



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



2044 106 420 706

L758-p v. 2

W. G. FARLOW



Facsimile-Edition. Ed.: W. Junk. No. 11.

C. LINNAEUS

SPECIES PLANTARUM.

TOMUS II.

HOLMIAE 1753.

EXEMPL. NO.....



W. JUNK

BERLIN W., KURFÜRSTENDAMM 201.

1907.

CAROLI LINNÆI

SÆ R:GIÆ M:ITIS SVETIÆ ARCHIATRI; MEDIC. & BOTAN.
PROFESS. UPSAL; EQUITIS DE STELLA POLARI;

SPECIES
PLANTARUM,

EXHIBENTES

PLANTAS RITE COGNITAS.

AD

GENERA RELATAS,

CUM

DIFFERENTIIS SPECIFICIS,
NOMINIBUS TRIVIALIBUS,
SYNONYMIS SELECTIS,
LOCIS NATALIBUS,

SECUNDUM

SYSTEMA SEXUALE

DIGESTAS.

TOMUS II.

Cum Privilegio S. R. Mitis Sueciæ & S. R. Mitis Polonia ac Electoris Saxon.

HOLMIÆ,
IMPENSIS LAURENTII SALVII.
1753.

Paulson 8/2

L758sp
v.2

Claff. XIV. p. 561

DIDYNAMIA,

GYMNOSPERMIA.

AJUGA. 613

1. AJUGA floribus resupinatis. orientalis.
Teucrium flaminibus tubo corollæ brevioribus. *Roy. lugdb.* 306.
Bugula orientalis villosa, flore inverso candido cum or-
 nis purpureis. *Tournef. cor.* 14.
 2. *Bugula* orientalis villosa, flore inverso cæruleo. alba
 macula notato. *Tournef. cor.* 14. *Dill. elib.* 60. t.
 43. f. 61.
Habitat in Oriente.
2. AJUGA tetragono-pyramidalis. *Fl. suec.* 475. *Mat. pyramidalis,*
med. 306. *Dalib. paris.* 169.
Consolida media pratensis cærulea. *Baub. pin.* 260. *non*
synon.
Consolida media genevensis. *Baub. hist.* 3. p. 432.
Bugula foliis angulose dentatis, caule simplici. *Hall.*
helv. 635.
Phyllochnois. *Reneal. spec.* 125. *
Habitat in Suecia, Helvetia, Germania.
3. AJUGA foliis tomentosis, calycibus hirsutis genevensis.
Bugula carneo flore. *Cins. hist.* 2. p. 43.
Habitat in Europa australiori.
4. AJUGA stolonibus reptatricibus. reptans.
Ajuga foliis obovatis crenatis, caule simplicissimo, sto-
 lonibus reptatricibus. *Gort. gelr.* 330.
Teucrium foliis obverse ovatis crenatis, caule simpli-
 cissimo, stolonibus reptatricibus. *Hort. cliff.* 301.
Roy. lugdb. 306. *Dalib. paris.* 170.
Consolida media pratensis cærulea. *Baub. pin.* 260.
Bugula. *Dod. pempt.* 135.
Habitat in Europa australiori.

N n

TE-

TEUCRIUM

614
*campanula-
 tum.*

1. **TEUCRIUM** foliis multifidis, floribus solitariis.
Teucrium foliis cuneliformibus trifidis: lacinis divisis, floribus solitariis. *Roy. lugdb.* 306.
Teucrium, calyce campanulato, laciniatum, flore magno & parvo subcæruleo. *Bærb. lugdb.* 1. p. 131.
Habitat in Oriente.

orientalis.

2. **TEUCRIUM** foliis multifidis, floribus racemosis.
Teucrium foliis pinnatifidis, laciniis linearibus, infimis divisis, notibus racemosis. *Hort. cliff.* 301. *Roy. lugdb.* 307.
Teucrium orientale angustifolium laciniatum, flore magno suaverubente. *Tournef. cor.* 14.
Habitat in Oriente.

Botrys.

3. **TEUCRIUM** foliis multifidis, floribus verticillatis utrinque ternis.
Teucrium foliis aliquoties trifidis, floribus verticillatis. *Hort. cliff.* 301. *Roy. lugdb.* 306. *Dalib. parif.* 171. *Sauv. monsp.* 248.
Botrys chamædryoides. *Bauh. pin.* 138.
Chamæpitys altera. *Dod. pempt.* 46.
Habitat in Germaniæ, Galliæ, Helvetiæ, Italiæ aprieti cultis. ☉

Pseudochamæpitys.

4. **TEUCRIUM** foliis trifidis linearibus integerrimis, floribus pedunculatis subracemosis, caule erecto.
Chamæpitys spuria, multifido folio, lamii flore. *Bauh. pin.* 250.
Pseudochamæpitys. *Clas. hist.* 2. p. 185.
Chamæpitys alia. *Cam. epit.* 680.
Habitat in Hispania.

Chamæpitys.

5. **TEUCRIUM** foliis trifidis linearibus integerrimis, floribus sessilibus axillaribus. *Hort. ups.* 160. *Mat. med.* 287.
Teucrium foliis simpliciter trifidis. *Hort. cliff.* 301. *Roy. lugdb.* 306. *Guet. stamp.* 2. p. 243. *Gron. virg.* 64. *Dalib. parif.* 173. *Sauv. monsp.* 175.
Chamæpitys lutea vulgaris L. folio trifido. *Bauh. pin.* 249.
Chamæpitys prima. *Dod. pempt.* 46.
Habitat in Italiæ, Galliæ, Angliæ, Hungariæ, Helvetiæ arvis. ☉

6.TEU-

6. **TEUCRIUM** foliis tripartitis linearibus, racemo terminali, caule repente.

Teucrium repens, folio trifido. *Nissol.*

Habitat - - -

Folia tripartita, linearia, glabra. Flores in racemo terminali, oppositi, pedunculati.

7. **TEUCRIUM** foliis subtridentatis linearibus, floribus lvs, sessilibus.

Teucrium foliis lanceolatis obtusis dentatis, floribus solitariis sessilibus. *Roy. lugdb.* 306. *Sauv. monsp.* 141.

Chamæpitys moschata, foliis serratis. *Bauh. pin.* 249.

Chamæpitys spuria prior. *Dod. pempt.* 47.

Anthyllis altera. *Clus. hist.* 2. p. 186.

Habitat in Lusitania, G. Narbonensi, Monspelii.

8. **TEUCRIUM** foliis pinnato-multifidis, caule simplicissimo erecto, bracteis subulato-palmatis.

Teucrium, delphinii folio, non ramosum. *Shaw. afric.*

575. t. 575.

Habitat in Mauritania.

9. **TEUCRIUM** foliis integerrimis oblongo-ovatis petiolatis: supra glabris subtus tomentosis.

Teucrium foliis ovato-oblongis integerrimis petiolatis, floribus solitariis pedunculatis. *Hort. cliff.* 302. *Roy. lugdb.* 305.

Teucrium peregrinum, folio sinuoso. *Bauh. pin.* 247.

Teucrium fruticans bæticum. *Clus. hist.* 1. p. 348.

Habitat in Bætica, Sicillæ maritimis. ♀

10. **TEUCRIUM** foliis integerrimis rhombeis acutis villosis: subtus tomentosis.

Teucrium fruticans bæticum, ampliore folio. *Dill. elsh.* 397. t. 284. f. 367.

Teucrium bæticum humilium & ramosius, folio subrotundo. *Hort. angl.* t. 3.

Habitat in regione Bætica. ♀

11. **TEUCRIUM** foliis lanceolato-linearibus integerrimis, racemo subspicato.

Teucrium foliis lanceolato-linearibus integerrimis sessilibus, floribus solitariis pedunculatis. *Hort. cliff.*

303. *Hort. ups.* 150. *Mat. med.* 186. *Roy. lugdb.* 307.

Polium angustifolium creticum. *Bauh. pin.* 221.

Rosmarinum stœchados facie. *Alp. exot.* 103. t. 102.

Habitat in Creta. ♀

564 DIDYNAMIA GYMNOSPERMIA.

- Marum.* 12. TEUCRIUM foliis integerrimis ovatis utrinque acutis, racemis secundis villosis.
Teucrium foliis ovatis utrinque acutis integerrimis, floribus solitariis utrinque digestis. *Hort. cliff.* 303. *Hort. ups.* 159. *Mat. med.* 285. *Roy. lugdb.* 305.
Marum cortusi. *Bauh. hist.* 3. p. 242.
Habitat in regno Valentino. 5
- multiflorum.* 13. TEUCRIUM foliis ovatis utrinque acutis; superne serrato-dentatis, floribus racemosis. *Roy. lugdb.* 307.
Teucrium foliis ovatis utrinque acutis utrinque bidentatis, verticillis laxis. *Hort. cliff.* 303.
Chamædrys multiflora tenuifolia Hispanica. *Tournef. inst.* 205. *Barb. lugdb.* 1. p. 182.
Habitat in Hispania.
- Sibiricum.* 14. TEUCRIUM foliis serratis ovatis, pedunculis trifloris, bracteis lineari-lanceolatis.
Teucrium chamædryos folio, flore parvo purpureo. *Gmel. mss.*
Habitat in Sibiria. 2
- canadense.* 15. TEUCRIUM foliis ovato-lanceolatis serratis, caule erecto, racemo terminali, verticillis hexaphyllis.
Teucrium foliis lanceolatis serratis petiolatis, floribus solitariis. *Gron. virg.* 64.?
Chamædrys canadensis, urticæ folio subtus incano. *Tournef. inst.* 205.
Habitat in Canada. 2
Planta e Canada Foliis gaudet angustioribus, æqualiter serratis, subtus tomentosis, planis; Spica ex verticillis aut sparsis floribus bractea minima singulis subjecta; at Hortensis Foliis majoribus rugosis, inæqualiter serratis, vix subtus pubescentibus & Floribus in Racemum subspicatum ex verticillis sexfloris, hexaphyllis: bracteis serratis.
- virginicum.* 16. TEUCRIUM foliis ovatis inæqualiter serratis, racemis terminalibus. *Gron. virg.* 64.
Habitat in Virginia.
- Scorodonia.* 17. TEUCRIUM foliis cordatis serratis petiolatis, racemis lateralibus secundis, caule erecto.
Teucrium foliis cordatis crenatis petiolatis, spicis laxis secundis. *Hort. cliff.* 301. *Roy. lugdb.* 306. *Dalib. parif.* 171. *Gort. gelr.* 331.
Scorodonia. Riv. mon. t. 12.

Scor-

Scordium alterum f. *Salvia sylvestris*. *Bauh. pin.* 247.
Habitat in arenosis castris Germaniæ, Helvetiæ, Gal-
liæ, Angliæ, Belgii: in dunis prædi Martecampi, be-
ati quondam mibi Paradisi. &

18. TEUCRIUM foliis oblongis sessilibus dentato-ser- *Scordium*.
 ratis, floribus geminis axillaribus, caule diffuso.

Teucrium foliis ovato-lanceolatis serrulatis sessilibus,
 floribus sæpius binatis. *Hort. cliff.* 302. *Fl. Suec.* 476.

Mat. med. 289. *Roy. lugdb.* 306. *Dalib. paris.* 171.

Scordium. *Bauh. pin.* 247. *Cam. epit.* 588. *Dod. pempt.*
 126.

Habitat in Europæ paludosis. &

19. TEUCRIUM foliis ovatis inciso-crenatis petiolatis, *Chamædry.*
 floribus subverticillatis. *Hort. cliff.* 302. *Mat. med.*
 288. *Roy. lugdb.* 307. *Dalib. paris.* 170. *Sauv. monsp.*
 145.

*Chamædry*s major repens. *Bauh. pin.* 248.

*Chamædry*s. *Riv. mon.* 10.

- β. *Chamædry*s minor repens. *Bauh. pin.* 248.

Habitat in Germania, Helvetia, Gallia. &

20. TEUCRIUM foliis cordatis undulatis obtuse serra- *florum*.
 tis: floralibus integerrimis concavis, caule fruticoso.

Teucrium foliis subovatis crenatis, floribus laxè spicatis
 pedunculatis, caule frutescente. *Hort. cliff.* 302. *Hort.*

upf. 160. *Roy. lugdb.* 307. *Sauv. monsp.* 145.

Teucrium. *Bauh. pin.* 247.

*Chamædry*s fruticosior, flore ochroleuco. *Morif. hist.* 3.

p. 421. *f.* 11. *t.* 22. *f.* 1.

Teucrium vulgare fruticans 1. *Clas. hist.* 1. *p.* 348.

- β. *Teucrium* lucidum, parvo folio, flore venuste purpu-
 reo. *Pluk. alm.* 367. *t.* 65. *f.* 2.

Habitat in Italia, Sicilia, Maltha, Hispania, Monspel-
lii. &

21. TEUCRIUM corymbo terminali, foliis lanceolatis *montanum*.
 integerrimis.

Teucrium foliis lanceolatis integerrimis petiolatis, spicis
 laxè subrotundis. *Hort. cliff.* 302. *Roy. lugdb.* 305.

Dalib. paris. 171. *Sauv. monsp.* 141.

Polium, lavendulæ folio. *Bauh. pin.* 220.

Polium majus. *Cord. hist.* 124. *b.*

Ajuga folio integro. *Riv. mon.* 15

Habitat in siccis Germaniæ, Monspellii, Genevæ, Hel-
vetiæ, Parisiis. &

- supinum.* 22. TEUCRIUM corymbo terminali, foliis linearibus.
 Polium montanum repens. *Baub. pin.* 220.
 Polium montanum supinum minimum. *Lab. ic.* 488.
Habitat in agro Viennensi.
Præcedenti maxime affinis est.
- pyrenaicum.* 23. TEUCRIUM corymbo terminali, foliis cuneiformi-orbieulatis crenatis.
 Teucrium foliis cuneiformibus crenatis, corymbo terminali, foliolis floralibus linearibus. *Vir. cliff.* 55.
Roy. lugdb. 305.
 Teucrium foliis suborbiculatis crenatis, spica laxa orbiculata depressa. *Hort. cliff.* 301.
 Polium pyrenaicum supinum, hederæ terrestris folio. *Tournef. inst.* 206.
 Polium saxatile purpureum chamædryoides, ampla coma. *Bocc. musf.* 2. t. 61. *Barr. rar.* 315. t. 1086. *folia male.*
Habitat in Pyrenæis. 2
- Polium.* 24. TEUCRIUM spicis subrotundis, foliis oblongis obtusis crenatis tomentosis sessilibus. *Hort. cliff.* 302.
Roy. lugdb. 305. *Sauv. monsp.* 141.
 Polium montanum luteum. *Baub. pin.* 220.
 β. Polium montanum album. *Baub. pin.* 221.
 γ. Polium montanum supinum alterum. *Baub. pin.* 221.
 δ. Polium maritimum supinum venetum. *Baub. pin.* 221.
 Polium montanum 5. *Clus. hist.* 1. p. 362.
Habitat in Italia, Hispania, Lusitania, Narbona. 2
Species Polii, a varietatibus distinctas, hodie confusas, inquirant solidi Botanici, in solo natali.
- capitatum.* 25. TEUCRIUM capitulis pedunculatis, foliis lanceolatis crenatis tomentosis, caule erecto.
 Polium montanum album serratum latifolium, *Barr. rar.* 325. t. 1079. 1078.
Habitat in Hispania.
Habitus Organi majoranæ, sed tomentosum & angustifolium.
- spinosum.* 26. TEUCRIUM spinosum, calycibus labio superiore ovatis, corollis resupinatis. *Læfl. descr.* VII. 38.
 Chamædrys multifida spinosa odorata. *Grisl. l'hist.*
 Chamædrys spinosa. *Baub. pin.* 248. *prodr.* 117. f. 117.
Morif. hist. 3. p. 423. f. 11. t. 22. f. 17.
Habitat in Lusitania. ☉
- onatum* 27. TEUCRIUM foliis serratis, spinis terminalibus, corollis rectis. Scor-

DIDYNAMIA GYMNOSPERMIA. 567

Scordium spinosum odoratum. Corn. cau. 123. Barr. rar. 319. t. 102.

Habitat — — — — — †

SATUREJA. 415

1. **SATUREJA** capitulis terminalibus, foliis lanceolatis. *virginiana*.
Satureja virginiana. Herin. par. 218. t. 218.

Pulegium crectum virginianum angustifolium, floribus in cymis dispositis. Moris. hist. 3. p. 371. f. 11. t. 7. f. 8.

Clinopodium foliis lanceolatis acuminatis, capitulis terminalibus. Hort. cliff. 304. Gron. virg. 65. Roy. lugdb. 314.

Clinopodium, pulegii angusto rigidoque folio, virginianum, flosculis in cymis dispositis. Pluk. alm. 110. t. 54. f. 2.

Serpentaria virginiana. Bocc. mus. 2. p. 161. t. 115.

Habitat in Virginia. 2

Stamina in fundo corollae latent Antheris minimis effatis, ut fere posset genere proprio tradi.

2. **SATUREJA** verticillis fastigiatis concatenatis, foliis *juliano*.
lineari-lanceolatis.

Satureja floribus verticillatis, foliis lanceolatis glabris. Roy. lugdb. 324.

Satureja spicata. Bauh. pin. 218.

Satureja perennis, verticillis spicatis & densius dispositis. Moris. hist. 3. p. 412. f. 11. t. 17. f. 4.

Habitat in Etruria, in Tyrrheni maris asperis, Florentiae. 2

3. **SATUREJA** floribus verticillatis, foliis ovatis acutis. *Thymbra*.
Roy. lugdb. 224.

Thymus frutescens, verticillis fere nudis globosis, foliis ovato-lanceolatis. Hort. cliff. 306.

Thymbra legitima. Clus. hist. 1. p. 358.

Thymum creticum pone verticillatum. Barr. rar. 279. t. 898.

Habitat in Creta.

4. **SATUREIA** verticillis lanuginosis, dentibus calycinis setaceis pilosis. *maffichtua*.
Hort. cliff. 306. Hort. upf. 160.
sub Thymo.

Samplicus s. Marum maffichen redolens. Bauh. pin. 224.

Marum vulgare s. Clinopodium. Dod. pempt. 471.

Habitat in Hispaniae petrosis. 2 **f. SA-**

- græca.* 5. SATUREJA pedunculis corymbosis lateralibus geminis, bracteis calyce brevioribus.
 Satureja annua orientalis tenuior, ad singulos nodos horifera. *Morif. hist.* 3. p. 411. f. 11. t. 17. f. 2.
 Clinopodium creticum. *Alp. exot.* 265. t. 264.
Habitat ad Archipelagum. ☉
Facie convenit cum S. montana, æque differt foliis minus mucronatis; sed precipue corymbis florum, qui utrinque non soli sed bini; accedit quod bractea f. involucrum in hac minimum est; in montana vero longitudine corymbi.
- montana.* 6. SATUREJA pedunculis dichotomis lateralibus solitariis, foliis mucronatis.
 Satureja pedunculis dichotomis, verticillis concatenatis. *Vir. cliff.* 57. *Hort. upf.* 161. *Roy. lugdb.* 324. *Sauv. monsp.* 142.
 Satureja montana. *Baub. pin.* 218.
 Satureja perennis. *Riv. mon.* 41.
 Melissa foliis linearibus integerrimis. *Hort. cliff.* 308.
Habitat in Hetruria, Narbona. 2
- hortensis.* 7. SATUREJA pedunculis bifloris. *Vir. cliff.* 87. *Hort. upf.* 161. *Mat. med.* 284. *Roy. lugdb.* 324. *Sauv. monsp.* 142.
 Satureja hortensis. *Baub. pin.* 218.
 Satureja altera. *Cam. epit.* 487.
 Thymus erectus annuus, foliis lanceolato-linearibus. *Hort. cliff.* 306.
Habitat in G. Narbonensi & Italia. ☉
- cretica.* 8. SATUREJA floribus spicatis, foliis carinatis punctatis ciliatis. *Mat. med.* 283.
 Thymus capitatus qui dioscoridis. *Baub. pin.* 219.
 Thymum legitimum. *Clus. hist.* 1. p. 357.
Habitat in Creta, Bætica, Hispali, Græcia. 2
- origanoides.* 9. SATUREJA foliis ovatis serratis, corymbis terminalibus dichotomis.
 Thymus foliis ovatis acuminatis serratis, corymbis lateralibus terminalibus pedunculatis. *Grou. virg.* 64.
 Calamintha mariana, innecronatis rigidioribus & crenatis foliis, flosculorum calyculis villis argenteis summo margine fimbriatis. *Pluk. mant.* 35. t. 344. f. 2.
 Calamintha erecta virginiana, mucronato folio glabro. *Morif. hist.* 3. p. 413. f. 11. t. 19. f. 7.
Habitat in Virginia.

THYM-

THYMBRA. 6 16

1. THYMBRA floribus spicatis.

spicata.

Thymbra spicata verior hispanica, *Barr. rar.* 281. t. 1230.

Thymum majus longifolium, stoecnadis foliaceo capite purpurascens, pilosum, *Pluk. alm.* 368. t. 116. f. 5.

Hyssopum montanum macedonicum valerandi dourez. *Baub. hist.* 3. p. 276.

Hyssopus capitata africana, saturejæ rigido hirsuto folio. *Morif. hist.* 3. p. 361.

Satureja hirsuta purpurea olibani. *Dodart. mem.*

Habitat in Macedonia: Libano.

2. THYMBRA floribus verticillatis.

verticillata.

Hyssopus angustifolia montana aspera. *Baub. pin.* 218.

Hyssopus montana. *Dalech. hist.* 934.

Habitat in Europa australi.

Planta 1. 2. similis Caules vix pedales. Folia opposita, lineari-lanceolata, integra, punctata: Floralia paulo latiora, acutiora, sed ciliata. Hæc omnino verticillata, floribus sessilibus: Præcedens obsolete spicata, spica oblonga bracteis s. foliis ciliatis imbricata. Calyx monophyllus, durus, utrinque latere linea hirsuta exaratus, bilabiatus: labio superiore latiore, semitrisido, æquali; L. inferiore angustiore, bipartito. Corolla ringens: labio superiore cuncavo, semibifido, obtuso; l. inferiore trisido, subæquali; Stamina didynamia sub labio superiore. Antheræ didymæ. Stylus semibifidus, filiformis. Semina 4

HYSSOPUS. 6 17

1. HYSSOPUS spicis secundis. *Hort. cliff.* 304. *Hort. officinale.*

apsf. 152. *Mat. med.* 290. *Roy. lugdb.* 323. *Gort. gelr.* 336.

Hyssopus officinarum cærulea s. spicata. *Baub. pin.* 217.

Hyssopus vulgaris. *Dod. pempt.* 287.

β Hyssopus rubro flore. *Baub. pin.* 217.

Habitat in - - - - - 2

2. HYSSOPUS corollis transversalibus, staminibus inferioribus corolla brevioribus. *Hort. apf.* 162. *

Habitat in Sibiria. 2

3. HYSSOPUS caule acuto quadrangulo. *Hort. apf.* 3. * nepe oider.

Nepeta caule acute quadrangulo glabro. *Vir. cliff.* 58.

Roy. lugdb. 316. *Gron. virg.* 66.

N 15

Bru-

Brunella bracteis lanceolatis Hort. cliff. 316.

Betonica virginiana elatior, foliis scrophulariæ glabris, dore ochroleuco. Pluk. alm. 67. t. 150. f. 3. Moris. hist. 3. p. 365. f. 11. t. 4. f. 11.

Habitat in Virginia, Canada. ♂

618

NEPETA.

- ontario.* 1. *NEPETA* floribus spicatis, verticillis subpedicellatis, foliis petiolatis cordatis dentato-serratis.
Nepeta floribus interrupte spicatis pedunculatis. Hort. cliff. 310. Hort. ups. 163. Fl. succ. 486. Roy. lugdb. 316. Gron. virg. 65. Mat. med. 291. Dalib. paraf. 116.
Mentha cataria vulgaris & major. Baub. pin. 228.
Cataria herba. Dod. pempt. 99.
Habitat in Europa. ♀
- pannonica.* 2. *NEPETA* floribus paniculatis, foliis cordatis petiolatis obsolete crenatis,
Nepeta caule paniculato, pedunculis partialibus ramosis multifloris. Roy. lugdb. 316.
Nepeta pannonica major & elatior. Moris. hist. 3. p. 415. f. 11. t. 6. f. 6.
Mentha montana verticillata. Baub. pin. 227.
Habitat in Pannonia. ♀
- violacea.* 3. *NEPETA* verticillis pedunculatis corymbosis, foliis petiolatis cordato-oblongis dentatis.
Nepeta foliis ovatis petiolatis glabris, verticillis diffuse spicatis glabris. Sauv. monsp. 151.
Nepeta montana purpurea major, sparsa spica. Barr. ic. 601. Bocch. mus. 2. p. 46. t. 36.
Cataria hispanica, betonica folio angustiore, flore cæruleo. Tournef. inst. 202.
Habitat in Hispania. ♀
- ucranica.* 4. *NEPETA* floribus paniculatis, foliis lanceolatis serratis sessilibus.
Cataria betonica folio angustiore, flore cærulescente. Gerb. tan 87.
Habitat in Ucraina.
- nuda.* 5. *NEPETA* foliis cordato-oblongis subsessilibus serratis, racemis verticillatis nudis.
Nepeta foliis sessilibus verticillatis nudis.
*Nepeta foliis sessilibus nudis, calycibus fructus ovatis æqualibus. Hort. ups. 164. **
Men-

DIDYNAMIA GYMNOSPERMIA. 571

Mentha cataria hispanica, olium *Sideritis altissima*, flore albo. *Baub. hist.* 3. p. 226
Habitat in Hispania. 2

NEPETA floribus sessilibus verticillato-spicatis, verticillis tomento obvolutis. *Hort. cliff.* 311. *Roy. lugdb.* 316.

Horminum spicatum, lavendulæ flore & odore. *Bocc. fec.* 48. t. 25. f. 2.
Habitat in Sicilia.

NEPETA floribus sessilibus verticillato-spicatis, bracteis lanceolatis longitudine calycis, foliis petiolatis.
Mentha Cataria minor alpina. *Baub. pin.* 228. *prodr.* 110.

Habitat in Italia. *D. Ratgeb.* 2
Radix perennis. *Caulis pedalis.* *Folia cordata, obtusa, crenata, petiolata.* *Spicæ terminales, verticillatæ: verticilli remoti, sessiles, obvaliati bracteis lanceolatis, viridibus, lateribus albidioribus, longitudine calycum, acuminatis.* *Calyces erecti, æquales.* *Corollæ alba.*

NEPETA floribus spicatis sessilibus, bracteis ovatis tuberosa, coloratis verticilla excipientibus, foliis sessilibus. *Hort. cliff.* 311. *Roy. lugdb.* 316.

Nepeta tuberosa spicata hispanica. *Bocc. mus.* 2. p. 44. t. 36. *Barr. ic.* 1131.
Habitat in Hispania, Lusitania. 2

NEPETA foliis lanceolatis, capitulis terminalibus, staminibus flore longioribus.

Clinopodium foliis lanceolatis, capitulis terminalibus. *Hort. cliff.* 305. *Gron. virg.* 65.

Clinopodium amaraci folio, floribus albis. *Pluk. alm.* 110. t. 85. f. 2.

Clinopodium, flore albo, ramosius, angustioribus foliis glabris; virginianum. *Morif. hist.* 3. p. 374. f. 11. t. 8. f. ult.

Habitat in Virginia. 2

Lobium inferius corollæ serratum, at non concavum.

NEPETA corollarum labio superiore integerrimo indiviso brevissimo.

Leucas foliis subrotundis serratis, flore albo. *Burm. zeyl.* 140. t. 63. an?

Nepeta indica, rotundior folio. *Morif. hist.* 3. p. 415. f. 11. t. 6. f. 7.

Ka-

572 DIDYNAMIA GYMNOSPERMIA.

Katu-kurka. *Rheed. mal.* 10. p. . . . f. 90.

Habitat in India.

*Labium corollæ superius brevissimum omnium, integrum
filamina autem, longitudine labii inferioris. Flori
verticillati.*

- florica.* 11. NEPETA corymbis geminis pedunculatis axillaribus
foliis cordato oblongis acuminatis serratis. *Hort. upf.*
164. *

Cataria montana, foliis veronicæ pratensis. *Buxb. cen.*
3. p. 27. t. 50. f. 1.

Habitat in Sibiria. &

- multifida.* 12. NEPETA floribus spicatis, foliis pinnatifidis integre
rimis.

Lavendula montana, folio valerianellæ varie secto, ca-
rulea. *Amm. ruth.* 69.

Betonica sibirientis. *Walsh. hort.* 122. t. 20.

Habitat in Dauria.

Caulis erectus absque ramis; Folia inferiora petiolata
superiora sessilia, non crenata, oblonga, triloba, quinqueloba,
driloba, quinqueloba: lobis alternis, oblongis. Flori
valde parvi; Antheræ bipartitæ.

LAVANDULA

- Spica.* 1. LAVANDULA foliis lanceolatis integris, spicis n-
dis. *Hort. cliff.* 303. *Hort. upf.* 162. *Mat. med.* 29
Roy. lugdb. 322. *Sauv. monsp.* 143.
Lavandula angustifolia. *Bauh. pin.* 216.
Lavandula altera. *Dnd. pempt.* 273.
β. Lavandula latifolia, *Bauh. pin.* 216.
Habitat in Europa australi. h

- multifida.* 2. LAVANDULA foliis duplicato pinnatifidis, *V.*
cliff. 56. *Hort. cliff.* 303. *Roy. lugdb.* 322. *Hort.*
upf. 162.
Lavandula folio dissecto. *Bauh. pin.* 216.
Lavandula multifido folio. *Clus. hist.* 1. p. 345.
Habitat in regione Bætica. δ

- dentata.* 3. LAVANDULA foliis pinnato-dentatis. *Hort. cliff.*
303. *Roy. lugdb.* 323.
Stæchas folio serrato. *Buxb. pin.* 216.
Stæchas crispa folio. *Clus. hist.* 1. p. 346.
Habitat in Hispaniæ Calape. h

4. *LAVANDULA* foliis lanceolato-linearibus, spica co-*Stachas*.
mosa. *Hort. cliff.* 303. *Hort. upf.* 162. *Mas. med.*
197. *Roy. lugdb.* 323. *Sauv. inensp.* 143.
Stachas purpurea. *Baub. pin.* 216.
Stachas brevioribus ligulis. *Clus. hist.* 1. p. 344.
5. *Stachas* cauliculis non foliatis. *Baub. pin.* 216.
Stachas longioribus ligulis. *Clus. hist.* 1. p. 344.
Habitat in Europa australi. 5

BETONICA. 6 2 3

1. *BETONICA* spica interrupta, corollarum lacinia la-^{officinalis.}
bii intermedia emarginata. *Roy. lugdb.* 316. *Fl. succ.*
487. *Hort. upf.* 164.
Betonica Dod. *pempt.* 40. *Hort. cliff.* 310.
Betonica purpurea. *Baub. pin.* 235.
6. *Betonica* alba. *Baub. pin.* 235.
Habitat in Europa. 2
1. *BETONICA* spica integra, corollarum lacinia labii ^{orientalis,}
intermedia integerrima. *Roy. lugdb.* 316.
Betonica orientalis, angustissimo & longissimo folio,
spica florum crassiori. *Tournef. cor.* 13.
Habitat in Oriente. 2
3. *BETONICA* spica basi foliosa. ^{alopecurus,}
Horminum minus album, *betonicæ* facie. *Baub. pin.*
239. *prodr.* 114. *Burs. XIII:* 119.
Betonica montana lutca. *Barr. icon.* 339? *Bocc. mus.*
2. p. 82. t. 7.?
Betonica folio, capitulo *alopecuri.* *Baub. pin.* 235.
Betonica folio *Alopecuros* quorundam. *Baub. hist.* 3.
p. 303.
Alopecuros alpina, *betonicæ* folio, capitulo *alopecuri.*
Pont. anth. 253. *Segu. ver.* 318.
Horminum alpinum luteum, *betonicæ* spica. *Raj. hist.*
547.
Habitat in montibus Sabaudicis, Austriæ superioris,
Lessanensibus, Italicis, Galloprovinciæ. 2
4. *BETONICA* verticillata, calycibus spinosis. *Hort. upf. annua.*
165.
Stachys foliis lanceolatis sessilibus basi attenuatis. *Hort.*
cliff. 310. *Gron. virg.* 66. *β. Dalib. parif.* 180.
Betonica arvensis annua, flore ex albo flavescente. *glabra.*
Tournef. inst. 203.

Be-

574 DIDYNAMIA GYMNOSPERMIA.

Betonica foliis inter diffitos verticillas longioribus ellipticis serratis. *Hall. helv.* 645.

Sideritis arvensis latifolia glabra. *Bauh. pin.* 233.

Alyssum majus. *Tabern. hist.* 929.

hirsuta.

β. Sideritis vulgaris hirsuta erecta. *Bauh. pin.* 233.

Sideritis vulgaris 1. *Clus. hist.* 2. p. 39.

Betonica foliis inter verticillos diffitos brevibus in aestivam terminatis integris. *Hall. helv.* 646.

Habitat ad agrorum margines Helvetiæ, Germaniæ, Galliæ. ☉

De Genere dubium, utrum ad Betonicas vel Stachydas pertineat; prius statuit Tournefort, posterius spinosæ calycis suadent.

621

SIDERITIS.

canariensis.

1. SIDERITIS fruticosa tomentosa, foliis cordatis, pedunculis ante florescentiam nutantibus.

Stachys fruticosa, foliis cordato-oblongis crenatis petiolatis. *Hort. cliff.* 310.

Stachys canariensis frutescens, verbasci folio. *Compt. hort.* 2. p. 197. t. 99.

Stachys amplifolius verbasci foliis, floribus albis parvis non galeatis, spica betonicæ. *Pluk. alm.* 356. t. 31.

Habitat in Canariis. ♀

cretica.

2. SIDERITIS fruticosa villosa, foliis cordato-oblongo-petiolatis, ramis divaricatis

Stachys cretica latifolia. *Tournef. inst.* 186.?

Habitat in Creta. ♀

syriaca.

3. SIDERITIS fruticosa tomentoso-lanata, foliis linearilanceolatis.

Cunila calycibus inermibus lanigeris. *Hort. cliff.* 311.

Sideritis cretica tomentosa candidissima, flore luteo. *Tournef. cor.* 12.

β. Stachys fruticosa, foliis lanceolato-linearibus integerrimis sessilibus. *Hort. cliff.* 310.

Stachys minor italica. *Bauh. pin.* 236.

Stachys lychnoides incana angustifolia, flore aureo. *Bauh. pin.* 1187.

Pilosella syriaca. *Bauh. pin.* 262.

Habitat in Creta. ♀

Fruticulus facie Salviæ, sed longior. Folia salviæ, tomentosa, lanata, alba: inferiora petiolata, superiora sessilia. Flores ex albo flavescentes.

4. S

4. **SIDERITIS** herbacea hispido-pilosa, foliis superiori-*persollata*.
bus amplexicaulibus.
Sideritis orientalis, phlomidis folio. *Tournef. cor.* 12.?
Stachys foliis oblongis acutis amplexicaulibus serratis:
florallibus cordatis acutis integerrimis. *Roy. lugdb.* 318.
Habitat in Oriente?
Caulis *erectus*, herbaceus. Folia *inferiora* caulis *præ-*
oblata; *superiora* sessilia, opposita, amplexicaulia, hinc
singularia. Corollæ albae *venis* aliquot rufis.
5. **SIDERITIS** herbacea decumbens, calycibus spinosis, *montana*.
labio superiore trifido.
Cunila calycum labio spinoso: superiore trifido, inferi-
ore bipartito. *Hort. cliff.* 313. *Hort. ups.* 169. *Roy.*
lugdb. 329.
Sideritis montana, parvo varioque flore. *Bauh. pin.* 233.
Sideritis montana, parvo flore nigro-purpureo, capite
medio croceo. *Col. exphr.* 1. p. 198. 2. 196.
Habitat in Italia ☉
Omnes enumeratas species esse genuinas Sideritides, patet
ex styli altero ramulo amplexante ramulum longiorem;
ut & ex staminibus intra corollæ tubum, limbo vix
ringentem; Confirmat hoc floris color parsim quasi fer-
ragineus.
6. **SIDERITIS** herbacea decumbens, calycibus spinosis: *romana*,
labio superiori indiviso.
Cunila calycum lacinia superiore latiori ovata trinervi.
Hort. cliff. 313. *Roy. lugdb.* 321.
Siceritis genus verticillis spinosis. *Bauh. hist.* 3. p. 428.
Sideritis, verticillis spinosis, minor procumbens *Morif.*
hist. 3. p. 388. f. 11. t. 12. f. 5.
Habitat in agris Europæ australis. ☉
Calycis lobus 5:tus s. superior ovatus, reliquis duplo la-
tior.
7. **SIDERITIS** foliis lanceolatis glabris integerrimis, bra-*hyslopifolia*,
cteis cordatis dentato-spinosis, calycibus æqualibus.
Sideritis foliis lanceolato-linearibus sessilibus glabris.
Hort. cliff. 313. *Roy. lugdb.* 322.
Sideritis alpina hyslopifolia. *Bauh. pin.* 233.
Sideritis VII. Clus. hist. 2. p. 41.
Habitat in Hetruria, Pyrenæis, Thuri. &
8. **SIDERITIS** foliis lanceolatis subdentatis, bracteis cor-*hirsuta*.
datis dentato-spinosis, calycibus æqualibus.

Side-

576 DIDYNAMIA GYMNOSPERMIA.

Sideritis foliis hirsutis profunde crenatis. *Bauh. pin.* 232.
 Sideritis montana scordioides altera. *Barr. ic.* 1160.
Habitat in G. Narbonensi.

622

MENTHA.

crispa.

1. MENTHA floribus spicatis, foliis cordatis dentatis undulatis sessilibus. *Hort. cliff.* 306. *Hort. upf.* 168.
Mat. med. 278.
 Mentha crispa danica s. germanica speciosa. *Morif. hist.* 3. p. 367. f. 11. t. 6. f. 5.
 Mentha crispa. *Riv. moss.* 47
Habitat in Sibiria. 2

spicata.

2. MENTHA floribus spicatis, foliis oblongis serratis
Hort. upf. 168.

Mentha sylvestris, longioribus nigrioribus & minus incanis foliis. *Bauh. pin.* 227.

viridis.

- a. Mentha spicis solitariis interruptis, foliis lanceolatis serratis sessilibus. *Hort. cliff.* 306. *Roy. lugdb.* 325
Dalib. paris. 177.

Mentha angustifolia spicata. *Bauh. pin.* 227,

Mentha III. IV. *Dod. pempt.* 95.

Mentha aquatica spicata, foliis oblongis viridibus serratis acuminatis. *Gron. virg.* 167.

longifolia.

- β. Mentha spicis confertis, foliis serratis tomentosis sessilibus. *Hort. cliff.* 306. *Roy. lugdb.* 325. *Dalib. paris.* 177. *Mat. med.* 279. *Gort. gelr.* 341.

Mentha sylvestris, folio longiore. *Bauh. pin.* 227.

Menthastrum spicatum, folio longiore candicante.
Bauh. hist. 3. p. 321.

rotundifolia.

- γ. Mentha sylvestris, rotundiore folio. *Bauh. pin.* 227.
Habitat in Dania, Germania, Anglia, Gallia. 2

aquatica.

3. MENTHA floribus capitatis, foliis ovatis serratis petiolatis, staminibus corolla longioribus. *Hort. cliff.* 306. *Fl. suet.* 482. *Roy. lugdb.* 325. *Dalib. paris.* 177.
 Mentha rotundifolia palustris s. aquatica major. *Bauh. pin.* 227.

Sisymbrium sylvestre. *Dalech. hist.* 677.

Habitat in Europa ad aquas. 2

Planta non hirta. Verticilli terminales in caputulum s. spicam obtusam conferti; Stamina corollâ longiora.

superita.

4. MENTHA floribus capitatis, foliis lanceolatis serratis subpetiolatis.

Men.

Acutia spicis brevioribus & habitioribus, foliis menthae ferratis, sapore fervido piperis. Raj. angl. 3. p. 234. t. 10. f. 2.

Habitat in Anglia. 2

MENTHA floribus verticillatis, foliis ovatis acutis serratis, staminibus corolla brevioribus.

Mentha hortensis verticillata, ocyni odore. Bauh. pin. 227.

Mentha verticillata minor acuta non crispa odore ocyni. Bauh. hist. 3. p. 216.

Mentha quarta. Fuchf. hist. 291.

Habitat in Europa australiore. 2

Similima M. arvensi, at minus hirta, Caulibus rubris; Stamina tubo corollae non longiora, Verticilli laterales.

MENTHA floribus verticillatis, foliis ovatis acutis serratis, staminibus corollam aequantibus. *Hort. cliff. 307. Fl. paec. 481. Roy. lugdb. 326. Hall. belv. 658. Dalib. parif. 178.*

Mentha arvensis verticillata procumbens. Morif. hist. 3. p. 369. f. 11. t. 7. f. 5

Calamintha arvensis verticillata. Bauh. pin. 229.

acutata in Europae agris frequens post messem. 2

Caules diffusi; Verticilli laterales; Planta tota hirta; Stamina corollae limbo longitudine equalia.

MENTHA floribus verticillatis, foliis lanceolatis serratis petiolatis.

Habitat in Canada. Kalm.

Differt a precedenti Caule altiore, non ramoso, multum piloso. Foliis dimidio angustioribus, acutioribus, profundius serratis, obscurioribus pilis vagis adspersis, uti etiam calyces. Calyces quinqueidentati, subaequales. Corolla asida: lobo superiore oblongo, longiore, integro; reliquis equalibus. Stamina 4, intra tubum corollae. Stylus corollae longior, apice bifidus.

MENTHA floribus verticillatis, foliis ovatis obtusis subcrenatis, caulibus subteretibus repentibus. *Hort. cliff. 307. Roy. lugdb. 326. Dalib. parif. 178. Mat. med. 280. Gort. gelr. 344.*

Pulegium latifolium. Bauh. pin. 222.

Pulegium. Fuchf. hist. 199.

Habitat in Angliae, Galliae, Helvetiae inundatis. 2

578 DIDYNAMIA GYMNOSPERMIA.

- Cervina.* 9. MENTHA floribus verticillatis, foliis linearibus. *Hort. cliff.* 307. *Hort. ups.* 168. *Sauv. monsp.* 143.
Pulegium angustifolium. Bauh. pin. 222. *Morif. bist.* 3. p. 371. f. 11. t. 7. f. 7. *Riv. mon.* 22.
Habitat Monspelii & ad Rhodanum. &
- canariensis* 10. MENTHA floribus capitatis axillaribus dichotomis, foliis ovatis crenatis, caule arborecente. *Hort. cliff.* 307.
Mentha canariensis frutescens, foliis subtus lanugine candidissima villosis, floribus glomeratis e sinu foliorum longioribus pediculis insidentibus. *Pluk. alm.* 248. t. 307. f. 2.
Heliotropium canariense arborecens, scorodonix folio. *Comm. hort.* 2. p. 129. t. 65.
Habitat in Canariis, &

GLECOMA.

- hed-rapen.* 1. GLECOMA foliis reniformibus crenatis. *Hort. cliff.* 307. *Fl. succ.* 483. *Roy. lugdb.* 320. *Mat. med.* 303.
Hedera terrestris vulgaris. Bauh. pin. 306.
Chamæcistus, Fuchf. bist. 876.
Habitat in Europæ septentrionalioris sepibus. &
- arvensis* 2. GLECOMA foliis cordatis oblongis crenatis, caule hispido erecto.
Sideritis calycibus subinermibus hispidis. Fl. succ. 488.
Sideritis alpine trizaginis folio. Bauh. pin. 233. *prodr.* 111.
Habitat in arvis Europæ. ☉
- belgica* 3. GLECOMA foliis cordato-oblongis crenatis. *Hort. cliff.* 307. *Roy. lugdb.* 320. *Gort. gelr.* 339.
Lamium paludosum belgicum, melissæ folio. Herm. lugdb. 351. t. 353.
Habitat in Belgii subhumidis. ☉

ORVALA.

- leganica.* 1. ORVALA.
Papia garganica, foliis urticæ alternis & eleganter incis, flore purpureo. Mich. geir. 20. t. 17.
Mentha hirsutior brachyplatyphyllos exotica. Pluk. alm. 248. t. 45. f. 2.
Habitat in valle Garganica montis Apuliæ.

LA-

LAMIUM. 625

1. LAMIUM foliis cordatis acuminatis serratis petiolo album.
Hort. cliff. 314. Fl. succ. 493. Mat. med. 307.
Roy. lugdb. 319. Dalib. paris. 178.

Lamium album non foetens, folio oblongo. Bauh. pin.
231.

Galeopsis. Cam. epit. 865.

3. Lamium maculatum. Bauh. pin. 231.

Habitat in Europæ cultis. ☉

1. LAMIUM foliis cordatis obtusis petiolatis. Hort. cliff. purpureum.
314. Fl. succ. 494. Roy. lugdb. 319. Dalib. paris.
179.

Lamium purpureum foetidum, folio subrotundo. Bauh.
pin. 230.

Urtica iners altera. Dod. pempt. 153.

Habitat in Europæ cultis. ☉

1. LAMIUM foliis floralibus sessilibus amplexicaulibus amplexicau-
obtusis. Hort. cliff. 314. Fl. succ. 495. Roy. lugdb. 1c.
319. Gron. virg. 66. Dalib. paris. 179.

Lamium, folio caulem ambiente, minus. Bauh. pin.
231.

Morus gallinæ, folio hederulæ, alterum. Lob. ic 463.

1. Lamium, folio caulem ambiente, majus. Bauh. pin. 231.

Habitat in Europæ cultis. ☉

LAMIUM foliis multipartitis Hort. cliff. 315. multitudin.

Lamium orientale, foliis eleganter laciniatis. Tournef.

cn. 11. Comm. rar. 26. t. 26.

Habitat in Oriente.

GALEOPSIS. 626

GALEOPSIS internodiis caulinis æqualibus: verticillis Ladanum,
lis omnibus remotis.

Galeopsis ramis summis pubescentibus. Hort. cliff. 314.

Fl. succ. 492. Roy. lugdb. 319. Dalib. paris. 181.

Sideritis arvensis angustifolia rubra. Bauh. pin. 233.

Ladanum segetum folio latiore. Riv. mon. 24.

Habitat in Europæ arvis & campis sabulosis. ☉

GALEOPSIS internodiis superne incrassatis, verticillis Tetrakt.
summis subconjugis.

Galeopsis ramis summis strigosis. Hort. cliff. 314. Fl.

succ. 491. Roy. lugdb. 319 Dalib. paris. 181.

003

Gale-

580 DIDYNAMIA GYMNOSPERMIA.

Galeopsis corolla rubra aut alba. *Fl. lapp.* 237.
 Urtica aculeata, foliis ferratis. *Bauh. pin.* 232.
 Cannabis spuria. *Riv. mon.* 44.

β. Galeopsis corolla flava: labio inferiore maculato. *Fl. lapp.* 193.

Lamium cannabinum aculeatum, flore specioso luteo
 labiis purpureis. *Pluk. alm.* 204. t. 41. f. 4.
 Cannabis spuria, flore majore. *Riv. mon.* 45.
Habitat inter Europæ segetes & olera. ☉

trifida. 3. GALEOPSIS caule piloso, calycibus labio corollæ
 superiore longioribus.

Marrubium nigrum rotundifolium. *Bauh. pin.* 230.
 Ocymastrum valentinum. *Clus. hist.* 2. p. 42.
Habitat in Hispania. ☉

Galeobdolon 4. GALEOPSIS verticillis sexfloris: involucri tetra-
 phyllo.

Leonurus foliis ovatis ferratis acutis. *Hort. cliff.* 317.
Roy. lugdb. 310. *Fl. suec.* 479. *Dalib. parif.* 180.
It. scan. 64.

Lamium folio oblongo luteum. *Bauh. pin.* 231.
 Urtica iners tertia s. Lamium flore luteo. *Do-
 pempt.* 153.

Habitat in Europæ nemoribus. 2

Frustrificatio parum recedit a congeneribus, vix tam
 distincto genere nominari meretur.

STACHYS.

sylvatica. 1. STACHYS verticillis sexfloris, foliis cordatis petio-
 latis. *Fl. suec.* 489. *Dalib. parif.* 180.

Stachys foliis cordatis acuminatis petiolatis, floribus la-
 spicatis. *Hort. cliff.* 309. *Roy. lugdb.* 317. *Hall. herb.*
 642.

Lamium maximum sylvaticum foetidum. *Bauh. pin.*
 231.

Galeopsis legitima. *Clus. hist.* 2. p. 35.

Habitat in Europæ nemoribus umbrosis. ☉

palustris. 2. STACHYS verticillis sexfloris, foliis lineari-lanceo-
 latis semi-amplexicaulibus. *Fl. suec.* 490. *Dalib. pa-
 ris.* 180.

Stachys foliis lineari-lanceolatis sessilibus basi emargina-
 tis. *Hort. cliff.* 309. *Roy. lugdb.* 317.

Stachys palustris foetida. *Bauh. pin.* 236.

Stachys aquatica. *Tabern. ic.* 577.

Hab

DIDYNAMIA GYMNOSPERMIA 581

Habitat in Europa ad ripas, inque cultis humidiusculis. 2

- 3 STACHYS verticillis multifloris, foliorum serraturis *alpina*.
apice cartilagineis.

Stachys verticillis duodecim-floris, caule villosa. *Hort.*
sp. 170 *

Stachys foliis lanceolato-ovatis basi emarginatis, verticillis tomentosis. *Hort. cliff.* 301. *Roy. lugdb.* 317.
Dalib. parif. 180.

Stachys latifolia major, foliis obscure virentibus, flore
galeato ferrugineo. *Pluk. alm.* 356. *t.* 317. *f.* 4.

Pseudo-stachys alpina. *Bauh. pin.* 236. *prodr.* 113.

Habitat in Germania, Helvetia. 2

S. alpina *minor*, colore obscura, corollarum galea horizontali, foliorum dentibus apice cartilagineo fusco. S.
germanica altior, lana alba, corollarum galea erecta.
Foliorum serraturis subimbricatis.

- 4 STACHYS verticillis multifloris, foliorum serraturis imbricatis.

Stachys verticillis quadraginta-floris, caule lanato. *Hort.*
sp. 170. *

Stachys foliis oblongo-cordatis, floribus verticillatis.
Hort. cliff. 309. *Roy. lugdb.* 317. *Dalib. parif.* 179.

Stachys major germanica. *Bauh. pin.* 236.

Stachys. *Fuchs. hist.* 766.

Habitat in Germania, Anglia, Gallia. 2

- 5 STACHYS verticillis triginta-floris, calycibus pungentibus. *Hort. sp.* 170. *

Stachys folio obscure virente, flore purpurascente.
Waltb. hort. 108. *t.* 19.

Stachys cretica. *Bauh. pin.* 236. *prodr.* 113.

Habitat in Creta. 2

- 6 STACHYS ramis ramosissimis, foliis lanceolatis glabris. *Hort. cliff.* 310. *Roy. lugdb.* 318.

Sideritis glutinosa bitumen redolens. *Marif. hist.* 3. *p.*
389.

Sideritis viscosa cretica bitumen redolens. *Zan. hist.* 136.

Habitat in Creta.

- 7 STACHYS ramulis spina terminatis. *Hort. cliff.* 310. *spinosa*.
Roy. lugdb. 318.

Stachys spinosa cretica. *Bauh. pin.* 236. *Morif. hist.*
3. *p.* 283. *f.* 11. *t.* 10. *f.* 9.

582 DIDYNAMIA GYMNOSPERMIA.

Gaidarothymo. *Clus. hist.* 2. p. 311.

Habitat in occidentali Cretæ.

- orientalis.* 8. STACHYS foliis tomentosis ovato-lanceolatis: flore
libus verticillo brevioribus. *Roy. lugdb.* 318.
Stachys orientalis altissima foetidissima. *Tournef. cor.* 13
Boerb. lugdb. 1. p. 154.
Habitat in Oriente.

628

BALLOTA.

- nigra* 1. BALLOTA foliis cordatis indivisis serratis, *Fl. suec.*
484.
Ballota, *Hort. cliff.* 311. *Roy. lugdb.* 315.
Ballote. *Fuchf. hist.* 154. *Cam. epit.* 572.
Marrubium nigrum foetidum. *Bauh. pin.* 230.
Habitat in Europæ ruderatis. &
- lanata.* 2. BALLOTA foliis palmatis dentatis, caule lanato.
Ballote foliis geranii batrachoides. *Amm. rusb.* 47.
Habitat in Sibiria.
Caules, lana alba induti. Verticilli hirsutissimi. Folia
grossularia, semizfida s. semisfida, lobis obtusis tri-
dentatis, supra glabra, subtus hirsuta. Calyces 5den-
tati denticulis setaceis. Corollæ albæ, ventricosa,
magna in hac gente.

629

MARRUBIUM.

* *Calycibus 5dentatis.*

- Alysson.* 1. MARRUBIUM foliis cuneiformibus, verticillis in-
volucro destitutis, *Hort. cliff.* 311. * *Roy. lugdb.*
314. *Sauv. monsp.* 151.
Marrubium album, foliis profunde incis, flore cæru-
leo. *Morif. hist.* 3. p. 377. f. 11. t. 10. f. 12.
Alysson verticillatum, foliis profunde incis. *Bauh. pin.*
232.
Alysson galeni. *Clus. hist.* 2. p. 35.
Habitat in Hispania. &
- peregrinum.* 2. MARRUBIUM foliis ovato-lanceolatis serratis, ca-
lycum denticulis setaceis. *Hort. cliff.* 311. n. 3. *Roy.*
lugdb. 314.
Marrubium album latifolium peregrinum. *Bauh. pin.*
230. *Morif. hist.* 3. p. 577. f. 11. t. 9. f. 8.
Marrubium alterum paunicum. *Clus. hist.* 2. p. 34.
β. Marrubium album angustifolium peregrinum. *Bauh.*
pin. 230. *Mæ-*

DIDYNAMIA GYMNOSPERMIA. 583

Marrubium creticum. *Dalech. hist.* 962.

Habitat in Sicilia, Cretæ, Austriæ siccis. 2

3. MARRUBIUM foliis subovatis lanatis superne emarginato-crenatis, denticulis calycinis subulatis. *Hort. cliff.* 312. n. 4. MARR.

Marrubium album candidissimum & villosum. *Tournef.*

cor. 12.

Habitat in Creta?

4. MARRUBIUM dentibus calycinis setaceis rectis villosis. *Hort. cliff.* 312. *Roy. lugdb.* 315. *Sauv. monsp.* 151. supinum.

Marrubium album, sericeo parvo & rotundo folio. *Barr.* ic. 685. *Bocc. mus.* 2. p. 78. t. 69.

Marrubium album hispanicum majus. *Barr.* ic. 686.

Habitat in Hispania, G. Narbonensi. 2

* *Calycibus 10dentatis.*

5. MARRUBIUM dentibus calycinis setaceis uncinatis. *vulgare*
Hort. cliff. 342. *Fl. suec.* 485. *Mat. med.* 305. *Roy. lugdb.* 315. *Hall. helv.* 648. *Dalib. parif.* 182.

Marrubium album vulgare. *Bauh. pin.* 230.

Marrubium vulgare. *Clus. hist.* 2. p. 34.

Habitat in Europæ borealis ruderatis. 2

6. MARRUBIUM foliis cordatis subrotundis emarginato-crenatis. *Hort. cliff.* 311. *Roy. lugdb.* 314. africanum.

Pseudo-Dictamnus africanus, foliis subrotundis subtruncatis. *Comm. hort.* 2. p. 179. t. 90.

Habitat ad Cap. b. Spei. 2

7. MARRUBIUM calycum limbis patentibus: denticulis acutis. *Hort. cliff.* 312. *Hort. ups.* 169. * *Roy. lugdb.* 315. hispanicum.

Marrubium album rotundifolium hispanicum *Herm.* par. 201. t. 201.

Marrubium subrotundo folio. *Barr.* ic. 767. *Bocc. mus.* 2. p. 167. t. 122.

Habitat in Hispania. 2

8. MARRUBIUM calycum limbis planis villosis, foliis pseudo-Dictamnus cordatis, caule fruticoso. *Hort. cliff.* 312. * *Hort. stamnis.* *ups.* 169. *Roy lugdb.* 315.

Pseudo-Dictamnus verticillatus inodorus. *Bauh. pin.* 222.

Pseudo-Dictamnus. *Dod. pempt.* 281

Habitat in. Creta. 5

O o 4

9. MAR-

acetabulo-
sum,

9. MARRUBIUM calycum limbis tubo longioribus membranaceis: angulis majoribus rotundatis.

Pseudo Dictamnus acetabulis moluccæ. *Bauh. pin.* 222.

Dictamnus falsus verticillatus, pericarpio conoide, baccatus. *Barr. ic.* 129. *

Habitat in Creta. &

Limbus calycis absoluta florentia excrescit, fit tubo duplo longior, nudus & membranaceus, nec villosus at in præcedentibus; limbi margo 20 angulis: minoribus scilicet acutis, latioribus obtusissimis rotundatis.

630

LEONURUS.

Cardiaca.

1. LEONURUS foliis caulinis lanceolatis trilobis. *Hort.*

cliff. 313. *Fl. sæc.* 496. *Mat. med.* 308. *It. scan.*

231. *Roy. lugab.* 312. *Dalib. parif.* 182.

Marrubium Cardiaca dictum. *Bauh. pin.* 230.

Cardiaca. *Fuchf. hist.* 395.

β. Cardiaca crispæ. *Raj. hist.* 572.

Habitat in Europæ ruderatis. &

Marrubia-
strum.

2. LEONURUS foliis ovatis lanceolatisque serratis, calycibus sessilibus spinosis.

Sideritis foliis ovatis & lanceolatis inciso-serratis. *Hort. cliff.* 313. *Sauv. monsp.* 279.

Marrubialtrum foliis cardiacæ. *Boett. mnsf.* 2. p. 120. t. 98.

Habitat in Bohemia, Ucraina.

catarrhus.

3. LEONURUS foliis tripartitis laciniatis, calycibus villosis. *Hort. upf.* 171. *

Sideritis foliis tripartito-multifidis. *Hort. cliff.* 313. *Hort. upf.* 169.

Cardiaca, foliis tenuius & profundius incisus, glabra s. villosa. *Amm. ruth.* 49. 50. ♂

Habitat in Tataria. ♂

libricus.

4. LEONURUS foliis tripartitis multifidis linearibus obtusiusculis *Hort. upf.* 170. *

Ballote inodora, foliis coronopi. *Amm. ruth.* 48. t. 8.

Habitat in Sibiria, China. ♂

631

PHLOMIS.

fruticosa.

1. PHLOMIS foliis subrotundis tomentosis crenatis, involucris lanceolatis.

Phlomis involucris tomentosis lanceolatis, foliis cordatis villosis, caule fruticoso. *Hort. upf.* 171.

Phlo-

DIDYNAMIA GYMNOSPERMIA. 585

Phlomis involucri radiis lanceolatis villosis. Hort. cliff.

315.

Verbascum latis salviæ foliis. Bauh. pin. 240.

Verbascum sylvestre alterum. Dod. pempt. 146.

2. *Phlomis cretica fruticosa, folio subrotundo, flore luteo.*

Tournef. cor. 10.

Phlomis latifolia capitata lutea grandiflora. Dill. elth. 316.

t. 237. f. 306.

Habitat in Sicilia, Hispania. 5

3. *PHLOMIS involucri lanceolatis, foliis ovatis subtus purpurea, tomentosis.*

Verbascum subrotundo salviæ folio. Bauh. pin. 240.

Salvia fruticosa, cisti folio haud incano, floribus purpureis. Pluk. alm. 329. t. 57. f. 6.

Phlomis fruticosa lusitanica, flore purpurascente, foliis acutioribus, Tournef. inst. 178.

Habitat in Lusitania. 5

3. *PHLOMIS foliis radicalibus cordatis utrinque tomentosis villosis.*

Phlomis orientalis, foliis auriculatis incanis, flore luteo. Nissol.

Habitat in Oriente.

Folia radicalia profunde cordata, oblonga, obtusiuscula; floralia ovata. Calyces hirsuti.

4. *PHLOMIS foliis lanceolatis tomentosis: floralibus o-lychnitis, varis, involucri setaceis lanatis.*

Phlomis foliis ligulatis utrinque tomentosis, radiis involucri setaceis villosis. Sauv. monsp. 143.

Verbascum angustis salviæ foliis. Bauh. pin. 240.

Verbascum sylvestre, foliis salviæ tenuifoliæ. Lob. ic. 558.

Habitat in Europa australi. D. Sauvages.

Habitus speciei 1, sed Folia angustiora. Corollæ vix calyce majores. Involucra linearia, pilis longis crinita.

5. *PHLOMIS foliis alternatim pinnatis: foliolis lacinia- laciniatis, calycibus lanatis.*

Phlomis orientalis, foliis laciniatis. Tournef. cor. 10.

Habitat in Oriente.

6. *PHLOMIS foliis ovatis subtus tomentosis, involu-cri subulatis strictis tripartitis.*

Phlomis radiis involucri setaceis trifidis, foliis cordatis.

Sauv. meth. 152.

O o 5

Phlo-

586 DIDYNAMIA GYMNOSPERMIA.

Phlomis involucri radiis subulatis strictis. *Hort. cliff.*

315. *

Phlomis samia herbacea, folio lunariæ. *Tournef. cor.* 10.

Habitat in Samo.

herba venti. 7. PHLOMIS involucris setaceis hispidis, foliis ovato-oblongis scabris, caule herbaceo. *Hort. upf.* 171.

Sauv. monsp. 152.

Phlomis involucri radiis setaceis hispidis. *Hort. cliff.*

315.

Marrubium nigrum longifolium. *Bauh. pin.* 230.

Habitat in Persia, Tataria, Narbona ad aggeres, &

tuberosa. 8. PHLOMIS involucris hispidis subulatis, foliis cordatis scabris, caule herbaceo. *Hort. upf.* 171.

Phlomis foliis cordiformibus, galea lacera. *Act. goss.*

211. t. 12.

Phlomis, urticæ foliis, glabra. *Amm. ruth.* 40. 39.

Galeopsis maxima, foliis hormini. *Buxb. cent.* 1. p. 4. t. 6.

Habitat in Sibiria campestribus. &

Involucra subulata & Calycis dentes patentes, hispidi.

Corollæ galea dentata. Radix tuberosa.

seylanica. 9. PHLOMIS foliis lanceolatis subserratis, capitulis terminalibus, calycibus octodentatis.

Leonurus foliis lanceolatis subserratis obtusiusculis, calycibus octodentatis. *Fl. zeyl.* 227. *

Cardiaca f. Leonurus, hyssopi hirsutis foliis, maderaspatana, floribus dilute rubris, unco tantum verticillo in ramulorum cymis. *Pluk. alm.* 81. t. 118. f. 4.

Habitat in India.

indica. 10. PHLOMIS involucris linearibus, calycibus unilabatis obliquis, foliis lanceolato-ovatis pilosis.

Habitat in India.

Folia ovata, serrata, basi attenuata, petiolata, pilis mollibus vestita, viridia. Verticilli versus apicem duo vel tres, crassi, cincti Involucris erectis, linearibus, carinatis, villosis. Calyces tubulosi, oblongi, decagoni, exteriori margine oris producti in labium 3 dentatum; a tergo, ubi margo brevissimus, dentes duo minutissimi. Corolla Leonuri Cardiaca, galea hirsutissima pilis albis, longior, incana. Labium inferius parvum.

nepetaefolia. 11. PHLOMIS foliis ovatis, calycibus decagonis septendentatis inæqualibus. *Hort. cliff.* 312. sub *Leonuro*

Car-

DIDYNAMIA GYMNOSPERMIA 587

Cardiaca americana annua, nepetæ folio, floribus brevibus phœniceis villosis. *Herm. lugdb.* 115. t. 113.

Habitat Surinami? &

12. PHLOMIS foliis lanceolatis serratis, involucris lineari-*Leonurus* ribus nudis, calycibus decagonis decemdentatis.

Leonurus foliis lanceolatis obtuse serratis. *Hort. cliff.* 312.

Leonurus capitis bonæ spei. *Brevn. cent.* 171. t. 86.

Stachys africana frutescens angustifolia, flore longissimo phœniceo, *Leonurus* dicta, *Morif. hist.* 3. p. 383.

f. 11. t. 19. f. 17.

Habitat ad Cap. b. Spei. &

6 3 2 MOLUCCELLA.

1. MOLUCCELLA calycibus quinquedentatis: denticulis æqualibus. *Roy. lugdb.* 314. *Hort. upf.* 172.

Molucca lævis. *Dod. pempt.* 92.

Melissa moluccana odorata. *Baub, pin.* 229.

Habitat in Syria. ☉

2. MOLUCCELLA calycibus septemdentatis. *Roy, spinosa.* *lugdb.* 314. *Hort. upf.* 172.

Molucca spinosa. *Dod. pempt.* 92.

Melissa moluccana foetida. *Baub, pin.* 229.

Habitat in Moluccis. ☉

3. MOLUCCELLA calycibus quinquefidis, corollis calyce longioribus.

Scordium spinosum, floris labio superiore f. galea lanuginosa f. villosa. *Raj. suppl.* 311.

Habitat in Pedemontio inter Tende & Braille e rupibus. *D. Sauvages.* &

Fruticulus Ramulis dichotomis, Folia ovata, pollicaria, pilis raris pubescentia, utrinque 2. f. 3. profundis serraturis notata, petiolata. Spinæ axillares, utrinque ternatae ut in Berberide, sed tantum ad florum verticillos. Calyces infundibuliformes, parvi, decagoni, limbo quinquesido, spinis terminalibus 12 ad 15. Corolla cardiaca, galea lanata. Media inter Balloten, Leonurum & Moluccellam.

CLINOPODIUM 6 3 3

1. CLINOPODIUM capitulis subrotundis hispidis, bracteis fetacis.

Climo-

588 DIDYNAMIA GYMNOSPERMIA.

Clinopodium foliis ovatis, capitulis verticillatis. Hort. cliff. 305. Fl. succ. 479. Roy. lugdb. 313.
Clinopodium origano simile. Bauh. pin. 224.
Clinopodium. Cam. epit. 563.
Habitat in rupestribus Europæ, Canadæ. &
Americana ex seminibus duplo minores gerit corollas, cæ-
terum eadem.

634
 incanum.

2. CLINOPODIUM foliis subtus tomentosis, verticillis explanatis, bracteis lanceolatis.
Clinopodium foliis lanceolatis serratis, verticillis pedunculatis. Roy. lugdb. 313.
Clinopodium, menthæ folio, incanum & odoratum. Dill. elsh. 87. t. 74. f. 85.
Clinopodium majus virginianum non ramosum, verticillis majoribus, floribus brevioribus carnis. Moris. hist. 3. p. 374. f. 11. t. 8. f. 4.
Clinopodium Serpentaria dictum, latiori folio, capitulis grandioribus. Pluk. mant. 51. t. 344. f. 7. Raj. suppl. 298.
Origanum foliis ad summum caulium canis. Raj. hist. 1229.
Habitat in Europa boreali. &

rugosum.

3. CLINOPODIUM foliis rugosis, capitulis axillaribus pedunculatis explanatis radiatis.
Clinopodium rugosum, capitulis scabiosæ. Dill. elsh. 88. t. 75. f. 86.
Scabiosæ affinis, chrysanthemi facie, lamii foliis, americana. Pluk. alm. 335. t. 222. f. 1.
Sideritis spicata, scrophulariæ folio, flore albo, spicis brevibus habitioribus rotundis pediculis intidentibus. Sloan. jam. 65. hist. 1. p. 174. t. 109. f. 2.
Mentha americana inodora, foliis subincanis. Raj. suppl. 284.
Mentha melissoides americana. Pluk. mant. 129.
Melissa altissima globularia. Plum. spec. 6.
Habitat in Carolina, Jamaica, Gallia æquinoctiali. &

634

ORIGANUM.

ægyptiacum.

1. ORIGANUM foliis carnosis tomentosis.
Majorana rotundifolia scutellata exotica. Tournef. inst. 199.
Origano cognata Zatarhendi. Bauh. pin. 223. Moris. hist. 3. p. 360. f. 11. t. 3. f. ult.

Zatar-

Zatarhendi. *Alp. ægypt.* 94. t. 95.

Habitat in Ægypto. 2

2. **ORIGANUM** foliis inferioribus tomentosis, spicis nutantibus. *Hort. cliff.* 304. *Mat. med.* 106. *Roy. lugdb.* 323.

Dictamnus creticus. Bauh. pin. 222.

Dictamnus cretense. Cam. epit. 472

Dictamnus. Lob. ic. 502.

Habitat in Cretæ monte Ida. 3

3. **ORIGANUM** foliis omnibus glabris, spicis nutanti-
bus. *Hort. cliff.* 304. *Roy. lugdb.* 323.

Origanum montis sipyli. Herm. lugdb. 462. t. 463.

Dictamnus sipyleus, majoranæ foliis. Moris. hist. 3.

p. 457. f. 11. t. 4. f. 2.

Habitat in Phrygiæ monte Sipulo.

4. **ORIGANUM** spicis aggregatis longis prismaticis re-
ctis, bracteis membranaceis, calyce duplo longioribus.

*Origanum foliis ovatis acutis glabris: venis scabris, spi-
cis tetragonis. Mat. med.* 300.

*Origanum foliis ovatis obtusis scabris integerrimis, spi-
cis confertis compactis glabris. Amœn. acad.* 1. p. 160.

Origanum creticum. Bauh. pin. 223.

- β. *Origanum folio subrotundo. Bauh. pin.* 223.

Origanum monspeliense pulchrum. Cam. epit. 468.

Habitat in Europa australi. 2

5. **ORIGANUM** foliis ovatis acutis serratis, spicis con-
gestis umbellatim fastigiatis. *Hort. cliff.* 304. *Roy.*

lugdb. 324.

Origanum smyrnæum. Wheel. itin. - - *Raj. hist.* 450.

*Majorana cretica, origani foliis, villosa, saturejæ odo-
re, corymbis majoribus albis. Tournes. cor.* 13.

*Habitat in Creta & Smyrnæ in monte cui arx inædi-
ficata.*

6. **ORIGANUM** spicis longis pedunculatis aggregatis, *heracleoti-
cum.*
bracteis longitudine calycum.

Origanum heracleoticum, Cunila gallinacea Plinii.

Bauh. pin. 223.

Orega, Origanum heracleoticum, Cunila. Lob. ic. 492.

Habitat in Europa australi, Græcia. 2

*Hæbitus O. sylvestris, sed spicæ digitales, laxè imbrica-
tæ; at differt ab O. cretico, quod bractæ vix calyci-
bus longiores & demum quod spicæ pedunculatæ.*

7. ORI-

590 DIDYNAMIA GYMNOSPERMIA.

- vulgare.* 7. **ORIGANUM** spicis subrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis.
Origanum foliis ovatis, spicis laxis erectis confertis paniculatis. *Hort. cliff.* 305. *Fl. suec.* 480. *Mat. med.* 299. *Roy. lugdb.* 323. *Gron. virg.* 65.
Origanum sylvestre. *Baub. pin.* 223.
Origanum sylvestre f. *vulgare.* *Fuchs. hist.* 522.
Habitat in Europæ, Canadæ rupestribus.
Americanum bracteis coloratissimis gaudet.
- mitet.* 8. **ORIGANUM** spicis oblongis aggregatis hirsutis, foliis cordatis tomentosis.
Origanum lignosum syracusanum perenne, unibella amplissima brevi, lato & nervoso folio. *Bocc. mus.* 2. p. 45. t. 38.
Origanum onites. *Baub. pin.* 222. an?
Habitat Syracusæ. 5
Habitus Majoranæ sed lignosus. Caules pilis longis patulis. Folia parva, cordata, subsessilia, acuta, rarius serrata, utrinque tomentosa, ex alis ramorum rudimenta. Spicæ congestæ, ut in Majorana, sed oblongæ, in singulo pedunculo ternæ, intermedio sessili, villosæ; Flores albi.
- Syriacum.* 9. **ORIGANUM** spicis longis ternatis pedunculatis villosis, foliis ovatis villosis.
Majorana syriaca vel cretica. *Baub. pin.* 224.
Marum syriacum. *Lob. ic.* 499. *
Habitat - - - - -
Habitus Majoranæ. Folia subrotundo-ovata, acuta, pilis vagis adpersa. Rami ex axillis. Corymbus terminalis, ex ramis longioribus brachiatus. Spicæ sæpius ubique 3, terminales, longæ, erectæ, tetragonæ, villosæ.
- Majorana.* 10. **ORIGANUM** foliis ovalibus obtusis, spicis subrotundis compactis pubescentibus. *Hort. cliff.* 304. *Hort. ups.* 151. *Mat. med.* 298. *Roy. lugdb.* 324.
Majorana vulgaris. *Baub. pin.* 224.
Amaracus vulgator. *Lob. ic.* 498.
 11. *Majorana* tenuifolia. *Baub. pin.* 224.
Majorana hortensis odorata perennis. *Morif. hist.* 3. p. 359.
Habitat - - - - -

THYMUS.

- Scorpyllum.* 1. **THYMUS** floribus capitatis, caulibus repentibus, foliis planis obtusis basi ciliatis. *Fl. suec.* 477. *Mat. med.* 232. *Thy-*

Thymus repens, foliis planis, floribus verticillato-spicatis. *Hort. cliff.* 306. *Roy. lugdb.* 325.

Serpyllum vulgare minus. *Baub. pin.* 220.

Serpyllum vulgare. *Dod. pempt.* 277.

β. *Serpyllum vulgare majus*. *Baub. pin.* 220.

γ. *Serpyllum vulgare minus*, capitulis lanuginosis. *Tournef. inst.* 197. *It. goetl.* 219.

δ. *Serpyllum angustifolium hirsutum*. *Baub. pin.* 220.

ε. *Serpyllum foliis citri odore*. *Baub. pin.* 220.

Habitat in Europæ aridis apricis. θ

1. **THYMUS erectus**, foliis revolutis ovatis, floribus *vulgaris*. verticillato-spicatis. *Hort. cliff.* 305. *Hort. upf.* 160.

Mat. med. 281. *Roy. lugdb.* 325. *Sauv. monsp.* 148.

Thymus vulgaris, folio tenuiore. *Baub. pin.* 219.

β. *Thymus vulgaris*, folio latiore. *Baub. pin.* 219.

Thymum durius. *Dod. pempt.* 276.

Habitat in G. Narbonensis, Hispaniæ montosis saxosis. θ

3. **THYMUS** floribus verticillato-spicatis, caule suffru-Zygis. ticofo, foliis linearibus basi ciliatis. *Læsl.*

Thymo vulgatori rigidiori simile. *Bauh. hist.* 2. p. 271.

Thymum angusto longiorique folio. *Barr. ic.* 777.

Serpyllum sylvestre Zygis diofcoridis. *Clus. hist.* 358.

Serpyllum folio thyma. *Baub. pin.* 220?

Habitat in Hispania.

Facies T. vulgaris, at Folia basi ciliata.

4. **THYMUS** floribus verticillatis, pedunculis unifloris, *Acinet.* caulibus erectis subramosis, foliis acutis serratis.

Fl. succ. 478.

Thymus caulibus vix ramosis, foliis ovatis acutis, pedunculis plurimis unifloris. *Hort. cliff.* 306. *Roy. lugdb.* 325.

Clinopodium arvense, ocymi facie. *Baub. pin.* 225

Clinopodium vulgare. *Lob. ic.* 506.

Habitat in Europæ glareosis, cretaceis, ficcis. ☉

5. **THYMUS** verticillis sexfloris, foliis obtusiusculis *alpinus.* concavis subserratis.

Clinopodium verticillis paucifloris in spicam congestis.

Hall. belv. 653.

Clinopodium montanum. *Baub. pin.* 225. *Bocc. mus.*

2. p. 50. t. 45.

Acini pulchra species. *Bauh. hist.* 3. p. 620.

Ha-

592 DIDYNAMIA GYMNOSPERMIA

Habitat in Alpibus Helveticis, Austriacis, Monspeli.

D. Rathgeb. ☉

Habitus & affinitas cum antecedenti.

cephalotos. 6. THYMUS capitulis imbricatis magnis, bracteis ovalis, foliis lanceolatis.

Thymus lusitanicus cephalotos, squamis capitulorum amplioribus. *Tournef. inst.* 196.

Tragoriganum, dictamni capite, hispanicum. *Barr. ic.* 787. *Bocc. mus.* 2. p. 50. t. 43.

β. Thymus lusitanicus cephalotos, capitulo majori. T.

γ. Thymus lusitanicus cephalotos, capitulo minori. T.

Habitat in Hispania, Lusitania. ♀

villosus. 7. THYMUS capitulis imbricatis magnis, bracteis densatis, foliis setaceis pilosis.

Thymus lusitanicus, folio capillaceo villoso, capite magno purpurascente oblongo. *Tournef. inst.* 196.

β. Thymus lusitanicus, folio capillaceo villoso, capite magno purpurascente rotundo. *Tournef. inst.* 190.

Habitat in Lusitania. ♀

pulegioides. 8. THYMUS ramis verticillatis longis, foliis ovalibus striatis.

Acinos thymi folio & facie, floribus inexpandis. *Morif.*

hist. 3. 404. f. 11. t. 19. f. 6.

Habitat Monspeli. D. Sauvages. ☉

Habitus *Melissae pulegioidis* f. *Menthae pulegii*, at

Caryocylus bilabiatus a *Satureja*, ore barbati a *Melissa* &

Mentha eam distinguunt. Caulis spithameus, erectus.

Rami pauci, simplices, breves. Folia ovalia, obtusa,

glabra, subius striata. Rami & planta maxima parti

verticillis obvallata.

MELISSA.

officinalis.

1. MELISSA racemis axillaribus verticillatis: pedicellis simplicibus.

Melissa floribus ex alis inferioribus subsessilibus. *Hort. cliff.* 307. *Hort. upf.* 163. *Mat. med.* 294. *Roy. lugd.*

320. *Sauv. monsp.* 149

Melissa hortensis. *Baub pin.* 229.

Apialtrum f. *Melissophyllum* *Lob. hist.* 227.

Habitat in montibus Geneventibus, Allobrogicis, Italicis. 2

grandiflora. 2. MELISSA pedunculis axillaribus dichotomis longitudine florum. Calg.

DIDYNAMIA GYMNOSPERMIA 593.

Calamintha magno flore. Bauh. pin. 229. Riv. mon. 43.

Calamintha montana præstantior. Besl. cyst. æst. 7. t.

7. f. 1.

Habitat in Hetruriæ montosis. 2

5. MELISSA pedunculis axillaribus dichotomis longitu- *Calamintha*,
dine foliorum.

*Melissa floribus ex alis superioribus pedunculo commu-
ni conertis. Hort. cliff. 307. Roy. lugdb. 320. Mat.
med. 295. Sauv. monsp. 149.*

*Calamintha vulgaris & officinarum germaniæ. Bauh.
pin. 228.*

Calamintha montana. Dod. pempt. 98.

Habitat in Italiæ, Hispaniæ, Galliæ clivis saxosis. 2

6. MELISSA pedunculis axillaribus dichotomis folio lon- *Nepeta*,
gioribus, caule decumbente.

*Melissa floribus ex alis superioribus, pedunculo dichoto-
mo, caule procumbente. Hort. cliff. 308. Gron. virg. 65.*

Roy. lugdb. 321. Sauv. monsp. 149. Gort. gelr. 337.

Calamintha pulegii odore f. Nepeta. Bauh. pin. 228.

6. *Calamintha montana præalta, pulegii odore. Bocc. mus.
2. p. 45. t. 40. f. 38.*

*Habitat in Italiæ, Galliæ, Angliæ, Helvetiæ aggeribus
glareosis.*

5. MELISSA racemis terminalibus, pedunculis solitariis *cretica*,
brevissimis.

Calamintha incana, ocyni foliis. Bauh. pin. 569.

Calamintha, pulegii odore, minor. Barr. ic. 1166.

Calamintha clinopodii austriaci foliis, odore pulegii.

Pluk. alm. 75. t. 163. f. 4

Habitat Monspelii.

6. MELISSA fruticosa, ramis attenuatis virgatis, foliis *fruticosa*,
subtus tomentosis.

*Melissa caule fruticoso, racemis verticillatis. Roy. lugdb.
321.*

*Calamintha hispanica frutescens, mari folio. Tournef.
inst. 194.*

Habitat in Hispania. 8

7. MELISSA floribus verticillatis glomeratis secundum *Pulegioid.*
longitudinem caulis, foliis tomentosis. *Gron. virg.*

107.
Habitat in Virginia, Canada. ☉

*Planta spithamea, brachiata: Ramis brevioribus, simpli-
P p cibus.*

cibus. Folia lanceolata, glabra. Verticilli laxi florum a radice ad summitates plantæ conferti. Differt a Thymo pulegioides, cui simillima, non tantum floribus minime spicatis ut in thymo, verum lateralibus per caulem sparsis, sed præcipue fauce calycis vix manifeste villosa, minime pilis albis clausa. Folia in hac utrinque uno alterove dente notata, in Thymo autem sæpius integerrima.

637

DRACOCEPHALUM.

* *Spicata.*

virginianum 1. DRACOCEPHALUM floribus spicatis, foliis lanceolatis serratis.

Dracocephalum foliis simplicibus, floribus spicatis. *Hort. cliff.* 308. *Roy. lugab.* 311.

Dracocephalum. *Breyn. ic.* 33. t. 27. *Hir. æt.* 1712. p. 276. t. 11.

Dracocephalus angustifolius, folio glabro serrato. *Morif. hist.* 3. p. 407. f. 11. t. 4. f. 1.

Pseudodigitalis perficæ foliis. *Bocc. sic.* 12. t. 6. f. 3.

Lysimachia galericulata spicata purpurea canadensis. *Barr. ic.* 1152.

Habitat in America septentrionali. 2

canariense. 2. DRACOCEPHALUM floribus spicatis, foliis compositis. *Hort. cliff.* 308. *Mat. med.* 293.

Dracocephalo affinis americana trifoliata, terebinthinæ odore. *Volk. norib.* 145. t. 145.

Camphorosina. *Morif. hist.* 3. p. 366. f. 11. t. 11. f. ult.

Melissa forte canarina triphyllis, odorem camphoræ spirans penetrantissimum. *Pluk. alm.* 401. t. 325. f. 5.

Cedronella canariensis viscosa, foliis plerumque ex eodem pedicello ternis. *Comm. hort.* 2. p. 81. t. 41.

Habitat in Canariis. 3

pinnatum. 3. DRACOCEPHALUM floribus spicatis, foliis cordatis pinnato-sinuatis.

Dracocephalum foliis pinnato-sinuatis obtusis, floralibus spicatis villosis coloratis. *Hort. upf.* 165. *

Habitat ad Jerkatsch. Steller, D. Gmelin.

Caules fruticosi, decumbentes. Folia cordata, transversim pinnatifida, obtusa lobis remotis, longe petiolata. Spica Melampyri arvensis: bracteis lanceolatis, denticato-setaceis, rubris sæpe. Corollæ parvæ; Styli floribus duplo longiores

4. DRA-

4. DRACOCEPHALUM floribus spicatis, foliis bracteatis. *confertis*.
 Etisque linearibus partitis spinosis.
 Hyssopus spicis interruptis *Hort. cliff.* 364.
 Hyssopus austriacus, magno flore, folio chamæpithyos.
Herm. lugdb. 330.
 Ruyschiana hirsuta, foliis laciniatis. *Amm. rusb.* 50.
 Chamæpithys cærulea austriaca *Bamb. pin.* 250.
 Chamæpithys austriaca. *Clus. hist.* 2. p. 185.
Habitat in Austria. 2
5. DRACOCEPHALUM floribus spicatis, foliis bracteatis. *Ruyschiana*.
 Etisque lanceolatis indivisis muticis.
 Dracocephalum foliis linearibus indivisis integerrimis,
 floribus spicatis, *Hort. nps.* 165. *
 Ruyschiana glabra, foliis integris. *Amm. rusb.* 50.
 Ruyschiana flore cæruleo magno. *Boerb. lugdb.* 1 p. 172.
 PseudoChamæpithys austriaca. *Riv. mon.* 106.
 Prunella hyssopi folio viridi, amplo flore cæruleo. *Mor-
 ris. hist.* 3. p. 364. f. 11. t. 5. f. 9.
Habitat in Sibiria, Suecia. 2
 * *Verticillata.*
6. DRACOCEPHALUM floribus verticillatis, foliis grandifloris.
 ovatis inciso-crenatis, bracteis lanceolatis integerrimis. *rum.*
 Dracocephalum floribus verticillatis, foliis oblongis ob-
 tusis sinuato-crenatis, bracteis oblongis. *Gmel. nss.*
Habitat in Sibiria. ☉
 Folia ovata s. ovalia, profunde & obtuse serrata, nuda:
 canina trimeria, floralia integerrima. Bractea lan-
 ceolata, integerrima, parva, pilis adspersa. Corol-
 la maxima, cærulea.
7. DRACOCEPHALUM floribus verticillatis, bracteis *Moldavica*.
 lanceolatis: serraturis capillaceis. *Hort. nps.* 166. *
Mat. med. 292.
 Dracocephalum floribus verticillatis, foliis ovato-lan-
 ceolatis. *Hort. cliff.* 308. *Roy. lugdb.* 312.
 Melissa peregrina, folio oblongo. *Bamb. pin.* 229.
 Melissa moldavica. *Cam. epit.* 576.
Habitat in Moldavia. ☉
8. DRACOCEPHALUM floribus verticillatis, bracteis *crucicatis*.
 is oblongis: serraturis spinosis, foliis subtoментosis.
Hort. nps. 66. *
 Dracocephalum floribus verticillatis, foliis lanceolatis,
 floribus oblongis. *Hort. cliff.* 308. *Roy. lugdb.* 312.
 Moldavica orientalis, betonica folio, flore magno vio-
 laceo. *Tournef. cor.* 11. *Comm. rar.* 28. t. 28.
 P p 2 Si-

596 DIDYNAMIA GYMNOSPERMIA.

Sideritis annua, flore luteo, utriculis & foliis longioribus. *Morif. hist.* 3. p. 389. f. 11. t. 8. f. 18.
Habitat in Oriente. ☉

- pelantum.* 9. DRACOCEPHALUM floribus verticillatis, bracteis orbiculatis serrato-ciliatis. *Hort. upf.* 166. *
Dracocephalum floribus verticillatis, foliis floralibus orbiculatis. *Hort. cliff.* 309. * *Key lugdb.* 312.
Moldavica orientalis, falcis folio, parvo flore cæruleo.
Tournef. cor. 41.
Habitat in Oriente. ☉

- nuttans.* 10. DRACOCEPHALUM floribus verticillatis, bracteis oblongis ovatis integerrimis, corollis majusculis nutantibus. *Hort. upf.* 167. n. 6. *
Moldavica betonica folio, floribus minoribus cæruleis pendulis. *Amm. ruth.* 44.
Habitat in Sibiria. ☉

- thymiflora.* 11. DRACOCEPHALUM floribus verticillatis, bracteis oblongis integerrimis, corollis vix calyce majoribus. *Hort. upf.* 167. *
Cedronella tatarica perennis, urticæ foliis, flosculis minoribus ex cæruleo rubentibus. *Key. lugdb.* 537.
Moldavica betonica folio, floribus minimis pallide cæruleis. *Amm. ruth.* 46.
Habitat in Sibiria. ☉

HORMINUM.

- 638
pyrenaicum. 1. HORMINUM foliis cordatis obtusis, caule nudo. *Horminum.* *Hort. cliff.* 309.
Callitrichum folio rotundo, flore magno violaceo. *Bauh. hist.* 3. p. 313. *Morif. hist.* 3. p. 395. *Raj. hist.* 546.
Melissa pyrenaica, caule brevi, plantaginis folio. *T. Magn. hort.* 133. t. 133.
Habitat in altis montium Tyrolensium, Pyrenæorum. 2
virginicum. 2. HORMINUM foliis cuneiformi-oblongis, caule bifolio.
Melissa atrorubens, pugulæ folio. *Dill. elsb.* 219. t. 175. f. 216.
Sideritis, bugulæ folio, mariana, floribus purpureis longo tubulo donatis. *Pluk. mant.* 171.
Habitat in Virginia, Carolina.

MELIT-

MELITTIS. 639

1. MELITTIS. *Hort. cliff.* 390. *Roy. lugdb.* 319. *Sauv. melissophyl.*
monsp. 150.

Melissa fuchii. *Cam. hort.* 99. 30.

Lamium monanum, *melissæ folio.* *Bauh. pin.* 231.

Habitat in subalpinis Germaniæ, Helvetiæ, Angliæ,
Monspeli. &

OCIMUM. 640

1. OCIMUM racemis secundis lateralibus, caule erecto. *frutescens.*

Ocimum foliis lanceolato-ovatis, radice perenni. *Hort.*

cliff. 315. *Fl. zeyl.* 228. * *Roy. lugdb.* 222.

Ocimum zeylanicum perenne odoratissimum latifoli-
um. *Burm. zeyl.* 174. t. 80. f. 1.

Cottam. Rheed. mal. 10. p. 153. t. 77.

Habitat in India ☉

Habet & hoc Stamina propria generi.

2. OCIMUM foliis ovatis glabris, calycibus ciliatis. *Hort. Basilicum.*

cliff. 315. *Hort. upf.* 168. *Roy. lugdb.* 322.

Ocimum caryophyllatum majus. *Bauh. pin.* 226.

Ocimum. Cam. epit. 308.

3. *Ocimum caryophyllatum maximum.* *Bauh. pin.* 226.

7. *Ocimum latifolium maculatum f. crispum.* *Bauh. pin.*
226.

8. *Ocimum viride, foliis bullatis.* *Bauh. pin.* 225.

Habitat in India, Persia. ☉

3. OCIMUM foliis ovatis integerrimis. *Hort. upf.* 169. *minimum.*

Ocimum foliis ovatis incanis. *Hort. cliff.* 315. *Roy.*

lugdb. 322

Ocimum minimum. *Bauh. pin.* 226. *Morif. hist.* 3.

p. 407. f. 11. t. 10. f. 17.

Habitat in Zeylona. ☉

4. OCIMUM foliis ovato-oblongis serratis, bracteis cor-
datis reflexis concavis. *tenuiflorum.*

Habitat in Malabar. ☉-b

Caulis pedalis f. bipedalis, teretiusculus, purpurascens,

brachiatus, pilis adpersus patulis; Rami breviores. Fo-

lia ovato-oblonga, obtuse serrata, mollia, longe petiola-

ta. Spicæ terminales, ternæ, longæ, angustæ, peduncu-

lata, Bracteis oppositis, glabris, cordatis, concavis, ar-

de reflexis. Flores tres ex singula bractea, subsessiles.

Calyces glabri ssidi: labio superiore plano, majore, re-

niformi, acuto. Corolla parva, purpurascens: Labio

P p 3

supe-

593 DIDYNAMIA GYMNOSPERMIA.

superiore 4fida; Labio inferiore simplici. Stamina 4, quorum 2 basi appendiculata. Stylus flore longior, bifidus. Flores tam parvi, ut vix nudis oculis conspiciantur, qui incipiunt fieri ab apice spicarum.

- meniboides.* 5. OCIMUM foliis lineari-lanceolatis serratis. *Fl. zeyl.* 229.
Mentha zeylanica, angustissimo folio dentato. *Raj. suppl.* 284.
Habitat in Zeylona.

641

TRICHOSTEMA.

- dichotomum.* 1. TRICHOSTEMA staminibus longissimis exsertis.
Trichostema. *Hort. cliff.* 493, *Gron. virg.* 64. *Roy. lugdb.* 310.
Scutellaria cærulea, majoranæ folio, americana. *Raj. suppl.* 317.
Cassida mariana, majoranæ folio. *Pet. jicc.* 243.
Habitat in Virginia, Pennsylvania. ☉
- brachiatum.* 2. TRICHOSTEMA staminibus brevibus inclusis.
Teucrium virginianum, origani folio. *Dill. elth.* 380.
t. 285. *f.* 369.
Habitat in America septentrionali.

642

SCUTELLARIA.

- orientalis.* 1. SCUTELLARIA foliis incisus subtus tomentosis.
Hort. nupl. 173.
Scutellaria foliis pinnatifidis. *Hort. cliff.* 316. * *Roy. lugdb.* 310.
Cassida orientalis, folio chamædrycos, flore luteo. *Tournef. cor.* 11 *itin.* 3. *p.* 306 *t.* 306. *Comm. rar.* 30. *t.* 30. *Mart. cent.* 18. *t.* 18.
β. Scutellaria orientalis incana, foliis laciniatis, flore luteo.
Tournef. cor. 11.
Habitat in America circa Tephlin. 2
- alpina.* 2. SCUTELLARIA foliis inciso-serratis utrinque glabris, spica rotundato-tetragona. *Hort. nupl.* 173. *
Cassida alpina lupina, magno flore albido. *Tournef. inst.* 182.
Habitat in Sibiria, Tartaria. 2
- laxiflora.* 3. SCUTELLARIA floribus lævibus carina scabris, necem lateralibus foliolis.
Scutellaria foliis cordato-lanceolatis serratis, pedunculis multifloris. *Roy. lugdb.* 311. *Scu-*

Scutellaria foliis ovato-lanceolatis petiolatis, racemis foliosis. *Gron. virg.* 67.

Scutellaria palustris repens virginiana major, flore minore. *Morif. hist.* 3. p. 416.

Habitat in Canada, virginia. &

Racemi laterales Foliis minoribus, lanceolatis, serratis; insuper singuli pedicelli seta utrinque notati sunt, ut in *S. galericulata* & *peregrina*.

4. *SCUTELLARIA* foliis cordato lanceolatis crenatis, *galericulata*. floribus axillaribus. *Fl. lapp.* 239. *Fl. succ.* 499. *Hort. cliff.* 316. *Roy. lugdb.* 310.

Lyfimachia cærulea *galericulata* f. *Gratiola* cærulea.

Baub. pin. 246.

Lyfimachia *galericulata*. *Dalech. hist.* 1060.

Habitat in Europæ littoribus. &

5. *SCUTELLARIA* foliis integerrimis: inferioribus hastifolia. statis, superioribus sagittatis. *It. gotl.* 212. *Fl. succ.* 500. *

Scutellaria folio non serrato. *Riv. mon.* 77.

Habitat ad littora Sueciæ rarius. &

An varietas precedentis? certe maxime affinis.

6. *SCUTELLARIA* foliis sessilibus ovatis: inferioribus *integrifolia*. obsolete serratis; superioribus integerrimis.

Scutellaria foliis integerrimis. *Gron. virg.* 67.

Scutellaria cærulea virginiana, lamell aut potius *teucriti* folio, minor. *Pluk. alm.* 338. t. 313. f. 4.

Scutellaria, *teucriti* folio, marilandica. *Raj. suppl.* 310.

Habitat in Virginia, Canada.

7. *SCUTELLARIA* foliis lanceolatis. *Gron. virg.* 167. *hystroptifolia*. *Cassida* inariana *hystroptifolia*. *Pet. act. angl.*

Habitat in Virginia.

8. *SCUTELLARIA* foliis subcordatis serratis, spicis *peregrina*. interruptis. *Hort. cliff.* 317. *Hort. nps* 172. *Roy.*

lugdb. 311.

Cassida. *Colum. ecphr.* 1. p. 187. t. 189.

Lanum peregrinum f. *Scutellaria* *Baub. pin.* 231.

Habitat in nemoribus Florentiæ, Liburni.

Alia huic simillima Calycibus lanatis, pilis griseis, Corollæ galea apice pilosa. Bracteis ovatis, brevibus.

9. *SCUTELLARIA* foliis ovatis crenatis, spicis imbricatis. *alpina*. *Hort. cliff.* 317. *Hort. nps* 173. *Roy. lugdb.*

311.

600 DIDYNAMIA GYMNOSPERMIA.

Teucrium alpinum inodorum, magno flore. *Baub. pin.* 247. *prodr.* 116. *Burf.*
Habitat in Alpibus Helvetiæ. 2

indica.

10. SCUTELLARIA foliis subovatis crenatis petiolatis, florum racemis nudiusculis.

Habitat in China.

Planta minor; Folia suborbiculata, subovata, obsolete serrata, petiolata. Racemus terminalis. Bracteis linearibus, brevissimis. Calyces brevissimi absque tubo oris. Corollæ angustæ, longiusculæ.

altissima.

11. SCUTELLARIA foliis cordato-oblongis acuminatis serratis, spicis subnudis.

Scutellaria orientalis altissima, urticæ folio. Tournef. cor. 11.

Habitat in Oriente.

cretica.

12. SCUTELLARIA foliis cordatis obtusis obtuseque serratis, spicis foliosis. *Roy. lugdb.* 311.

Cassida cretica fruticosa, folio catariæ, flore albo. Tournef. cor. 11.

Habitat in Creta.

643
vulgaris.

PRUNELLA.

1. PRUNELLA foliis ovato-oblongis petiolatis.

Prunella bracteis cordatis. Hort. cliff. 316. *Fl. succ.* 498. *Mat. med.* 309. *Roy. lugdb.* 314.

Brunella major, folio non dissecto. Baub. pin. 260.

Brunella. Dod. pempt. 136.

- grandiflora.* β. *Brunella cærulea, magno flore. Baub. pin.* 261. *It. gotl.* 219. 228.

Brunella 1. non vulgaris. Clus. hist. 2. p. 42.

laciniata.

- γ. *Brunella folio laciniato. Baub. pin.* 261.

Brunella 11. non vulgaris. Clus. hist. 2. p. 43.

- δ. *Brunella minor alba laciniata. Baub. pin.* 261.

- ε. *Brunella verbenulæ folio, flore cæruleo. Vaill. paris.*

22. t. 5. f. 1.

Habitat in Europæ pascuis. 2

Planta α in apricis procumbens, pollicaris; in nemoribus erecta pedalis. β grandiflora in rupium fissuris plerumque nascitur; γ Laciniata tam multa habet in fructificatione communia, ut vix videatur distincta.

- byssopifolia.* 2. PRUNELLA foliis lanceolato-linearibus ciliatis subsessilibus. *Sanv. monsp.* 141.

Bru-

DIDYNAMIA GYMNOSPERMIA. 601

Brunella hyssopifolia. Baub. pin. 261.

*Brunella angustifolia integra hirsutior. Moris. hist. 3.
p. 364. f. 11. t. 5. f. 7.*

Habitat Monspelii. 2

3. PRUNELLA bracteis pinnato-dentatis ciliatis. *Læf. lusitanica, descr. 31.*

Prumella odorata lusitanica, flore violaceo. Barr. ic. 561.

*Clinopodium lusitanicum spicatum & verticillatum.
Tournef. inst. 195.*

*Bugula odorata lusitanica. Comm. canad. 46. Moris. hist.
3. p. 394. f. 11. t. 5. f. 4.*

Habitat in Lusitania, Hispania.

PRASIUM. 644

1. PRASIUM foliis ovato-oblongis serratis. *Hort. cliff. majus, 309. Hort. upf. 173.*

*Lamium arboreum perenne creticum. Zan. hist. 112.
t. 46. Raj. hist. 561.*

Melissa fruticosa cretica sempervirens, teucarii facie, flore albo. Moris. hist. 3. p. 408. f. 11. t. 21. f. 3.

Teucrium fruticans, amplo & albo flore, italicum. Barr. ic. 895.

Habitat in Sicilia, Romæ & in agro Tingitano. 5

2. PRASIUM foliis ovatis crena utrinque duplici. *Hort. minus, cliff. 309. **

Lamium fruticans, teucarii folio lucido, calyce & flore magno candido: tantilla purpura varie notato. Gup. cathol. 106.

Habitat in Sicilia. 5

An hac sufficienter specie a priori distincta ulterius indagandum.

PHRYMA. 645

1. PHRYMA. *Gen. nov. 1092.*

Leptostachya.

Leptostachya. Mitch. gen. 9.

Verbena racemo simplicissimo, floribus sessilibus, calycibus fructus reflexis racemoque appressis. Gron. virg.

7. Roy. lugdb. 536.

Habitat in America septentrionali.

ANGIOSPERMIA.

646

BARTSIA.

- coccinea.* 1. BARTSIA foliis alternis linearibus utrinque bidentatis. *Barttia* foliis alternis. *Hort. cliff.* 235. *Gron. virg.* 68. *Amœn. acad.* 1. p. 160.
Pedicularis f. cristæ galli æstivis virginiana, ajugæ multifido folio: apicibus coccineis, floribus pallidis in spicam congestis. *Pluk. alm.* 283. t. 102. f. 5. *Raj. suppl.* 400.
Horminum, teuui coronopi folio, virginianum, *Morif. hist.* 3. p. 395. f. 11. t. 13. f. 28.
Habitat in Virginia, Noveboraco. 2
- pallida.* 2. BARTSIA foliis alternis lanceolatis integerrimis, floralibus ovatis dentatis.
Barttia foliis alternis bidentatis. *Amœn. acad.* 2. p. 356. *
Habitat in Sibiria D. *Gmelin. Demidoff.*
- viscosa.* 3. BARTSIA foliis superioribus alternis ferratis, floribus lateralibus.
Euphrasia latifolia viscata ferrata. *Morif. blæs.* 264.
Euphrasia major lutea latifolia palustris. *Raj. hist.* 772. *angl.* 3. p. 285.
Euphrasia lutea palustris. *Pluk. alm.* 142. t. 27. f. 5.
Euphrasia lutea latifolia palustris. *Morif. hist.* 3. p. 432.
Alectorolophos italica luteo-pallida. *Barr. rar.* 209. t. 665.
Habitat in Angliæ, Galliæ, Italiæ paludibus adriaticis. ☉
- alpina.* 4. BARTSIA foliis oppositis cordatis obtuse ferratis. *Hort. cliff.* 325. *Fl. suec.* 515. *Roy. lugdb.* 298.
Euphrasia caule simplici, foliis cordatis sessilibus obtuse ferratis. *Fl. lapp.* 246.
Clinopodium alpinum hirsutum. *Baub. pin.* 225. *Pluk. alm.* 110. t. 163. f. 5.
Clinopodium alpinum. *Pou. bald.* 343.
Teucrium alpinum, coma purpureo-cærulea. *Baub. pin.* 247.
Stachellna. *Hall. helv.* 624.
Habitat in Alpibus Lapponicis, Helveticis, Allobrogicis, Baldo, Vallicis. 2
- Trixago.* 5. BARTSIA foliis oppositis lanceolatis obtuse ferratis.
Trixago apula unicaulis. *Colum. cephr.* 1. p. 199. *Morif. hist.* 3. p. 427. f. 11. t. 2. 4. f. 8.

Chamæ-

Chamædrys unicaulis spicata. Baub. pin. 248.

Antirrhinum folio dissecto. Baub. pin. 211.

Habitat in Italiæ maritimis, bumentibus. ☉

RHINANTHUS. 647

1. RHINANTHUS corollis labio superiore subulato *orientalis*, incurvo.

Elephas orientalis, flore magno proboscide incurva.

Tournef. cor. 48. itin. 2. p. 299. t. 299.

Habitat in Oriente.

Calyx quadrifidus: lacinia prima (sub rostro corollæ) ovata; lateralibus utrinque reflexis; quarta, recta obtusa.

2. RHINANTHUS corollis labio superiore subulato *re-Elephas*.
Ro. Roy. lugdb. 298.

Elephas italica, flore magno proboscide surrecta. Tournef. cor. 48.

Elephas campoclarensium. Col. ecphr. 1. p. 186. t. 188.

Euphrasia lutea, floribus elephantis caput cum proboscide referentibus. Moris. hist. 3. p. 432. f. 11. t. 24. f. 14.

Scordio affinis Elephas ob florem. Baub. pin. 248.

- β. *Elephas orientalis, flore parvo, proboscide surrecta.*

*Tournef. cor. 48. itin. 3. p. 54. **

Habitat in umbrosis Italiæ. ☉

Habitus præcedentis, at flos diversus calyce trifido: laciniis duabus reflexis: tertia majore erecta, plicata, acuta.

3. RHINANTHUS corollarum labio superiore com-Crista galli, presso brevior. *Fl. lapp. 248. Fl. suec. 503. Hort. cliff. 325. Roy. lugdb. 298.*

Pedicularis pratensis lutea f. Crista galli. Baub. pin. 163.

- β. *Crista galli angustifolia montana. Baub. pin. 163. prodr. 86.*

- γ. *Crista galli mas. Baub. hist. 3. p. 436.*

Habitat in Europæ pratis. ☉

4. RHINANTHUS foliis sub lanceolatis pilosis integer-
rimis. Fl. zeyl. 238,

Hylsopus zeylanicus tenellus pratensis. Burm. zeyl. 122.

Habitat in Zeylona.

5. RHINANTHUS corollis fauce patente, foliis sinu-
to dentatis. Gron. virg. 168. †

Habitat in Virginia.

EU.

648

EUPHRASIA.

- latifolia.* 1. EUPHRASIA foliis dentato-palmatis, floribus subcapitatis.
 Euphrasia foliis ovatis, floribus spicatis. *Sauv. monsp.* 139.
 Euphrasia purpurea minor. *Bauh. pin.* 111. *Magn. monsp.* 95. t. 94.
 Euphrasia pratensis latifolia italica. *Bauh. pin.* 234. *Morif. hist.* 3. p. 430. f. 11. t. 24. f. 8.
 Euphrasia tertia latifolia pratensis. *Col. ecphr.* 200. t. 202. f. 2.
Habitat in Apulia, Italia, Monspelii. ☉
- officinalis.* 2. EUPHRASIA foliis ovatis argute dentatis. *Fl. lapp.* 247. *Fl. suec.* 516. *Mat. med.* 315. *Roy. lugdb.* 299. *Hall. belv.* 628.
 Euphrasia caule ramoso, foliis ovatis acute dentatis. *Hort. cliff.* 325.
 Euphrasia officinarum. *Bauh. pin.* 233.
 Euphrasia minor. *Dill. app.* 53.
 Euphrasia. *Cam. epit.* 767.
Habitat in Europæ pascuis aridis. ☉
- aricuspidata* 3. EUPHRASIA foliis linearibus tricuspидatis.
 Euphrasia angustis & tricuspидatis foliis, floribus ex albo purpureis. *Pluk. alm.* 142. t. 177. f. 1. *Zan. hist.* 110. t. 76.
Habitat in Italia. ☉
- Odontites.* 4. EUPHRASIA foliis linearibus: omnibus ferratis.
Hort. cliff. 346. *Fl. suec.* 517. *Roy. lugdb.* 299.
 Euphrasia pratensis rubra. *Bauh. pin.* 234.
 Euphrasia altera. *Dod. pempt.* 55.
 β. Euphrasia sylvestris major purpurea latifolia. *Col. ecphr.* 1. p. 201. t. 202. f. 1.
Habitat in Europæ arvis pascuisque sterilibus. ☉
- lutea* 5. EUPHRASIA foliis linearibus ferratis: superioribus integerrimis.
 Euphrasia pratensis lutea. *Bauh. pin.* 234. *Morif. hist.* 3. p. 432. f. 11. t. 24. f. 16.
 Euphrasia lutea montana angustifolia major altera. *Col. ecphr.* 1. p. 204. t. 203.
Habitat in Europa australi. ☉
- limifolia.* 6. EUPHRASIA foliis linearibus: omnibus integerrimis.
 Euphrasia foliis linearibus. *Sauv. monsp.* 138.

Eu-

Euphrasia foliis lini angustioribus. Bauh. pin. 234.

Euphrasia limifolia. Col. ecphr. 2. p. 68. t. 69.

Habitat in Italia, Helvetia, Gallia. ☉

MELAMPYRUM. 649

1. MELAMPYRUM spicis quadrangularibus compa-*cristatum.*
etis obtusis imbricatis. *Fl. suec. 510.*

Melampyrum luteum angustifolium. Bauh. pin. 234.

Melampyrum luteum, linariæ folio. Bauh. prodr. 112.

Melampyrum angustifolium cristatum. Pluk. alm.

249. t. 99. f. 2.

Habitat in Europæ borealis pratis asperis. ☉

2. MELAMPYRUM spicis conicis laxis laceris. *Fl. suec. arvensis.*
511.

Melampyrum foliis laciniatis laxè spicatis. Roy. lugdb.

298.

Melampyrum purpurascens coma. Bauh. pin. 234.

Triticum vaccinum. Dod. pempt. 541.

Melampyrum lanuginosum bæticum. Bauh. pin. 234.

Habitat in Europæ agris. ☉

3. MELAMPYRUM bracteis cordato-lanceolatis: sum-*nemorosum,*
mis coloratis sterilibus, calycibus lanatis. *Fl. suec. 512.*

Melampyrum coma cærulea. Bauh. pin. 234.

Parietaria sylvestris 1. Clus. hist. 2. p. 44.

Habitat in Europæ borealis nemoribus. ☉

4. MELAMPYRUM florum conjugationibus remotis, *pratense.*
corollis clausis. *Fl. suec. 513.*

Melampyrum foliis lanceolatis, florum paribus remotis.

Fl. lapp. 240.

Melampyrum luteum latifolium. Bauh. pin. 243.

Habitat in Europæ borealis pratis siccis. ☉

5. MELAMPYRUM corollis hiantibus. *Fl. suec. 514. sylvaticum.*

Melampyrum floribus spatis sessilibus binatis, foliis o-
mnibus indivisis. Hall. belv. 627.

Melampyrum latifolium, floribus parvis luteis. Cels.

apf. 32. Fl. lapp. 240.

Habitat in Europæ borealis sylvis ☉

Differt a precedente corollis dimidio brevioribus, limbo
hiantibus.

LATHRÆA. 650

1. LATHRÆA caule ramoso subterrestri, floribus ere-*Clandesti-*
ctis. *La-*

Lathræa foliis cordatis, floribus fasciculatis. *Guett. stamp.* 454. *Dalib. parisi.* 191.

Clandestina flore subcæruleo. *Tournef. inst.* 652.

Dentaria aphyllis, flore purpureo. *Raj. hist.* 1230.

Orobanche montana. *Rudb. elys.* 2. p. 233. f. 14.

Orobanche flore majore ex cæruleo-purpurascens
Bamb. pin. 87. *prodr.* 31. *Burf. III.* 155. *Rudb. elys.* 2. p. 229. f. 2. *Raj. hist.* 1227. *Morif. hist.* 3. p. 502. f. 12. t. 16. f. 15.

Orobanche f. *Dentaria* aphyllis purpurea, cespite denso
Morif. hist. 3. p. 503. f. 12. t. 16. f. 15.

Planta clandestina, f. *Madrona*. *Dalech. hist.* 960.

β. *Clandestina*, *Madronna*, *Occulta*. *Borel. obs.* 38.

Habitat in umbrifis Galliæ, Pyrenæorum, Italiæ. &
Caulis infra terram corymbosus, ut fructificationes sola
supra terram emineant solitariae in singulo ramo.

β. *elypæa*. 2. *LATHRÆA* corollis campanulato-patentibus.

Phælypæa lusitanica, flore luteo. *Tournef. cor.* 47.

Orobanche elegantissima verna, flore luteo. *Griff. luss.*
Morif. hist. 3. p. 502. *

β. *Orobanche* orientalis, flore coccineo. *Tournef. cor.* 47.
Habitat in Lusitanie umbrifis. &

Anblatum. 3. *LATHRÆA* corollarum labiis indivisiis. †

Anblatum orientale, flore purpurascens. *Tournef. cor.* 48.

Habitat in Oriente. &

Squamata. 4. *LATHRÆA* caule simplicissimo, corollis pendulis
labio inferiore trifido. *Fl. suec.* 518. *Dalib. parisi.* 191

Orobanche, radice dentata, major. *Bamb. pin.* 88.

Orobanche radice squamata, foliis rotundis, flore pendente & suave rubente, fundboensis. *Rudb. elys.* 2. p. 234. f. 17.

Orobanche radice dentata altius radicata, foliis & floribus albo purpureis. *Mentz. pug.* t. 3. f. 3. *Morif. hist.* 3. p. 503. f. 12. t. 16. f. 14.

Habitat in Europæ frigidioris umbrosissimis. &

651

SCHWALBEA.

americana. 1. *SCHWALBEA*. *Gron. virg.* 71. *

Euphrasia major mariana, floribus spicatis amplis: tubi longioribus. summis oris profunde incisis. *Pluk. man.* 73. t. 348. f. 2. ?

Habitat in America septentrionali.

TOZ

TOZZIA. 652

1. TOZZIA.

Tozzia alpina lutea, alfines folio, radice squamata. *alpina.*

Mich. gen. 20. t. 16.

Euphrasia lutea alfinifolia, radice squamata. *Bauh. pin.*

234. *prodr.* 111.

Dentaria buguloides, radice globosa: squamulis myon-
toideis, alpina. *Mentz. pug. t. 9. f. 3. 4.*

Orobanche buglossoides, radice rotunda, squamulis
myontoideis, alpina. *Morif. hist. 3. f. 12. t. 16. f. an-
tepenult.*

Anonyma f. gregorii, radice dentariæ. *Col. ecphr. 2. p.*
49. t. 50.

*Habitat in Alpibus Helveticis, Austriacis, Italicis, Py-
renæis, locis asperis, humidis. &*

PEDICULARIS. 653

* *Caule ramoso.*

1. PEDICULARIS caule ramoso, calycibus calloso-*palustris.*
punctatis, corollis labio obliquis. *It. w-gorb. 40. Mat.*
med. 314.

Pedicularis caule ramoso erecto, calycibus bifidis cre-
natis. *Hall. belv. 621. Fl. succ. 505.*

Pedicularis pratensis rubra elatior. *Raj. hist. 700.*

Pedicularis danica maxima. *Lob. illust. 147.*

Habitat in Europæ septentrionalis paludibus. ☉

2. PEDICULARIS caule ramoso, calycibus oblongis *sylvarum.*
angulatis lævibus, corollis labio cordato. *It. w-gorb.*
40.

Pedicularis caule ramoso, floribus solitariis, calycibus
quinquefidis. *Fl. succ. 504.*

Pedicularis caule ramoso, floribus solitariis remotis. *Fl.*
lapp. 241. Hort. cliff. 326.

Pedicularis pratensis purpurea. *Bauh. pin. 163.*

Fistularia. *Dod. pempt. 556.*

Habitat in Europæ sylvis paludosis. ☉

3. PEDICULARIS caule subramoso, corollis galea ro-*rostrata.*
strato-acuminatis, calycibus subhirtutis.

Pedicularis alpina, foliis alternis: pinnulis incisfis, flori-
bus sparsis rostratis purpureis. *Hall. belv. 621. t. 16.*
f. 1.

Pedicularis alpina, filicis folio, minor. *Bauh. pin. 163.*

Ale-

Alectorolophus alpina 3 minor. *Clas. bist.* 2. p. 210.
Habitat in Alpibus Helvetiæ, Austriæ.

* *Caule simplicissimo.*

- Sceptrum Carolin.* 4. **PEDICULARIS** caule simplici, floribus terno-verticillatis, corollis clausis, calycibus crenatis, capsulis regularibus. *Fl. suec.* 506.

Pedicularis capsulis subrotundis regularibus. *Roy. lugdb.* 209.

Pedicularis alpina, folio ceterach. *Helv. flor.* 39. t. 39.
Sceptrum carolinum. *Rudb. it.* 1. p. 1. *Fl. lapp.* 243.
 t. 4. f. 4. *Hort. cliff.* 326.

Habitat in Sueciæ, Borussia, Rutheni *spongiosis, sylvaticis, rignis.* &

Scapus nudus. Spica interrupta ex *Verticillis trifloris, foliolo exceptis.* Corolla non bians. Folia pinnatifida, crenata.

- verticillata.* 5. **PEDICULARIS** caule simplici, foliis quaternis. *Pedicularis alpina*, floribus purpureis spicatis, foliis pinnatis verticillatis. *Hall. belv.* 623. t. 17. f. 1.
Alectorolophus alpina 2, floribus purpureis. *Clas. bist.* 2. p. 210.

Filipendula montana altera. *Bauh. pin.* 163.

Habitat in Sibiria, Helvetia, Austria. &

- resupinata.* 6. **PEDICULARIS** caule simplici, foliis lanceolatis serratis crenulatis, floribus resupinatis.

Habitat in Sibiria *D. Gmelin, D. Demidoff.*

Caulis pedalis, levis, simplicissimus. Folia lanceolata, indivisa, profunde serrata serraturis crenulatis, alterna. Flores ex alis superioribus solitarii, sessiles. Calyx levis, obtusus, bifidus, integer. Corolla saturatissime purpurea, resupinata: Galea adunca, compressa; Vexillum amplum, rotundatum.

- reclita.* 7. **PEDICULARIS** caule simplici, staminibus floris galea longioribus.

Pedicularis alpina, foliis alternis pinnatis, floribus ex atrorubentibus in spicam densam congestis. *Hall. belv.* 623. t. 16. f. 2.

Habitat in Helvetiæ *summis Alpibus.*

- villosa.* 8. **PEDICULARIS** caule simplici, corollarum galeæ margine villotis.

Pedicularis caule simplici, foliis semipinnatis: pinnulis obtusis acute serratis. *Gmelin.*

Habitat in Sibiria.

Cau

Caulis *simplicissimus, spithameus*. Folia *alterna, pinnatifida, serrulata*. Flores *in spicam s. capitulum congesti*. Corollæ *obtusæ, flavæ; galea parum incurva, margine villosa*.

PEDICULARIS caule simplici, foliis pinnatis *retrofloratis, imbricatis*.

Pedicularis caule simplici, foliis semipinnatis obtusis: laciniis imbricatis crenatis.. *Fl. lapp. 244. t. 11. f. 2. Fl. succ. 509.*

Pedicularis caule erecto non ramoso, pinnis foliorum retroversis imbricatis. *Hall. helv. 622.*

Pedicularis alpina, folio ceterach. *Bauh. pin. 163.*

Habitat in Alpibus Lapponiæ, Helvetiæ. &

PEDICULARIS caule simplici, foliis dentato-pinnatis linearibus, calycibus hirsutis.

Pedicularis caule simplici, calycibus villosis, foliis linearibus dentatis crenatis. *Fl. lapp. 245. t. 4. f. 3. Fl. succ. 508.*

Habitat in Lapponiæ Alpibus.

PEDICULARIS caule simplici, foliis pinnatis *seriatim, internatis*, calycibus rotundatis glabris, corollis *galea uncinatis acutis*.

Habitat in Siberia.

Folia *Radicalia pinnata, pedalis; pinnis pinnatifidis, serratis*. Caulium *pinnatifida fere ad basin, serrata*. Caulis *lævis, plusquam pedalis, foliis alternis*. Spica *nuda, calycibus semiglobosis, quinquedentatis, glabris*. Corolla *incarnata; Galea falcata & uncinata, acuta, valde adunca.*

PEDICULARIS caule simplici, foliis pinnatifidis *lapponica*, serratis, calycibus bifidis obtusis.

Pedicularis caule simplici, foliis lanceolatis semipinnatis serratis acutis. *Fl. lapp. 242. t. 4. f. 1. Fl. succ. 507. Hort. cliff. 326. Roy. lugdb. 399.*

Habitat in Alpibus Lapponicis frequens. &

PEDICULARIS caule simplici, spica foliosa, corollæ *tomato*, galea acutis emarginatis, calycibus quinquedentatis.

Pedicularis foliis alternis: pinnis pinnatis, foliis ex spica florifera longe eminentibus. *Hall. helv. 623. t. 17 f. 3.*

Pedicularis alpina, filicis folio, major. *Bauh. pin. 163.*

Habitat in Alpibus Helveticis. &

Calyx *densatus, tubulosus, integerrimus, parum pilosus.*

610 DIDYNAMIA ANGIOSPERMIA.

Corolla labro superiore acutoped emarginato dentib.
duobus, acutis, nutantibus.

- tuberosa.* 14. **PEDICULARIS** caule simplici, calycibus crenatis, corollis galea rostrato aduncis.
Pedicularis alpina, foliis alternis pinnatis: pinnulis pinnatis, floribus ochroleucis rostratis in spicam congestis. *Hall. belv.* 620. *t.* 15. *Sauv. monsp.* 247.
Filipendula montana, flore pediculariæ. *Bauh. pin.* 16
Alectorolophus montana, flore luteo. *Barr. sc.* 46
Bocc. maf. 315. *t.* 8.
Habitat in Alpibus Helveticis, Italicis. 2

GERARDIA.

- tuberosa.* 1. **GERARDIA** foliis subovatis tomentosis repandis, longitudine caulis.
Gerardia humilis, bugulæ foliis, asphodeli radice. *Pin. gen.* 31.
Habitat in America calidior.
- purpurea.* 2. **GERARDIA** foliis linearibus,
Digitalis foliis linearibus, floribus remotis. *Gron. vi.* 169,
Digitalis rubra minor, labiis florum patulis, foliis parvis angustis. *Banist. virg.* 1926. *Gron. virg.* 72. *
Digitalis virginiana rubra, foliis & facie antirrhini vulgaris. *Pluk. mant.* 65. *t.* 388. *f.* 1.
Habitat in Virginia, Canada. ☉
Caulis pedales, filiformes, simplicissimi aut brachiales. Folia linearia, integerrima, opposita, sæpe alterna. Flores oppositi: Pedunculis unifloris, filiformibus, aut sessilibus. Calyces læves, parvi, campanulati quinque-dentati. Corollæ saturatissime purpureæ, rotatae aut campanulatae aut tubulosæ. Sic variat corollæ figura.
- læva.* 3. **GERARDIA** foliis lanceolatis pinnato-dentatis, caule simplicissimo.
Anonymos floribus flavis speciosis digitali æqualibus. *Gron. virg.* 74.
Digitalis lutea elatior, jaceæ nigrae foliis. *Banist. vi.* 1926. *Pluk. mant.* 64. *t.* 368.
Habitat in Virginia, Canada.
Caulis pedalis f. altior. Folia opposita, lanceolata, petiolata (Lycopi aut jaceæ), basi incisapinnatifidius patulis. Spica terminalis, laxa, ex
opp

DIDYNAMIA ANGIOSPERMIA. 681

oppositis, magnis, flavis. Antheræ in duas spinas deorsum tendentes. Stamen quintum deest.

GERARDIA foliis oblongis duplicato-ferratis, caule *pedicularis*. paniculato, calycibus crenatis.

Pedicularis foliis lanceolatis pinnatifidis ferratis, floribus peduncularis. *Gran. virg. 68.*

Digitalis, verhesinæ foliis e regione binis, americana, capsularum apicibus longissimis filamentis donata. *Pluk. mant. 64.*

Digitalis mariana, filipendulæ folio. *Raj. suppl. 397.*

Habitat in Virginia, Canada.

Corollæ extus pubescentes, oblongæ. Calycis dentes crenati, exsiccatione nigra evadit, ut Pedicularis; sed folia opposita & corollæ patula.

GERARDIA foliis ovatis ferratis, bracteis linearibus glutinosis, hispida.

Habitat in China

Caulis teretissimus, erectus; ramis brevioribus. Folia opposita, petiolata, ovata, acuta, pollice latiora, profunde ferrata, subvillosa. Racemi terminales, solitarii ex floribus oppositis brevissime peduncularis. Calyces acuti, 5partiti, lacinia superiore majore. Bractea filiformis una alterave calyci adjacet ejusdem longitudinis, pilis glutinosis, uti calyx hispida.

CHELONE. 655

CHELONE foliis lanceolatis ferratis: summis opposi- *glabra*. tis. *Hort. cliff. 493. Gran. virg. 70.*

Chelone acadientis, flore albo. *Tournef. art. 1706.*

Digitalis mariana, persicæ folio. *Raj. suppl. 397*

Digitalis mariana, ferratis densioribus rigidis & angustis foliis, semine tegopyri triquetro. *Pluk. mant. 64. 3. 348. f. 3.*

Habitat in Virginia, Canada. 2

CHELONE caule foliisque hirsutis.

hirsuta.

Anonymos flore pallide cæruleo digitalis instar in summis caulis disposito, foliis villosis acuminiatis. *Gran. virg. 71.*

Digitalis virginiana, panacis colonii foliis, flore amplo pallescente. *Pluk. mant. 64.*

Digitalis flore pallido transparentii, foliis & caule molli hirsutie imbutis. *Banist. virg. 1928.*

Habitat in Virginia.

Q q 2

Huic

612 DIDYNAMIA ANGIOSPERMIA.

*Haec flamen quintum longum cum Anthera villosissim
D. Jussieu.*

Pentstemon. 3. CHELONE foliis amplexicaulibus, panicula dichotoma.

Pentstemon. Mitch. gen. 14.

Habitat in Virginia.

Caulis herbaceus, brachiatus. Folia ovato-lanceolata, opposita, amplexicaulia, denticulata, acuminata, utrinque viridia. Caules et Ramuli terminantur pedunculo dichotomo. Calyces quinquepartiti. In prima dichotomia flores communiter bini, propriis pedicellis.

656

GESNERIA.

humilis.

1. GESNERIA foliis lanceolatis serratis sessilibus, pedunculis ramosis multifloris.

Gesneria humilis, flore flavescente. Plum. gen. 27.

Habitat in America australi.

tomentosa.

2. GESNERIA foliis lanceolatis crenatis hirsutis, pedunculis lateralibus longissimis corymbiferis. *Hort. cliff. 318.*

Gesneria amplo digitalis folio tomentoso. Plum. gen. 27.

Digitalis folio oblongo serrato, ad foliorum alas floridi Sloan. jam. 60. hist. 1. p. 162. t. 104. f. 2. Rq suppl. 396.

Habitat ad Jamaicae ripas. b

ANTIRRHINUM.

* *Foliis angulatis.*

Asarina.

1. ANTIRRHINUM foliis oppositis cordatis crenatis

Hort. cliff. 313. Roy. lugdb. 295.

Hedera saxatilis, magno flore. Baub. pin. 306.

Asarina s. Hederula saxatilis. Lob. ic. 601.

Habitat in rupibus Vegenio conterminis Genevae.

Cymbalaria.

2. ANTIRRHINUM foliis cordatis quinquelobis alternis. *Hort. cliff. 32. Hort. nps. 175. Roy. lugdb. 295.*

Guett. stamp. 2. p. 210. Gort. gehr. 358.

Cymbalaria. Baub. pin. 306. Cam. epit. 860. Dalech. hist. 1322.

Habitat in rupibus & muris antiquis Basileae, Parisii Harlemi. ☉

Elatine.

3. ANTIRRHINUM foliis hastatis alternis, caule flaccido. *Hort. cliff. 323. Hort. nps. 175. Gron. virg. 68. Roy. lugdb. 295.*

Elatine

Elatine folio acuminato in basi auriculato, flore luteo.

Bauh. pin. 253.

Elatine. Cam. epit. 754.

Elatine folio acuminato, flore cæruleo. *Bauh. pin.* 253.

Habitat in Germaniæ, Angliæ, Galliæ, Italiæ *arvis.* ☉

ANTIRRHINUM foliis ovatis alternis, caule flacci-*spurius*,

do. Hort. upf. 175 *Hort. cliff.* 323. *Roy. lugdb.* 295.

Elatine folio subrotundo. *Bauh. pin.* 253.

Veronica fernina. *Dad. pempt.* 42.

Habitat in Germaniæ, Angliæ, Galliæ, Italiæ *arvis.* ☉

ANTIRRHINUM foliis hastatis, caule erecto ramo-*egyptianum.*
sissimo.

Linaria memphitica pumila, hastato folio tricuspidi, flo-
re luteo. *Lipp.*

Habitat in Ægypto.

* *Foliis oppositis.*

ANTIRRHINUM foliis ternis ovatis. *Hort. cliff. triphyllum.*

324. *Hort. upf.* 174. *Roy. lugdb.* 295. *Sauv. monsp.*

165.

Linaria triphylla minor lutea. *Bauh. pin.* 212.

Linaria hispanica. *Clus. hist.* 1. p. 320.

Linaria triphylla cærulea. *Bauh. pin.* 312.

Habitat in umbrosis montibus Valentini & Hyblæis *circa.*

Syracusas. ☉

ANTIRRHINUM foliis quaternis lanceolatis, caule triornitho-

erecto ramoso, floribus pedunculatis. *Hort. cliff.* 324. *phorum.*

Roy. lugdb. 296.

Linaria americana maxima, purpureo flore. *Herm. lugdb.*

376. *Raj. bist.* 1884.

Linaria, flore purpureo, americana. *Riv. mon.* 81.

Habitat in Lusitania, America.

ANTIRRHINUM foliis lanceolatis sparsis: inferiori-*triste.*

ribus oppositis, nectariis subulatis, floribus subsessilibus.

Hort. cliff. 498. *Roy. lugdb.* 296.

Linaria trillis hispanica. *Dist. elsb.* 201. t. 164. f. 199.

Habitat Gibraltariæ.

ANTIRRHINUM foliis quaternis linearilanceolatis, *purpureum*

Antirrhinum foliis quaternis linearilanceolatis, caule flo-

rifero erecto spicato. *Hort. upf.* 174.

Antirrhinum foliis linearibus sparsis, nectariis subulatis

recurvis, floribus laxè spicatis. *Hort. cliff.* 498. *Roy.*

lugdb. 296.

Li-

624 DIDYNAMIA ANGIOSPERMIA.

Linaria purpurea major odorata. Baub. pin. 213.

Linaria altera purpurea. Dod. pempt. 183.

Habitat ad radices Vesuvii.

bipunctatum 10. ANTIRRHINUM foliis lanceolatis glabris: inferioribus quaternis, caule erecto paniculato, floribus spicato-capitatis.

Antirrhinum foliis lanceolato-linearibus, floribus in summitate caulis confertis parvis. *Gvett. stamp. 2. p. 205*

Antirrhinum erectum, foliis linearibus: imis verticillatis, pauciflorum. *Sauv. monsp. 165.*

Linaria lutea parva annua. Baub. hist. 3. p. 457.

Habitat in Hispania. Læsing.

Caulis spithameus, Ramis strictis ad angulum rectum excurrentibus. Folia lanceolata, sessilia, plana, patentia vix carnosae, fere laevia: inferiora quaternaria. Flores subspicati, pauci, sessiles. Corollæ flavae, punctis duobus fulvis in palato glabro versus labium superius. Labium superius bipartitum, replicatum, lanceolatum, erectum.

sepena. 11. ANTIRRHINUM foliis sublinearibus: inferne quaternis, calycibus capsulae aequalibus.

Antirrhinum foliis linearibus alternis saepius oppositis acuminatis, floribus laxè spicatis. *Roy. lugdb. 296.*

Antirrhinum foliis lanceolato-linearibus sparsis, calycinis laciniis capsulae aequalibus. *Gvett. stamp. 2. p. 204. Dab. parif. 186.*

Linaria angustifolia, uore cinereo striato. Dill. elth. 198. t. 163. f. 197.

Habitat in Anglia, Gallia. &

arvensis. 12. ANTIRRHINUM foliis sublinearibus: inferioribus verticillatis, calycibus piloso-viscidis, floribus spicatis, caule erecto, *Hort. upf. 174.*

Antirrhinum foliis linearibus alternis recurvis, rami caule brevioribus. *Roy. lugdb. 297.*

Linaria arvensis caerulea. Baub. pin. 213. prodr. 107. Dill. elth. 199. t. 163. f. 198.

β. *Linaria pumila, foliolis carnosis, flosculis minimis flavis. Baub. pin. 213.*

γ. *Linaria quadrifolia lutea. Baub. pin. 213.*

Linaria tetraphylla lutea. Col. ecphr. 1. p. 299. t. 300 f. 1.

Habitat in Angliæ, Galliæ, Italiæ arvis. ☉

Flores spicati sunt.

DIDYNAMIA ANGIOSPERMIA. 615

13. **ANTIRRHINUM** foliis caulinis lanceolato-linearibus sparsis: radicalibus rotundis ternis. *Guest. num stamp. 2. p. 204. Dalib. parif. 186.*
Antirrhinum foliis ad caulem linearibus, ad radicem ovatis sæpe ternis. *Sauv. monsp. 166.*
Linaria annua purpureo-violacea, calcaribus longis, foliis imis rotundioribus. *Vaill. parif. 118. Magn. monsp. 159.*
Linaria cærulea minor. *Lab. illustr. 103.*
Linaria cærulea calcaribus longis. *Bauh. hist. 3. p. 461.*
Habitat in Gallia.
14. **ANTIRRHINUM** foliis oppositis lanceolatis, floribus laxe racemosis. *Roy. lugdb. 297. origanifolium.*
Antirrhinum saxatile, foliis serpilli. *Bauh. pin. 212. prodr. 106. Barf. X: 7.*
Linaria folio crasso origani minore, floribus variegatis ex pallido & cæruleo rubescente. *Boerb. lugdb. 1. p. 232.?*
Habitat Massiliæ in muris vetustis, inque Pyrenæis major. ☉
An hoc varietas A. minoris?
15. **ANTIRRHINUM** foliis quinis linearibus. *Hort. multicaule. cliff. 324. Roy. lugdb. 296.*
Linaria sicula multicaulis, molliginis folio. *Bocc. sic. 38. t. 19. f. 1.*
Habitat in Sicilia. &
Caules juncei, laxi, debiles. Folia linearia, obtusa.
16. **ANTIRRHINUM** foliis subquaternis linearibus, caule diffuso, floribus racemosis. *Roy. lugdb. 297. alpinum.*
Sauv. monsp. 165.
Linaria quadrifolia supina. *Bauh. pin. 213.*
Linaria 3 stiriaca. *Clus. hist. 1. p. 322.*
Linaria cærulea repens. *Bauh. pin. 213.*
Linaria foliis carnosis cinereis. *Bauh. pin. 213. ex Hall.*
Habitat in Helvetia, Austria, Baldo.
- * *Foliis alternis.*
17. **ANTIRRHINUM** foliis linearibus acuminatis alternis, floribus racemosis, caule diffuso. *Roy. lugdb. 296. Dalib. parif. 186. Sauv. monsp. 48.*
Linaria pumila supina lutea. *Bauh. pin. 213*
Linaria hispanica. *Clus. hist. 1. p. 321.*
Habitat in Gallia, Hispaniæ arenosis.

dalmaticum. 18. ANTIRRHINUM foliis lanceolatis alternis, caule fruticoso.

Linaria latifolia dalmatica, magno flore. *Bauh. pin.* 212. *prodr.* 106.

Linaria maxima, folio lauri. *Bauh. hist.* 3. p. 458.

Habitat in Creta, &

hirtum 19. ANTIRRHINUM foliis lanceolatis hirtis alternis, floribus spicatis, foliolo calycino supremo maximo. *Læst.*

Habitat in Hispania. ☉

Caules aliquot simplices, spithamei, non ramosi. Folia lanceolata (Lavendulae), alterna. Flores dense spicati, colore & magnitudine A. Linariae. Calycis foliolum superius duplo majus quam reliqua. Capsula subrotunda.

genistifolium 20. ANTIRRHINUM foliis lanceolatis acuminatis, panicula virgata.

Linaria genistae folio glauco, flore luteo. *Herm. par. app.* 9. *Dili. elth.* 202.

Habitat in Sibiria. D. Gmelin. &

Caules subdecumbentes vel erectiusculi. Folia alterna, lanceolata, glabra, acuminata, unguina Genistae similis. Racemi longi, in paniculam dispositi: Floribus alternis, distantibus, pedicellis longitudine calycis. Calyx quinquepartitus, glaber, acutus, longitudine capsulae. Corollae luteae, magnitudine A. Linariae.

Linaria 21. ANTIRRHINUM foliis lanceolato-linearibus confertis, caule erecto, spicis terminalibus sessilibus, floribus imbricatis. *Fl. suec.* 501. *Mat. med.* 313. *Roy. lugdb.* 297.

Antirrhinum foliis linearibus sparsis. *Hart. cliff.* 325.

Antirrhinum foliis lanceolato-linearibus sparsis, calycis laciniis capsula dimidio brevioribus. *Guett. stamp.* 2. p. 203.

Linaria vulgaris lutea, flore majore. *Bauh. pin.* 212.

Osyris. *Fuchs. hist.* 543. *Cam. epit.* 930.

Habitat in Europae ruderalis. &

Palatum villosum fulvum.

monspessula-nou. 22. ANTIRRHINUM foliis linearibus confertis, caule nitido paniculato, pedunculis spicatis nudis. *Roy. lugdb.* 297. *Sauv. monsp.* 47.

Linaria, capillaceo folio, odora. *Bauh. pin.* 213. *prodr.* 106. n. 4.

Li.

DIDYNAMIA ANGIOSPERMIA. 617

Linaria odorata monspessulana. Bauh. hist. 3. p. 459.
Habitat in Gallia.

23. ANTIRRHINUM foliis linear-lanceolatis alternis, *chalepensis*
 floribus racemosis, calycibus corolla longioribus,
 caule erecto. *Roy. lugdb. 296. Hort. upf. 174.*
Linaria chalcensis minor erecta, flore albo lineis vio-
laceis. Moris. hist. 3. p. 502. f. 5. t. 35. f. 9.
Linaria annua angustifolia, flosculis albis longis cauda-
tis. Trinius. obs. 87. t. 87. f. 2. Raj. hist. 1884.
Habitat in Italia. ☉

24. ANTIRRHINUM foliis lanceolatis obtusis alter-*minus*,
 nis, caule ramosissimo diffuso. *Hort. cliff. 324. Fl.*
succ. 502. Roy. lugdb. 295.
Antirrhinum arvense minus. Bauh. pin. 212.
Antirrhinum tertium. Cam. epit. 922.
Habitat in glareosis Europæ ☉

* *Corollis biantibus aut ecaudatis, foliis alternis*
integris.

25. ANTIRRHINUM corollis ecaudatis, calycibus *majus*,
 rotundatis. *Hort. upf. 175. Mat. med. 312.*
Antirrhinum foliis lanceolatis petiolatis, calycibus bre-
vissimis, racemo terminali. Vir. cliff. 61. Hort. cliff.
324. n. 4. Roy. lugdb. 294.
Antirrhinum majus alterum, folio longiore. Bauh. pin.
211.
Antirrhinum luteo flore. Bauh. pin. 211.
Antirrhinum. Dod. pempt. 182. Lob. ic. 404.
Habitat in Europæ australis maceris, sepibus. ♂

26. ANTIRRHINUM corollis ecaudatis, calycibus *co-orientum*,
 rolla longioribus. *Hort. upf. 176.*
Antirrhinum foliis lanceolatis petiolatis, calycibus flo-
res superantibus. Hort. cliff. 324. Roy. lugdb. 294.
Antirrhinum arvense majus. Bauh. pin. 212.
Antirrhinum sylvestre, Phyteuma. Dod. pempt. 212.
Habitat in Europæ agris & arvis. ☉

27. ANTIRRHINUM foliis ovatis serratis, corollis *bellidifolium*
 patulis. *Hort. cliff. 498. Roy. lugdb. 295.*
Antirrhinum foliis imis spathulatis, summis digitatis.
Sauv. monsp. 67.
Linaria bellidis folio. Bauh. pin. 212. prodr. 106. t.
106.
Linaria odorata. Dod. pempt. 184.
 Q 9 5

Ha-

618 DIDYNAMIA ANGIOSPERMIA.

Habitat in agris inter Lugdunum & Viennam, interque Lugdunum et Gratianopolin. &

- maadense.* 28. ANTIRRHINUM foliis linearibus alternis, corollis hiantibus: labio inferiore explanato.

Antirrhinum caule simplicissimo longissimo, foliis caulinis linearibus; stolonum procumbentium lanceolatis minimis. *Gron. virg.* 67.

Habitat in Virginia, Canada. ☉

Radix fibrosa. Caulis erectus, filiformis, pedalis, vix ramosus. Folia alterna, remota, linearia, glabra. Racemus terminalis, rarior: floribus alternis, pedicellatis. Calyx quinquepartitus, acutus. Corollæ labium superius breve, reflexum. Lab. inferius magnum, patens, dependens.

658

CYMBARIA.

- daurica.* 1. CYMBARIA.

Cymbaria daurica pumila incana, linariæ folio, magno flore luteo guttato. *Amm. ruth.* 47. t. 1. f. 2.

Habitat in Dauriæ montanis, apricis, saxosis.

659

CRANIOLARIA.

- annua.* 1. CRANIOLARIA foliis cordatis angulatis.

Martynia annua villosa & viscosa, aceris follo, flore albo tubo longissimo. *Ehresh. pict.* 1. f. 2.

Habitat in America. ☉

- fruticosa.* 2. CRANIOLARIA foliis lanceolatis dentatis.

Gesnera arborescens, amplo flore fimbriato & maculoso. *Plum. gen.* 27.

Habitat in America calidiore.

Folia lanceolata, serrata. Pedunculi laterales multisfiori. Corolla oblonga, interne punctata margine ciliato.

660

MARTYNIA.

- perennis.* 1. MARTYNIA caule simplici, foliis serratis.

Martynia foliis serratis. *Hort. cliff.* 322. t. 18.*

Habitat in America. &

- annua.* 2. MARTYNIA caule ramoso, foliis angulatis.

Martynia foliis dentatis. *Hort. cliff.* 312.

Martynia annua villosa & viscosa, folio subrotundo, flore magno rubro. *Mort. cent.* 42. t. 42. *Ehresh. pict.* t. 1. f. 1.

Habitat in Americæ Vera Cruce. ☉

TO.

TORENIA. 661

1. TORENIA. *Gen. nov.* 1108. *asiatica.*
Hederæ terrestris species. Com. mal. 35.
Kaka-pu. Rheed. mal. 9. p. 103. t. 53.
Habitat in India. Toren.
Variat Foliis glabris & villosis: Floribus majoribus & minoribus.

BESLERIA. 662

1. BESLERIA pedunculis ramosis, foliis ovatis. *Melissifolia.*
Besleria melissæ tragi folio. Plum. gen. 29.
Habitat in America.
2. BESLERIA pedunculis simplicibus confertis, foliis lanceolatis.
Besleria, virgæ aureæ foliis, flore luteo, minor. Plum. gen. 29.
 3. *Besleria, virgæ aureæ foliis, flore luteo, major. Plum. gen.* 29.
Habitat in America.
3. BESLERIA pedunculis simplicibus solitariis, involucris pentaphyllis propriis.
Besleria scandens cristata, fructu nigro. Plum. gen. 29.
Habitat in America.
Radix repens, Folia ovata, serrata, petiolata. Pedunculi laterales, uniflori, terminati Involucro pentaphyllo: foliolis cordatis, dentatis, sessilibus. Corolla solitaria, externe hirsuta. Stamina & Pistilla mire impleta, forte aut Martinia aut distincti generis planta.

SCROPHULARIA. 663

1. SCROPHULARIA foliis cordatis serratis acutis basi marilandica, rotundatis, caule obtusangulo. *Hort. ups.* 177.
Scrophularia foliis cordatis oppositis, racemo terminali. Gron. virg. 71.
Scrophularia marilandica, longo profunde serrato utricæ folio. Raj. suppl. 396.
Habitat in Virginia. 2
Affinis maxime sequentis, ut vix distingui mereatur.
2. SCROPHULARIA foliis cordatis basi transversis, nodosa, caule obtusangulo.
Scrophularia foliis cordatis, racemo terminali. Hort. cliff. 322. *Fl. suec.* 520. *Mat. med.* 310. *Roy. lugdb.* 293. *Gron. virg.* 71. *Dalib. parif.* 191.

Scro-

Scrophularia nodosa fœtida. Bauh. pin. 235.

Scrophularia. Cam. epit. 866.

Habitat in Europæ succulentis. 2.

- aquatica.* 3. SCROPHULARIA foliis cordatis petiolatis decurrentibus, caule membranis angulato, racemis terminalibus. *Hort. ups. 177. Mat. med. 319.*

Scrophularia foliis cordatis. petiolorum alis in caulem decurrentibus. Guett. stamp. 2. p. 201.

Scrophularia aquatica major. Bauh. pin. 235. caule fimbriato. Læs. pruss. 248. t. 13.

Scrophularia femina. Cam. epit. 867.

Habitat in Angliæ, Helvetiæ, Galliæ humidis. 3.

- auriculata.* 4. SCROPHULARIA foliis cordato-oblongis basi appendiculatis, racemis terminalibus. *Læs.*

Scrophularia betonicæ folio. Tournef. inst. 166.

Scrophularia aquatica montana mollior. Barr. rar. 189. t. 274.

Betonica aquatica septentrionalium. Lab. ic. 533. fig.

Habitat in Hispania,

Caules vix ramosi, acutè tetragani, nudis. Folia opposita, petiolata, cordata, oblonga, serrata, subtus villosa; infra basin auriculata utrinque foliola ovato, parvo, sessili. Racemus terminalis.

- Scorodonia.* 5. SCROPHULARIA foliis cordatis duplicato-serratis, racemo composito. *Roy. lugdb. 296.*

Scrophularia melissæ folio. Tournef. inst. 166.

Scrophularia scorodonix folio. Pluk. alm. 333. t. 59. f. 5. Raj. hist. 766.

Habitat in Lusitania & Jersea insula Angliæ.

- orientalis.* 6. SCROPHULARIA foliis lanceolato-linearibus acutè serratis: inferne incis, racemo composito. *Roy. lugdb. 294.*

Scrophularia orientalis, foliis cannabinis. Tournef. cor. 9.

Habitat in Oriente.

- pernaltis.* 7. SCROPHULARIA foliis cordatis, pedunculis axillaribus solitariis dichotomis. *Hort. cliff. 322. Hort. ups. 177. Roy. lugdb. 293.*

Scrophularia flore luteo Bauh. pin. 236. prodr. 112. t. 112.

Lamium pannonicum & exoticum. Clus. hist. 2. p. 38.

Habitat in Italia, Helvetia. 3.

- fimbriatifolia* 8. SCROPHULARIA foliis difformibus, pedunculis axillaribus aggregatis. *Roy. lugdb. 292.* *Scro-*

- Scrophularia* foliis laciniatis. *Bauh. pin.* 236. *prodr.* 173.
Scrophularia sambucifolia, flore rubro luteo vario pulchro. *Pluk. alm.* 338. t. 313. f. 6.
Scrophularia sambuci foliis, capsulis maximis. *Morif. hist.* 2. p. 483. f. 5. t. 8. f. 6.
 9. *Scrophularia lusitanica*, maximo flore, foliis dissectis. *Raj. suppl.* 396.
Habitat in Hispania, Lusitania. ☉
9. SCROPHULARIA foliis pinnatis, racemo terminata-canina. li mudo, pedunculis bifidis. *Vir. cliff.* 61. *Hort. cliff.* 322. *Roy. lugdb.* 294. *Sauv. mousp.* 247.
Scrophularia Ruta canina dicta. *Bauh. pin.* 236.
Ruta canina. *Clus. hist.* 2. p. 109. *Lob. ic.* 2. p. 55.
Habitat in Helvetia, Narbōna, Italia. ☉
10. SCROPHULARIA foliis cordatis: superioribus al-peregrina. ternis, pedunculis axillaribus bifloris. *Hort. cliff.* 322. *Roy. lugdb.* 293.
Scrophularia urticæ folio. *Bauh. pin.* 236.
Scrophularia peregrina. *Cam. hort.* 157. t. 43.
Habitat in Italia. ☉
11. SCROPHULARIA foliis lanceolatis obtusis ferru-frutescens, to-dentatis, pedunculis bifidis. *Roy. lugdb.* 294.
Scrophularia lusitanica frutescens, verbenacæ foliis. *Tournef. inst.* 167. *Raj. suppl.* 395.
Habitat in Lusitania.
12. SCROPHULARIA foliis quaternis ovatis, floribus coccineis, verticillatis spicatis. *Roy. lugdb.* 294.
Scrophularia flore coccineo, foliis urticæ quaternis caulem ambientibus. *Houft. mss.*
Habitat in Vera Cruce.

CELSIA. 664

1. CELSIA foliis duplicato-pinnatis. *Hort. cliff.* 321 *orientalis.* *Hort. mss.* 179. t. 1. *Roy. lugdb.* 301.
Verbalcum orientale, sophiæ folio. *Tournef. cor.* 8. *Buxb. cent.* 5. p. 17.
Blattaria orientalis, agrimonix folio. *Buxb. cent.* 1. p. 14. t. 20.
Habitat in Cappadocia, Armenia. ☉

DIGITALIS. 665

1. DIGITALIS calycinis foliolis ovatis acutis, corollis purpureo-obtusis: labio superiore integro. *Hort. mss.* 178.

Di-

622 DIDYNAMIA ANGIOSPERMIA.

- Digitalis foliolis calycinis ovatis. Hort. cliff. 318. Dalib. parif. 192. acuminatis. Roy. lugdb. 292. Gort. gelr. 371.*
Digitalis purpurea, folio aspero. Baub. pin. 243.
Digitalis purpurea. Dod. pempt. 168.
β. *Digitalis alba, folio aspero. Baub. pin. 229.*
Habitat in Europa australiore. 3
- alba.* 2. **DIGITALIS** calycinis foliolis lanceolatis, corollis acutis; labio superiore bifido. *Hort. upf. 178.*
Digitalis foliis calycinis subulatis, floribus imbricatis. Hort. cliff. 318. Roy. lugdb. 293. Dalib. parif. 192.
Digitalis major lutea f. pallida, parvo flore. Baub. pin. 224.
Digitalis lutea, minore flore. Morif. hist. 2. p. 479. f. 5. t. 8. f. 5.
Habitat in Gallia, Italiae sabulosis. 2
- ferruginea.* 3. **DIGITALIS** calycinis foliolis ovatis obtusis, corollae labio inferiore longitudine floris.
Digitalis foliolis calycinis ovatis obtusis, corollis villosis; labio longissimo. Buttn. canon. 226.
Digitalis foliolis calycinis ovatis obtusis. Roy. lugdb. 292.
Digitalis angustifolia, flore ferrugineo. Baub. pin. 244. Riv. mon. 98.
Digitalis latifolia, flore ferrugineo. Morif. hist. 2. p. 478. f. 5. t. 8. f. 2. 3.
Digitalis lutea non ramosa, scorzonerae folio. Baxx. cent. 5. p. 25. t. 49.
Habitat in Italia, Constantinopoli. 2
- canariensis.* 4. **DIGITALIS** calycinis foliolis lanceolatis, corollis bilabiatis acutis, caule fruticoso.
Gesneria foliis lanceolatis serratis, pedunculo terminali laxo spicato. Hort. cliff. 318.
Digitalis acanthoides canariensis frutescens, flore aureo. Comm. hort. 2. p. 105. t. 53.
Digitali affinis canariensis, solidaginis acutis foliis leviter pilosis, flore aureo cucullato. Pluk. alm. 400. t. 325. f. 2.
Habitat in Canariis. 5

BIGNONIA.

- Carap. a.* 1. **BIGNONIA** foliis simplicibus cordatis, caule erecto, floribus diandris.
Bignonia foliis simplicibus cordatis. Hort. cliff. 317. Roy. lugdb. 289.
Bigno-

Bignonia urueu foliis, flore sordide albo intus maculis purpureis & luteis adperso, siliqua longissima & angustissima. *Catesb. car.* 1. p. 49. t. 49.

Kawata Filagi. *Kempf. jap.* 841. t. 842.

Habitat in Japonia, Carolina. h

Arbor erecta. Folia *terna*. Cirrhi *nulli*. Flores *dianthri*: *staminibus* duobus *antheriferis*; reliquis *tribus* *castratis*.

1. BIGNONIA foliis simplicibus lanceolatis, caule *vo-* *scuperu-*
luoli. *lucis*.

Bignonia foliis ovato-lanceolatis conjugatis integerri-
mis. *Roy. lugdb.* 291.

Gelseminum f. *Jasminum* luteum odoratum virginia-
num scandens sempervirens. *Catesb. car.* 1. p. 53.
t. 53. *Raj. hist.* 1769.

Syringa volubilis virginiana, myrti majoris folio, alato
semine, floribus odoratis luteis. *Pluk. alm.* 359. t.
112. f. 5.

Habitat in Virginia.

Folia *opposita*. Siliquæ *cordata*, *simplices*.

3. BIGNONIA foliis conjugatis: cirrho brevissimo arcu- *unguis* *cast.*
ato tripartito.

Bignonia americana. capiteolis aduncis donata, siliqua
longissima. *Tournef. inst.* 164.

Gelseminum iudicum hederaceum tetraphyllum, folio
subrotundo acuminato. *Sloan. jam.* 90. *hist.* 1. p. 208

Clematis quadrifolia, flore digitalis luteo, claviculis a-
duncis. *Plum. amer.* 80. t. 94.

Clematis myrsinites, amplioribus foliis, americana te-
traphyllos. *Pluk. alm.* 109. t. 163. f. 2.

Habitat in Barbados, Domingo.

4. BIGNONIA foliis conjugatis cirrhosis: foliolis ova- *agrimonia-*
to-lanceolatis, pedunculis bifloris, siliquis linearibus. *lis*.

Bignonia bifolia scandens, siliquis latis & longioribus,
semine lato. *Plum. spec.* 5.

Habitat in Cayenna.

5. BIGNONIA foliis conjugatis cirrhosis: foliolis cor- *paniculata*,
dato-ovatis, floribus racemoso-paniculatis.

Bignonia bifolia scandens, flore violaceo odore, fructu
ovato duro. *Plum. spec.* 5.

- β *Bignonia* bifolia scandens, flore e luteo purpurascente.
inodoro, fructu ovato duro. *Plum. spec.* 5.

Habitat in America calidure.

6. BIG-

624 DIDYNAMIA ANGIOSPERMIA.

- ernigera.* 6. BIGNONIA foliis conjugatis cirrhosis: foliolis cordatis; foliis imis ternatis. *Vir. cliff.* 60. *Hort. cliff.* 317. *Roy. lugdb.* 289.
Bignonia scandens, flore atro flavo minori subtus albidante. *Gron. virg.* 73.
PseudoApocynum folliculis maximis obtusis, seminibus amplissimis alis membranaceis. *Morif. hist.* 3. p. 672. f. 15. t. 3. f. 16.
Habitat in Virginia & australiori America.
Caulis volubilis, scaber, transversim dissectus crucem representat.
- capreolata.* 7. BIGNONIA foliis conjugatis cirrhosis: foliolis cordato-lanceolatis; foliis imis simplicibus. *Vir. cliff.* 59. *Hort. cliff.* 317. *Roy. lugdb.* 289.
Bignonia americana capreolis donata, siliqua brevior. *T. Breyn. ic.* 33. t. 25.
Clematis americana siliquosa tetraphyllos. *Dod. act.* 71.
Clematis tetraphylla americana. *Bocc. sic.* 31. t. 15. f. 31. *Zan. hist.* 74. t. 2. *Raj. hist.* 1329.
Habitat in America.
- Leucoxylon.* 8. BIGNONIA foliis digitatis integerrimis. *Hort. cliff.* 497.
Bignonia arbor pentaphylla, flore roseo, major & minor, siliquis planis. *Plum. spec.* 5. *Catesb. car.* 1. p. 37. t. 37.
Leucoxylon arbor siliquosa, quinque foliis, floribus neriis, alato semine. *Plum. alm.* 215. t. 200. f. 4.
Nerio astinis arbor siliquosa, folio palmato s. digitato, flore albo. *Steen. jum.* 154. *hist.* 2. p. 62. *Raj. dendr.* 114.
Habitat in Jamaica. Caribæis, agris humidioribus, & ad ripas. b
- radinta.* 9. BIGNONIA foliis digitatis pinnatifidis.
Bignonia flore luteo, foliis radiatis & elegantissime dissectis. *Ferr. peru.* 1. p. 731. t. 22. *Roy. lugdb.* 289.
Habitat in Peru.
- radicans.* 10. BIGNONIA foliis pinnatis, foliolis incis, geniculis radicatis. *Hort. cliff.* 217. *Hort. ups.* 178. *Gron. virg.* 73.
PseudoGelsemium siliquosum. *Riv. mon.* 101.
PseudoApocynum hederaceum americanum, tubuloso flore phoeniceo, fraxini folio. *Morif. hist.* 3. p. 612. f. 15. t. 3. f. 1.

Gelse-

DIDYNAMIA ANGIOSPERMIA. 625

Gelsemium hederaceum indicum. *Corn. can.* 102. t. 103.

β. *Bignonia fraxini* foliis, coccineo flore minore. *Catesb. car.* 1. p. 65. t. 65.

Habitat in America. 5

11. *BIGNONIA* foliis decompositis: foliolis incis, *ge-peruviana*,
nuculis cirrhosis. *Hort. cliff.* 317. *Roy. lugdb.* 290.

Bignonia arbor, flore luteo, fraxini folio. *Plum. spec.* 5.

Clematis peruana, digitalis flore, folio fraxini. *Pluk. alm.* 108. t. 162. f. 4.

Habitat in Peru.

12. *BIGNONIA* foliis bipinnatis: foliolis integris sub-indica
cordatis acutis. *Fl. zeyl.* 236.

Arbor filiquosa malabarica, cordato folio, fructu maximo oblongo plano. *Raj. hist.* 1741.

Palega pajanelli. *Rheed. mal.* 1. p. 177. t. 43.

β. *Arbor filiquosa malabarica*, folio majore mucronato,
fructu maximo oblongo plano. *Raj. hist.* 1741.

Pajanel *Rheed. mal.* 1. p. 179. t. 45.

Habitat in India.

13. *BIGNONIA* foliis bipinnatis: foliolis lanceolatis in-
tegris.

Arbor guajaci latiore folio, *bignoniæ* flore cæruleo,
fructu duro in duas partes dissiliente, seminibus alatis
imbricatis politis. *Catesb. car.* 1. p. 42. t. 42.

Habitat in Carolina. 5.

CITHAREXYLUM. 667

1. *CITHAREXYLUM*. *Amœn. acad.* 1. p. 406. *spinosum.*

Citharexylon americanum alterum, foliis ad marginem
dentatis. *Pluk. alm.* 108. t. 161. f. 5.

Habitat in Barbados. 5

HALLERIA. 668

1. *HALLERIA*. *Hort. cliff.* 323.

Halleria foliis ovatis longitudinaliter serratis. *Roy. lugdb.* *lucida.*
289.

Lonicera foliis lucidis acuminatis dentatis, fructu rotundo. *Burm. afr.* 244. t. 89. f. 2.

Solanum flore periclymeni. *Amm. herb.* 591.

β. *Halleria* foliis lanceolato-ovatis superne serratis. *Roy. lugdb.* 289.

R 1

Loni-

626 DIDYNAMIA ANGIOSPERMIA.

Lonicera folio acuto serrato, flore pendulo, fructu oblongo. Burm. afr. 243. t. 89. f. 1.
Habitat in Æthiopia.

669

CRESCENTIA.

Cujete.

1. CRESCENTIA.

Crescentia foliis lanceolatis utrinque attenuatis. Hort. cliff. 327.

Cujete foliis oblongis & angustis, magno fructu ovato Plum. gen. 23.

Arbor americana cucurbitifera, folio longo mucronato fructu oblongo. Comm. hort. 1. p. 137. t. 71.

β. *Cujete angustifolia, fructu minori globoso aut ovato Plum. gen. 23.*

γ. *Cujete minima, fructu duro. Plum. gen. 23.*

δ. *Cujete latifolia, fructu putamine fragili. Plum. gen. 23.*
Cucurbitifera arbor, subrotundis foliis confertis, fructuovali. Pluk. alm. 124. t. 171. f. 2.

Arbor cucurbitifera americana, folio subrotundo. Sloan. jam. 206. hist. 2. p. 172. Raj. hist. 1667.

Habitat in Virginiae, Jamaicae, Brasiliae apricis. b

670

statice.

GMELINA.

1. GMELINA. Fl. zeyl. 230. *

Michelia spinosa, floribus luteis. Amm. act. petrop. 1. p. 218. t. 18.

Lycium maderaspatanum indici alpino putati æmulum foliis minoribus & majoribus bijugis, & grandioribus aculeis horridum. Pluk. alm. 234. t. 303. f. 3. t. 97. f. 2.

Habitat in India. b

671

volubilis.

PETREA.

1. PETREA. Hort. cliff. 319. *

Habitat in America. b

Periit egregius Petreus 1742. ætatis 32. variolis corruptus, longiori vita dignissimus, utpote qui Floræ indici domicilia extruxit in Europa omnium splendidissim

672

retrofolia.

LANTANA.

1. LANTANA foliis ternis, spicis oblongis imbricatis

Camara trifolia, purpurascens flore. Plum. gen. 32.

Myrobatindum spicatum, viburni foliis ex adverso ternis. Vaill. act. 1722. p. 277.

H

Habitat in America calidiore.

An sequentis sola varietas?

2. LANTANA foliis oppositis, caule inermi, spicis oblongis. *Hort. upf.* 380.
Periclymenum rectum humilius, salviæ folio rugosiore, flore purpureo, fructu oblongo esculento purpureo. *Sloan. jam.* 164. *bist.* 2. p. 82. t. 195. *Raj. dendr.* 30.
 3. Viburnum americanum odoratum, folio parvo oviculato, floribus & baccis folio interceptis. *Pluk. alm.* 386. t. 114. f. 5. *Herm. par.* 384.
Habitat in America calidiore. ☉
3. LANTANA foliis oppositis, caule inermi ramoso, floribus capitato-umbellatis. *Camara.*
Lantana foliis oppositis petiolatis, floribus capitatis. *Hort. cliff.* 349. *Roy. lugdb.* 290.
Viburnum americanum non spinosum, melissæ foliis, floribus coccineis. *Pluk. alm.* 385. t. 114. f. 4.
Viburnum americanum, urticæ foliis, lamii odore, floribus miniatis. *Comm. hort.* 1. p. 151. t. 78.
 4. *Periclymenum rectum*, urticæ folio hirsuto majore, flore flavo. *Sloan. jam.* 163.
Habitat in America calidiore. ♀
4. LANTANA foliis oppositis, caule aculeato ramoso, aculeata, floribus capitato-umbellatis.
Lantana foliis oppositis petiolatis, caule aculeato. *Hort. cliff.* 498. *Hort. upf.* 380. *Roy. lugdb.* 290.
Myrobatindum viburnifolium spinosum, floribus coccineis. *Vaill. act.* 1722. p. 276.
Viburnum americanum odoratum, urticæ foliis latioribus, spinosum, floribus miniatis. *Pluk. alm.* 285. t. 223. f. 5.
Habitat in America calidiore. ♀
5. LANTANA foliis alternis, spicis oblongis. *bullata.*
Salvia barbadensis dicta, spica florum compactiori. *Pluk. alm.* 329. t. 221. f. 3.
Periclymenum rectum, salviæ folio rugoso majore oblongo bullato, flore albo, fructu longiore. *Sloan. jam.* 163. *bist.* 2. p. 81. *Raj. dendr.* 31.
 6. *Periclymenum rectum*, salviæ folio rugoso majore subrotundo bullato. *Sloan. jam.* 163. *bist.* 2. p. 81. t. 195. f. 1. *Raj. dendr.* 31.
Habitat in Jamaica. ♀

- corymbosa*. 6. LANTANA foliis alternis, floribus corymbosis †
 Periclymenum rectum, salviae foliis majoribus oblongis
 mucronatis subtus villosis alternatim sitis, flore & fru-
 ctu minoribus. *Sloan. jam.* 164. *hist.* 2. p. 83. t.
 194. f. 3. *Raj. dendr.* 30.
 Ulmi angustifoliae facie baccifera jamaicensis, foliis su-
 perne scabris, subtus villosis alternatim sitis, floribus
 parvis perpusillis, fructu botryoide monospermo. *Pluk*
alm. 393. t. 329. f. 5.
Habitat in Jamaica. †
Refertur ex fide Sloanei; De floris structura nul-
la mihi certitudo.
- fricana*. 7. LANTANA foliis alternis sessilibus, floribus solitariis
Hort. cliff. 320. *Roy. lugdb.* 290.
 Lantana floribus solitariis. *Gen. plant.* 632.
 Jasminum africanum, ilicis folio, flore solitario ex ali-
 foliorum proveniente. *Comm. rar.* 6. t. 6.
Habitat in Aethiopia. †

CORNUTIA

- pyramidata*. 1. CORNUTIA. *Hort. cliff.* 319.
 Cornutia flore pyramidato, foliis incanis. *Plum. gen.* 31.
 Agnanthus viburni folio. *Vaill. act.* 1722. p. 273.
Habitat in Caribæis, Campechia. †

LOESELIA.

- climata*. 1. LOESELIA. *Roy. lugdb.* 299.
 Royena foliis acute dentatis & veluti spinosis, flore luteo
Hoult. mss.
Habitat in Vera Cruce.

CAPRARIA.

- biflora*. 2. CAPRARIA.
 Capraria foliis alternis, corollis quinquefidis *Hort. cliff.*
 320. *Roy. lugdb.* 301.
 Capraria curassavica. *Herm. parad.* 110. t. 110.
 Gratiolæ affinis frutescens americana, foliis agerati f. vi-
 ronicæ erectæ majoris. *Comm. hort.* 1. p. 79. t. 4.
 Lysimachiae peruvianæ affinis americana procumben-
 ononidis vernæ frutescentis folio singulari glabro. *Plu*
alm. 237. t. 98. f. 4.
Habitat in Curassao. †
 Conferatur *Capraria peruviana*, agerati foliis absque
 dicalis. *Fewill. per.* 1. p. 764. t. 48. St

SELAGO. 676

1. SELAGO corymbo multiplici. corymbosa

Selago caule erecto, corymbo terminali. *Hort. cliff.* 321.

Roy. lugdb. 300.

Camphorata africana umbellata frutescens. *Comm. hort.*

2. p. 79. t. 40. *Raj. suppl.* 130.

Millefolio affinis maderaspatana, camphorata foliis radiatum nascentibus. *Morif. hist.* 3. p. 39. *Raj. suppl.* 219.

Habitat in Æthiopia. &

2. SELAGO spicis imbricatis. † spuria.

Melampyrum africanum spicatum, foliis angustissimis.

Burm. afric. 115. t. 42. f. 3.

Habitat in Æthiopia. †

Caulis suffrutex. Folia linearia, alterna. Spica ovato-oblonga, obtusa, arcte imbricata; bracteis subovatis, membranaceis. Fructificatio hodie ignota.

3. SELAGO racemo simplici. † dubia.

Thymelæa foliis angustissimis linearibus, flosculis spicatis. *Burm. afr.* 130. t. 47. f. 3.

Thymelæa æthiopica spicata glabra, foliis longioribus angustis. *Pluk. mant.* 180. t. 445. f. 6

Valerianoides parva, foliis in caule raris, spica in summo caule oblonga ex foliolis & flosculis confertis composita. *Raj. suppl.* 245.

Habitat in Æthiopia.

Caulis erectus. Folia conferta, linearia. Racemus simplex, terminalis, longissimus, erectus. Corollæ tubus filiformis, longissimus. Stamina modo 2 intra tubum corollæ; antheris incumbens, linearibus. Fructus ignotus; hinc genere dubio.

HEBENSTRETIA. 677

1. HEBENSTRETIA foliis dentatis. *Hort. cliff.* 497. *dentata.*

326. *Roy. lugdb.* 300.

Valerianella africana, foliis angustis, flore macula rubicante notato. *Comm. hort.* 2. p. 247. t. 109.

Valerianoides flore monopetalo, semine unico oblongo. *Raj. suppl.* 245.

Pedicularis foliis angustissimis dentatis, floribus spicatis.

Burm. afric. 114. t. 42. f. 2

Habitat in Æthiopia.

2. HEBENSTRETIA foliis integerrimis. *Hort. cliff.* 497. *integrifolia.*

Roy. lugdb. 300.

Habitat in Æthiopia.

ERI-

678

ERINUS

- europæus.* 1. ERINUS floribus racemosis
 Erinus. *Sauv. monsp.* 116.
 Ageratum serratum alpinum. *Bauh. pin.* 221.
 Ageratum purpureum. *Dalech. hist.* 1184.
 β. Ageratum minus saxatile, flore albo. *Barr. rar.* 23. s. 1192.
Habitat in Alpibus Helveticis, Pyrenæicis, Montpelii. &
- africanus.* 2. ERINUS floribus lateralibus sessilibus, foliis lanceo-
 latis subdentatis.
 Buchnera foliis obtusis serratis. *Hort. cliff.* 501. * *Roy.*
lugdb. 300.
 Lychnidea villosa, foliis ex alis floriferis, florum petala-
 lis cordatis. *Burm. afric.* 139. t. 50. f. 1.
 Euphrasia æthiopica, drabæ foliis, summis oris floscu-
 lorum altius divitis. *Pluk. mant.* 73. *Raj. suppl.* 401.
 β. Buchnera foliis acutis dentatis. *Hort. cliff.* 501. * *Roy.*
lugdb. 300.
Habitat in Æthiopia.
- peruvianus.* 3. ERINUS foliis lanceolato-ovatis serratis.
 Lychnidea veronicæ folio, flore coccineo. *Few. peruv.*
 3. p. 25. f. 3.
Habitat in Peru.
- laciniatus.* 4. ERINUS foliis laciniatis.
 Lychnidea verbena tenuifoliæ folio. *Few. peruv.* 3. p.
 35. t. 25.
Habitat in Peru.

679

BUCHNERA.

- americana.* 1. BUCHNERA foliis densis oppositis.
 Cortusæ f. Verbasci species, caule non ramoso, flori-
 bus violaceis. *Gran. virg.* 74. *
Habitat in Virginia, Canada.
- asiatica.* 2. BUCHNERA foliis integerrimis alternis.
Habitat in Zeylona, China.
 Caulis ramis alternis, habitu Euphrasiæ, obtuse tetra-
 gonus. Folia lineari-lanceolata, scabra, integerrima,
 alterna, versus basin opposita. Spicæ longæ floribus al-
 ternis remotis. Calyces quinquepartiti, scabri. Tubus
 corollæ filiformis, calyce duplo longior, cervice incur-
 vato, stamina 4 includente; Limbus quinquesidus,
 purpureus; laciniis 2 superioribus brevissimis reflexis,
 tribus reliquis obovatis subæqualibus.

BRO-

DIDYNAMIA ANGIOSPERMIA. 631

BROWALLIA. 680

1. BROWALLIA. *Hort. cliff.* 318. t. 17. *Hort. nps. americana.* 179. *.
Habitat in America australi. ☉

LINNÆA. 681

1. LINNÆA floribus geminatis. *Fl. lapp.* 250. t. 12. f. borealls.
4. *Hort. cliff.* 320. *Fl. succ.* 522. t. 1. *Mat. med.*
517. *Roy. lugdb.* 236. *Hall. helv.* 608. *Segu. veron.*
183. *Sauv. meth.* 137.
Serpillifolia. *Buxb. act.* 2. p. 346. t. 21.
Campanula serpillifolia. *Baub. pin.* 93. *prodr.* 35. *Act.*
nps. 1720. p. 96. t. 1.
Nummularia norvegica, flore purpureo. *Kyll. act. dan.*
2. p. 346. f. C.
Habitat in Sueciæ, Sibiria, Helvetiæ, Canadæ sylvis an-
tiquis, muscicis, acerosis, sterilibus, umbrosis. ☿

SIBTHORPIA. 682

1. SIBTHORPIA foliis reniformi-subpeltatis crenatis. *europæa.*
Gen. nov. 1099.
Chrysosplenium cornubiense. *Pet. herb.* t. 6. f. 11.
Alini spuria pusilla repens, foliis saxifragiæ aureæ. *Raj.*
angl. 3. p. 352. *Pluk. alm.* 23. t. 7. f. 6.
Habitat in Cornubiæ, Devonix, Lusitaniæ udis agge-
ribus.
2. SIBTHORPIA foliis orbiculatis integris crenatis, pe- *africana.*
dunculis solitariis. *Gen. nov.* 1099.
Chrysosplenii foliis planta aquatica, flore flavo pentape-
taio. *Shaw. afric.* 149. f. 149.
Habitat in Africa.
3. SIBTHORPIA foliis reniformibus crenatis, peduncu- *peregrina.*
lis geminis. *Gen. nov.* 1099.
Planta. *Pluk. phyt.* 257. f. 5.
Habitat - - -

LIMOSELLA. 683

1. LIMOSELLA. *Fl. lapp.* 249. *Fl. succ.* 521. *Hall. aquatica.*
helv. 629. *Dalib. parif.* 193.
Limosella annua. *Lind. alif.* 156. t. 5.
Plantaginella. *Hall. jen.* 23. t. 6. f. 2.
Plantaginella palustris. *Baub. pin.* 190. *Meris. hist.* 3.
p. 605. f. 15. t. 2. f. 1

R r 4

Sper-

Spergula perpusilla, lanceolatis foliis. *Loef. pruss.* 261. t. 81.

Alfina palustris repens, foliis lanceolatis. *Pluk. alm.* 20. t. 74. f. 4.

Alfina palustris exigua, foliis lanceolatis. *Menz.* pag. 2. t. 7. f. 6.

Habitat in Europa septentrionalis inundatis. ☉

684
indica.

ÆGINETIA.

1. ÆGINETIA.

Tiliacum-cumulu. *Rheed. mal.* 11. p. 97. t. 47.

Habitat in Malabar.

Folia nulla. Caules *simplicissimi*, teretes, *stricti* *ovis* *triflori*, *uniflori*. Flos *examinandus*.

685
virginica.

OBOLARIA.

1. OBOLARIA. *Hort. cliff.* 323. *Gran. virg.* 74.

Orobanche virginiana, radice corallibide summo caule foliis subrotundis. *Morif. hist.* 3. p. 504. f. 12. t. 16. f. 23. *Pluk. alm.* 273. t. 209. f. 6. *Raj. suppl.* 595.

Habitat in Virginia.

686
lævis.

OROBANCHE.

1. OROBANCHE caule simplicissimo lævi, staminibus exsertis.

Orobanche caule simplici cæruleo, bracteis brevibus. *Sauv. monsp.* 4.

Orobanche majore flore. *Bauh. pin.* 88. *Morif. hist.* 3. f. 12. t. 16. f. 2.

Orobanche magna purpurea monspeliulana. *Bauh. hist.* 2. p. 782.

Habitat Monspelii.

major.

2. OROBANCHE caule simplicissimo pubescente, staminibus inclusis.

Orobanche caule simplicissimo. *Hort. cliff.* 321. *Fl. Suec.* 519. *Roy. lugab.* 299. *Dalib. paris.* 190. *Gert. gelr.* 368.

Orobanche caule simplici, bracteis longioribus. *Sauv. monsp.* 5.

Orobanche major garyophyllum olens. *Bauh. pin.* 87.

Orobanche 1. *Clus. hist.* 1. p. 270.

Rapum genistæ. *Lob. ic.* 2. p. 89.

Habitat in Europæ agris, pratis ficcis.

DIDYNAMIA ANGIOSPERMIA. 623

3. OROBANCHE caule ramoso, corollis quinquefidis. *ramosa*.
 Orobanche caule ramoso. *Hort. cliff.* 321. *Roy. lugdb.*
299. Hall. belv. 610. *Dalib. parisi.* 191. *Gort. gelr.*
369.
 Orobanche ramosa. *Bauh. pin.* 88.
 Orobanche. *Cam. epit.* 311.
Habitat in Europæ siccis.
4. OROBANCHE caule ramoso, corollis quadridentatis. *virginiana*.
 Orobanche caule ramoso, floribus distantibus. *Gron. virg.*
168.
 Orobanche minor virginiana lignosior, per totum cau-
 lem floribus minoribus onusta. *Morisi, hist. 3. p.* 502.
f. 12. t. 16. f. 9. Raj. suppl. 595.
Habitat in Virginia.
5. OROBANCHE caule unifloro. *uniflora*.
 Orobanche aut Helleborine affinis marilandica, caule
 nudo, unico in summitate flore. *Raj. suppl.* 595.
 Dentaria s. Anblato cordi affinis, flore pallide cæruleo.
Gron. virg. 70.
 Gentiana minor aurea, flore simplici amplo deflexo pal-
 lide flavescente. *Pluk. mant.* 89. *t.* 348. *f.* 3.
 Aphyllon. *Mitch.* 25.
Habitat in Virginia.

DODARTIA. 687

1. DODARTIA foliis linearibus integerrimis glabris. *orientalis*.
 Dodartia *Hort. cliff.* 326. *Roy. lugdb.* 297.
 Dodartia orientalis, flore purpurascente. *Tournef. cor.*
47. itin. 3. *p.* 208. *t.* 208.
Habitat in monte Ararat, Tartaria. &
2. DODARTIA foliis ovatis serratis villosis. *indica*.
Habitat in India,
 Caules teretiusculi, villosi, subramosi. Folia ovata, pe-
 tiolata, obtuse serrata, villosa, pollicis extimo latiora.
 Racemus terminalis foliis minoribus. Flores oppositi,
 secundi, subsessiles. Calyx obtusus, villosus. Corolla
 flava, labio exteriori recto brevi.

LIPPIA. 688

1. LIPPIA. *american.*
Habitat in Vera Cruce. &
 Frutex Caulibus teretibus, pallidis, ad genicula com-
 pressis. Folia lanceolato-ovata, extra medium serrata.
R r 5 *petio-*

petiolata, scabra, opposita. Pedunculi axillares, oppositi, brevissimi, capitati. Capitula retrorsum imbricata squamis innumeris, tetragona, ovali, magnitudine pili, flosculo inter singulam squamam.

689

orientale.

SESAMUM.

1. SESAMUM foliis ovato-oblongis integris. *Hort. cliff.* 318. *Fl. zeyl.* 318.

Sesamum pedunculo inter duas glandulas. Vir. cliff. 60. *Roy. lugdb.* 292.

Digitalis orientalis Sesamum dicta. T. Burm. zeyl. 87. t. 38. f. 1.

Schit-elu. Rheed. mal. 9. p. 105. t. 54.

Habitat in Zeylona, Malabar. ☉

indianum.

2. SESAMUM foliis inferioribus trifidis. *Roy. lugdb.* 292.

Sesamum alterum, foliis trifidis, orientale, semine obscuro. Pluk. alm. 344. t. 109. f. 4.

Habitat in India. ☉

690

ringens.

MIMULUS

1. MIMULUS. *Hart. upf.* 176. t. 2. * *Aët. upf.* 1741. p. 82.

Lythimachia galericulata f. *Gratiola elatior* non tamosa. *Gron. virg.* 69.

Gratiola canadensis latifolia, flore magno cæruleo. Berk. lugdb. 2. p. 265.

Digitalis perfoliata glabra, flore violaceo minore. Meris. hist. 2. p. 479. f. 5. t. 8. f. 6. Raj. hist. 769.

Habitat in Virginia, Canada. ☿

691

strepens.

RUELLIA.

1. RUELLIA foliis petiolatis, floribus verticillatis subsessilibus. *Hort. upf.* 178.

Ruellia foliis petiolatis, fructu sessili conferto. Hort. cliff. 318. *Gron. virg.* 73. *Roy. lugdb.* 291.

Ruellia strepens, capitulis comosis. Dill. elieb. 300. t. 249. f. 321.

Habitat in Virginia, Carolina. ☿

clandestina.

2. RUELLIA foliis petiolatis, pedunculis longis subdivisis nudis. *Hort. upf.* 179.

Ruellia foliis sessilibus, pedunculis trifloris. Hort. cliff. 308. *Roy. lugdb.* 291.

Ruel-

DIDYNAMIA ANGIOSPERMIA. 635

Ruellia capsulis teretibus. Dill. elsh. 328. t. 248. f. 320.
Habitat in Barbados.

3. **RUELLIA** foliis petiolatis, pedunculis dichotomis *pa-paniculato*,
 nicularis.

Ruellia pedunculis multifloris dichotomis folio longioribus. Hort. cliff. 313. Roy. langb. 292.

Speculum veneris majus impatiens. Sloan. iam. 59. hist. 1. p. 158. t. 100. f. 2. Raj. suppl. 389.

Habitat in Jamaica. 2

4. **RUELLIA** foliis ovatis crenatis, pedunculis unifloris. *tuberosa*.

Ruellia capsulis angulosis. Dill. elsh. 328.

Ruellia numilis, flore cæruleo, asphodeli radice. Plum. gen. 12.

Gentianella flore cæruleo, integro vaseulo feminali ex humidi contactu impatiens. Sloan. jam. 52. hist. 1. p. 149. t. 95. f. 1. Raj. suppl. 370.

Habitat in Jamaica. 2

5. **RUELLIA** floribus geminis.

biflora.

Ruellia minore folio, floribus gemellis. Dill. elsh. 331.

Habitat in Carolina.

6. **RUELLIA** foliis subcrenatis lanceolato-ovatis, capis *crispa*,
 tulis ovatis foliosis hispulis.

Achatoda luzanensis, spica plana. Pet. gaz. t. 73. f. 6.?

Habitat in India. Osbeck. 2

Radix lignosa, repens. Caules digitales, simplicissimi, articulati, teretiusculi. Folia lanceolato-ovata, subpetiolata, pilis adpersa, margine reflexo undulato crispa f. crenata. Capitula ad genicula lateraliter, solitaria, ovata, imbricata. Foliis ovatis, acuminatis, piloso-scabris, intra quæ flores Calyce 5-partito hirsuto lineari. Corolla flava. Stamina. 4.

7. **RUELLIA** foliis oblongis integerrimis, floribus *foli-ringens*,
 tuis sessilibus, caule procumbente. *Fl. zeyl. 234. **

Upudali. Rheed. mal. 10. p. 125. t. 64.

Habitat in India.

8. **RUELLIA** foliis mucronato-ferratis, caule repente, *antipoda*,
 floribus subspicatis terminalibus quinis terulisve. *Fl.*

*zeyl. 235. **

Gentianella utriusque Indiæ impatiens, foliis agerati Sinop gralis barbadensibus. Pluk. mant. 167. t. 186. f. 2.

Pectanga-pulpani. Rheed. mal. 9. p. 115. t. 59.

Habitat in Indiis. ☉

BAR.

BARLERIA.

- 692
foliatis. 1. BARLERIA spinis axillaribus, foliis lanceolatis denticulatis.
 Barleria aculeata, folani folio angustiore, flore cæruleo.
Plum. gen. 31.
Habitat in America australi.
- Prionitis.* 2. BARLERIA spinis axillaribus quaternis foliis integerrimis.
 Barleria foliis integerrimis, spinis lateralibus. *Roy. lugdb.* 231. *Fl. zeyl.* 231. *
Prionitis. Hort. cliff. 486.
 Lycium indicum, spinis quaternis ad foliorum singulorum exortum. *Seb. thes.* 1. p. 21. t. 13. f. 1.
 Melampyro cognata maderaspatana, spinis horrida. *Pluk. alm.* 243. t. 119. f. 5.
 Coletta-Veetla. *Rheed. mal.* 9. p. 77. t. 41.
Habitat in India. 2
- maxifolia.* 3. BARLERIA spinis axillaribus oppositis solitariis foliis subrotundis integerrimis.
 Barleria americana spinosissima frutescens, buxi folio parvo. *Amm. herb.* 104.
 Caraschulli. *Rheed. mal.* 2. p. 91. t. 47.
Habitat in Indiis.
- erigata.* 4. BARLERIA foliis oblongis integerrimis, calycis foliolis duobus latioribus ciliatis, duobusque linearibus acutis.
Habitat in India.
 Caulis teres. Folia oblongo-ovata, utrinque acuta, integerrima. Flores axillares, sessiles. Calyx quadripartitus foliolis 2 ovatis, acuminatis, serrato-spinosis; 2 alternis brevioribus, linearibus, acutis, integerrimis, patentibus. Corolla cærulea, monopetala, erecta, infundibuliformis, patens, quinquefida: lobis ovatis subæqualibus: quinto profundius diviso. Stam. filamenta 4, quorum 2 tubo corollæ longiora, filiformia: Antheris oblongis; 2 brevissima in fundo corollæ: Anthæris emarcidis. Pistilli Germen ovatum. Stylus filiformis, longitudine majorum staminum. Stigma obtusum. Capsula longitudine calycis, compressa, acuta, bilocularis, bivalvis; valvulis cymbiformibus dissepimento longitudinali, margine altero spina elastica. Semina in singulo loculo 2, compressa.

5. BAR-

5. **BARLERIA** inermis, foliis ovatis denticulatis petio-
latis.
Barleria solani folio, flore coccineo. *Plum. gen.* 31.
Habitat in America meridionali.

DURANTA. 693

1. **DURANTA** spinosa. *repens.*
Caltorea repens spinosa. *Plum. gen.* 30.
Habitat in America.
2. **DURANTA** inermis. *erecta.*
Caltorea racemosa, flore caeruleo, fructu amaro. *Plum*
gen. 30.
Habitat in America
Caulis erectus, ramosus. Folia oblonga: inferiora opposi-
ta, suprema quaterna s. cruciatim posita.

OVIDEA. 694

1. **OVIDEA.** *spinosa.*
Valdia cardui folio, fructu subcaeruleo. *Plum. gen.* 14.
Habitat in America meridionali.
Folia opposita, petiolata, oblonga, dentata. Flores co-
rymbosi.

VOLKAMERIA. 695

1. **VOLKAMERIA** spinis petiolorum rudimentis. *aculeata.*
Ligustroides. *Hort. cliff.* 489.
Duglassia spinosa, ligustri folio. *Amm. herb.* 576.
Ligustrum aculeatum, fructu testiculato. *Plum. spec.* 489.
Paliuro affinis ligustrifolia spinosa, flore monopetalo dif-
formi. *Sloan. jam.* 137. *bist.* 2. p. 137. t. 166. f.
2. 3. *Raj. dendr.* 95.
Habitat in Jamaica, Barbados. b
2. **VOLKAMERIA** ramis inermibus. *Fl. zeyl.* 231. * *inermis.*
Periclymeni similis myrtifolia arbor maderaspatensis.
Pluk. alm. 287. t. 211. f. 4.
Habitat in India. b

CLERODENDRUM. 696

1. **CLERODENDRUM** foliis simplicibus cordatis to-
mentos. *Fl. zeyl.* 232. * *informata*
Peragu. *Rheed. mal.* 2. p. 41. t. 25. *Raj. bist.* 1571.
β. Clerodendrum folio lato & acuminato. *Burm. zeyl.* 66.
t. 29.
Habitat in India VI.

VITEX.

697
Agnus Castus.

1. VITEX foliis digitatis, spicis verticillatis.
Vitex. Cam. epit. 105. Hort. cliff. 327. Roy. lugdb. 291. Gron. virg. 169. Mat. med. 319.
Vitex foliis angustioribus cannabis modo dispositis.
Bauh. pin. 475.
- β. *Vitex latiore folio. Bauh. pin. 475.*
Vitex folio serrato. Bauh. hist. 1. p. 205.
Habitat in Sicilia et Neapolis paludosis. ♀
- trifolius.* 2. VITEX foliis ternatis quinatisque, paniculis dichotomis.
Vitex trifolia minor indica. Pluk. alm. 390. t. 206. f. 5. Fl. zeyl. 413.
Piperi similis, fructus striatus, semina. Bauh. pin. 412.
Caranosi. Rheed. mal. 2. p. 13. t. 10. Raj. hist. 1575.
Habitat in India.
Foliola ovata, acuta, integerrima, subtus tomentosa.
- Negundo.* 3. VITEX foliis quinatis ternatisque serratis, floribus racemoso-paniculatis.
Vitex trifolia minor indica serrata. Pluk. alm. 390. Fl. zeyl. 414.
Negundo arbor mas. Bauh. hist. 1. p. 189.
Bernnosi. Rheed. mal. 2. p. 15. t. 11. Raj. hist. 1575.
Habitat in India. ♀
Foliola 3. f. 5, ovato-lanceolata, sinuato-serrata, subtus nuda venosa.
- pinnata.* 4. VITEX foliis pinnatis, paniculis trichotomis.
*Pistacio-vitex. Fl. zeyl. 415. **
Habitat in Zeylona. ♀

BONTIA.

698
daphnoides.

1. BONTIA.
Bontia arborescens, thymelææ facie. Plum. gen. 32.
Bontia laureolæ facie. Dill. elth. 57. t. 49. f. 57.
Olea sylvestris barbadensis, folio angusto piligui leviter crenato, Pluk. alm. 269. t. 209. f. 3. Raj. dendr. 47.
Habitat in Barbados. ♀

COLUMNEA.

699
scandens.

1. COLUMNEA.
Columnnea scandens, phœniceo flore, fructu albo. Plum. gen. 28.
- β. *Columnnea scandens, flore lutescente, fructu albo. Plum. gen. 28.*
Ille

Habitat in Gallia æquinoctiali.

Canlis, Folia, Corollæ villosa. Folia ovata, serrata, petiolata.

ACANTHUS. 700

1. ACANTHUS foliis sinuatis inermibus. *Hort. cliff. mollis.*

326. *Hort. upf.* 181. *Mat. med.* 318. *Roy. lugdb.*

304. *Sauv. monsp.* 278.

Acanthus sativus f. *mollis virgilii*. *Baub. pin.* 383.

Carduus Acanthus f. *Branca ursi*. *Baub. hist.* 3. p. 75.

Habitat in Italiæ, Siciliæ humentibus, duris. &

2. ACANTHUS foliis pinnatifidis spinosis. *Hort. cliff. spinosus.*

326. *Roy. lugdb.* 304.

Acanthus aculeatus. *Baub. pin.* 383.

Acanthus sylvestris. *Dod. pempt.* 719.

Habitat in Italiæ humentibus. &

3. ACANTHUS caule fruticoso aculeato.

tricuspidatus.

Acanthus malabaricus, agrifolii folio. *Pet. succ.* 10. *gaz.*

t. 94. f. 15.

Frutex indicus spinosus, foliis agrifolii, filiqua geminata brevi. *Raj. hist.* 1766. *Pluk. phyt.* 261. f. 4.

Carduus aquaticus f. *indorum* *Dilivaria*. *Camell. lux.*

G. n. 16.

Paiba schylli. *Rheed. mal.* 2. p. 93. t. 48.

Habitat in India. &

4. ACANTHUS foliis quaternis obsolete dentatis inermibus, calycibus ciliatis. *maderaspatis.*

Adharoda maderaspatis, calyce barbato, foliis subtus villosis. *Pet. musf.* 23.

Habitat in India.

MELIANTHUS. 701

1. MELIANTHUS stipulis solitariis petiolo adnatis. *major.*

Hort. cliff. 492. *Hort. upf.* 181. *Roy. lugdb.* 402.

Melanthus africanus. *Herm. lugdb.* 414. t. 415.

Habitat in Æthiopiæ succulentis. &

2. MELIANTHUS stipulis geminis distinctis. *Hort. minor.*

cliff. 492. *Roy. lugdb.* 402.

Melanthus africanus minor foetidus. *Comm. rar.* 4. t. 4.

Habitat in Æthiopia. &

Clas.

TETRADYNAMIA.

SILICULOSA.

702

MYAGRUM.

- perenne.* 1. MYAGRUM siliculis biarticulatis dispermis, foliis extrorsum sinuatis denticulatis. *Hort. nps.* 182. *
Crambe foliis lanceolatis dentato-sinuatis. Hort. cliff.
340. Roy. lugdb. 329. Gort. getr. 404.
Rapistrum monospermum. Baub. pin. 95. prodr. 37.
t. 37. Mapp. asat. 266. t. 266.
Habitat in Asia. 4
- orientale.* 2. MYAGRUM siliculis sulcatis lævibus, foliis oblongis dentato-sinuatis. *Hort. nps.* 182. *
Rapistrum orientale, folio raphani, capsulis rugosis. Baub.
lugdb. 2. p. 2.
Habitat in Oriente. 5
- rugosum.* 3. MYAGRUM siliculis sulcatis pilosis rugosis, foliis oblongis obtusis dentatis. *Hort. nps.* 103. *
Habitat in Europa australi. 5
- hispanicum.* 4. MYAGRUM siliculis lævibus subtorosis, foliis lyratis.
Sinapi hispanicum minus, raphani folio. Tournef. inst.
227
Habitat in Hispania. 3
Caulis pilis raris reflexis scaber. Racemi virgati, longi
Folia lyrata, dentata, petiolata, oblonga. Siliculæ cy-
lindricæ, laeves, nec sulcatæ rugosæ aut striatæ; con-
stantes Corpore oblongo, pedicellato, cylindrico, sterili, cu-
Silicula ovata insidet collo angustiore: articulo inferiori
biloculari inani, superiore monospermo. Corollæ luteæ
- perfoliatum* 5. MYAGRUM siliculis cordatis subsessilibus foliis, amplexicaulibus. *Hort. nps.* 182. * *Hort. cliff.* 328. *Roy.*
lugdb. 330. Savv. monsp. 77.
Myagrum monospermum latifolium. Baub. pin. 104
prodr. 52. t. 51. Moris. hist. 2. p. 267. f. 3. t. 21.
antepenult.
Habitat inter Gallie segetes. 5

Silicul

TETRADYNAMIA SILICULOSA 641

Siliculae triloculares sed monospermae, f. in medio loculo tantum semine facta.

6. MYAGRUM filiculis ovatis pedunculatis polyspermis. *Jatium*.
Hort. cliff. 328. *Fl. suec.* 197. *Roy. lugdb.* 330.
Dalib. paris. 193.
Myagr. sylvestre. *Bauh. pin.* 109.
β. Myagr. sativum. *Bauh. pin.* 109.
Camelina f. Myagrion. *Dod. pempt.* 532.
Habitat in Europa inter linum. ☉
7. MYAGRUM filiculis globosis compressis punctato-paniculatum rugosis. *Hort. cliff.* 328. *Fl. suec.* 542. *Roy. lugdb.* 330. *Hall. belv.* 536. *Dalib. paris.* 193.
Myagr. monospermum hirsutum, filiquis rotundis
Lef. pruss. 174. t. 56.
Myagro similis filiqua rotunda. *Bauh. pin.* 109. *prodr.* 52. t. 52.
Habitat in Europa, juxta agros. ☉

8. MYAGRUM filiculis angulatis, foliis tripartitis. *Hasselt.* *egyptium.*
Habitat in Aegypto.
Caulis divaricatus, Racemis virgatis. Folia petiolata, tripartita vel quinquepartita, dentata. Capsulae duobus articulis tetragonis. Flores lutei. An potius Bunias?

VELLA. 703

1. VELLA foliis pinnatifidis, filiculis pendulis, *annua.*
Vella. Hort. cliff. 329. *Hort. ups.* 186. *Roy. lugdb.* 331.
Nasturtium sylvestre, erucae facie. *Bauh. pin.* 105.
Nasturtium sylvestre valentinum. *Clus. hist.* 2. p. 130.
Habitat in Hispania ☉
2. VELLA foliis integris, filiculis erectis. *PseudoCytisus.*
PseudoCytisus flore leucoji luteo. *Bauh. pin.* 230.
Cytisi facie Alysson fruticans quorundam. *Lob. ic.* 2. p. 49. *Bauh. hist.* 1. p. 374.
Habitat in Hispania circa oppidum Aranjuez. D. Minuart. ☉
Planta frutescens, Foliis integris, sessilibus, birtis. obverse ovatis, Folliculis erectis. *Minnart.*

ANASTATICA. 704

1. ANASTATICA. *Hort. cliff.* 328. *Hort. ups.* 183. *hierocuntica.*
Roy. lugdb. 331. S s Thlaspi

642 TETRADYNAMIA SILICULOSA,

- Thlaspi Rosa de Hiericho dictum. Moris. hist. 2. p. 228.*
f. 5. t. 25. f. 2. 3.
Rosa hierochuntea vulgo dicta. Bauh. pin. 484.
Rosa hierochuntica. Com. hort. t. 41.
Habitat in littoribus maris rubri. ☉

SUBULARIA.

- 705*
aquatica. 1. **SUBULARIA.** *Fl. lapp. 253. Fl. succ. 527.*
Alyssum palustre, folio junci. Buxb. act. 2. p. 369. t.
23. f. 1.
Graminifolia aquatica, thlaspeos capitulis rotundis: sepi-
mento siliculam dirimente. Pluk. alm. 180. t. 188. f. 5.
Habitat in Europæ borealis inundatis lacustribus sin-
viis. ☉

DRABA.

- 706*
alpina. 1. **DRABA** scapo nudo simplici, foliis lanceolatis integer-
rimis. Fl. lapp. 255. Fl. succ. 524. Hort. cliff. 333.
Roy. lugdb. 333. Dalib. parisi. 197.
Sedum alpinum hirsutum luteum. Bauh. pin. 284.
Leucojum luteum aizoides montanum. Col. ecph. 2.
p. 62.
Bursa pastoris alpina rosca lutea. Moris. hist. 2. p. 306.
f. 3. t. 20. f. 6.
Habitat in Alpibus Europæ. 2
- pyrena.* 2. **DRABA** caulibus nudis, foliis incis. *Hort. cliff. 333.*
Fl. succ. 523. Roy. lugdb. 333. Hall. belv. 538.
Gron. virg. 76.
Bursa pastoris minor, loculo oblongo. Bauh. pin. 108.
Alysson vulgare, polygoni folio, caule nudo T. Segu.
veron. 1. p. 575. t. 4. f. 3.
Pilosella minor. Thal. barc. 84. t. 7. f. E
Habitat in Europæ aridis. ☉
- pyrenæica.* 3. **DRABA** caule nudo, foliis pinnatis ternatisque. *Læf.*
Alysson pyrenaicum perenne minimum, foliis trifidis.
Tournef. inst. 217.
Habitat in Pyrenæis. 2
- muralis.* 4. **DRABA** caule ramoso, foliis cordatis dentatis amplexi-
*caulibus. Roy. lugdb. 33. It. gotl. 192. * Fl. succ*
525. Hall. belv. 539.
Bursa pastoris major loculo oblongo. Bauh. pin. 108
prodr. 50. t. 50.
Habitat in Europæ nemoribus. ☉

5. **DRA**

TETRADYNAMIA SILICULOSA. 643

5. DRABA caule ramoso, foliis ovatis sessilibus dentatis. *nemorosa*, Bursa pastoris alpina hirsuta. *Bauh. pin.* 108. *prodr.* 51. t. 51?

Draba minima muralis discoides. *Col. ecpbr.* 1. p. 274. t. 272.

Habitat in Sueciæ nemoribus. ☉.

Præcedenti simillima, sed folia majora, minus acuta, magis hirsuta, sessilia nec amplexicaulia. Flores lutei, nec ut in præcedenti albi.

6. DRABA foliis caulinis numerosis incauis, siliculis oblongis. *Fl. succ.* 526. *It. gotl.* 192. *

Draba caule ramoso folioso, foliis dentatis. *Fl. tapp.* 254. *Hort. cliff.* 334.

Leucojum f. Lunaria vasculo oblongo intorto. *Pluk. alm.* 215. t. 42. f. 1.

Habitat in Alpibus & alpinis Europæ. ♂

LEPIDIUM. 707

1. LEPIDIUM foliis caulinis pinnato-multifidis, rami-*perfoliatum*, feris cordatis amplexicaulibus integris. *Hort. cliff.* 331.

Hort. nsp. 183. *Roy. lugdb.* 335.

Nasturtium spicatum persicum perfoliatum maximum.

Morif. hist. 2. p. 294. f. 3. t. 25. f. 17.

Thlaspi verum dioscoridis 1. *Zan. hist.* 193.

Habitat in Persia, Syria. ☉

2. LEPIDIUM cauliumgeniculis inflatis.

vesicarium.

Lepidium orientale, nasturtii folio, caule vesicario. *Tournef. cor.* 15. *Buxb. cent.* 1. p. 17. t. 26.

Habitat in Iberiæ, Mediæ aridis. ☉

3. LEPIDIUM scapo nudo simplicissimo, floribus te-*nudiculis*, trandris.

Lepidium foliis filiformibus apice pinnatifidis, caule nudo. *Sauv. monsp.* 228. 281.

Nasturtium minimum vernum, foliis tantum circa radicem. *Magn. monsp.* 187. t. 186.

Habitat Monspeli. ☉

Folia radicalia multa, linearia, apice pinnatifida vel *dentata*. Siliculæ emarginatæ. Petala equalia. Stamina, 4.

4. LEPIDIUM foliis pinnatis: impari majore, caule pro-*procumbens*, cumbente. *Sauv. monsp.* 228.

Nasturtium foliis omnibus alternatim pinnatis. *Dalib. parif.* 194.

644 TETRADYNAMIA SILICULOSA.

Nasturtium pumilum vernalium. *Magn. monsp.* 185. t. 184.

Habitat Montpelii. ☉

petraeum.

5. LEPIDIUM foliis pinnatis integerrimis, petalis emarginatis calyce minoribus. *It. al.* 52. *Fl. succ.* 535. Nasturtium alpinum tenuissime divifum. *Bauh. pin.* 105. Nasturtium petraeum annuum nostras. *Pluk. alm.* 162. t. 206. f. 4. *mal.*

Cardamine pusilla saxatilis montana discoides. *Col. ceph.* 1. p. 274. t. 273. *bon.*

Nasturtium pumilum vernalium. *Bauh. pin.* 105.

Habitat in Lapidosis Oelandiae, Angliae. ☉

futuum.

6. LEPIDIUM floribus tetradynamis, foliis oblongis multifidis. *Vir. cliff.* 63. *Hort. upf.* 183. *Mat. med.* 322. *Roy. lugdb.* 335.

Lepidium folliis varie divifis incisisque. *Hort. cliff.* 331.

Nasturtium sylvestre vulgatum. *Bauh. pin.* 103.

Nasturtium hortense. *Dod. pempt.* 771.

- β. Nasturtium hortense crispum. *Bauh. pin.* 104. *prodr.* 44. t. 43.

Habitat - - - ☉

lyratum.

7. LEPIDIUM foliis lyratis crispis.

Lepidium orientale, nasturtii crispi folio. *Tournef. cor.* 15. *itin.* 339. t. 339.

Habitat in Oriente.

latifolium.

8. LEPIDIUM foliis ovato-lanceolatis integris serratis. *Hort. cliff.* 330. *Fl. succ.* 533. *Roy. lugdb.* 334. *Dalib. parif.* 194.

Lepidium latifolium. *Bauh. pin.* 97.

Lepidium plinii. *Dod. pempt.* 716.

Habitat in Galliae, Angliae umbrosis, succulentis. &

subulatum.

9. LEPIDIUM foliis subulatis indivifis fparfis, caule fuf-fruticofio.

Lepidium, capillaceo folio, fruticofum hispanicum. *Tournef. inf.* 216.

Habitat in Hispania. Laff. &

Radix perennis, cespitem promens foliorum & ramulorum persistentium. Caules vix spithamei, simplices, plures. Folia fparfa, fubulata, linearia, acuta, fupra plana, fubtus femitretia. Racemi terminales & ex alis fupremis foliorum, simplices. Petala alba. Siliculae omnino L. ruderalis. Stamina sex.

10. LE-

TETRADYNAMIA SILICULOSA. 645

10. LEPIDIUM floribus diandris apetalis, foliis radicalibus dentato-pinnatis; ramiferis linearibus integerrimis. *Fl. suec.* 534. *Hort. cliff.* 331. *Roy. lugdb.* 335. Nasturtium sylvestre, osyridis folio. *Bauh. pin.* 105. Nasturtium angustifolium. *Fuchf. hist.* 307. *Habitat in Europæ ruderatis & ad vias.* ☉
11. LEPIDIUM floribus subtriandris tetrapetalis, foliis linearibus pinnatis. *Virgati- cum.*
Lepidium foliis lanceolato-linearibus serratis. *Gron. virg.* 76.
Lepidium foliis omnibus lineari-lanceolatis serratis. *Roy. lugdb.* 334.
Iberis humilior annua virginiana ramosior. *Morif. hist.* 2. p. 311. f. 3. t. 21. f. 2. *Sloan. jam.* 80. *hist.* 1. p. 195. t. 123. f. 3. *Raj. hist.* 827.
Iberis virginiana latifolia ramosa. *Morif. præl.* 277.
Habitat in Virginia, Jamaicae glareosis. ☉
12. LEPIDIUM floribus diandris tetrapetalis, foliis inferioribus lanceolatis serratis; superioribus linearibus integerrimis. *Roy. lugdb.* 334.
Lepidium foliis lanceolato-linearibus serratis. *Hort. cliff.* 331.
Iberis latiore folio. *Bauh. pin.* 97.
Iberis. *Dod. pempt.* 714.
Habitat in Germania, Gallia, Italia secus vias. ☉
13. LEPIDIUM floribus diandris tetrapetalis, foliis omnibus pinnato-multifidis.
Lepidium foliis pinnatifidis incis. *Roy. lugdb.* 335.
Thlaspi bonariense multifidum, flore invilibili. *Dill. elsb.* 281. t. 286. f. 370.
Habitat in Bonaria. ☉
14. LEPIDIUM foliis lanceolatis amplexicaulibus dentatis. *Hort. cliff.* 331. *Sauv. monsp.* 78. 120.
Draba umbellata f. Draba major capitulis donata. *Bauh. pin.* 109. *Morif. hist.* 2. p. 313. t. 21. f. 1.
Draba 1. vulgaris. *Clus. hist.* 2. p. 123.
Draba chalepensis repens humilior, foliis minus cinereis & quasi viridibus. *Morif. hist.* 2. p. 314.
Habitat in Germania, praesertim Austria, Gallia, Italia. 2
- THLASPI. 708
1. THLASPI filiculis suborbiculatis, foliis lanceolatis peregrinum integerrimis. *Hort. cliff.* 330. * *Roy. lugdb.* 334.
S s 3 Thla-

646 TETRADYNAMIA SILICULOSA

Thlaspi, capsula cordata, peregrinum. *Bauh. hist.* 2. p. 927. *Morif. hist.* 2. p. 297. f. 3. t. 18. f. 30.
Habitat - - - - -

- arvense.* 2. THLASPI siliculis orbiculatis, foliis oblongis dentatis glabris. *Fl. lapp.* 251. *Fl. succ.* 530. *Hort. cliff* 330. *Mat. med.* 323. *Roy. lugdb.* 333.
 Thlaspi arvense, siliquis latis. *Bauh. pin.* 105.
 Thlaspi latius. *Dod. pempt.* 712.
Habitat in Europæ agris. ☉
- alliacum.* 3. THLASPI siliculis subovatis ventricosis, foliis oblongis obtusis dentatis glabris. *Roy. lugdb.* 334.
 Thlaspi allium redolens. *Morif. hist.* 2. p. 297. f. 3. t. 18. f. 28.
 Scorodo-Thlaspi ulyffis aldrovandi. *Bauh. hist.* 2. p. 932.
Habitat in Europa australi.
Antecedenti similis, sed Silicula ovato-ventricosa, margine minore.
- saxatile.* 4. THLASPI siliculis ellipticis, foliis lanceolato-linearibus integerrimis. *Sauv. monsp.* 53. 71. †
 Thlaspi parvum saxatile, flore rubente. *Bauh. pin.* 107.
 Lithonthlaspi quartum, carnosio rotundo follo. *Col. ecphr.* 1. p. 279. t. 277. f. 2
Habitat in Italiæ, G. Narbonensis, Provinciæ saxoss.
- perfoliatum.* 5. THLASPI siliculis subrotundis, foliis amplexicaulibus cordatis subserratis.
 Thlaspi foliis radicalibus ovatis, caulinis amplexicaulibus lanceolatis. *Guett. stamp.* 142. *Dalib. paris.* 196
 Thlaspi arvense perfoliatum majus. *Bauh. pin.* 106.
 Thlaspi oleraceum. *Tabern. ic.* 462.
 β. Thlaspi perfoliatum minus. *Bauh. pin.* 106.
 Thlaspi 3. pumilum. *Clus. hist.* 2. p. 131.
Habitat in Germaniæ, Helvetiæ, Galliæ apricis, agris.
- hirtum.* 6. THLASPI siliculis subrotundis pilosis, foliis sagittatis hirsutis. *Roy. lugdb.* 333. *Sauv. monsp.* 120.
 Thlaspi capsulis hirsutis. *Bauh. hist.* 2. p. 922.
 Thlaspi villosum, capsulis hirsutis. *Bauh. pin.* 106.
prodr. 212. t. 212.
Habitat in Italia, Narbona.
- campestre.* 7. THLASPI siliculis subrotundis, foliis sagittatis dentatis incanis. *Hort. cliff.* 330. *Fl. succ.* 531. *Roy. lugdb.* 333. *Hall. belv.* 544. *Dalib. paris.* 196.

Thla-

TETRADYNAMIA SILICULOSA. 647

- Thlaspi arvense*, vaccariæ incano folio, majus. *Bauh.*
pin. 106.
Thlaspi latifolium. *Fuchf. hist.* 306.
Habitat in Europæ arvis, vis argillofis, apricis. &
8. **THLASPI** filiculis obcordatis, foliis radicalibus cu-
neiformibus integerrimis.
Thlaspi filiculis obverse cordatis, foliis imis spathula-
tis; summis amplexicaulibus sagittatis. *Sauv. monsp.*
121.
Thlaspi alpinum, bellidis cœruleæ folio. *Bauh. pin.*
106.
Thlaspi montanum, burfæ pastoris folio. *Col. ecphr.* 1.
p. 275. t. 276.
Thlaspi montanum 2. *Clus. hist. 2. p. 131.*
Habitat in Helvetiæ, Austriæ, Italiæ, Monspelii petrosis.
9. **THLASPI** filiculis obcordatis, foliis radicalibus pin-
natifidis. *Fl. lapp. 252. Fl. succ. 532. Hort. cliff. 330.*
Mat. med. 324. Roy. lugdb. 334. Hall. belv. 145.
Dalib. parif. 196.
Bursa pastoris major, folio sinuato. *Bauh. pin. 108.*
Bursa pastoris. *Fuchf. hist. 611.*
β. Bursa pastoris media. *Bauh. pin. 108.*
Habitat in Europæ cultis ruderatis. ☉

COCHLEARIA. 709

1. **COCHLEARIA** foliis radicalibus subrotundis, cau-
linis oblongis subsinuatis. *Fl. lapp. 256. Fl. succ. 537.*
Hort. cliff. 332. Mat. med. 320. Roy. lugdb. 335.
Cochlearia folio subrotundo. *Bauh. pin. 110.*
Cochlearia. *Dod. pempt. 594.*
Habitat in Europæ borealis littoribus marinis. ☉
2. **COCHLEARIA** foliis hastatis angulatis. *Fl. succ. 196. danica.*
Cochlearia aremorica. *Tournef. inst. 215.*
Thlaspi hederaceum. *Lob. ic. 615.*
β. Cochlearia danica repens. *Bauh. pin. 110. prodr. 53.*
γ. Cochlearia minor erecta. *Bauh. pin. 110. prodr. 53.*
Habitat in Danicæ, Sueciæ littoribus marinis. ☉
3. **COCHLEARIA** foliis reniformibus integris. *Hort. grœnlandica.*
*cliff. 498. * Roy. lugdb. 335.*
Cochlearia minima repens insulæ alholmiæ. *Barth. act.*
3. p. 143. t. 144.
Cochlearia minima. *Boerb. lugdb. 2. p. 10.*
Habitat in Norvegia, Islandia, Grœnlandia. ☉

648 TETRADYNAMIA SILICULOSA.

- foetilis.* 4. COCHLEARIA foliis radicalibus lanceolatis serratis, caulinis linearibus. *Sauv. monsp.* 122.
Cochlearia perennis saxatilis minima. *Magn. hort.* 59.
Thlaspi alpinum minus, capitulo rotundo. *Bauh. pin.* 107. *prodr.* 48. t. 48.
Thlaspi tertium saxatile. *Cam. epit.* 338.
β. Thlaspi alpinum majus, capitulis rotundis. *Bauh. prodr.* 49. t. 49. *pin.* 106. n. 6.
Habitat in Alpibus Helvetiæ, Monspeli. 2
- Coronopus.* 5. COCHLEARIA foliis pinnatifidis. *Hort. cliff.* 331.
Fl. succ. 539. *Roy. lugdb.* 335. *Dalib. parif.* 195.
Sauv. monsp. 254.
Ambrosia campestris repens. *Bauh. pin.* 138.
PseudoAmbrosia. *Cam. epit.* 596.
Habitat in Europæ apricis, nudis. ☉
- Armoracia.* 6. COCHLEARIA foliis radicalibus lanceolatis crenatis, caulinis incilis. *Hort. cliff.* 332. *Fl. succ.* 540. *Mat. med.* 321. *Roy. lugdb.* 336.
Raphanus rusticus. *Bauh. pin.* 96.
Raphanis magna. *Dod. pempt.* 678.
Habitat in Europæ fossis & ad rivulos. 2
- glastifolia.* 7. COCHLEARIA foliis caulinis cordato-sagittatis amplexicaulibus. *Hort. cliff.* 332. *Hort. ups.* 184. *Roy. lugdb.* 335.
Lepidium glastifolium. *Bauh. pin.* 97. *Morif. hist.* 2. p. 312. f. 3. t. 21. f. 3.
Lepidium annuum. *Dalech. hist.* 1297.
Habitat - - - - - ♂

716

IBERIS.

- semperflorens.* 1. IBERIS frutescens, foliis cuneiformibus integerrimis obtusis. *Hort. cliff.* 330. *Hort. ups.* 184. *Roy. lugdb.* 336.
Leucojum fruticosum umbellatum persicum, foliis leucoji intar sempervirentibus. *Morif. hist.* 2. p. 296.
Thlaspi latifolium polycarpon, leucoji foliis. *Bocc. sic.* 55. t. 22. f. a-1.
Thlaspi fruticosum, folio leucoji, semperflorens. *Seb. xbof.* 1. p. 2. t. 13. f. 4.
Habitat in Sicilia, Persia. 5
- sempervirens.* 2. IBERIS frutescens, foliis linearibus acutis integerrimis. *Hort. cliff.* 330. *Roy. lugdb.* 336.

Thla-

TETRADYNAMIA SILICULOSA. 649

Thlaspi montanum sempervirens. *Baub. pin.* 106.

Thlaspi montanum candidum. *Dalech. hist.* 1180.

Thlaspi creticum perenne, flore albo. *Barr. ic.* 214. ☞

724.

Habitat in Cretæ rupestribus. ☞

3. IBERIS frutescens foliis apice dentatis.

gibraltarica

Thlaspidium hispanicum, ampliore flore, folio crasso
dentato. *Dill. elib.* 382. t. 287. f. 37.

Habitat ad Gibraltariam Hispaniæ. ☞

4. IBERIS foliis subrotundis crenatis. *Roy.*

rotundifolia.

Thlaspi montanum, serrato cepeæ folio, flore purpura-
scente umbellato. *Barr. ic.* 848.

Thlaspi alpinum, folio rotundiore carnosio, flore pur-
purescente. *Tournef. inst.* 212. *Schenck. alp.* 50. 143.

Lepidium caule repente, foliis ovatis amplicaulibus.

Hall. helv. 546.

Habitat in Helvetia. ☞

5. IBERIS foliis lanceolatis acuminatis: inferioribus ser-
natis; superioribus integerrimis. *Hort. cliff.* 330. *Hort.*

ups. 184. *Roy. lugdb.* 336

Thlaspi umbellatum creticum, iberidis folio. *Baub. pin.*
106.

Draba f. Arabis f. Thlaspi candiæ. *Dod. pempt.* 713.

Habitat in Metturgia, Creta. ☉

Fructus hujus corymbosus hemisphericus persistit, f. a-
mare vero racemosus evadit.

6. IBERIS foliis lanceolatis indivisis, corymbis hemisphæ-
ricis. *Sauv. monsp.* 72. †

Thlaspi creticum, foliis indivisis, f. lini, amarum. *Baub.*
hist. 2. p. 924.

Habitat Monspeli. ☉

7. IBERIS foliis lanceolatis acutis subdentatis, floribus
racemosis. *Hort. ups.* 184.

Iberis foliis obverse lanceolatis, floribus umbellatis.

Guss. stamp. 2. p. 146.

Thlaspi umbellatum arvense, iberidis folio. *Baub. pin.*
106.

Thlaspidium folio iberidis. *Riv. test.* 112.

Habitat in Helvetia ☉

8. IBERIS foliis linearibus superne dilatatis serratis. *Roy. odorata.*
lugdb. 336. *Hort. cliff.* 330.

Thlaspi umbellatum creticum, flore albo odoro, minus.
Baub. pin. 106.

Thla-

650 TETRADYNAMIA SILICULOLA.

Thlaspi parvum 4, odorato flore. *Glas. hist.* 2. p. 132.
Habitat in Alpibus Allobrogicis. ☉

- nadicantis.* 9. *IBERIS* foliis sinuatis, caule nudo simplici. *Hort. cliff.* 328. *Fl. suec.* 536. *It. el.* 139. *Roy. lugdb.* 336.
Bursa pastoris minor, foliis incis. *Baub. pin.* 108.
Bursa pastoris minor. *Dod. pempt.* 103.
Nasturtium petræum. *Tabern. sc.* 451.
Habitat in Europæ arenosis, sylvaticis, nudis, sterilibus. ☉

711 ALYSSUM.

* *Suffruticosa.*

- spinosum.* 1. *ALYSSUM* ramis senilibus spiniformibus nudis
Hort. cliff. 332. *Roy. lugdb.* 331. *Dalib. paris.* 197.
Thlaspi fruticolum spinosum. *Baub. pin.* 108.
Leucojum f. *Thlaspi spinosum.* *Baub. hist.* 2. p. 931.
Thlaspi spinosum hispanicum. *Borr. ic.* 808.
Habitat in Hispaniæ, Galliæ cantibus. ☉

- montanum.* 2. *ALYSSUM* ramulis suffruticosis diffusis, foliis punctato echinatis. *Hort. ups.* 185.
Alyssum foliis lanceolatis obtusis incanis, caulibus procumbentibus, radice perenni. *Hort. cliff.* 332. *Roy. lugdb.* 331.
Thlaspi montanum luteum. *Baub. hist.* 2. p. 928. t. 929.
Habitat in Helvetia. ☉

- halimifolium.* 3. *ALYSSUM* foliis lanceolato-linearibus acutis integerrimis, caulibus procumbentibus perennantibus. *Hort. cliff.* 333. *Roy. lugdb.* 331.
Thlaspi, halimi folio, sempervirens. *Herm. lugdb.* 59. t. 595.

- β. *Thlaspi parvum, halimi angusto incano folio.* *Boerh. mss.* 2. p. 45. t. 39.
Habitat in Europæ australis aridis. ☉

- saxatile.* 4. *ALYSSUM* caulibus frutescentibus paniculatis, foliis lanceolatis undulatis integris. *Roy. lugdb.* 331. n. 1.
Alysson folio leucoji incano, flore luteo. *Boerh. lugdb.* 2. p. 3.
Alyssum creticum saxatile, foliis undulatis incanis. *Tonnesf. cor.* 15.
Habitat in Creta. ☉

* *Herbacea.*

- incanum.* 5. *ALYSSUM* caule erecto, foliis lanceolatis incanis i

TETRADYNAMIA SILICULOSA. 651

tegerimis, floribus corymbosis. *Hort. cliff.* 332. *Fl. succ.* 528. *Roy. lugdb.* 331.

Thlaspi fruticosum incanum. Bauh. pin. 108.

Thlaspi incanum machliniense. Clus. hist. 2. p. 132.

Habitat in Europæ septentrionalioris arenosis, apricis. 2 ♂

8. ALYSSUM caulibus herbaceis diffusis, foliis lineari-*minimum.* bus tomentosis, siliculis compressis.

Alyssum procumbens annuum, foliis lineari-lanceolatis subtomentosis. Samv. meth. 70.

Lunaria annua minima hispanica, folio leucoji maritimi. Borrh. lugdb. 2. p. 6.

Habitat in Hispania. ☉

9. ALYSSUM caule erecto herbaceo, siliculis sessilibus *clypeatum.* ovalibus compresso-planis, petalis acuminatis.

Lunaria siliculis subsessilibus incanis lateralibus. Roy. lugdb. 333.

Lunaria leucoji folio, siliqua oblonga majori. Tournef. inst. 218.

Leucojum alyssoides clypeatum majus. Bauh. pin. 201.

Alysson dioscoridis. Doct. pempt. 89.

Habitat in Europa australi. ☉

10. ALYSSUM caule herbaceo, foliis lanceolatis *denta- sinuatum.* tis, siliculis inflatis

Alyssum foliis lanceolatis repando-dentatis. Hort. cliff. 333. *Roy. lugdb.* 332.

Alyssoides incanum, foliis sinuatis. Tournef. inst. 213.

Leucojum incanum, siliquis rotundis. Bauh. pin. 201.

Erica peregrina. Clus. hist. 2. p. 134.

Habitat in Hispaniæ incultis, ad vias. ☉ ♂

Petala bifida. Filamenta lateralibus obsolete dentata.

11. ALYSSUM caule herbaceo erecto, foliis incanis lanceolatis integerrimis, siliculis inflatis. *eroticum.*

Alyssoides fruticosum creticum, leucoji folio incano. Tournef. cor. 15.

Leucojum luteum, utriculato semine. Alp. exot. 117. t. 118.

Habitat in Creta.

12. ALYSSUM foliis linearibus dentatis, siliculis inflatis *Vesicaria.* angulatis acutis.

Vesicaria orientalis, foliis dentatis. Tournef. cor. 49. *itin.* 2. p. 97. t. 14.

Habitat in Oriente.

13. ALYSSUM foliis incanis dentatis, staminibus quatuor *hyperboreum.* bifurcatis. Lu-

652 TETRADYNAMIA SILICULOSA.

Lunaria foliis ellipticis incondite dentatis. *Kraschen ad. petr.* 1747. p. 381. t. 15. f. 1.

Habitat in America septentrionali. D. Krascheninnikof
Stamina 2 breviora ungue acuto notata, reliqua 4 ad
basin squama emarginata.

712

CLYPEOLA.

Thlaspi. 1. CLYPEOLA filiculis unilocularibus monospermis.

Hort. cliff. 329. *Hort. ups.* 185. *Roy. lugdb.* 332.

Thlaspi clypeatum, serpilli folio. *Baub. pin.* 107.

Thlaspi minimum spicatum lunatum. *Col. exphr.* 1
p. 281. t. 284.

Habitat in Italiæ, Narbonæ sabulosis. ☉

Alyssoides. 2. CLYPEOLA filiculis bilocularibus tetraspermis *Hort cliff.* 329. *Roy. lugdb.* 332.

Thlaspi Alysson dictum campestre majus. *Baub. pin.* 107.

Alysson minimum. *Clus. hist.* 2. p. 133.

Habitat in Austria, Gallia. ☉

3. CLYPEOLA annua, filiculis bilocularibus dispermis, calycibus persistentibus. *Sauv. monsp.* 71. †

Thlaspi Alysson dictum campestre minus. *Baub. pin.* 107.

Habitat in Gallia. ☉

maritima. 4. CLYPEOLA perennis, filiculis bilocularibus ovatis dispermis. *Sauv. monsp.* 71. †

Thlaspi Alysson dictum maritimum. *Baub. pin.* 107.

Thlaspi narbonense, centunculi angusto folio. *Tabern. ic.* 461.

Habitat in G. Narbonensi. 2

3

BISCUTELLA.

auriculata. 1. BISCUTELLA calycibus nectario utrinque gibbis, filiculis in stylum coeuntibus. *Hort. cliff.* 329. *Hort. ups.* 185. *Roy. lugdb.* 336.

Thlaspidium biscutatum villosum, flore calcari donato. *Baub. pin.* 107. *prodr.* 49.

Leucojum montanum, flore pedato. *Col. exphr.* 2. p. 59. t. 61.

JonDraba alyssoides lutea angustifolia. *Barr. ic.* 230.

Habitat in Italia, Galloprovincia. ☉

2. BI-

TETRADYNAMIA SILIQUOSA. 653

2. BISCUTELLA filiculis orbiculato-didymis a stylo *indym.* divergentibus. *Hort. cliff.* 329. *Hort. xps.* 185. *Roy. lugdb.* 337. *Suvv monsp.* 72. 282.
Thlaspi bicutatum asperum hieracifolium majus. Bauh. pin. 107.
Thlaspi clypeatum. Clus. hist. 2. p. 133.
 β Jon Draba alyssoides apula spicava. *Col. ecphr.* 1. p. 283. t. 285.
Habitat in Germania, Gallia, Italia. ☉
Vulgaris fructu majori: disco glabro; Rarior minori fructu: disco scabro.

LUNARIA. 714

1. LUNARIA filiculis oblongis. *rediviva,*
Lunaria foliis cordatis. Hort. cliff. 333. *Fl. suec.* 529. *Roy. lugdb.* 332.
Viola lunaria major, filiqua oblonga. Bauh. pin. 203.
Viola latifolia. Lunaria odorata. Clus. hist. 1. p. 297.
Habitat in Europa septentrionaliore. &
2. LUNARIA filiculis subrotundis. *minna,*
Lunaria major, filiqua rotundiore. Bauh. hist. 2. p. 881.
Viola latifolia. Dod. pempt. 161. *Dalech. hist.* 805.
Habitat in Germania.
Ita affinis precedentis, ut otium dubium utrum vere distincta.

SILIQUOSA.

DENTARIA. 715

1. DENTARIA foliis ternis ternatis. *cinephyllus*
Dentaria foliis omnibus ternatis. Roy. lugdb. 340.
Dentaria triphyllus. Bauh. pin. 322. *Clus. hist.* 2. p. 121. n. 5.
Ceraria plinii. Col. ecphr. 1. p. 308. t. 307.
Habitat in Austria, Italia. &
2. DENTARIA foliis inferioribus pinnatis, summis simplicibus. *Hort. cliff.* 335. *Fl. suec.* 565. *Roy. lugdb.* 340. *Hall. helv.* 557.
Dentaria heptaphyllos baccifera. Bauh. pin. 322.
Dentaria heptaphyllos baccifera. Clus. hist. 2. p. 121.
 β *Dentaria baccifera, foliis ptarmicæ. Bauh. pin.* 322.
Habitat in Europa australi ad radices montium umbrosar. &
3. DEN-

654 TETRADYNAMIA SILIQUOSA.

- pentaphyllos* 3. DENTARIA foliis summis digitatis.
 α. Dentaria foliis septenis, superioribus quinatis *Hall. belv.* 556.
 Dentaria heptaphyllos. *Baub. pin.* 322.
 Dentaria 8. heptaphyllos. *Clus. hist.* 2. p. 123.
 β. Dentaria pentaphyllos, foliis mollibus. *Garid. prov.* 152
 t. 29.
 Dentaria pentaphyllos. *Baub. pin.* 322.
 γ. Dentaria pentaphyllos, foliis asperis. *Tournef. inst.* 225
 Dentaria 6. pentaphyllos 1. *Clus. hist.* 2. p. 122.
Habitat in Alpibus Helveticis, Allobrogicis, aliisque.

CARDAMINE.

* *Foliis simplicibus.*

- bellidifolia* 1. CARDAMINE foliis simplicibus ovatis integerrimis
 petiolis longis. *Fl. lapp.* 206. t. 9. f. 2. *Fl. succ.* 564
Hall. belv. 566.
 Nasturtium alpinum, bellidis folio, minus. *Baub. pin.* 105.
 Plantula cardamines æmula. *Clus. hist.* 2. p. 129.
Habitat in Alpibus Lapponiæ, Helvetiæ, Britannicæ.
- asartifolia.* 2. CARDAMINE foliis simplicibus subcordatis.
 Nasturtium montanum, asari folio. *Bocc. sic.* 5. t. 3.
Herm. par. 203. t. 203. *Raj. hist.* 846.
 Nasturtium alpinum palustre rotundifolium. radice
 pente. *Morif. hist.* 2. p. 224.
 Nasturtium alpinum, bellidis folio, majus. *Bamb. pin.* 105. *prodr.* 46?
Habitat in Alpibus Italicis.
- nudicaulis.* 3. CARDAMINE foliis simplicibus lanceolatis sinuato
 dentatis, caulibus nudis.
Habitat in Sibiria. D. Gmelin.
- petræa.* 4. CARDAMINE foliis simplicibus oblongis dentatis.
 Cardamine petræa cambrica, nasturtii facie. *Dill. el.* 70. t. 61. f. 71.
 Nasturtium petræum. *Pluk. alm.* 261. t. 201. f. 3.
Pet. herb. 50. f. 3.
Habitat in Angliæ, Arvonæ, Merviniæ, Sueciæ rupibus excelsis.

* *Foliis ternatis.*

- trifolia.* 5. CARDAMINE foliis ternatis obtusis, caule subnudo.

Ci

TETRADYNAMIA SILIQUOSA. 655

Cardamine foliis ternatis. Hall. belv. 559. Fl. suec. 563.

Cardamine foliis ternatis crenatis, caule simplici. Fl. lapp. 257. Hort. cliff. 336. Roy. lugd. 346.

Cardamine alpina trifolia. Clus. hist. 2. p. 127.

Nasturtium alpinum trifolium. Baub. pin. 104.

Habitat in Alpibus Helveticis, Lapponicis.

6. CARDAMINE foliis ternatis acuminatis, caule ra- mosissimo.

*Nasturtium africanum, foliis ternis, facie christophoria-
næ. Herm. par. 202. t. 202.*

*Nasturtium africanum, floribus albis spicatis, foliis ter-
nis christophorianæ facie. Pluk. alm. 152. t. 101. f.*

5. Raj. suppl. 414.

Habitat in Africa.

* *Foliis pinnatis.*

7. CARDAMINE foliis pinnatis: foliolis quinis incis. (*chelidonia.*)

Cardamine glabra, chelidonii folio. T. Barr. sc. 156.

*Nasturtium pyrenæorum aquaticum latifolium, purpu-
rascente flore. Herm. par. 203. t. 204.*

Habitat in Sibiria, Italia.

8. CARDAMINE foliis pinnatis, floribus apetalis. (*impatiens.*)

*Cardamine foliis pinnatis: pinnis laciniatis. Hort. cliff.
336. Fl. suec. 561. Roy. lugd. 345. Hall. belv. 557.*

Sauv. monsp. 255.

*Sisymbrii cardamines species quædam insipida. Baub.
hist. 2. p. 886.*

Habitat in Europæ nemoribus ad radices montium. ☉

9. CARDAMINE foliis pinnatis: foliolis palmatis æqua- libus petiolatis. (*græca.*)

*Nasturtium montanum nanum, rotundo thalictri folio,
cyrenæum. Bocc. mus. 2. p. 171. t. 166.*

*Sio minimo prosperi alpini affinis, foliolis latis. Bocc.
sc. 84. t. 44. f. 2.*

Habitat in Sicilia, Corsica, insulis Græciæ. ☉

10. CARDAMINE foliis pinnatis, floribus tetrandris. (*hirsuta.*)

*Hort. cliff. 336. Fl. suec. 562. Roy. lugd. 345.
virg. 77.*

Nasturtium aquaticum minus. Baub. pin. 104.

Sisymbrium aquaticum alterum. Cam. epit. 270.

Habitat in Europæ arvis, hortis, arvis.

11. CAR-

656 TETRADYNAMIA SILIQUOSA.

- pratense.* 11. **CARDAMINE** foliis pinnatis: foliolis radicalibus subrotundis: caulinis lanceolatis.
Cardamine foliis pinnatis: foliolis subrotundis integris.
Fl. suec. 559. *Dalib. paris.* 198.
Cardamine foliis pinnatis, caule erecto. *Fl. lapp.* 258.
Hort. cliff. 336. *Roy. lugdb.* 345.
Nasturtium pratense, magno flore. *Bauh. pin.* 104.
Flos cuculi. Dod. pempt. 592.
 β. *Nasturtium* pratense, folio rotundiore, flore majore.
Bauh. pin. 104.
Habitat in Europæ pascuis aquosis. &
- emars.* 12. **CARDAMINE** foliis pinnatis, axillis stoloniferis. *L. W. goth.* 76. *
Cardamine foliis pinnatis, foliolis subrotundis angulosis.
Hall. belv. 558. *Fl. suec.* 560. *Dalib. paris.* 199.
Nasturtium aquaticum majus & amarum. *Bauh. pin.* 104.
Habitat in Europæ septentrionalioris nemoribus. &
- resedifolia* 13. **CARDAMINE** foliis pinnatis: foliolis decurrentibus confluentibus.
Nasturtium alpinum minus, resedæ folio. *Bauh. pin.* 104. *prodr.* 45. t. 45. *Bocc. mus.* 2. p. 41. t. 46.
Habitat in Alpibus Helveticis, Pyrenæis.
- Virginica.* 14. **CARDAMINE** foliis pinnatis: foliolis lanceolatis basi unidentatis,
Alyssum foliis radicalibus pinnatis in orbem positis; caulinis lanceolatis, filiculis compressis. *Gron. virg.* 170.
Nasturtium, burse pastoris folio, virginianum, flore albo, siliqua compressa. *Pluk. alm.* 251. t. 101. f. 4.
Habitat in Virginia.
Folia radicalia in orbem digesta, pinnata: foliolis numerosis, fere imbricatis, sublanceolatis, a latere postico denticulo uno brevi juxta basin. Caulis minus foliosus: foliis raris, sæpius linearibus integris. Habitus Burse pastoris.
- * *Foliis supradecompositis.*
- Lunaria.* 15. **CARDAMINE** foliis supradecompositis, siliquis unilocularibus pendulis.
Habitat in Ægypto. D. B. Jussieu.
Folia supradecomposita. Siliquæ late, membranaceo-compressæ, fere Lunariæ. Semina orbiculata, emarginata.

SISYM.

SISYMBRIUM. 717

* *Siliquis declinatis brevibus.*

1. SISYMBRIUM siliquis declinatis. foliis pinnatis: fo-*Nasturtium*
liolis subcordatis. *Hort. cliff.* 336. *Fl. succ.* 552. *It.* *¶ m.*
Scan. 278. * *Mat. med.* 331. *Roy. lugdb.* 340. *Gron.*
virg. 170. *Dalib. parif.* 203.
Sisymbria Cardamine. *Fuchf. hist.* 723.
Nasturtium aquaticum supinum. *Baub. pin.* 104.
Habitat in Europa & America septentrionali ad fontes.
2. SISYMBRIUM siliquis declinatis, foliis pinnatis: fo-*sylvestre*.
liolis lanceolatis serratis. *Hort. cliff.* 336. *Roy. lugdb.*
341. *Dalib. parif.* 204. *Sauv. monsp.* 228. *Bæhm.*
lapp. 118. *Gort. gelr.* 399.
Eruca palustris, nasturtii folio, siliqua oblonga. *Baub.*
pin. 98.
Eruca quibusdam sylvestris repens, flosculo luteo. *Baub.*
hist. 2. p. 866.
Eruca sylvestris minor, luteo parvoque flore. *Baub. pin.*
98.
Eruca sylvestris *Fuchf. hist.* 263.
Habitat in Helvetiæ, Germaniæ, Galliæ ruderatis. &
3. SISYMBRIUM siliquis, declinatis oblongo ovatis, *ambibium*.
foliis pinnatifidis serratis.
- a. *Sisymbrium* foliis pinuatifidis serratis. *Fl. succ.* 551. palustre.
Sisymbrium foliis infimis capillaceis, summis pinnatifidis.
Hort. cliff. 337. *Roy. lugdb.* 341. *Dalib. parif.* 203.
Raphanus aquaticus, foliis in profundas lacinias divisus.
Baub. pin. 97. *prodr.* 38. t. 38. *Fl. lapp.* 262.
- β. *Sisymbrium* foliis simplicibus dentatis serratis. *Hort. aquaticum.*
cliff. 336. *Fl. succ.* 350. *Roy. lugdb.* 341. *Hall. belv.*
548.
Raphanus aquaticus, rapistri folio. *Baub. pin.* 97.
Raphanus aquaticus alter. *Baub. prodr.* 38. t. 38.
γ. *Sisymbrium* aquaticum, foliis variis. *Vaill. parif.* 185. t. *restre.*
Sisymbrium foliis imis integris ovatis serratis, superio-
ribus pinnatis. *Hall. belv.* 548.
Habitat in Europæ septentrionalioris aquis.
- * *Siliquis sessilibus axillaribus.*
4. SISYMBRIUM siliquis axillaribus sessilibus solita-*supinum*.
riis, foliis dentato-*annuatis*. *Hort. ups.* 196.
Erysimum caule decumbente, siliqua laterali sessili.
Roy. lugdb. 343. *Dalib. parif.* 201.

T t

Eruca

658 TETRADYNAMIA SILIQUOSA.

Eruca supina alba, siliqua singulæ foliorum alis erumpente. Isnard. æst. 1724. p. 295. t. 18.

Habitat Parisiis ad agrorum margines, inque Hispania.

polyceratium 5. **SISYMBRIUM** siliquis axillaribus sessilibus subulatis aggregatis, foliis repando-dentatis. *Hort. ups. 193. Samu. monsp. 283.*

Erysimum siliquis in alis foliorum sessilibus. Hort. cliff. 337.

Erysimum polyceratium f. corniculatum Banb. pin. 101.

Iris altera Dalech. hist. 653.

Habitat in Helvetiæ, Italiæ ruderalis. ☉

montale, 6. **SISYMBRIUM** foliis lyratis denticulatis, siliquis sessilibus.

Sisymbrium foliis sinuato-pinnatis. Gnett. stamp. 150. Dalib. parif. 205.

Eruca sylvestris minor lutea, burse pastoris folio. Banb. pin. 98. prodr. 39. Barr. ic. 1016.

Eruca ficula, burse pastoris folio. Bocca sic. 18.

Habitat Gallia, Sicilia.

** Caule nudo.*

vimineum. 7. **SISYMBRIUM** foliis lanceolatis sinuato-ferratis, caule subnudo erectiusculo. *Hort. cliff. 337. Roy. Lugdb. 341. Dalib. parif. 204.*

Eruca, minimo flore, monspeliensis. Banb. hist. 2. p. 862.

Bellis viminea, ibiridis folio, luteo flore. Barr. ic. 131.

Habitat Parisiis & ad Tyberis margines, inque Moravia.

arenosum, 8. **SISYMBRIUM** foliis lyrato-ferratis scabris radicalibus, caule subnudo ramoso.

Eruca cærulea in arenosis proveniens. Banb. pin. 99. prodr. 40. fig. mal.

Eruca sylvestris major minorque, foliis subasperis in orbem sparsis. Læf. pruss. 68. t. 13. bon.

Habitat in Germania, Helvetia.

monensis. 9. **SISYMBRIUM** foliis pinnatifidis integerrimis, caulibus subnudis simplicibus procumbentibus.

Eruca monensis laciniata, flore luteo majore. Dill. elsh. 135. t. 111. f. 135.

Habitat in Mona, insula Angliæ.

** Bipin-*

* *Foliis bipinnatis.*

10. **SISYMBRIUM** siliquis scabris, foliis pinnatifidis: *asperum*.
pinnis linearilanceolatis subdentatis, corollis calyce
longioribus. *Sauv. monsp.* 283. †
Sisymbrium foliis pinnatifidis subdentatis angustissimis.
Sauv. monsp. 255.
Sinapi parvum, siliqua aspera. *Baub. pin.* 99. *prodr.* 41.
Sinapi montepellulanum, siliqua aspera hirsuta. *Baub. hist.*
2. p. 857.
Habitat Monspelii.
11. **SISYMBRIUM** petalis calyce minoribus, foliis de *Sophia*.
composito-pinnatis. *Fl. suec.* 553. *Dalib. parif.* 204.
Mat. med. 332.
Sisymbrium corolla calyce minore, foliis multifidis line-
aribus. *Fl. lapp.* 261. *Hall. belv.* 337. *Key. lugdb.* 341.
Deiscurea. *Guss. stamp.* 2. p. 164.
Nasturtium sylvestre tenuissime divisum. *Baub. pin.* 105.
Scripium Abinthium. *Fuchs. hist.* 2.
Habitat in Europæ maceris, muris, tectis. ☉
12. **SISYMBRIUM** foliis pinnatis: foliolis lanceolatis *tanacetifo-*
pinnatifidis. *Key. lugdb.* 342. *Hum.*
Eruca fruticosa, foliis tanaceti, indica. *Zan. hist.* 86. t.
33.
Eruca tanaceti foliis. *Morif. hist.* 2. p. 231. f. 2. t.
6. f. 19.
Habitat in Sabaudia?
13. **SISYMBRIUM** foliis pinnato-hastatis flaccidis: foli- *altissimum*.
olis sublinearibus integerrimis pedunculis laxis. *Hort.*
upf. 193. * *Sauv. monsp.* 229.
Erysimum siliquis laxis, foliis hastato-pinnatis. *Hort. cliff.*
338. *Sauv. monsp.* 284.
Rapistrum italicum, siliquis longissimis. *Baub. pin.* 95.
Rapistrum montanum f. sylvestre, irionis folio. *Col.*
ecphr. 1. p. 266. t. 265.
Erysimum montepellulanum, sinapios folio. *Walsh. hort.*
55 t. 22.
Habitat in Italia, Gallia, Sibiria. ☉
14. **SISYMBRIUM** foliis pinnato-hastatis dentatis, sili- *iride*.
quis erectis.
Erysimum foliis lyrato-pinnatis: extimo hastato. *Fl. suec.*
556. *Dalib. parif.* 201. *Sauv. monsp.* 284.
Erysimum latifolium majus glabrum. *Baub. pin.* 103.
Irioneis apulus, erucæ folio. *Col. ecphr.* 1. p. 264. t. 265.
T t 2 *Habi-*

660 TETRADYNAMIA SILIQUOSA.

Habitat in Europæ cultis.

Differt ab S. altissimo Foliis summis non in pinnas lineares longissimas sectis; Foliis omnibus magis dentatis, Caule brevior, minus glabro; Siliquis magis erectis nec laxis subpendulis.

* *Foliis lanceolatis integris.*

- strictissimum* 15. **SISYMBRIUM** foliis lanceolatis dentato-ferratis caulinis. *Hort. cliff.* 337. *Hort. nps.* 192. *Roy. lugdb.* 341.

Draba lutea, siliquis strictissimis. Baub. pin. 110.

Arabis quibusdam dicta planta. Cam. epit. 342.

Habitat in Helvetiæ montibus. &

- integrifolium* 16. **SISYMBRIUM** foliis linearibus integerrimis, pedunculis glutinoso-hispidis.

Hesperis caule ramolissimo, foliis linearibus, pedunculis hispidis. Gmel.

Habitat in Sibiria.

Caulis digiti longitudine. Folia linearia, glabra, obtusiuscula. Racemi terminales, pubescentes, visci. Flores majusculi, albi. Siliquæ lineares.

ERYSIMUM.

- officinale.* 1. **ERYSIMUM** siliquis spicæ adpressis. *Hort. cliff.* 337. *Fl. suec.* 554. *Mat. med.* 333. *Roy. lugdb.* 342. *Dalib. parif.* 200.

Erysimum vulgare. Baub. pin. 100.

Verbena mas. Fuchs. hist. 592.

Habitat in Europæ ruderatis. ☉

- Barbarea.* 2. **ERYSIMUM** foliis lyratis extimo subrotundo. *Fl. suec.* 557. *Mat. med.*

Erysimum foliis basi pinnato-dentatis apice subrotundis.

Fl. lapp. 264. *Hort. cliff.* 338. *Roy. lugdb.* 342.

Dalib. parif. 202.

Eruca lutea latifolia f. Barbarea. Baub. pin. 98.

β. *Sisymbrium, erucæ folio glabro, minus & procerius. Tournef. inst.* 226.

γ. *Sisymbrium orientale, barbareae facie, folio rotundo. Tournef. cor.* 16.

Habitat in Europa.

- Alliaria.* 3. **ERYSIMUM** foliis cordatis. *Hort. cliff.* 338. *Fl. suec.* 558. *Mat. med.* 334. *Roy. lugdb.* 342. *Dalib. parif.* 201.

Alliaria.

TETRADYNAMIA SILIQUOSA. 661

Alliaria. *Bauh. pin.* 110. *Fuchs. hist.* 104. *Cam. epit.* 589.

Habitat in Europæ sepibus, cultis, umbrosis. ♂

4. ERYSIMUM foliis integris lanceolatis. *Fl. lapp.* 263. *Cheiranthoid.* *Hort. cliff.* 337. *Fl. succ.* 555. *Roy. lugdb.* 342. *Dalib. parif.* 202.

Turritis foliis integris lanceolatis. *Guet. stamp.* 2. p. 165.

Myagrum siliqua longa. *Bauh. pin.* 109.

Camelina, Myagrum alterum thlaspi effigie. *Lob. ic.* 225.

Habitat ubique in Europæ arvis. ☉

Habitus *simillimus Cheiranthæ Erysimoidæ præter flores minores & siliquas magis patulas. Stigmata in hac parva rix divisa, in sequenti quasi ex nodis duobus divaricatis confecta*

CHEIRANTHUS. 719

1. CHEIRANTHUS foliis lanceolatis dentatis, caule re- *erysimoides.*

cto, siliquis tetragonis.

Erysimum foliis integris lanceolatis margine repando-dentatis. *Hort. ups.* 192.

Leucojum luteum sylvestre angustifolium. *Bauh. pin.* 202.

Leucojum sylvestre. *Clus. hist.* 1. p. 299. *Bauh. hist.* 2. p. 873.

Habitat in Hungaria, Gallia. ♂

2. CHEIRANTHUS foliis lanceolatis acutis glabris. *Cheiri.*

Hort. cliff. 334. *Hort. ups.* 187. *Mat. med.* 335.

Roy. lugdb. 337. *Guet. stamp.* 2. p. 156. *Dalib. parif.* 197. *Gort. gelr.* 382.

Leucojum luteum vulgare. *Bauh. pin.* 202.

Leucojum luteum. *Dod. pempt.* 160.

- β. Leucojum luteum, magno flore. *Bauh. pin.* 202.

- γ. Leucojum luteum, serrato folio, flore grandiore. *Bauh.*

pin. 202. *prodr.* 102.

- δ. Leucojum luteum, pleno flore, majus. *Bauh. pin.* 202.

- ε. Leucojum, pleno flore, minus. *Bauh. pin.* 202.

Habitat in Angliæ, Helvetiæ, Galliæ, Hispaniæ muris. rectis. ♂. ♀. h.

3. CHEIRANTHUS foliis lanceolatis subdentatis retu- *chius.*

lis, siliquis apice subulatis. *Hort. ups.* 187.

Hesperis caule ramosissimo, foliis lanceolatis dentatis

Hort. cliff. 335. *n.* 3. *Roy. lugdb.* 337. *n.* 3.

T 13

Hesperis

662 TETRADYNAMIA SILIQUOSA.

Hesperis filiquis hirsutis, flore parvo rubello. *Dill. elsh.* 180. t. 147. j. 178.

Leucojum thlaspece facie, flore albo. *Herm. par.* 192. s. 193.

Habitat in Chio. ☉

incanum.

4. CHEIRANTHUS foliis lanceolatis integerrimis obtusis incanis, filiquis apice truncatis compressis, caule suffruticoso. *Hort. ups.* 187. *Hort. cliff.* 334.

Leucojum, incano folio, hortense. *Baub. pin.* 200. n. 1. 2. 4. 5. 6. 7.

Viola alba & purpurea. *Lob. ic.* 329.

Habitat in Hispaniæ maritimis. 2. 5.

annuum.

5. CHEIRANTHUS foliis lanceolatis subdentatis obtusis incanis, filiquis cylindricis apice acutis, caule herbaceo.

Leucojum incanum minus. *Baub. pin.* 200.

Habitat in Europæ australis maritimis. ☉

Fortè sola varietas antecedentis?

fruticulosus, 6. CHEIRANTHUS foliis linearibus subsinuatis, floribus sessilibus petalis undatis, caule suffruticoso. *Læst.*

Leucojum maritimum angustifolium. *Baub. pin.* 201.

Leucojum minus, breviori folio, obsoleto flore. *Barr. ic.* 99. n. 2.

Leucojum minus, lavendulæ folio, obsoleto flore. *Bucc. mus.* 148. s. 111.

Habitat in Hispania, Italia, Monspelii. 5

trilobus.

7. CHEIRANTHUS foliis dentatis obtusis, calycibus lævibus, filiquis nodosis mucronatis.

Leucojum martimum minimum hispanicum vernum, foliis erucæ. *Tournef. inst.* 221?

Leucojum maritimum minimum. *Baub. pin. Burf.* XI. 28.

Habitat in Hispania & prope insulas Stœrachadum. ☉

Caules ramosi, patuli, spithamei, incani. Folia lanceolata, obtusa, unico vel duobus utrinque dentibus profundioribus. Calyces non pilosi. Corolla purpurea, majuscula. Siliquæ lineares, teretiusculæ, suturis carinatae, lateribus torulis nodosæ, apice longo subulato acuminatae.

lacernus.

8. CHEIRANTHUS foliis lacero-dentatis acuminatis, calycibus pilosis, filiquis nodosis mucronatis.

Leu-

TETRADYNAMIA SILIQUOSA. 663

Leucojum lusitanicum purpureum, foliis eleganter dentatis. *Herm. parad.* 193. t. 193.

Habitat in Lusitania. ☉

9. CHEIRANTHUS siliquarum apicibus tridentatis. *tricuspidatus*

Hort. cliff. 335. *Roy. lugdb.* 338.

Leucojum marimum. Cam. hort. 87. t. 24. *Meris. hist.*

2. p. 242. f. 3. t. 8. f. 13.

Habitat in Europæ australis maritimis? ☉

HESPERIS. 720

1. HESPERIS caule hispido ramoso patente. *Hors. aff.* 187. *tristis.*

Hesperis caule hispido procumbente. Hort. cliff. 335.

Roy. lugdb. 338.

Hesperis montana pallida odoratissima. Baub. pin. 202.

Hesperis pannonica. Com. hort. t. 18.

Habitat in Hungaria. ♂

2. HESPERIS caule simplici erecto, foliis ovato-lanceo-*matronalis.*

latis denticulatis, petalis mucrone emarginatis. *Hort.*

cliff. 188. *Roy. lugdb.* 338. *Dalib. parif.* 197.

Hesperis hortensis. Baub. pin. 202.

Viola matronalis. Dod. pempt. 161.

Habitat in Italia. ♂

3. HESPERIS caule simplici, foliis lanceolatis dentato-*sibirica.*

ternatis, petalis obtusissimis integris.

Habitat in Sibiria. ♀

Habitus omnibus partibus H. matronalis, ut facile pro

varietate haberem; Differt vero Caule duplo altiore;

Corollis purpureis nec albidis; Ungvibus petalorum

calyce duplo longioribus; Petalorum bracteis obtusis

vix ac ne vix emarginatis, cum præcedentis emargi-

nata ungue intermedio parvo prominulo. Folia longi-

ora, angustiora, magis serrata. Ceterum Sibiria pro-

ducit varias plantas majores: Ononidem, Polemonium,

Serratulam.

4. HESPERIS caule ramosissimo diffuso, foliis lanceola-*africana.*

tis serratis scabris, siliquis sessilibus.

Hesperis caule ramosissimo, foliis lanceolatis sæpius den-

tatis. Hort. cliff. 335. *Roy. lugdb.* 338.

Hesperis africana, hieracii folio hirsuto, flore minimo

purpurascente. Niss. act.

Hesperis sylvestris, parvo flore. Baub. pin. 202. *prodr.*

103. *Burf.*

Habitat in Africa. ☉

T t 4

5. HESPE-

664 TETRADYNAMIA SILIQUOSA.

- provincialis.* 5. HESPERIS caule ramoso diffuso, filiquis teretibus.
Hort. ups. 188.
 Cheiranthus foliis lanceolato-linearibus acutis glabris utrinque dentatis. *Hort. cliff.* 331. *Roy. lugdb.* 337.
 Leucojum minus, lavendulæ folio, obsoleto flore, angustifolium. *Bocc. mus.* 2. p. 148. t. 111. *Barr. ic* 803.
Habitat in Galloprovincia. 2.
- dentata.* 6. HESPERIS foliis dentato-pinnatifidis, caule lævi.
 Hesperis foliis multifidis. *Roy. lugdb.* 338.
 Hesperis flore albo minimo, filiqua longa, folio profunde dentato. *Dill. elsh.* 179. t. 148. f. 177.
Habitat in Sicilia. ☉
- verna.* 7. HESPERIS caule erecto ramoso, foliis cordatis amplexicaulibus serratis villosis.
 Turrilis annua verna, flore purpurascens. *Tournef. inst.* 224.
 Leucojum minus rotundifolium, flore purpureo. *Barr. ic.* 876.
 Leucojum maritimum latifolium. *Morif. hist.* 2. p. 241. f. 3. t. 8. f. 5.
 Leucojum marinum alterum latifolium purpureo-violeum. *Lob. ic.* 333.
 Leucojum maritimum latifolium. *Bauh. pin.* 201.
 Rapistrum floribus leucoji marini. *Bauh. pin.* 95. *prodr.* 37. *Bursf. IV:* 47.
Habitat in Galloprovinciæ maritimis.

721
alpina.

ARABIS.

1. ARABIS foliis amplexicaulibus dentatis. *Hort. cliff.* 335.
 α. Arabis foliis amplexicaulibus dentatis, caule erecto simplici. *Roy. lugdb.* 338. *Hall. helv.* 561.
 Arabis caule simplici, foliis ovatis utrinque dentatis. *Fl. lapp.* 257.
 Draba alba siliquosa repens. *Bauh. pin.* 109.
 Draba 3, succulento folio. *Clusf. hist.* 2. p. 125.
 β. Draba foliis amplexicaulibus dentatis, caule diffuso ramoso. *Roy. lugdb.* 339.
 Draba alba siliquosa. *Bauh. pin.* 109.
 Draba 2. *Clusf. hist.* 2. p. 24.
Habitat in Alpibus Helveticis, Lapponicis.
 Mire varians ex loco planta. α. brevis erecta alpina; β altera, inferne procumbens magna; hinc facie diversissima.

2. ARA-

TETRADYNAMIA SILIQUOSA. 669

2. ARABIS caule nudo. *Amœn. acad.* 358. t. 4. f. 20. * *grandiflora*.
Habitat in Sibiria. 2

3. ARABIS foliis petiolatis lanceolatis integerrimis. *Vir. thaliana*.
cliff. 64. *Fl. succ.* 567. *Roy. lugdb.* 339. *Dalib. parif.*
 200.

Turritis foliis lanceolatis integris petiolatis, ad exor-
 tum ramorum solitariis. *Hort. cliff.* 339.

Bursæ pastoris similis siliquosa major *Baub. pin.*
 108.

Pilosella siliquosa minor. *Thal. barc.* 84. t. 7. f. D.

Habitat in Europæ septentrionalioris sabulosis. ☉

4. ARABIS foliis glabris radicalibus lyratis: caulinis li- *lyrata*.
nearibus.

Cheiranthus caule filiformi lævi, foliis lanceolatis: infi-
 mis incis. *Gron. virg.* 76.

Habitat in Canada. D. *Kalm.* ☉

Statura & habitus A. thaliana, præter quod folia radi-
calia lyrata, glabra, & flores majores.

5. ARABIS foliis caulinis lanceolatis dentatis glabris. *canadensis*.
Eruca virginiana, bellidis majoris folio. *Pluk. alm.* 136.
 t. 86. f. 8.

Turritis foliis lanceolatis dentatis: radicalibus maximis,
 siliquis compressis sulcatis. *Gron. virg.* 77?

Planta *erecta, pedalis bipedalis, lævis. Folia lato-lanceo-*
lata, 4 vel 5 denticulis crassis remotis serrata utrinque.
Racemus terminalis nudus, & ex summis alis duæ
vel tres laterales racemi.

6. ARABIS foliis amplexicaulibus, siliquis ancipitibus li- *pendula*.
nearibus, calycibus subpilosis. *Hort. upf.* 191. *

Turritis latifolia hirsuta, siliquis pendulis. *Amm. ruth.*
 58.

Habitat in Sibiria. ☉

7. ARABIS foliis amplexicaulibus, siliquis decurvis pla- *Turrita*.
nis linearibus, calycibus subrugosis. *Hort. upf.* 192.

Turritis caule simplici, foliis lanceolatis subdentatis gla-
 bris. *Sauv. monsp.* 73.

Turritis perraro ramifera, foliis lanceolatis obscure den-
 tatis. *Guett. stamp.* 2. p. 162. *Dalib. parif.* 202.

Leucojum helperidis folio. *Tournef. inst.* 221.

Brassica sylvestris, albido flore, nutante siliqua. *Bocc.*
mus. 2. p. 81. t. 72.

T t 5

Tur-

666 TETRADYNAMIA SILIQUOSA.

Turrita major, platea. *Clus. hist.* 2. p. 126.

Habitat in Helvetia, Hungaria, Gallia, Sicilia. ☉

722

TURRITIS.

glabra

1. **TURRITIS** foliis radicalibus dentatis hispidis, caulinis integerrimis amplexicaulibus glabris. *Hort. cliff.* 339. *Fl. suec.* 544. *Roy. lugdb.* 339. *Hall. belv.* 500. *Dalib. parif.* 203

Brassica sylvestris, foliis circa radicem cichoraceis. *Bauh. pin.* 112.

Habitat in Europæ pascuis siccis apricis.

hirsuta.

2. **TURRITIS** foliis omnibus hispidis, caulinis amplexicaulibus. *Hort. cliff.* 339. *Fl. suec.* 545. *Roy. lugdb.* 339. *Dalib. parif.* 62.

Erysimo similis *hirsuta*, non laciniata, alba. *Bauh. pin.* 101. *prodr.* 42. t. 42.

Habitat in Sueciæ, Germaniæ, Angliæ pascuis sylvaticis,

723

BRASSICA.

orientalis.

1. **BRASSICA** foliis cordatis amplexicaulibus glabris: radicalibus scabris integerrimis, filiquis tetragonis. *Hort. ups.* 190.

Brassica orientalis perfoliata, flore albo, filiqua quadrangula. *Tournef. cor.* 16.

Habitat in Oriente. ☉

campestris.

2. **BRASSICA** radice cauleque tenui, foliis caulinis uniformibus cordatis sessilibus. *Hort. cliff.* 339. *Fl. suec.* 546. *Roy. lugdb.* 344.

Brassica campestris perfoliata, flore luteo. *Laf. pruss.* 29. *Fl. lipp.* 265.

Habitat in agris non argillofis Europæ. ☉

Napus.

- 3 **BRASSICA** radice caulescente fusiformi. *Hort. cliff.* 339. *Fl. suec.* 547. *Mat. med.* 328. *Roy. lugdb.* 344. *Dalib. parif.* 199.

Napus sylvestris. *Bauh. pin.* 95.

Napus. *Dod. pempt.* 674.

β. *Napus sativa.* *Bauh. pin.* 95.

Habitat in arenosis maritimis Gotlandiæ, Belgii, Angliæ. ♂

Rapa.

4. **BRASSICA** radice caulescente orbiculari depresso carosa. *Hort. cliff.* 339. *Hort. ups.* 190. *Mat. med.* 329. *Roy. lugdb.* 344. Ra-

Rapa sylvestris. *Raj. hist.* 800.

Rapa sativa rotunda. *Baub. pin.* 84.

Rapum. *Cam. epit.* 218.

β. Rapa sativa oblonga f. femina. *Bauh. pin.* 89.

Habitat in arvis Angliæ, Belgii. ♂

γ. BRASSICA radice caulescente tereti carnosa, *Hort. oleracea, cliff.* 338. *Hort. upf.* 189. *Mat. med.* 327. *Roy. lugdb.* 342.

Brassica maritima arborea f. procerior ramosa. *Morif. sylvestris, hist.* 2. p. 208. *Raj. hist.* 796.

β. Brassica alba vel viridis. *Bauh. pin.* 111.

viridis.

γ. Brassica capitata rubra. *Bauh. pin.* 111.

rubra.

δ. Brassica capitata alba. *Bauh. pin.* 111.

capitata.

ε. Brassica alba crispa. *Bauh. pin.* 111.

tabauda.

Brassica undulata. *Reencalm. spec.* 113.

γ. Brassica laciniata rubra. *Bauh. hist.* 2. p. 832.

laciniata.

η. Brassica angusto apii folio. *Bauh. pin.* 112.

felenisia.

θ. Brassica fimbriata. *Baub. pin.* 112.

fabellica.

Brassica fabellica. *Reencalm. spec.* 134. t. 133.

ι. Brassica cauliflora. *Bauh. pin.* 111. *Reencalm. spec.* 131 botrytis.

t. 133. *Cam. epit.* 253.

κ. Brassica radice napiformi. *Tournef. inst.* 219.

Napobrassica.

Napo-Brassica. *Baub. pin.* 111. *prodr.* 54.

gongylodes

λ. Brassica gongylodes. *Bauh. pin.* 111.

Brassica cauloRapa. *Bauh. hist.* 2. p. 829.

Caulo-Rapum. *Cam. epit.* 251.

Habitat in maritimis Angliæ. ♂

6. BRASSICA foliis lanceolato-ovatis glabris indivisis *violacea, dentatis. Hort. upf.* 191.

Habitat in China. ♂

* *Eruca filiquis stylo ensiformi.*

7. BRASSICA foliis dentato pinnatifidis, caule hispido, *Erucastrum, filiquis lævibus.*

Sisymbrium foliis pinnato-dentatis. *Hort. cliff.* 337.

Roy. lugdb. 341. *Dalib. parit.* 205. sub Sisymbrio.

Eruca sylvestris major lutea, caule aspero. *Bauh. pin.*

98.

Eruca sativa. *Fuchf. hist.* 262.

Habitat in Europæ australioris ruderalis. ☉

8. BRASSICA foliis lyratis, caule hirsuto, foliis glabris. *Eruca, Hort. upf.* 190. *Mat. med.* 330.

Sisymbrium foliis pinnatifidis: laciniis exterioribus majoribus, caule hirsuto. *Hort. cliff.* 337.

Eru-

668 TETRADYNAMIA SILIQUOSA.

Eruca latifolia alba. Baub. pin. 98.
Sinapi alterum genus. Fuchf. hist. 539.
Habitat in Helvetia. ☉

vesicaria.

9. BRASSICA foliis dentato-pinnatifidis, siliquis hispida
 tectis calyce tumido.
*Brassica foliis lyratis, caule hirsuto, siliquis retrorsum
 hispidis. Hort. nps. 191.*
*Eruca chalepensis, caulibus & siliquis hirsutis. Morj
 hist. 2. p. 228.*

Habitat in Hispania. Læsing. ☉

*Radix fusiformis. Folia lanceolata, pinnatifido-dentata
 supra glabra. Caulis pilosus, ramosus. Racemi erecti
 terminales. Corollæ flavæ venis saturatioribus. Ca
 lyx floris cylindricus, in fructu vero ventricosus infla
 tus persistens evadit. Siliquæ venter calyce non longior
 pistillum remanens lanceolatum. Differt itaque ab eruci
 calyce, sub fructu, inflato vesicario, foliis non lyratis se
 lanceolatis; Siliquis hispidis retrorsum nec glabris
 Calycibus persistentibus ad maturitatem fructus ne
 deciduis*

724

SINAPIS.

arvensis.

1. SINAPIS siliquis multangulis toroso-turgidis rostro
 longioribus. *Hort. cliff. 338. Fl. succ. 548. Roy
 lugdb. 343. Hall. belv. 554. Dalib. paris. 200.*
Rapistrum flore luteo. Baub. pin. 95.
Irion. Fuchf. hist. 257.
Habitat in agris Europæ. ☉

alba.

2. SINAPIS siliquis hispidis: rostro obliquo longissimo
Hort. cliff. 338. Hort. nps. 191. Roy. lugdb. 343
Dalib. paris. 200.
Sinapi apii folio. Baub. pin. 99.
Sinapi sativum alterum. Dod. pempt. 707.
Habitat in agris Belgii, Angliæ, Galliæ. ☉

nigra.

3. SINAPIS siliquis glabris tetragonis. *Hort. cliff. 338*
*Fl. succ. 549. Mat. med. 326. Roy. lugdb. 343. Da
 lib. paris. 199.*
Sinapi rapi folio. Baub. pin. 99.
*Habitat in aggeribus ruderatis Europæ septentrionalis
 ris. ☉*

juncea

4. SINAPIS ramis fasciculatis, foliis summis lanceolatis
 integerrimis. *Hort. nps. 191. **

Si

TETRADYNAMIA SILIQUOSA. 669

Sinapi indicum maximum, lactucæ folio. *Herm. par.*

230. t. 230.

Habitat in Asia. ☉

5. SINAPIS foliis duplicato-pinnatis: laciniis linearibus. *hispanica.*

Hort. cliff. 338. *Roy. lugdb.* 343.

Sinapi hispanicum, nasturtii folio. *Tournef. inst.* 227.

Habitat in Hispania.

RAPHANUS. 725

1. RAPHANUS siliquis teretibus torosis bilocularibus. *fusion.*

Hort. cliff. 340. *Hort. wps.* 188. *Mat. med.* 325. *Roy.*

lugdb. 344.

Raphanus minor oblongus. *Bauh. pin.* 96.

Raphanus major orbicularis f. rotundus. *Bauh. pin.* 96.

β. Raphanus niger. *Bauh. pin.* 96.

Habitat. - - - ☉. ♂

2. RAPHANUS siliquis teretibus articulatis lævibus u- *Raphani-*

nilocularibus *Hort. cliff.* 340. *Fl. suec.* 568. *Roy. lugdb.* *strum.*

344. *Hall. belv.* 555. *Dalib. parif.* 198. *Sauv. monsp.*

285.

Rapistrum flore albo, siliqua articulata. *Bauh. pin.* 95.

Rapistrum flore albo striato. *Bauh. hist.* 2. p. 851.

Habitat inter segetes Europæ.

3. RAPHANUS siliquis teretibus torulosis villosis, foliis *sibiricus.*

linearibus pinnatifidis.

Raphanus foliis pinnatis: pinnis confluentibus, siliquis

teretibus articulatis unilocularibus. *Gmel.*

Habitat in Sibiria.

Habitus Erucicæ; planta *pygmæa*. Caules maxima ex par-

te nudi. Folia radicalia et caulina pinnatifida, linea-

ria. Corollæ magnæ, flavae. Legumina teretia, pedi-

cellata, moniliformia, uti caulis villosa, rostro oblon-

gu: Figura *Cardamines alpinae insipidae*. *Col. ecphr.*

2. p. 69. hanc refert.

BUNIAS. 726

1. BUNIAS siliculis bicornibus divaricatis basi spinosis. *cornuta.*

Habitat in Sibiria. D. *Gmelin.*

Planta maxime singularis. Silicula transversim ovalis,

utrinque desinens in cornu f. spicam longissimam vali-

diffimam, ut referat silicula duo cornua in medio sili-

cule spinæ parvæ 4 undique directæ.

1. BU.

- Erucago.* 2. **BUNIAS** siliculis tetragonis: angulis bicristatis.
Bunias foliis antrorsum lineatis. *Hort. apf.* 132. *Sauv. monsp.* 285
Bunias. Hort. cliff. 340. *Roy. lugdb.* 344.
Erucago monspeliaca, siliqua quadraligula echinata.
Bamb. pin. 97. prodr. 41. t. 41.
Habitat Monspehi in agris humidiusculis. ☉
- orientalis.* 3. **BUNIAS** siliculis ovatis gibbis verrucosis.
Bunias foliis retrorsum sinuatis. *Hort. npsj.* 186.
Crambe foliis pinnato-hastatis. *Roy. lugdb.* 330.
Crambe orientalis, dentis leonis folio, crucaginis facie.
Tournef. cor. 14.
Habitat in Russia. &
- Cakile.* 4. **BUNIAS** siliculis ovatis lævibus ancipitibus.
Crambe foliis succulentis linearibus pinnato-dentatis.
Sauv. monsp. 280.
Raphanus siliquis ovatis angulatis monospermis. *Hort. cliff.* 340. *Fl. suec.* 569. *Roy. lugdb.* 344. *Gron. virg.* 170. *Sauv. monsp.* 285.
Eruca maritima italica, siliqua halix cuspidi simili. *Bamb. pin.* 99.
Cakile serapionis. Lob. ic. 223.
Habitat in Europæ, Africæ, Americæ maritimis. ☉

7 2 7
 ISATIS.

- timforia.* 1. **ISATIS** foliis radicalibus crenatis; caulinis sagittatis, siliculis oblongis. *Fl. suec.* 543. *Dalib. parij.* 205.
Isatis. Hort. cliff. 341. *Roy. lugdb.* 330.
Isatis sylvestris f. angustifolia. *Bamb. pin.* 113.
Isatis sylvestris. Fuchf. bist. 330.
 β. *Isatis sativa* f. latifolia. *Bamb. pin.* 113.
Habitat ad litora maris Baltici & Oceani Europæ. ♂
- lusitanica.* 2. **Isatis** foliis radicalibus crenatis; caulinis sagittatis, pedunculis subtoментosis.
Isatis sylvestris minor lusitanica. *Herm. lugdb.* 678.
Isatis orientalis maritima canescens. *Tournef. cor.* 14.
Bamb. cent. 1. p. 4. t. 5.
Habitat in Hispania & Oriente. ☉
Vix a præcedente sufficienter distincta, sed annua et minor.
- armena.* 3. **ISATIS** foliis integerrimis cordatis posticè obtusis, siliculis cordatis.

Isa-

TETRADYNAMIA SILIQUOSA. 671

Isatis armena, foliis brassicæ perfoliatæ, fructu cordiformi canescente. *Tournef. cor.* 14. *Buxb. cent.* 1. p. 3. t. 4. *Habitat in Armeniæ pratis siccioribus ad rivulos.*

4. ISATIS foliis omnibus dentatis.

ægyptica.

Habitat in Ægypto.

CRAMBE. 728

1. CRAMBE foliis cauleque glabris. *Fl. suec.* 570. *maritima.*
Crambe foliis cordatis crispis carnosiss. *Hort. cliff.* 340.

Roy. lugdb. 329.

Brassica maritima monospermos. *Bauh. pin.* 112.

Brassica marina sylvestris multicaulis monospermos.

Lob. ic. 245.

Habitat ad littora Oceani septentrionalis. &

2. CRAMBE foliis scabris, caule glabro.

orientalis.

Crambe foliis & foliolis alternatim pinnatifidis. *Roy.*

lugdb. 330.

Rapistrum orientale, acanthi folio. *Tournef. cor.* 14.

Habitat in Oriente.

3. CRAMBE foliis cauleque scabris. *Hort. ups.* 193. *hispanica.*

Crambe foliis subcordatis crenatis scabris. *Hort. cliff.*

340. *Roy. lugdb.* 329.

Rapistrum maximum rotundifolium monospermum.

Corn. canad. 147. t. 148. *Moriz. hist.* 2. p. 266. f.

3. t. 13. f. 1.

Habitat in Hispania. ☉

CLEOME. 729

1. CLEOME floribus gynandris, foliis simplicibus.

fruticosa.

Habitat in India. ☿

Caulis frutescens, teres, ramosus. Folia ovali oblonga, integerrima, glabra, petiolata, alterna. Racemi terminales, simplices. Calyces tetraphylli, patentes, ovato-lanceolati. Petala 4, unguibus oblongis, subrotunda, ascendentia. Germen pedicello longissimo, in cuius medio stamina 4, saepe duo inferiora basi coalita. Antheræ ovales. Siliqua oblonga.

- CLEOME floribus gynandris, foliis digitatis. *Hort. cliff. gynandra.*

341. *Hort. ups.* 193. *Fl. zeyl.* 239 * *Roy. lugdb.* 339.

Quinquefolium lupini folio. *Bauh. pin.* 326.

Cara-vecla. *Kbeed. mal.* 9. p. 43. t. 24.

Habitat in Alia, Atrica, America calidissimis. ☉

3. CLE-

672 TETRADYNAMIA SILIQUOSA.

- icosandra.* 3. CLEOME floribus icosi-tetrandris, foliis quinatis. *Fl. zeyl.* 240. *
Sinapistrum zeylanicum pentaphyllum viscosum, flore flavo. *Burm. zeyl.* 215. t. 99.
Habitat in Zeylona. ☉
- viscosa.* 4. CLEOME floribus dodecandris, foliis quinatis ternatisque. *Fl. zeyl.* 241. *
Sinapistrum zeylanicum triphyllum & pentaphyllum viscosum, flore flavo. *Mart. cent.* 25. t. 25.
Aria-veela. Rheed. mal. 9. p. 41. t. 23.
Habitat in Zeylona, Malabaria. ☉
- dodecandra.* 5. CLEOME floribus dodecandris, foliis ternatis. *Fl. zeyl.* 242. *
Sinapistrum triphyllum pumilum glabrum, flosculo purpureo, siliqua membranacea. *Burm. zeyl.* 216. t. 100. f. 1
Sinapistrum indicum triphyllum, flore carneo, non spinosum. *Sloan. jam.* 80. *hist.* 1. p. 194. t. 124. f. 1.
Habitat in Indiis. ☉
- ornithopodioides.* 6 CLEOME floribus hexandris, foliis ternatis: foliolis ovali lanceolatis.
Cleome floribus hexandris, genitalibus declinatis, siliquis teretibus torosis. *Hort. cliff.* 341. * *Hort. mps.* 194.
Roy. lugdb. 340.
Sinapistrum orientale triphyllum, ornithopodii siliquis. *T. Dill. elth.* 359. t. 266. f. 345. *Buxb. cent.* 1. p. 6. t. 9. f. 2.
Habitat circa Peram in arvis. ☉
- violacea.* 7. CLEOME floribus hexandris, foliis ternatis solitariisque: foliolis lineari-lanceolatis.
Cleome floribus hexandris, genitalibus declinatis, siliquis subulatis. *Hort. cliff.* 341. *Roy. lugdb.* 340,
Trifolium siliquosum, flore violaceo, lusitanicum. *Barr. ic.* 866.
β. Sinapistrum peregrinum alterum, flore luteo. *Barr. ic.* 865. *melior.*
Habitat in Lusitania. ☉
- monophylla.* 8. CLEOME floribus hexandris, foliis simplicibus ovato-lanceolatis. *Fl. zeyl.* 243. *
Sinapistrum zeylanicum viscosum, folio solitario, flore flavo, siliqua tenui. *Burm. zeyl.* 217. t. 100. f. 2.
Tsien-veela. Rheed. mal. 9. p. 63. t. 34.
Habitat in India. ☉

Cl

Classis XVI.

MONADELPHIA.

PENTANDRIA.

WALTHERIA.

730

1. **WALTHERIA** foliis planis serrato dentatis. *americana.*
Waltheria foliis cordato-ovatis serratis. *Hort. cliff.*
342. Roy. lugdb. 348.
Monosperm-Althea arborescens villosa, folio majore.
Isnard. act. 1721. p. 362. t. 14.
Althææ similis americana, flore luteo. *Herm. lugdb.*
*2. p. 267. **
Betonica arborescens, foliis amplioribus. *Pluk. alm. 67.*
t. 150. f. 6.
Althæa americana pumila, flore luteo spicato. *Breyn.*
cent. - t. 57
Habitat in Bahama, Barbiches, Surinamo. *h*
2. **WALTHERIA** foliis ovatis serratis undulatis. *Roy. indica.*
*lugdb. 348. n. 1. Fl. zeyl. 244. **
Melochia foliis oblongis obtusis serratis tomentosis, floribus confertis. *Hort. cliff. 343.*
Betonica arborescens maderaspatana villosa, foliis profunde venosis. *Pluk. alm. 67. t. 50. f. 5.*
Habitat in India. *h*

HERMANNIA

731

1. **HERMANNIA** foliis ovatis plicatis crenatis tomentosis. *althæaefolia.*
Hort. cliff. 343. Roy. lugav. 347.
Hermannia capensis, althææ folio. *Pet. gaz. 53. t. 34.*
f. 2.
Ketmia africana frutescens, foliis mollibus & incanis.
Comm. hort. 2. p. 151. t. 79.
Habitat in Æthiopia. *h*
Pedunculi ex axillis foliorum, sæpius terni, apice flores in singulo duos, sub foliolis tribus lanceolatis proferentes.
2. **HERMANNIA** foliis obovatis acute-incisis, pedunculis bidloris. *Roy. lugdb. 347.*
grossularifolia.

U u

Kut-

174 MONADELPHIA PENTANDRIA.

Ketmia africana vesicaria, uvæ crispæ foliis. *Comm. rar*
7. t. 7.

Habitat in Æthiopia. h

- alnifolia.* 3. *HERMANNIA* foliis cuneiformibus plicatis crenato-
emarginatis. *Hort. cliff.* 342. *Roy. lugdb.* 347.
Ketmia africana vesicaria fruticans & erecta, alni foliis
latis & majoribus. *Comm. hort.* 2. p. 155. t. 78.
Habitat in Æthiopia. h

- Hysopifolia.* 4. *HERMANNIA* foliis lanceolatis obtusis serratis. *Hort*
cliff. 342. *Hort. nps.* 195. *Roy. lugdb.* 347.
Althæa africana frutescens, floribus aureis cochlearis
pendulis minoribus. *Volk. norib.* 24. t. 24.
Habitat in Æthiopia. h

- lavendulifolia.* 5. *HERMANNIA* foliis lanceolatis obtusis integerrimis
Hort. cliff. 342. *Roy. lugdb.* 347.
Hermannia frutescens, folio lavendulæ latiore & obru-
so, flore parvo aureo. *Boerb. lugdb.* 1. p. 273.
Habitat in Æthiopia. h

- trifoliata.* 6. *HERMANNIA* foliis ternatis sessilibus. *Hort. cliff*
342. *Roy. lugdb.* 347.
Hermannia frutescens, folio oblongo molli cordato
hirsuto. *Boerb. lugdb.* 1. p. 273.
Habitat in Æthiopia. h

- pinnata.* 7. *HERMANNIA* foliis pinnatifidis linearibus. *Hort*
cliff. 342. *Roy. lugdb.* 348.
Hermannia frutescens, folio multifido tenui, caule ru-
bro. *Boerb. lugdb.* 1. p. 273.
Cistoides frutex æthiopicus, parvis coronopi foliis, ac-
nodos caulem radiatim ambientibus. *Pluk. mant.* 50
t. 46. f. 3
Habitat in Æthiopia. h

732 MELOCHIA.

- pyramidata.* 1. *MELOCHIA* floribus solitariis, capsulis pyramidatis
pentagonis: angulis acutis. *Hort. cliff.* 343. *Fl. zeyl.*
245.
Althæa brasiliensis frutescens, incarnato flore, sagopyri
semine. *Pluk. alm.* 25. t. 131. f. 3.
Habitat in Brasilia. 2

- depressa.* 2. *MELOCHIA* floribus solitariis, capsulis depressis pen-
tagonis: angulis obtusis. *Key. lugdb.* 348.

Abu

MONADELPHIA DECANDRIA. 675

Abutilon americanum, ribesii foliis, flore carneo, fructu pentagono aspero. *Houst. mss.*
Habitat in Havana.

3. MELOCHIA racemis confertis terminalibus, capsu-*cconcatenata*,
 lis globosis sessilibus. *Fl. zeyl.* 247. *

Alcea indica. flosculis parvis, fasciculatim ramulis ad-
 fixis. *Pluk. alm.* 26. t. 9. f. 5.

Habitat in India utraque. 2

4. MELOCHIA floribus capitatis sessilibus, capsulis sub-*corchorifolia*,
 rotundis. *Fl. zeyl.* 246.

Melochia corchori folio. *Dill. elsb.* 221. t. 176. f. 217.

Althæa peregrina, longiore betonicæ folio, floribus
 albis perexiguis, capitulis arcte conglomeratis. *Pluk.*
alm. 25. t. 44. f. 5.

Tieru-uren. *Rheed. mal.* 9. p. 143. t. 73.

Habitat in India.

5. MELOCHIA floribus capitatis, foliis ovatis ferratis, *supina*.
 caulibus procumbentibus.

Alcea supina pusilla, geranii exigui maritimi folio & fa-
 cie, maderaspatensis, fructu in summo caule glome-
 raro, pericarpio duro. *Pluk. alm.* 14. t. 132. f. 4.

Habitat in India.

DECANDRIA. 733

CONNARUS.

1. CONNARUS. *Fl. zeyl.* 248. *

Rhus zeylonicus trifoliatus, phaseoli facie, floribus co-
 piofis spicatis. *Burm. zeyl.* 199. t. 89.

Phaseolus arborescens zeylanicus monocarpos. *Raj.*
suppl. 438.

Habitat in India. b

HUGONIA. 734

1. HUGONIA spinis oppositis revolutis. *Fl. zeyl.* 249. *Myxstrax.*

Ægoceros arbor zeylanica, lauri fructu, cortice co-
 litum olente. *Raj. dendr.* 45.

Myxstar frutex baccifer malabaricus, fructu calycu-
 lato rotundo monopireno. *Raj. hist.* 1570.

Modira-canni. *Rheed. mal.* 2. p. 29. t. 19.

Habitat in India.

735

GERANIUM.

* *Staminibus 7 fertilibus.**triste.*

1. GERANIUM calycibus monophyllis: tubis longissimis subsessilibus, radice subrotunda. *Hort. cliff.* 344.

Hort. upf. 197.

Geranium calycibus monophyllis: tubis longissimis subsessilibus, radice subrotunda, foliis lobatis. *Roy. lugdb.* 352.

Geranium africanum noctu olens tuberosum, vitis foliis hirsutis. *Comm. hort.* 2. p. 123. t. 62.

- β. Geranium calycibus monophyllis: tubis longissimis sessilibus, radice subrotunda, foliis pinnatifidis. *Roy. lugdb.* 352.

Geranium triste. *Corn. canad.* 109. t. 110.

- γ. Geranium noctu olens æthiopicum, radice tuberosa, foliis myrrhidis latioribus & angustioribus. *Breyn. cent.* 126. t. 58.

Habitat in Æthiopia & forte in India. 2

fulgidum.

2. GERANIUM calycibus monophyllis, foliis tripartitis incis: intermedia majore, pedunculis umbelliferis geminis, caule carnos. *Vir. cliff.* 67. *Hort. cliff.* 498. *Roy. lugdb.* 353.

Geranium africanum, folio alceæ, flore coccineo fulgidissimo. *Dill. elsb.* 156. t. 130. f. 137.

Geranium surinamense, chelidonii folio, flore coccineo, petalis æqualibus. *Till. pis.* 68. t. 26.

Habitat in Æthiopia. 5

inguinat.

3. GERANIUM calycibus monophyllis, foliis orbiculato-reniformibus tomentosis crenatis integrifusculis. *Hort. upf.* 195.

Geranium calycibus monophyllis, foliis florentibus erectis, foliis subcordatis. *Hort. cliff.* 345. *Roy. lugdb.* 353.

Geranium africanum arborescens, malvæ folio pingui, flore coccineo. *Dill. elsb.* 151. t. 125. f. 151. 152.

Geranium africanum arborescens, malvæ folio planolucido, flore elegantissimo kermesino. *Mart. cent.* 3. f. 3.

Habitat in Africa. 5

Folia digitis tacta inquinant colore ferrugineo.

papilionaceum.

4. GERANIUM calycibus monophyllis, corollis papilionaceis: vexillo dipetalò maximo, foliis angulatis. *Hort. cliff.* 345. *Hort. upf.* 197. *Roy. lugdb.* 353.

Ger-

Geranium africanum arborescens, flore veluti dipetalo eleganter variegato. *Dill. elth.* 154. t. 128. f. 155.

Geranium africanum arborescens, malvæ folio mucronato, petalis florum inferioribus vix conspicuis. *Mart. cent.* 15. t. 15.

Habitat in Africa. 5

5. GERANIUM calycibus monophyllis, corollis papi-^{myrrhisfoli-} lionaceis, vexillo dipetalo, foliis bipinnatis. *Hort. um. cliff.* 345. *Roy. lugdb.* 353.

Geranium africanum, coriandri folio, floribus incarnatis, majus. *Herm. lugdb.* 279. t. 280.

β. *Geranium africanum* tuberosum, anemones folio, incarnato flore. *Herm. par.* 178. t. 178.

Geranium africanum, myrrhidis folio, flore albicante, radice rapacca. *Comm. hort.* 2. p. 125. t. 63.

Habitat in Africa. 2

6. GERANIUM calycibus monophyllis, foliis pinnatis: ^{pinnatum.} foliolis ovatis.

Geranium africanum, astragali folio. *Comm. præl.* 53. t. 53.

Habitat in Africa.

7. GERANIUM calycibus monophyllis, foliis cuculla-^{cucullatum.} tis dentatis. *Hort. cliff.* 345. *Hort. upf.* 196. *Roy. lugdb.* 353.

Geranium africanum arborescens, foliis cucullatis angulosis. *Dill. elth.* 155. t. 129. f. 156.

Geranium africanum arborescens, ibisci folio rotundo, carlinæ odore. *Herm. lugdb.* 274. t. 275.

Geranium africanum arborescens, ibisci folio anguloso, floribus amplis purpureis. *Mart. cent.* 28. t. 28.

Habitat in Africa.

8. GERANIUM calycibus monophyllis, caule carnosio ^{gibbosum.} gibboso, foliis subpinnatis.

Geranium calycibus monophyllis geniculis nodofo, foliis duplicato pinnatifidis. *Hort. upf.* 345. *Roy. lugdb.* 354.

Geranium africanum carnosum, petalis angustis albicantibus. *Dill. elth.* 153. t. 127. f. 154.

Geranium calycibus monophyllis, caule carnosio nodofo: internodiis filiformibus, foliis lobato-pinnatifidis.

Roy. lugdb. 354.

Geranium africanum noctu olens tuberosum & nodofo, aquilegiæ foliis. *Herm. lugdb.* 284. t. 285.

Seiff. bot. 111. t. 111.

Habitat in Africa.

U u 3.

9. GE-

678 MONADELPHIA DECANDRIA.

- peltatum.* 9. GERANIUM calycibus monophyllis, foliis quinquelobis integerrimis glabris peltatis. *Hort. cliff.* 345. *Roy. lugdb.* 352.
Geranium africanum, foliis inferioribus asari, superioribus staphidis agræ maculatis splendentibus & acetosæ sapore. *Comm. præl.* 52. t. 2.
Habitat in Africa. 5
- acetosum.* 10. GERANIUM calycibus monophyllis, foliis glabris subovatis carnosissimis crenatis. *Hort. cliff.* 345. *Roy. lugdb.* 352.
Geranium africanum frutescens, folio crasso & glauco, acetosæ sapore. *Comm. præl.* 54. t. 4.
Habitat in Africa. 5
- Zonale.* 11. GERANIUM calycibus monophyllis, foliis cordato orbiculatis incisissimis zona notatis. *Hort. ups.* 196.
Geranium africanum arborescens, alchemillæ hirsuto folio, floribus rubicundis. *Comm. præl.* 51. t. 1.
Habitat in Africa. 5
- vitifolium.* 12. GERANIUM calycibus monophyllis, foliis adscendentibus lobatis pubescentibus. *Hort. ups.* 196.
Geranium africanum arborescens, vitis folio, odore melissæ. *Dill. elth.* 152. t. 126. f. 153.
Habitat in Africa.
- capitatum.* 13. GERANIUM calycibus monophyllis, foliis lobatis undatis villosis. *Hort. ups.* 196.
Geranium calycibus monophyllis, floribus capitatis, foliis cordatis lobatis crenatis pilosis. *Hort. cliff.* 345. *Roy. lugdb.* 353.
Geranium africanum frutescens, malvæ folio laciniato odorato. *Herm. lugdb.* 277. t. 278.
Geranium americanum arborescens, vitis folio, odore melissæ. *Dill. elth.* 152. t. 126. f. 153.
Habitat in Africa.
- alchimilloides.* 14. GERANIUM calycibus monophyllis, foliis orbiculatis palmatis incisissimis pilosis, caule herbaceo. *Vir. cliff.* 345. *Hort. ups.* 197. * *Roy. lugdb.* 354.
Geranium calycibus monophyllis longissimis sessilibus fructu assurgente, foliis palmatis crenatis. *Hort. cliff.* 345.
Geranium africanum, alchemillæ hirsuto folio, floribus albidis. *Herm. lugdb.* 282. t. 283. *Stiff. bot.* 107. t. 107
Habitat in Africa.

15. GE

15. GERANIUM calycibus monophyllis, caule carno-^{odoratissi-}
 so brevissimo, ramis longis, foliis cordatis. *Hort. cliff.* ^{mum.}
 345. *Roy. lugdb.* 354.
 Geranium africanum humile, folio fragrantissimo mol-
 li. *Dill. elth.* 157. t. 131. f. 138.
Habitat in Africa. 2
16. GERANIUM calycibus monophyllis, pedunculis ^{grossulacini-}
 multifloris, foliis cordatis subrotundis lobatis crenatis. ^{des.}
 Geranium pedunculis bifloris, foliis cordatis incis glab-
 rris, caulibus filiformibus procumbentibus. *Roy. lugdb.*
 351.
 Geranium africanum, uvæ crispæ folio, calycibus pro-
 cumbentibus, floribus exiguis rubellis. *Herm. lugdb.*
 287. t. 289.
Habitat in Africa.
 Caules filiformes, procumbentes, Folia cordata, glabra, in-
 cisa. Pedunculi biflori cum calycibus monophyllis.
17. GERANIUM calycibus monophyllis, foliis oblon-^{auritum.}
 gis integris tripartitisque, pedunculis radicalibus.
 Geranium africanum, foliis plerumque auritis, floribus
 ex rubro purpurascens. *Comm. hort.* 2. p. 121.
 t. 61.
Habitat in Africa. 2
18. GERANIUM calycibus monophyllis, foliis corda-^{glaucophyl-}
 to-ovatis crenatis petiolo brevioribus, caule herbaceo. ^{lum.}
Roy. lugdb. 354.
 Geranium ægyptium glaucophyllum, rostris longissimis
 plumosis. *Dill. elth.* 150. t. 124. f. 150.
Habitat in Ægypto. 2
19. GERANIUM calycibus monophyllis, foliis ovatis ^{althæoides,}
 sinuatis crenatis, caule filiformi. *Hort. cliff.* 345. *Roy.*
lugdb. 354.
 Geranium, folio althææ, africanum, odore melissæ. *Boerb.*
lugdb. 1. p. 263.
Habitat in Africa. 2
- 20 GERANIUM calycibus monophyllis, foliis ovatis ^{betulinum,}
 inæqualiter serratis planis.
 Geranium frutescens, folio lato dentato, flore magno
 rubente. *Burm. afric.* 92. t. 33. f. 1.
 Geranium frutescens, folio subrotundo dentato, flore pur-
 pureo. *Burm. afric.* 93. t. 33. f. 2.
Habitat in Africa. 5

* *Staminibus quinque fertilibus*

- eichenarium*, 21. GERANIUM pedunculis multifloris, calycibus pentaphyllis, floribus pentandris, foliis pinnatis incisitis obtusis. *Hort. cliff.* 334. *Fl. suec.* 579. *Mat. med.* 337. *Roy. lugdb.* 352. *Hall. belv.* 369. *Dalib. parif.* 208. Geranium, cicutæ folio, minus & lupinum. *Bauh. pin.* 319. Geranium supinum. *Doa. pempt.* 63.
- moschatum*, 22. Geranium, cicutæ folio, moschatum. *Bauh. pin.* 319. *Habitat in Europæ sterilibus cultis.* ☉
- malacoides*, 22. GERANIUM pedunculis multifloris, calycibus pentaphyllis, floribus pentandris, foliis cordatis sublobatis. *Hort. cliff.* 344. *Roy. lugdb.* 352. *Samu. monsp.* 154. Geranium folio althææ. *Bauh. pin.* 318. Geranium malacoides. *Lob. ic.* 662. Geranium pedunculis bifloris, foliis ovatis hirsutis crenato-incisitis obtusis, caulibus procumbentibus. *Roy. lugdb.* 391. *Habitat in Italiæ G. narbonensis, Angliæ maritimis.* ☉
- gratum*, 23. GERANIUM pedunculis multifloris, calycibus pentaphyllis, floribus pentandris, foliis pinnatis acutis sinuatis. Geranium pedunculis multifloris pentandris. *Hort. cliff.* 498. Geranium pedunculis bifloris pentandris, radice annua. *Vir. cliff.* 66. *Roy. lugdb.* 352. Geranium acu longissima. *Bauh. pin.* 319. *prodr.* 138. Geranium speciosum annuum, longissimis rostris, creticum. *Bauh. bist.* 3. p. 479. *Habitat in Creta.* ☉
- Docet Fructificatio & habitus tres ultimas plantas ab initio unam fuisse, ortas procul dubio ex miscela diversarum, ut pro eadem haberi possent: at veropossesse etiam distingvi distinctas arbitrarium videtur.*
- * *Staminibus decem fertilibus.*
- tuberosum*, 24. GERANIUM pedunculis bifloris, foliis multipartitis: laciniis linearibus subdivisis obtusis. *Hort. cliff.* 343. *Roy. lugdb.* 350. Geranium tuberosum majus. *Bauh. pin.* 318. *Habitat in Anglia?* 2
- macrorrhizum*, 25. GERANIUM pedunculis bifloris, calycibus inflatis, pistillo longissimo. *Hort. cliff.* 343. * *Roy. lugdb.* 350. Geranium batrachioides odoratum. *Bauh. pin.* 318. *Habitat - - -* 2

26. GERANIUM pedunculis bifloris foliisque alternis, *phenon*, calycibus subaristatis, caule erecto.
 Geranium pedunculis bifloris alternatim cauli integro superne nudiusculo insidentibus. *Hort. cliff.* 343. *
Hort. upf. 198. *Roy. lugdb.* 350.
 Geranium montanum fuscum. *Bauh. pin.* 318.
 β. Geranium batrachioides hirsutum, flore atro-rubente.
Bauh. pin. 318.
 Geranium 1, pulio flore. *Clus. hist.* 2. p. 99.
Habitat in Alpibus Pannonicis, Helveticis, Styrlacis. &
 27. GERANIUM pedunculis bifloris, foliis caulinis trinodisum.
 lobis integris serratis: summis subsessilibus *Hort. cliff.*
 343. *Roy. lugdb.* 350.
 Geranium nodosum. *Bauh. pin.* 318.
Habitat in Delphinatu.
 28. GERANIUM pedunculis bifloris, foliis peltatis incisoferratis, caule erecto. *Fl. lapp.* 266. *Fl. succ.* 572.
Hort. cliff. 344. *Roy. lugdb.* 351.
 Geranium batrachioides, folio aconiti. *Bauh. pin.* 317.
 Geranium 2. batrachioides minus. *Clus. hist.* 2. p. 99.
Habitat in Europæ borealis sylvis. &
 29. GERANIUM pedunculis bifloris, foliis peltatis multipartitis rugosis pinnato-laciniatis acutis. *Hort. cliff.*
 344. *Fl. succ.* 573. *Roy. lugdb.* 350. *Hall. helv.* 367.
 Geranium batrachioides, Gratia dei germanorum. *Bauh. pin.* 318.
 Geranium 3 batrachioides majus. *Clus. hist.* 2. p. 100.
Habitat in Europæ borealis pratis. &
 30. GERANIUM pedunculis bifloris, caule dichotomo erecto, foliis quinquepartitis incisil: summis sessilibus. *Gron. virg.* 78.
 Geranium batrachioides americanum maculatum. floribus obsolete cæruleis. *Dill. elib.* 158. t. 132. f. 159.
Habitat in Carolina, Virginia, Sibiria. &
 31. GERANIUM pedunculis bifloris, calycibus pilosis rebertianum decem-angulatis.
 Geranium pedunculis bifloris, foliis quinque-tripartitis, lobis pinnatifidis, calycibus hirsutis. *Hort. cliff.*
 344. *Fl. succ.* 578. *Mat. med.* 336. *Dalib paris.*
 207. *Roy. lugdb.* 351.
 Geranium rebertianum primum. *Bauh. pin.* 319.

Geranium 3. *Fuchs, hist.* 206.

Habitat in Europæ borealis rupibus. 2

Calyx pilosus, decem carinatus, ex foliolis 1. 3. exterioribus tricarinatis, 2. 5. unicarinatis, 4. bistarinatis. Petala integra.

- lucidum.* 32. GERANIUM pedunculis bifloris, calycibus pyramidatis angulatis rugosis, foliis quinquelobis rotundatis. *It. gosl.* 128. * *Fl. suec.* 574. *Dalib. paris.* 208. *Sauv. monsp.* 207.

Geranium lucidum saxatile. Barb. pin. 318.

Geranium saxatile. Thal. herc. 44. t. 5.

Geranium rotundifolium saxatile montanum. Col. ecphr. 1. p. 138. t. 137.

Habitat in Europæ rupibus arborosis. ☉

- malle.* 33. GERANIUM pedunculis bifloris foliisque ramis alternis, caule ramoso diffuso, calycibus muticis.

Geranium pedunculis bifloris, foliis quinquepartito-multifidis rotundatis: laciniis obtusiusculis, capsulis hirtis. Fl. suec. 577. *Dalib. paris.* 206.

Geranium pedunculis bifloris, foliis sparsis reniformibus semiquinquefidis mollibus & latissimis. Hall. belv. 366. * *Sauv. monsp.* 207.

Geranium columbinum minus, majori flore & foliis bifidis. Magn. monsp.

Geranium columbinum villosum, petalis bifidis purpureis. Vaill. paris. 79. t. 15. f. 3.

Habitat in Europa ad plateas. ☉

- carolinianum.* 34. GERANIUM pedunculis bifloris, calycibus aristatis, foliis multifidis, pericarpis hirsutis. *Roy. Ingdh.* 351. *Gron. virg.* 78.

Geranium columbinum carolinum, capsulis nigris hirsutis. Diss. elth. 162. t. 135. f. 162.

Habitat in Carolina, Virginia. ☉

- columbinum.* 35. GERANIUM pedunculis bifloris, foliis quinquepartito-multifidis angulatis: laciniis acutis, capsulis glabris, calycibus aristatis. *Fl. suec.* 576. *Dalib. paris.* 207. *Geranium columbinum, foliis dissectis, pediculis florum longissimis. Vaill. paris.* 79. t. 15. f. 4. *Geranium foliis ad nervum quinquefidis, pediculo longissimo, caule prostrato. Hall. belv.* 367. *Habitat in Gallia, Helvetia, Germania.*

MONADELPHIA POLYANDRIA. 689

36. GERANIUM pedunculis bifloris, petalis integris longitudine calycis, caule prostrato, foliis reniformibus incis. *rotundifolium.*
 Geranium pedunculis bifloris, foliis subrotundis multifidis, caule procumbente. *Hort. cliff.* 344. *Fl. suec.* 575. *Roy. lugdb.* 351. *Hall. belv.* 365. *Dalib. paris.* 206.
 Geranium folio malvæ rotundo. *Bauh. pin.* 318.
 Geranium alterum. *Fuchs. hist.* 205.
 Geranium columbinum majus, flore minore cæruleo. *L'ail. paris.* 79. t. 15. f. 1.
Habitat in Europæ cultis. ☉
37. GERANIUM pedunculis subunifloris, foliis quinquepartitis acutis: foliolis pinnatifidis. *sibiricum.*
Habitat in Sibiria.
Caules pedales diffusi, divaricati. Folia quinquepartita: lobis pinnatifidis. Pedunculus e singula bifurcatum. Caulis longus, uniflorus, in medio geniculatus bracteis 2. oppositis subulatis. Corolla alba stris purpureis petalis vix emarginatis. Calyx 5-phyllus, aristatus. Capsula subvillosa.
38. GERANIUM pedunculis unifloris, foliis quinquepartitis trifidis orbicularis. *sanguineum.*
 Geranium pedunculis simplicibus unifloris. *Hort. cliff.* 343. *Fl. suec.* 571. *Roy. lugdb.* 350. *Hall. belv.* 365. *Dalib. paris.* 206.
 Geranium sanguineum, maximo flore. *Bauh. pin.* 318.
 Geranium VII hæmatodes. *Clus. hist.* 2. p. 201.
 β. Geranium hæmatodes laucastriensis, flore eleganter striato. *Dill. elsh.* 163. t. 136. f. 163.
Habitat in Europæ pratis siccis umbrosis. 2
39. GERANIUM pedunculis simplicibus unifloris, foliis tripartito-multifidis linearibus, caule arborco. *grandiflorum.*
 Geranium frutescens incanum, flore magno luteo dentato. *Burm. afric.* 88. t. 34. f. 1.
Habitat in Æthiopia. 3

POLYANDRIA.

SIDA. 736

1. SIDA foliis cordato-ovatis serratis, stipulis setaceis, 2-spinosis, villis subspinosis. *Hort. ups.* 199. *Fl. zeyl.* 254. *
Roy. lugdb. 349.

Sida

- Sida* foliis cordato-ovatis serratis, capsulis quinis bica-
spidatis. *Hort. cliff.* 346.
- Alcea*, carpini folio, americana frutescens, flosculis lu-
teis, semine duplici rostro. *Comm. hort.* 1. p. 3. t. 2.
- Althæa* carpini folio, flosculis luteis. *Bocc. sic.* 11. t. 2.
- Althæa virginiana* bidens, pimpinellæ majoris acutiore
folio, flosculis minimis luteis. *Pluk. alm.* 25. t. 9. f. 6.
- β. *Malvinda* bicornis, sangvisorbæ folio villosa & glabro.
Dill. elsb. 214. t. 171. f. 210.
- Althæa* rostrata cormandelienſis, pimpinellæ majoris
folio subrotundo, semine bicorni. *Pluk. alm.* 26. t.
9. f. 3.
- Habitat in India utraque.* ☉
- rhombifolia*, 2. *SIDA* foliis lanceolato-rhomboidibus serratis. *Vir. cliff.*
68. *Hort. cliff.* 346. *Hort. upf.* 199. *Fl. zeyl.* 252. *
Gran. virg. 78. *Roy. lugdb.* 349.
- Malvinda* unicornis, folio rhomboide, perennis, *Dill.*
elsb. 216. t. 172. f. 212.
- β. *Althæa* carpini folio, flosculis luteis. *Bocc. sic.* 11. t. 6.
f. 2.
- Alcea* pernambucana, carpini folio, flosculis minimis.
Morif. hist. 2. p. 528. f. 5. t. 19. f. ult. mal.
- Habitat in India utraque.* ☉
- alvifolia*, 3. *SIDA* foliis orbiculatis plicatis serratis. *Hort. cliff.* 346.
Hort. upf. 200. *Fl. zeyl.* 253. *. *Roy. lugdb.* 349.
- Malvinda* stellata alvifolia. *Dill. elsb.* 215. t. 172. f.
211.
- β. *Althæa* indica, latiori folio cordiformi ad summum ſi-
nuato *Pluk. alm.* 26. t. 9. f. 2.
- Habitat in India.* ☉
- cordifolia*, 4. *SIDA* foliis cordatis subangulatis serratis villosis.
Malva bicornis, ballotes folio molli. *Dill. elsb.* 218.
t. 171. f. 209.
- Althæa* maderaspathana, subrotundo folio molli & hir-
suto, multipilis. *Pluk. alm.* 25. t. 131. f. 2.
- Habitat in India.* ☉
- Radix annua.* Calyx simplex. Seminum rostra deciden-
te corolla, in fasciculum longitudine calycis prominen-
tia. Calyx pilosus.
- periplocaefolia* γ. *SIDA* foliis cordato-lanceolatis integerrimis. *Fl. zeyl.*
251.
- Sida* foliis ovato-lanceolatis, caule paniculato. *Roy.*
lugdb. 349.

Abu-

MONADELPHIA POLYANDRIA. 685

Abutilon periplocæ acutioris folio, fructu stellato. Dill.

elsh. 4. t. 3. f. 3.

Alcea orientalis, scammoniae folio, major, floribus albis.

Pluk. alm. 17. t. 74. f. 7.

Habitat in Zeylona. b. ☉

6. SIDA foliis subrotundo-cordatis acuminatis integer-*amplissima*, rimis. *Roy. lugdb. 350.*

Abutilon amplissimo folio, caule villosa. Plum. spec. 2.

Habitat in America calidiore.

7. SIDA foliis subrotundo-cordatis acuminatis crenatis. *Abutilon.*

Roy. lugdb. 349. Hort. cliff. 346. Hort. upf. 198.

Gron. virg. 78.

Althæa theophrasti, flore luteo. Bauh. pin. 316.

Althæa altera f. Abutilon. Cam. epit. 668.

β-*Abutilon indicum, flore luteo minore. Fl. zeyl. 520.*

Alceæ affinis indica Abutilon dicta, flore flavo, foliis mollibus profunde venosis, filiculis lanuginosis. Pluk.

phys. 126. f. 5.

Habitat in Indiis. ☉

8. SIDA foliis cordatis subrotundis crenatis tomentosis, *crispa*, pericarpis inflatis multilocularibus crenatis repandis.

Abutilon vesicarium crispum, floribus melinis parvis.

Dill. elsh. 6 t. 5. f. 5.

Abutilon americanum, fructu subrotundo pendulo e capsulis vesicariis crispis conflato. Mart. cent. 29.

t. 29.

Habitat in Carolina, Providentia, Bahama.

9. SIDA foliis angulatis: inferioribus cordatis, superiori-*crispa*, bus panduriformibus

Sida foliis crenatis: inferioribus dentatis obtusis, superioribus acuminatis. Roy. lugdb. 349.

Sida foliis cordatis obsolete ferratis. Hort. cliff. 346.

Abutilon lavateræ flore, fructu cristato. Dill. elsh. 3.

t. 2. f. 2.

β-*Sida foliis inferioribus cordato-angulatis, superioribus hastatis, petalis integerrimis. Roy. lugdb. 349. Hort.*

upf. 199.

Habitat in Mexico. ☉

10. SIDA capitulis pedunculatis triphyllis septemfloris. *capitata.*

Lin. a. f. upf. 1743. p. 137. t. 2.

Malva caule erecto hispido, foliis erectis crenatis difformibus. Roy. lugdb. 357.

Mal-

Malva aspera major aquatica ex hortensium rosarum genere, flore minore luteo, semine aculeato. *Sloan. Jam. 96. hist. 1. p. 217. Raj. suppl. 320.*
Habitat in Caribæis, locis paludosis. &

737

NAPÆA.

hermaphro-
dita.

1. NAPÆA pedunculis nudis lævibus, foliis glabris, floribus hermaphroditis. *Gen. nov. 1089. **
Sida foliis palmatis: laciniis lanceolato attenuatis. *Hort. cliff. 346. Hort. nps. 198. Roy. lugdb. 348.*
Malva virginiana, ricini folio. *Herm. lugdb. 22. s. 23.*
Habitat in Virginia. &

dioica.

2. NAPÆA pedunculis involucreatis angulatis, foliis scabris, floribus dioicis. *Gen. nov. 1089. **
Abutilon folio profunde dissecto, pedunculis multifloris, mas. *Ehret. pict. 8. f. distincta.*
Abutilon folio profunde dissecto, pedunculis multifloris, femina. *Ehret. pict. 7. f. 1.*
Althæa magna aceris folio, cortice cannabino, floribus parvis, semina rotatim in summitate caulium, singula singulis cuticulis rostratis cooperta ferens. *Banist. 1928.*
Habitat in Virginia. &
Flores feminae gaudent etiam staminibus, at anthera minima, effæta & steriles sunt.

738

ALTHÆA.

officinalis.

1. ALTHÆA foliis simplicibus tomentosis. *Hort. cliff. 348. Hort nps. 204. Mat. med. 338. Roy. lugdb. 357. Dalib. parif. 210. Gort. gelr. 410.*
Althæa dioscoridis & plinii. *Banb. pin. 315.*
Althæa. Fuchf. hist. 15. Cam. epit. 667.
 β. *Althæa* laciniato folio. *Banb. pin. 316.*
Habitat in Hollandiæ, Angliæ, Galliæ, Sibiriz sub humidis. &

cannabina.

2. ALTHÆA foliis inferioribus palmatis, superioribus digitatis, caule fruticoso *Hort. nps. 205. Sauv. monsp. 200.*
Althæa foliis compositis scabris. *Hort. cliff. 349. Roy. lugdb. 357.*
Alcea cannabina. Banb. pin. 316.
Alcea fruticosa, cannabino folio. *Clus. hist. 2. p. 25.*
Habitat in Hungaria, Italia, G. Narbonensi. &

3. AL-

3. *ALTHÆA* foliis trifidis piloso-hispidis. *Hort. cliff. hirsuta.*
 349. *Roy. lugdb.* 357. *Dalib. parif.* 210. *Sauv.*
monsp. 200.
Alcea hirsuta. *Baub. pin.* 317
Alcea hirsuta minima, flore cælio hispanica. *Barr. ic.* 1169.
Alcea villosa. *Dalecb. hist.* 594.
Habitat in Gallia, Italia, Hispania.

ALCEA. 739

1. *ALCEA* foliis sinuato angulosis. *Hort. cliff.* 348. *rosea.*
Hort. upf. 204. *Mat. med.* 340. *Roy. lugdb.* 357.
Malva rosea, folio subrotundo. *Baub. pin.* 315.
Malva hortensis. *Dod. pems.* 652.
Habitat - - - - δ
 2. *ALCEA* foliis palmatis. *Hort. cliff.* 348. *Hort. upf. ficifolia.*
 204. *Roy. lugdb.* 357.
Malva rosea, folio ficus. *Baub. pin.* 315.
Malva rosea simplex peregrina. *Tabern. hist.* 315.
Habitat - - - - δ
Vix sufficienter a precedente distincta.

MALVA. 740

* *Foliis indivisis.*

1. *MALVA* foliis cordatis crenatis tomentosis, floribus *tomentosa.*
 lateralibus congestis, caule fruticoso. *Fl. zeyl.* 255. *
Malva foliis cordatis crenatis villosis. *Hort. cliff.* 347.
Roy. lugdb. 357.
Althæa orientalis bidens, subrotundis foliis, floribus par-
vis luteis. *Pluk. amalib.* 11. t. 356. f. 1?
Habitat in India. 5.
Caulis frutescens, tomentosus. Folia sine lobis cordata, to-
mentosa. Calyx inferior triphyllus. Seminum rostra,
deciduo flore, brevissima, angulos fructus retusi consti-
tuentia.
Confer. Althæa indica, abutili subrotundo folio, flore lu-
teo spicato. *Pluk. phyt.* 74. f. 6.
 2. *MALVA* foliis ovato-oblongis acute serratis, capsulis *coromande-*
 tricuspatis. *Hort. cliff.* 346. * *Hort. upf.* 201. *liana.*
Althæa coromandeliana, angustis prælongis foliis, semine
bicorni. *Pluk. mant.* 10. t. 334. f. 2.
Habitat in America? ☉
 3. *MALVA* foliis cordatis crenatis, floribus lateralibus *so-*
litaris terminalibus spicatis. *Roy. lugdb.* 359. *americana.*

Al-

688 MONADELPHIA POLYANDRIA.

Althæa americana pumila, flore luteo spicato. *Breyn. cent.* 124. t. 57.

Habitat in America. ☉

* *foliis angulatis.*

peruviana. 4. MALVA caule erecto herbaceo, foliis lobatis, spicis secundis axillaribus.

Habitat in Peru. D. Jussieu. ☉

Caulis erectus, tripedalis, pilis rarioribus sæpius geminis. Folia septemlobata, plicata, glabra, venosa, acute serrata; petiolis longitudine foliis. Stipulæ ovato-lanceolatæ. Pedunculus axillaris, nudus, longus. Spica secunda, inversum versa, recurvata ante florescentiam. Corollæ parvæ purpureæ. Semina 10, supra exasperata dentibus alternis plurimis.

bryonifolia. 5. MALVA foliis palmatis scabris, caule tomentoso fruticoso, pedunculis multifloris. *Roy. lugdb.* 356.

Althæa frutescens, bryoniæ folio. Baub. pin. 316. p. 138.

Althæa profunde serrata l. dentato folio. Baub. hyst. 2. p. 955.

Habitat in Hispania. ☿

Planta belle echinata per discum foliorum.

capensis. 6. MALVA foliis subcordatis laciniatis glabris, caulcareborecente. *Hort. cliff.* 347. *Hort. upf.* 201.

Malva capensis frutescens, grossulariæ folio minori glabro, Dill. elth. 208. t. 169. f. 206.

β. *Malva capensis frutescens, grossulariæ folio majore hirsuto. Dill. elth.* 200. t. 169. f. 207.

γ. *Malva africana frutescens, flore rubro. Comm. hort.* 2. p. 171. t. 86.

Malva africana frutescens, flore parvo carneo unguiculis atrorubentibus. Pluk. amalsh. 140. t. 431. f. 5.

Habitat in Æthiopia. ☿

caroliniana. 7. MALVA caule repente, foliis multifidis. *Hort. cliff.* 347. *Hort. upf.* 201.

Abutilon repens, alceæ foliis, flore helvulo, Dill. elth. 5. t. 4. f. 4.

Abutilon carolinianum repens, alceæ foliis, gilvo flore, Mart. cent. 34. t. 34.

Habitat in Carolina. ☉

rotundifolia. 8. MALVA caule repente, foliis cordato-orbiculatis obsolete quinquelobis. *Hort. cliff.* 347. *Fl. jucc.* 580.

Mat.

MONADELPHIA POLYANDRIA. 689

Mat. med. 339. *Roy. lugdb.* 356. *Hall. belv.* 363. *Gren. virg.* 79. *Dalib. parif.* 209.

Malva sylvestris, folio subrotundo. *Bauh. pin.* 314.

Malva sylvestris pumila. *Fuchs. hist.* 508.

Habitat in Europæ ruderatis, viis, plateis. ☉

9. *MALVA* caule erecto herbaceo, foliis lobatis obtusis, *sylvestris.*
pedunculis petiolisque pilosis.

Malva caule erecto, foliis multipartitis. *Hort. cliff.* 347.

Fl. suec. 581. *Roy. lugdb.* 356. *Hall. belv.* 362. *Dalib. parif.* 208.

Malva sylvestris, folio sinuato *Bauh. pin.* 314.

Malva sylvestris recta. *Cord. hist.* 114.

Habitat in Europæ campestribus.

10. *MALVA* caule erecto herbaceo, foliis lobatis obtu- *marritana.*
lis, pedunculis petiolisque glabris.

Malva hederaceo folio. *Bauh. pin.* 315. *I. rt. upf.* 20

Malva major tertia. *Dalech. hist.* 586.

- β. *Malva folio ficus altera.* *Bauh. pin.* 315. *prodr.* 37.

Habitat in Italia, Lusitania, Hispania. ☉

11. *MALVA* caule erecto, foliis semiorbiculatis cr 12- *hispanica.*
tis, calyce exteriori diphyllo. *Læf.*

Malva rotundifolia glabra ex hispania, flore amplo rubello. *Pluk. alm.* 238. t. 44. f. 3.

Habitat in Hispania.

12. *MALVA* caule erecto, foliis angulatis, floribus axil- *verticillata,*
laribus glomeratis. *Vir cliff.* 356. *Hort. cliff.* 502.
Hort. upf. 200. *Roy. lugdb.* 356.

Malva annua rotundifolia, floribus omnium minimis albis pentaphyllis verticillatim genicula ambientibus.

Morif. hist. 2. p. 521. *Raj. hist.* 598.

- β. *Malva foliis angulatis crispis*, floribus axillaribus glo- *crispa,*
meratis. *Hort. cliff.* 347. *Hort. upf.* 200.

Malva foliis crispis. *Bauh. pin.* 315.

Malva crispa. *Dod. pempt.* 653.

Habitat a in China, Syria ☉

13. *MALVA* caule erecto, foliis multipartitis scabriuscu- *Alcea.*
lis. *Hort. cliff.* 347. *Fl. suec.* 582. *Roy. lugdb.* 356.

Hall. belv. 353. *Dalib. parif.* 209.

Alcea vulgaris major. *Bauh. pin.* 316.

Alcea vulgaris. *Clas. hist.* 2. p. 25.

Habitat in Germania, Anglia, Gallia. &

X x

14. MAL-

- meschata.* 14. MALVA foliis radicalibus reniformibus incisís, caulinis quinquepartitis pinnato-multifidis. *Hort. nps.* 202.
Malva montana f. Alcea rotundifolia laciniata. *Col. ecphr.* 1. p. 148. t. 147.
Alcea folio rotundo laciniato. *Bauh. pin.* 316.
Habitat in Italia, Gallia. &

- egyptia* 15. MALVA foliis palmatis dentatis, corollis calycemini-
noribus.
Alcea ægyptia, geranii folio. *Lipp. ægypt. mss.*
Habitat in Ægypto.
Habitus *antecedentis sed tenerior.* Caules filiformes, debiliores, scabri. Folia radicalia quinqueloba subdentata obtusa; superiora tripartita linearia dentata. Flores axillares solitarii, pedunculis longitudine petiolorum. Calyx exterior setis 3; interior latus, semiquinquesidus, acutus, Corollæ cærulescentes.

LAVATERA.

* *Caule fruticoso.*

- arbores.* 1. LAVATERA caule arboreo, foliis septemangul-
aribus tomentosis plicatis, pedunculis confertis unifloris axillaribus. *Hort. nps.* 202.
Lavatera foliis septemangularibus obtusis plicatis villosis, caule fruticoso, floribus ad alas confertis. *Hort. cliff.* 348. *Roy. lugdb.* 355.
Malva arborea veneta dicta, parvo flore. *Bauh. pin.* 315.
Malva arborefcens. *Dod. pempt.* 653.
Habitat inter Pisas & Liburnum. ♂

- micans,* 2. LAVATERA caule arboreo, foliis septemangul-
aribus acutis crenatis plicatis tomentosis, racemis terminalibus. *Roy. lugdb.* 356.
Malva foliis mollibus undulatis: in margine superi-
micis sulphureis ad solem splendentibus donatis. *Morif. hist.* 1. p. 523. f. 5. t. 17. f 9.
Malva hispanica. *Morif. pral.* 285.
Habitat in Hispania, Lusitania. ♂

- alba.* 3. LAVATERA caule fruticoso, foliis quinquelobo-
statis. *Hort. nps.* 202.
Lavatera foliis quinquangularibus acutis crenatis: lac-
inia media productiore. *Hort. cliff.* 348. *Roy. lugdb.* 355.
Althæa frutescens. folio acuto. parvo flore. *Bauh. pr.* 316.

Althæa

Althæa arborea olbia in galloprovincia. *Lob. ic.* 653.

Habitat in Olbia insula Galloprovinciæ. ♀

4. **LAVATERA** caule fruticoso, foliis subcordatis sub-trilobis trilobis rotundatis crenatis, stipulis cordatis, pedunculis unifloris.

Lavatera foliis subcordatis emarginatis obsoleto trilobis creuatis petiolo brevioribus, stipulis cordatis. *Roy.*

lugdb. 355

Malva foliis subcordatis trilobis obtusis serratis villosis.

Hort. cliff. 347.

Althæa frutescens, folio rotundiore incano. *Bauh. pin.* 316.

Althæa fruticans hispanica, aceris monspessulani incanis foliis, grandiflora saponem spirans. *Pluk. alm.* 24.

t. 8. *f.* 3.

Althæa frutex 1. *Clus. hist.* 2. *p.* 24.

Habitat in Hispania. ♀

Planta tomentosa: tota pilis minimis glutiniferis; eisque immixtis aliis pilis majoribus apice stellatis. Pedunculi tres pluresve ex alis petiolorum, uniflori. Flos magnus, pallide purpurascens.

5. **LAVATERA** caule fruticoso, foliis septemangulatis-lusitanica. bus tomentosis plicatis, racemis terminalibus. *Roy.*

lugdb. 355.

Althæa frutescens lusitanica, folio ampliore minus incano. *Tournef. inst.* 97.

Habitat in Lusitania. ♀

* *Caule herbaceo.*

6. **LAVATERA** caule herbaceo, fructibus denudatis, calycibus incisis. *Hort. upf.* 203.

Lavatera foliis inferioribus cordato-subrotundis, superioribus lobatis crenatis, calycibus utrinque incisis, caule herbaceo. *Roy. lugdb.* 355.

Althæa thuringiaca. *Cam. hort.* 12. *t.* 6.

Althæa thuringiaca grandiflora. *Dill. elch.* 9. *t.* 8. *f.* 8.

Althæa flore majore. *Bauh. pin.* 316.

Habitat in Paunonia, Thuringia, Tataria ad sepes. ♂

7. **LAVATERA** caule erecto, ramis inferioribus diffusis, pedunculis confertis unifloris axillaribus. *Hort.*

upf. 202. *

Malva cretica annua, flore parvo ad alas umbellato.

Tournef. cor. 2.

Habitat in Creta. ☉

X x 2

8. LA-

692 MONADELPHIA POLYANDRIA.

- trimestris.* 8. LAVATERA foliis glabris, caule scabro herbaceo, pedunculis unifloris, fructibus orbiculo tectis. *Hort. upf.* 203.
 Lavatera foliis trilobis: lacinia media productiore, caule herbaceo. *Hort. cliff.* 348.
 Lavatera foliis ovato-lanceolatis: inferne angulatis, pedunculis unifloris, caule herbaceo. *Roy. lugdb.* 355
Sanv. monsp. 101.
 Malva folio vario. *Bauh. pin.* 315, *prodr.* 137. t. 137.
 9. Lavatera foliis subcordatis obtusis angulatis serratis, caule herbaceo. *Roy. lugdb.* 355.
Habitat in Syria, Hispania, G. Narbonensi.

742
Malacoides.

MALOPE.

1. MALOPE foliis ovatis crenatis glabris. *Hort. cliff.* 347. *Hort. upf.* 206. *Roy. lugdb.* 357.
 Malva betonica folio. *Morif. bist.* 2. p. 522. f. 5. t. 17. f. 11. *Bocc. sic.* 15. t. 8. f. 2.
 Alcea betonica folio, flore purpureo-violaceo. *Barr. ic.* 1189.
Habitat in Hetruriæ pratis.

743
lobata.

URENA.

1. URENA foliis angulatis. *Hort. cliff.* 348. *Hort. upf.* 200. *Fl. zeyl.* 256. *Roy. lugdb.* 358.
 Urena finica, xanthii facie. *Dill. elsb.* 340. t. 319. f. 412.
 Trifolio adfinis indiæ orientalis, xanthii facie. *Breyn. cent.* 82. t. 35.
Habitat in China. b
sinuata. 2. URENA foliis sinuato multifidis villosis. *Fl. zeyl.* 257.
 Alcea indica frutescens, foliis in lacinias varie dissectis. *Pluk. alm.* 15. t. 74. f. 1.
 Malvinda foliis inferioribus multifidis, superioribus incisis, flore solitario. *Burm. zeyl.* 150. t. 69. f. 2.
Habitat in India. b
procumbens. 3. URENA foliis hastato-subcordatis indivisis serratis, caule procumbente.
Habitat in Chinæ monticulis. Osbeck. b
 Caulis frutescens, procumbens, repens, ramosissimus. Folia magnitudine origani, subcordata, integra nec lobata, ævia, argute serrata. Flores foliis majores sunt.

1. GOS-

GOSSYPIUM. 744

1. GOSSYPIUM foliis quinquelobis, caule herbaceo. *herbaceum*
Hort. upf. 203. *Mat. med.* 341.
Gossypium caule decumbente. *Hort. cliff.* 350. *Roy. lugdb.* 359.
Gossypium frutescens, semine albo. *Bauh. pin.* 430.
Gossypium. Cam. epit. 203.
Habitat in America. ☉
2. GOSSYPIUM foliis trilobis integerrimis. *Hort. upf. barbados.* 204.
Gossypium frutescens annuum, folio trilobo, barbadosense. *Pluk. alm.* 172. t. 188. f. 1.
Habitat in Barbados. ♂. ♀
3. GOSSYPIUM foliis palmatis: lobis lanceolatis, caule fruticoso. *arborescens.*
Gossypium caule erecto. *Hort. cliff.* 350. *Roy. lugdb.* 359.
Gossypium arborescens, caule lævi. *Bauh. pin.* 430.
Gossypium herbaceum f. *Xylon maderaspatense*, rubicundo flore, pentaphyllum. *Pluk. alm.* 172. t. 188. f. 3.
Cuduparini. Rheed. mal. 1. p. 55. t. 31.
Habitat in Indiæ arenosis. ♀

HIBISCUS. 745

1. HIBISCUS foliis ovatis acuminatis serratis, caule simplicissimo, petiolis floriferis. *Moscheutos.* *Hort upf.* 205.
Hibiscus foliis ovatis crenatis: angulis lateralibus obsolete. *Hort. cliff.* 349. *Gron. virg.* 79. *Roy lugdb.* 358.
Alcea rosea peregrina, forte *Rota moscheutos plinii.* *Corn. canad.* 144. t. 145. *Morif. bist.* 2. p. 532. f. 5. t. 19. f. 6.
Habitat in Canada, Virginia. 2
2. HIBISCUS caule herbaceo simplicissimo, foliis ovato-palmatis, subtus tomentosis, floribus axillaribus.
Althæa palustris. Bauh. pin. 316. (ex borto C. B. *Burserus*).
Althæa hortensis f. *peregrina.* *Dod. pempt.* 655.
Habitat in Virginia. Gronov. Canada Kalm. 2
Habitus H. Moscheutos. Caulis sesquipedales, non ramosi, annui. Folia lato-ovata, obtuse serrata, trimeria, acuminata, subtus tomentosa. Pedunculi ex axillis
X x 3 lis

694 MONADELPHIA POLYANDRIA.

- lis foliorum superiorum solitariis, petiolo longiores, ar-
nisi flori, non e petiolo enati, geniculati. Flos maximus.*
- populaceus.* 3. HIBISCUS foliis cordatis integerrimis. *Fl. zeyl.* 258. *
Alcea malabariensis, abutili folio, flore majore ex albo
flavescente. *Raj. bist.* 1069.
Bupariti. *Rheed. mal.* 1. p. 51. t. 29.
Habitat in India. h
Calyx exterior, in hac & sequente, monophyllus, de-
cumbentatus.
Conferatur *Keemia amplissimo tilia folio subtus argen-
teo. Catesb. car.* 2. p. 90. t. 90.
- tiliaceus.* 4. HIBISCUS foliis cordato-subrotundis acuminatis
crenatis, caule arboreo. *Roy. lugdb.* 532. *Fl. zeyl.*
259. *
Alcea malabarica, abutili folio, flore minore ex albo
flavescente exterius aspero. *Raj. bist.* 1070.
Pariti s. Tali-pariti. *Rheed. mal.* 1. p. 53. t. 30.
Habitat in India. h
- Rosa sinensis.* 5. HIBISCUS foliis ovatis acuminatis serratis glabris,
caule arboreo. *Fl. zeyl.* 260. *
Alcea javanica arborescens, flore pleno rubicundo.
Breyn. cent. 121. t. 56.
Scheru-pariti. *Rheed. mal.* 2. p. 25. t. 16.
Habitat in India. h
- hirtus.* 6. HIBISCUS foliis lanceolato-ovatis acuminatis serratis,
ramis scabris, caule herbaceo.
Habitat in India.
Planta parva. Caulis & praesertim Pedunculi, nec non
Folia et Calyces subtus pilis rigidiusculis hirti: Folia
lanceolata, argute serrata. Corolla purpurea.
- mutabilis.* 7. HIBISCUS foliis cordato-quinquangularibus obsolete
serratis, caule arboreo. *Hort. ups.* 205. *Hort. cliff.*
349. *Roy. lugdb.* 358.
Althæa arborea Rosa sinensis. *Marif. bist.* 2. p. 530.
f. 5. t. 18. f. 2.
Rosa sinensis. *Ferr. flor.* 493. t. 497. *Mer. sur.* 31. t.
31.
Hina-pariti. *Rheed. mal.* 6. p. 66. t. 38. 39. 40. 41.
Habitat in India. h
- Malavicus.* 8. HIBISCUS foliis cordatis crenatis: angulis lateralibus
extimis parvis, caule arboreo. *Hort. cliff.* 349. *Roy.*
lugdb. 358.

Mal-

MONADELPHIA POLYANDRIA. 695

Malvaviscus arborescens, flore miniato clauso. *Dill. elzb.* 210. t. 170. f. 208.

Alcea indica arborescens, folio molli, flore amplo eleganter coccineo. *Pluk. alm.* 14. t. 257. f. 1.

Habitat in Mexico. h

9. *HIBISCUS* foliis ovato-lanceolatis: superne inciso-ser-*fractus*.
ratis, caule arboreo. *Hort. cliff.* 350. *Hort. ups.* 205.

Alcea arborescens syriaca. *Baub. pin.* 316.

Alcea arborescens. *Cam. hort.* t. 3. 4.

Habitat in Syria. h

10. *HIBISCUS* foliis quinquefido-palmatis, caule actu-*ficulneus*.
leato. *Hort. cliff.* 498. *Roy. lugdb.* 359. *Fl. zeyl.* 269.*

Ketmia zeylanica, fici folio, perianthio oblongo integro.

Dill. elzb. 190. t. 157. f. 190.

Habitat in Zeylona.

11. *HIBISCUS* foliis serratis: inferioribus ovatis indivi-*Sabdariffa*.
fis; superioribus tripartitis, caule aculeato.

Hibiscus inermis, foliis serratis: inferioribus ovatis integris; superioribus trilobis. *Hort. cliff.* 350. *Fl. zeyl.*

262. *Roy. lugdb.* 359.

Athæa indica, gossypii folio, acetosæ sapore. *Raj. bist.* 1900.

Alcea acetosa indica mitis, foliis superioribus gossypil in modum tripartito-divisis. *Pluk. alm.* 15. t. 6. f. 2.

β. *Hibiscus inermis*, foliis serratis: infimis indivisis; mediis tripartitis; summis quinquepartitis. *Roy. lugdb.* 359.

Alcea indica, magno flore. *Baub. pin.* 317.

Alcea americana. *Clus. bist.* 2. p. 26.

γ. *Hibiscus* foliis palmato-digitatis quinquepartitis: laciniis lanceolatis, caule aculeato. *Fl. zeyl.* 264.

Alcea spinosa maderas. saporis acidi, foliis profunde laciniatis cannabis in modum pentaphyllea. *Pluk. alm.*

15. t. 5. f. 4.

Narinam-poulli. *Rheed. mal.* 6. p. 75. t. 44.

δ. *Alcea benghalensis spinosissima*, acetosæ sapore, flore luteo pallido: umbone purpurascente. *Comm. hort.* 1.

p. 35. t. 18.

Ketmia indica, foliis digitatis, flore magno sulphureo: umbone atro-purpureo, petiolis spinosis. *Ebret. t.*

6. f. 1.

Habitat in India, ☉

γ *Arcliquis diversa foliis magis palmatis; at Color florum,*
X x 4 *Aculei*

Aculei erecti, Locus, Sapor & omnia indicant has vel loco, vel hybrida partu degenerasse.

- surattensis.* 12. **HIBISCUS** recurvato-aculeata, foliis trilobis, calycibus exterioribus appendiculatis, stipulis semicordatis.

Habitat in India.

Habitus & statura antecedentis. Folia inferiora ovato-subtriloba, acuta, serrata: superiora hastato-triloba, latiora fere quam longa. Aculei recurvi & pili recti sparsi per caulem, ramos, petiolos, peduncullos. Stipulae dimidiato-cordatae s. parva stipularum cordata. Calyx pilis strigosus: exterior valde singularis, cujus radii in medio extrorsum aucti foliolo parvo ovato.

- Manihot.* 13. **HIBISCUS** foliis palmato-digitatis septempartitis. *Hort. cliff.* 350. *Hort. mps.* 206. *Roy. lugdb.* 358. *Ketmia* folio manihot ferrato, flore amplo sulphureo. *Dill. elth.* 189. t. 156. f. 189.

Alcea finica, manihot stellato folio, capsula longa pilosa pyramidata quinquefariam divisa. Pluk. amalib. 7. p. 355. f. 2.

Habitat in Indiis. b

- Abelmoschus.* 14. **HIBISCUS** foliis subpeltato-cordatis septemangulatis ferratis hispidis. *Hort. cliff.* 349. *Hort. mps.* 206. *Fl. zeyl.* 261. * *Mat. med.* 342. *Roy. lugdb.* 358.

Alcea hirsuta, flore flavo, semine moschato. Marsgr. bras. 45. t. 45.

Alcea aegyptiaca villosa. Baub. pin. 317.

Flos moschatus. Mcr. surin. 42. t. 42.

Cattu-gasturi. Rheed. mal. 2. p. 71. t. 38.

Habitat in Indiis. b

- stultentus.* 15. **HIBISCUS** foliis quinquepartito-pedatis, calycibus interioribus latere rumpentibus.

Alcea americana annua, flore albo (flavo potius) maximo, fructu pyramidalis sulcato. Comm. hort. 1. p. 37. t. 19. *Raj. suppl.* 518.

Alcea maxima, malvae rosae folio, fructu decagono recto crassiore brevioris esculento, Sloan. jam. 98. *biff.* 1. p. 223. t. 133. f. 3.

Habitat in Indiis.

- vitisfolia.* 16. **HIBISCUS** foliis quinquangularibus acutis serratis, caule inermi, floribus pendulis. *Fl. zeyl.* 265. * *Althaea indica, vitis folio, flore amplo flavo pendente. Herm. lugdb.* 26. t. 28.

Katou-

Katou-Beloeren. *Rheed. mal.* 6. p. 79. t. 46.

Habitat in India. ☉

17. **HIBISCUS** foliis cordato-hastatis, pedunculis alter-*seylonicus*.
nis unifloris geniculatis. *Fl. zeyl.* 266. *

Habitat in Zeylona. ☉

18. **HIBISCUS** foliis inferioribus cordatis acuminatis ter-*virginicus*.
ratis; superioribus hastatis. *Gron. virg.* 171.

Alcea maderaspatana, trilobis f. hastatis foliis glabris, pericarpio tantummodo villosa. *Pluk. aim.* 15. t. 127.
f. 2. *forte?*

Habitat in Virginie paludosis salis.

Caulis viridis, erectus, angustior. Folia caulina cordato-oblonga, demum hastata, viridia, subtomentosa, trinervia, tenuissime serrata. Pedunculi uniflori: ex alis solitarii, in summitate subracemosi, nudi.

19. **HIBISCUS** foliis inferioribus cordatis angulatis, su-*Pentacarpus*.
perioribus subhastatis, floribus subnutantibus, pitillo cernuo.

Hibiscus foliis cordatis angulatis serratis, stipulis setaceis divaricatis. *Hort. cliff.* 350. *Roy. lugdb.* 359.

Ketmia palustris minor, folio angusto, flore parvo purpurascente, fructu depresso pentagono. *Zannich. venet.* 155. t. 91.

Habitat in paludosis Venetie. 2

Planta antecedenti simillima; sed fructu diversa. Caulis tripedalis, angustus. Folia caulina coraeo-oblonga; superiora hastata, viridia, subtrinervia, subtomentosa, tenuissime serrata. Pedunculi axillares in summitate plantæ, solitarii, nudi, petiolo longiores, uniflori. Calyx inferior patens, radiis subulatis, vix longitudine interioris. Corolla patens, dilute rubra, subnutans. Stamina lutea tota; antheræ saturatiæ. Stylus cernuus cum staminibus, albus; Stigmata 5, capitata, incarnata.

20. **HIBISCUS** foliis tripartitis incisis, calycibus inflatis. *Trionum*,
Hort. upf. 206.

Trionum. *Hort. cliff.* 349. *Roy. lugdb.* 358.

Alcea vesicaria. *Bauh. pin.* 317.

Alcea peregrina solisequa. *Lob. ic* 656.

- β. *Ketmia vesicaria africana.* *Tournef. inst.* 101.

Alcea vesicaria capitis b. *spei. Moris. præl.* 227. *hist.* 2. p. 533.

Habitat in Italia, Africa. ☉

X x 5

PEN-

746

PENTAPETES.

- phoenicea*. 1. PENTAPETES foliis hastato-lanceolatis serratis.
Blattaria zeylonica, flore amplo coccineo. *Comm-hort.*
1. p. 11. t. 6.
Alcea indica, lucido hastato folio, flore blattariae phoeniceo. *Raj. suppl.* 523. *Pluk. phyt.* t. 255. f. 3.
Alcea fruticosi pentaphylloidis æmula, floribus amœnissime rubellis, calyce producto. *Pluk. alm.* t. 126. f. 4. mala.
Habitat in India. ☉
Involucrum uniflorum deciduum. Stamina 15. inter singula tria Filamentum petaloideum lanceolatum, coloratum, longitudine corollæ. Cotyledones bipartiti, uti in *Convolvulis*. Pedunculi biflori.
- suberifolia*. 2. PENTAPETES foliis ovatis repandis. *Amœn. acad.*
1. p. 407.
Pentapetes. *Fl. zeyl.* 250.
Pterospermadendrum suberis folio anguloso subtus incano, floribus albis, *Amm. act.* 8. p. 215. t. 14.
Habitat in India. ☿
- acerifolia*. 3. PENTAPETES foliis cordatis repandis. *Amœn. acad.*
1. p. 407.
Pterospermadendron foliis aceris, flore fructuque majore. *Amm. act.* 8. p. 216. t. 16. 17.
Habitat in India.

747

STEWARTIA.

- Malacodendron*. 1. STEWARTIA. *Act. upf.* 1741. p. 79. t. 2. *Catesb. car.* 3. p. 13. t. 13.
Malacodendron. *Mitch. gen.* 16.
Habitat in Virginia. ☿
Folia *Cerasi*, acutissime serrata, subtus villosa.

748

CAMELLIA.

- japonica*. 1. CAMELLIA
Tsubakki montanus s. sylvestris, flore roseo simplici.
Kempf. amœn. 850. t. 851.
β. Tsubakki hortensis, flore pleno maximo rosæ hortensis.
Kempf. amœn. 852.
Habitat in Japonia, China. ☿

Clas-

Classis XVII.

DIADELPHIA.

HEXANDRIA.

FUMARIA.

749

* *Corollis bicalcaratis.*

1. FUMARIA scapo nudo. *Hort. cliff.* 351. * *Gron. Cucullaria, virg.* 171. *Roy. lugdb.* 393.

Fumaria tuberosa insipida. *Corn. canad.* 127.

Fumaria siliquosa, radice grumosa, flore bicorporeo ad labia conjuncto, virginiana. *Pluk. alm.* 162. t. 90. f.

3. *Raj. suppl.* 475.

Cucullaria. *Juss. act. paris.* 1743.

Habitat in Virginia, Canada. 2

Radix tuberosa. Folium radicale tricompositum. Scapus nudus, Racemo simplici; bractea vix ulla; Nectarium duplex corollam basi bicornem efficiens.

2. FUMARIA floribus postice bilobis, caule folioso. *Spettabilis, Habitat in Sibiria. D. Demidoff.*

Planta eximia floribus speciosissimis, maximis. Habitus Fumariae bulbosae, sed majora omnia. Rami ex alis rariore. Caulis erectus. Racemus absque bracteis. Corollae magnitudine extimi articuli pollicis, pome in duos lobos aequales, rotundatos divisa.

* *Corollis unicalcaratis.*

3. FUMARIA caule simplici, bracteis longitudine flo- bulbosa, rum.

Fumaria caule simplicissimo diphylo, floribus calyce destitutis. *Hort. cliff.* 351. *Vir. cliff.* 70. *Fl. succ.* 585.

Mat. med. 344. *Roy. lugdb.* 394.

- a Fumaria bulbosa, radice cava, major. *Bauh. pin.* 142. cava.

Pistolochia. *Fuchf. hist.* 91.

- β Fumaria bulbosa, radice non cava, minor. *Bauh. pin.* intermedia, 244.

- γ Fumaria bulbosa, radice non cava, major. *Bauh. pin.* solida.

144.

Habitat in Europae nemoribus & umbrosis. 2

Conjunctae has plantas specie: Radix carnosa, *simphon,*
Gem-

Gemmæ radicales duæ, trivalves, Caulis simplex, subangulatus, unico rario, Foliorum structura decomposita: Racemus bracteis immixtus, terminalis, foliarius; Corolla, Calyx minimus. Semina, Stamina Siliquæ structura; Tempus vernum. Species reliquæ propriis notis distinctissimæ.

sempervirens.

4. *FUMARIA* siliquis linearibus paniculatis, caule erecto. *Hort. nosf.* 207.

Fumaria caule erecto ramoso, siliquis filiformibus corymbosis. *Hort. cliff.* 352. *Roy. lugdb.* 394.

Fumaria siliquosa sempervirens. *Corn. canad.* 57. t. 57. *Morif. b. fl.* 2. p. 259. f. 3. t. 12. f. 1.

Habitat in Canada, Virginia. ☉

capnoides.

5. *FUMARIA* siliquis linearibus tetragonis, caulibus diffusis acutangulis.

Fumaria caule ramoso diffuso, siliquis oblongis, radice perenni. *Hort. cliff.* 352. *Roy. lugdb.* 394.

Fumaria lutea. *Baub. pin.* 143.

Fumaria lutea montana. *Dalech. hist.* 1294.

Fumaria tingitana, radice fibrosa, perennis, flore ex albo flavescente, siliquis curtis. *Pluk. alm.* 262. t. 90. f. 2.

Habitat in Gallia, Italia, Mauritania. ☉

hennepbylla.

6. *FUMARIA* foliis triternatis: foliolis cordatis.

Fumaria hispanica saxatilis, foliis amplioribus cordiformibus, semine compresso. *Tournef. inst.* 422.

Fumaria hennepbyllos hispanica saxatilis. *Bocc. mus.* 2. p. 83. t. 73. *Raj. suppl.* 475. *Barr. ic.* 42.

Fumaria radice fibrosa, foliis ad petiolum sinuatis crassioribus. *Pluk. alm.* 162.

Habitat in Hispaniæ, Siciliæ saxosis.

officinaria.

7. *FUMARIA* pericarpis monospermis racemosis, caule diffuso.

Fumaria pericarpis monospermis. *Hort. cliff.* 252. *Fl. succ.* 584. *Mat. med.* 343. *Roy. lugdb.* 394. *Hall. herb.* 605.

Fumaria officinarum & dioscoridis. *Baub. pin.* 143.

Fumaria. *Fuchf. hist.* 338. *Cam. epit.* 890.

Habitat in Europæ agris, cultis. ☉

spicata.

8. *FUMARIA* pericarpis monospermis spicatis, caule erecto, foliolis filiformibus. *Sauv. monsp.* 263.

Fumaria minor tenuifolia. *Baub. pin.* 143. *Segn. ver.* 2. p. 111.

Fuma-

Fumaria tenuifolia erecta hispanica purpurea. Barr. ic. 41.

Fumaria minor tenuifolia præcox, semine lini. Moris. hist. 2. p. 262. f. 3. t. 12. f. 13.

Capnos tenuifolia. Clus. hist. 2. p. 208.

Habitat in Hispania, G. Narbonensi, Veronæ ad agros, vias.

Varietates, qui statuas hanc cum sequente esse Fumariæ officinalis vix erret, dictitante fructu. An hybrida?

Spica ex flosculis arte congestis est.

9. **FUMARIA** pericarpiis monospermis racemosis, foliis *capreolata*, scandentibus subcirrhis.

Fumaria viticulis & capreolis plantis vicinis adhærens.

Bauh. pin. 143.

Fumaria major scandens, flore pallidiore. Raj. hist. 405

Habitat in G. Narbonensi. Anglia. ☉

Pedunculi partiales recurvati sunt, & Foliola extrema in cirrhis transeunt.

10. **FUMARIA** filiquis linearibus, foliis cirrhiferis. *claviculata,*

Fumaria foliis cirrhiferis, floribus spicatis. Hort. cliff.

*351. * Roy. lugdb. 394. Gort. gelr. 413.*

Fumaria claviculis donata. Bauh. pin. 143. Moris. hist.

2. p. 266. f. 3. t. 12. f. 3.

Fumaria alba latifolia. Raj. angl. 3. p. 335.

Habitat in Angliæ locis uliginosis saxosis.

11. **FUMARIA** filiquis globosis inflatis. *Hort. ups. 207. vesicaria.*

Fumaria foliis cirrhiferis, filiquis ovatis inflatis pendulis. Hort. cliff. 351. Roy. lugdb. 394.

Fumaria alba vesicaria, capreolis donata, sub exitum autumnum florens, æthiopica. Pluk. alm. 400. t. 355. f. 3.

Cysticapnos africana scandens. Boerb. lugdb. 1. p. 310. t. 310.

Habitat in Æthiopia. ☉

OCTANDRIA.

POLYGALA.

* *Cristati (flores appendice penicelliformi.)*

1. **POLYGALA** floribus cristatis spicatis, caule herbaceo *incarnata*, ramoso erecto, foliis alternis subulatis.

Polygala marilandica, foliis tenuissimis, spica parva compacta in summo caule flosculis carneis. Raj. suppl. 639.

Habitat in Virginia, Canada. ☉

Caulis pithameus, pedalis, erectus, niliformis, angulatus. Ramis communiter tres, in alis summis, longi, angusti. Folia subulata, plana, acutissima, remota, sæpius vix in caule octo.

octo. Spicæ terminales, solitariae, oblongae, vividissime incarnatae; Corollae, quam alae calycinae triplo longiores, instructae pennis multifido, dilatato, incarnato, uti corollae erecto.

- vulgaris.* 2. **POLYGALA** floribus cristatis racemosis, caulibus herbaceis simplicibus procumbentibus, foliis lineari-lanceolatis. *Aman. acad. 2. p. 136.*
Polygala foliis lineari-lanceolatis, caulibus diffusis herbaceis. Vir. cliff. 70. Hort. cliff. 352. Fl. suec. 586. Roy. lugdb. 393.
Polygala major. Bauh. pin. 215.
Polygala vulgaris major. Clus. hist. 1. p. 324. Vaill. parif. 161. t. 32. f. 1.
 3. *Polygala vulgaris. Bauh. pin. 215.*
Polygala foliis imis subrotundis, superioribus angustis acutis. Hall. herb. 606.
 4. *Polygala minor, foliis circa radicem rotundiusculis. Vaill. parif. 161. t. 32. f. 2.*
Habitat in Europæ pratis, & pascuis siccis. &
- monspeliaca.* 3. **POLYGALA** floribus cristatis racemosis, caule erecto, foliis lanceolato-linearibus acutis. *Sauv. monsp. 53.*
Polygala annua erecta angustifolia, floribus coloris obsoleti carneis lineis virgatis. Magn. monsp. 208.
Polygala vulgaris coloris obsoleti, foliis angustissimis. Bauh. hist. 3. p. 338. Raj. hist. 1336.
Habitat Monspeli in collibus sterilibus. ☉
- sibirica.* 4. **POLYGALA** floribus cristatis, racemo laterali nudo, caulibus herbaceis, foliis lanceolatis.
Habitat in Sibiria. D. Gmelin.
Habitus antecedentis. Folia inferiora ovata, superiora lanceolata. Racemus unus (raro duo), lateralis, latus, absque bracteis, ad exortum pedicellorum partialium.
- bracteolata.* 5. **POLYGALA** floribus cristatis, racemo laterali, bracteis triphyllis, foliis lanceolatis, caule suffruticoso.
Polygala floribus cristatis racemosis, carina cristis longiore, caule suffruticoso, foliis lineari-lanceolatis. Aman. acad. 2. p. 137.
Polygala africana glabra, florum galea binis cristis simbratis ornata, ex involucrio magno dipetalo erumpente. Pluk. alm. 300. t. 53. f. 2.
Polygala frutescens angustifolia ramosa, floribus in summitate velut umbellatis. Raj. suppl. 640.

3. Poly

- β. *Polygala floribus cristatis racemosis: carina cristis brevior, caule infruticoso, foliis lineari-subulatis. Amæn. acad. 2. p. 137.*
Polygala foliis lineari-subulatis. Hort. cliff. 353.
Polygala trutecens, foliis linearibus, flore majore purpureo. Burm. afric. 202. t. 73. f.
- γ. *Polygala floribus cristatis racemosis, caule erecto suffruticoso simplicissimo, foliis subulatis. Amæn. acad. 2. p. 137.*
Polygala foliis linearibus acutis, flore purpureo minore. Burm. afric. 203. t. 73. f. 3.
Habitat Æthiopia. h
- δ. **POLYGALA** floribus cristatis alternis, caule erecto *scabra*, suffruticoso ramoso, foliis linearibus obtusis scabris.
Amæn. acad. 2. p. 138.
Polygala æthiopica, angustis hirsutis foliis, flore obsolete purpureo. Pluk. mant. 153. t. 404. f. 6.
Polygala folio lineari-obtuso, flore albente minimo. Burm. afric. 204. t. 73. f. 4.
Habitat in Æthiopia. h
- ε. **POLYGALA** floribus cristatis carina lunulata, caule *myrtifolia*, fruticoso, foliis lævibus oblongis obtusis. *Amæn. acad. 2. p. 138.*
Polygala foliis lanceolatis obtusis, caule frutescente. Hort. cliff. 353.
Polygala arborea myrtifolia, floribus albis: intus purpureis. Com. hort. 1. p. 87. t. 46. Raj. suppl. 640.
Polygala arborea, myrtilli subrotundis foliis, fructu magno tordylii. Pluk. mant. 153. t. 437. f. 4.
Polygala frutescens, foliis oblongis glabris, flore purpureo. Burm. afric. 200. t. 13. f. 1.
Habitat in Æthiopia. h
** Imberbes (flores absque penicillo carinatis) frutescentes.*
- ζ. **POLYGALA** floribus imberbibus lateralibus solitariis, *Penæa*, caule arboreo, foliis obtusis petiolatis. *Amæn. acad. 2. p. 140.*
Penæa arborescens, buxi folio aspero. Plum. spec. 22.
Habitat in America meridionali h
- η. **POLYGALA** floribus imberbibus racemosis, caule ar- *diversifolia*, boreo, foliis senioribus oblongo-ovatis, recentibus subovatis. *Amæn. acad. 140.*
Polygala foliis ramorum provectorum oblongo-ovatis,

tis, teneſſorum ſubovatis, caule arboreo. *Hort. cliff.* 353. *

Habitat in America calidiore. ♀

- obinervis.* 10. POLYGALA floribus imberbibus ſpicatis axillaribus, caulibus ſuffruticoſis, foliis ovalibus.

Habitat in India. ♀

Cauleſ pedales, decumbentes, lignoſi, non glabri. Folia ovalia, integerrima, magnitudine extimi pollicis, ſubpetiolata, nuda, obtuſa cum mucrone. Rami laterales, rariſſimi. Spicæ laterales, foliis ſæpe longiores, virides.

- Chamæbuxus.* 11. POLYGALA floribus imberbibus ſparſis: carina apice ſubrotundo, caule fruticoſo, foliis lanceolatis. *Amœn. acad.* 2. p. 140.

Chamæbuxus flore coluteæ. *Banb. pin.* 471.

Anonymos flore coluteæ. *Cluſ. hiſt.* 1. p. 105.

Habitat in Aultriæ, Helvetiæ, Alſatiæ montanis. ♀

- ſpiñoſa.* 12. POLYGALA floribus imberbibus lateralibus, caule arboreo ſpiñoſo, foliis ovalibus mucronatis. *Amœn. acad.* 2. p. 141. *

Habitat in Æthiopia. ♀

- Heiſteria.* 13. POLYGALA floribus imberbibus lateralibus, caule arboreſcente, foliis triquetris mucronatis.

Polygala aſtragaloides fruteſcens. *Breyn. prodr.* 2. p. 85.

Suffrutex ericæformis, flore rubro elegantiffimo. *Seb. theſ.* 2. p. 63. t. 62. f. 8.

Thymbra capenſis, nepetæ theophrasti foliis aculeatis, flore parvo purpureo. *Pluk. alm.* 366. t. 229. f. 5.

Heiſteria. *Hort. cliff.* 352. * *Roy. lugdb.* 393.

Habitat in Æthiopia. ♀

* *Imberbes herbaceæ, caule ſimpliciſſimo.*

- Senega.* 14. POLYGALA floribus imberbibus ſpicatis, caule erecto herbaceo ſimpliciſſimo foliis lato-lanceolatis. *Amœn. acad.* 2. p. 139.

Polygala caule ſimplici erecto, foliis ovato-lanceolatis alternis integerrimis, racemo terminali erecto. *Gran. virg.* 80. * *Mat. med.* 345.

Plantula marilandica, caule non ramoſo, ſpica in ſtigio ſingulari e thoſculis albis compoſita. *Raj. ſuppl.* 640.

Habitat in Virginia, Penſylvauia, Marilandia, &

15. POLYGALA floribus imberbibus oblongo-capitatis, *lucra*.
caule erecto herbaceo simplicissimo, foliis lanceolatis
acutis. *Amoen. acad. 2. p. 140.*
Polygala f. Flos Ambervalis floribus luteis in capitulum oblongum congestis. *Pluk. alm. 301. Raj. suppl. 639.*
Polygala virginiana, flore luteo capitato. Pet. ficc. 58.
Habitat in Virginia. ☉
16. POLYGALA floribus imberbibus globoso-capitatis, *viridescens*,
caule erecto herbaceo simplicissimo, foliis lanceolatis
obtusiusculis. *Amoen. acad. 2. p. 140.*
Polygala foliis lanceolatis alternis, caule simplicissimo,
corymbo terminali capitato. *Gron. virg. 80?*
Habitat in Virginia ☉
Caulis spithameus f. pedalis, angulatus, absque ramis, erectus. Folia alterna, linearia f. linearilanceolata, levia, integerrima. Capitulum terminale, subglobosum, ex albo viride, cum obsolete rubri mixtura in alis florum.
17. POLYGALA floribus imberbibus, caule herbaceo triflora.
erecto, foliis linearibus alternis, pedunculis subtrifloris. *Fl. zeyl. 269.*
Habitat in Zeylona. ☉
18. POLYGALA floribus imberbibus, pedunculis multi-glancoideis.
floris lateralibus, caulibus diffusis herbaceis, foliis acutis. *Fl. zeyl. 270. **
Habitat in Zeylona. ☉
- * *Imberbes herbaceæ ramosæ.*
19. POLYGALA floribus imberbibus, capsulis ciliato-ciliatis.
dentatis, caule herbaceo erecto. *Fl. zeyl. 268.*
Habitat in India. ☉
Variet Foliis cordatis, Caulibus semipedalibus, ramosis.
20. POLYGALA floribus imberbibus, pedunculis squarrosius, caule herbaceo ramoso erecto.
Habitat in Virginia Kalm. ☉
Caulis spithameus f. pedalis, firmus, angulatus, ramosus, erectus, ramis caule altioribus. Folia alterna, angustolanceolata. Spicæ terminales, laxæ, sanguineæ, ovatae, decidunt enim flosculi inferiores maturi, dum novelli accrescunt unde, Pedunculus lacerus f. squarrosus.

- verticillata.* 21. **POLYGALA** floribus imberbibus, spicis floribus remotis, foliis linearibus verticillatis, caule herbaceo ramoso.
Polygala caulibus filiformibus, foliis linearibus alternis pedunculis spicatis. *Gron. virg.* 172.
Polygala foliis imberbibus spicatis, caule erecto herbaceo filiformi ramoso, foliis linearibus. *Aman. acad.* 2. p. 159.
Polygala mariana quadrifolia minor, spica parva albicante. *Pluk. mant.* 153. t. 438. f. 4.
Polygala quadrifolia minima marilandica, spicis florum parvis albescentibus. *Raj. suppl.* 639.
Habitat in Virginia. ☉
 Folia sepius quina ad gemicula, interdum alterna. Spicæ albae angustissima flosculis remotis.
- cruciata.* 22. **POLYGALA** floribus imberbibus, foliis quaternis. *Aman. acad.* 2. p. 138.
Polygala foliis quaternis. *Gron. virg.* 80.
Polygala quadrifolia f. cruciata, floribus ex viridi rubentibus in globum compactis. *Pluk. alm.* 301.
Habitat in Virginia.

DECANDRIA

ERYTHRINA.

- herbacea.* 1. **ERYTHRINA** foliis ternatis, caule simplicissimo infermi. *Hort. cliff.* 354. *Roy. lugdb.* 373.
Coral carolinensis, hastato folio. *Dill. elieb.* 107. t. 90. f. 106.
Coralodendron humile, spica florum longissima, radice crassissima, *Catesb. car.* 49. t. 49.
Habitat in Carolina, Missipi. 2
- coralodendron.* 2. **ERYTHRINA** foliis ternatis, caule arboreo aculeato. *Hort. cliff.* 354. *Hort. mps.* 207. *Fl. zeyl.* 275. *Roy. lugdb.* 373.
- occidentalis.* *Ceratia* f. *Siliqua* sylvestris spinosa arbomindica. *Banks. pin.* 402.
Coral arbor americana. *Comm. hort.* 1. p. 211. t. 108
- orientalis.* 3. *Mouricou.* *Rheed. mal.* 6. p. 13. t. 7.
Habitat in Indiis. ☿
 Est quædam differentia inter occidentalem & orientalem, sed vix videtur sufficiens pro specie distincta; aculei enim in orientali nigrescentes sunt.

3. ERY.

3. ERYTHRINA foliis pinnatis, leguminibus membra- *Piscipula*
 uaceo-tetragonis. †
 Coral arbor polyphylla non spinosa, fraxini folio, siliqua
 alis foliaceis exstantibus rotæ molendinariz fluviati-
 lis aucta. *Sloan. jam. 143. hist. 2. p. 39. t. 176. f.*
45. Raj. dendr. 108.
Habitat in America calidiore. b

SECURIDACA. 752

1. SECURIDACA. *columbina.*
Spartium scandens, citri foliis, floribus albis ad nodos
confertim nascentibus. Plum. spec. 19.
Habitat in America meridionali. b

BORBONIA. 753

1. BORBONIA foliis lanceolatis trinerviis integerrimis. *trinervia.*
Borbonia foliis lanceolatis acuminatis trinerviis. Hort.
cliff. 494.
Frutex æthiopicus, rusci angusto & minore folio. Pluk.
alm. 158. t. 414. f. 1.
Habitat in Æthiopia. b
2. BORBONIA foliis lanceolatis multinerviis integerrimis. *lanceolata*
Frutex æthiopicus, rusci angusto & minore folio. Pluk.
alm. 158. t. 297. f. 4.
Genista africana frutescens, rusci nervosis foliis. Raj.
dendr. 107.
Habitat in Æthiopia. b
3. BORBONIA foliis cordatis multinerviis integerrimis. *cordata.*
Borbonia foliis cordatis acuminatis integerrimis multi-
nerviis. Hort. cliff. 494. Roy. lugdb. 372.
Spartium africanum frutescens, ruscitolio caulem am-
plexante. Comm. hort. 2. p. 195. t. 98. Raj. dendr. 107.
Genista africana frutescens, rusci foliis nervosis, flore
luteo. Seb. thes. 1. p. 38. t. 24. f. 3.
Frutex æthiopicus, foliis rusci, floribus papilionaceis se-
rica lanugine fusca villolis. Pluk. alm. 159. t. 297. f. 3
Habitat in Æthiopia. b
4. BORBONIA foliis ovatis tomentosis. *tomentosa.*
Genista africana tomentosa, folio cochleariformi, flore
luteo. Seb. thes. 1. p. 38. t. 24. f. 1.
5. BOR-
 5. BOR-
Genista africana arborescens, folio argentea lanugine ve-
stito. Seb. thes. 1. p. 38. t. 24. f. 2.
Habitat in Æthiopia. b

708. DIADELPHIA DECANDRIA.

crenata.

5. BORBONIA foliis cordatis multinerviis crenatis.
Frutex æthiopicus leguminosus, foliis rusci majoribus
ambitu spinulis fimbriatis. *Pluk. alm.* 159.
Planta leguminosa æthiopica, foliis rusci. *Breyn. sem.*
69. t. 28.
Habitat in Æthiopia. 5

754

SPARTIUM

* *Foliis simplicibus.*

fruticent.

1. SPARTIUM ramis oppositis teretibus apice floriferis
foliis lanceolatis. *Hort. cliff.* 356. *Hort. ups.* 208.
Roy. lugab. 370. *Sauv. monsp.* 60.
Spartium arborescens, seminibus lenti similibus. *Bauh.*
pin. 396.
Spartium macrolobium. *Renalm. spec.* 34. t. 33.
Habitat in G. Narbonenti, Italia, Sicilia, Turcia. 5

radicans.

2. SPARTIUM ramis oppositis angulatis, foliis opposi-
tis subulatis.
Spartium æquicolorum minimum montanum triphyllum
Col. cephr. 1. p. 295. t. 294.
Genista radiata f. stellaris. *Bauh. hist.* 1. p. 399.
Habitat in Italia. 5

*monosper-
mum.*

3. SPARTIUM ramis angulatis, racemis lateralibus, fo-
liis lanceolatis. *Hort. cliff.* 356. *Roy. lugab.* 370.
Sauv. monsp. 60.
Spartium alterum monospermum, semine reni simili
Bauh. pin. 396.
Spartium hypolphærolobium. *Ren. spec.* 35. t. 33.
β. Spartium tertium, flore albo. *Bauh. pin.* 396.
Spartium 3. hispanicum. *Clus. hist.* 1. p. 103.
Habitat in Hispaniæ sterilibus. 5

Scorpius

4. SPARTIUM ramis spinosis patentibus, foliis ovatis
Genista-spartium spinosum majus, flore luteo. *Bauh.*
pin. 394.
Aspalathus secunda monspeliensis. *Bauh. hist.* 1. p. 402.
β. Genista-spartium spinosum majus secundum, flore pal-
lido. *Bauh. pin.* 394.
γ. Genista-spartium spinosum majus tertium hirsutum
Bauh. pin. 394.
Aspalathus alter 111. *Clus. hist.* 1. p. 106.
Habitat in Hispania, G. Narbonensi. 5

• *Folia*

* *Foliis alternis.*

5. SPARTIUM foliis solitariis ternatisque, ramis sexan-
gularibus apice floriferis.

Spartium orientale, filiqua compressa glabra & annula-
ta. *Tournef. cor.* 44.

Habitat in Oriente, b

Anguli tres decurrunt per ramos a singulo folio.

6. SPARTIUM foliis ternatis: foliolis involutis, cauli-
bus inermibus prostratis glabris, leguminibus scabris.

Cytisus ramis humifusis albidis, floribus capitatis termi-
nalibus, foliolis ovalibus glabris aggestis. *Sauv. monsp.*

190

Cytisus foliis incanis angustis quasi complicatis. *Bauh.*
pin. 390. *Guett. stamp.* 2. p. 417.

Cytisus montis calcaris. *Bauh. hist.* 1. p. 370.

Cytisus 11. *Clas. hist.* 1. p. 94.

Habitat in Hispania, Gallia australiore. b

7. SPARTIUM foliis ternatis solitariisque, ramis iner-
mibus angulatis. *Hort. cliff.* 356. *Fl. succ.* 589. *Roy.*

lugdb. 370. *Gort. gelr.* 415.

Genista angulosa & scoparia. *Bauh. pin.* 395.

Genista. *Dod. pempt.* 761.

Habitat in Europæ australioris arenosis. b

8. SPARTIUM foliis ternatis, ramis angulatis spinosis. *spinosum.*
Hort. cliff. 356. *Hort. upf.* 208. *Roy. lugdb.* 370.

Sauv. monsp. 191.

Acacia trifolia. *Bauh. pin.* 392.

Aspalathus secunda trifolia. *Bauh. hist.* 1. p. 375.

Habitat in Europæ australis asperis maritimis. b

GENISTA.

755

* *Inermes.*

1. GENISTA foliis ternatis. *Hort. cliff.* 355. *Mat. med. canariens.*
347. *Roy. lugdb.* 371.

Cytisus minoribus foliis, ramulis tenellis villosis. *Bauh.*
pin. 390.

Cytisus 1. *Clas. hist.* 1. p. 94.

2. Cytisus canariensis sempervirens & incanus. *Comm.*
hort. 2. p. 103. t. 52.

Cytisus canariensis, flore candido & citrino. *Sch. thes.*
2. p. 6. t. 4. f. 6. 7.

Y 7 3

Cy

710 DIADELPHIA DECANDRIA.

Cytisus canariensis microphyllus angustifolius prorsus incanus. *Pluk. alm.* 128. t. 277. f. 6.
Habitat in Hispania, Canariis. ♀

- sagittalis.* 2. GENISTA ramis ancipitibus articulatis, foliis ovato-lanceolatis. *Hort. cliff.* 355. *Roy. lugab.* 371.
Genista herbacea f. *Chamæ-spartium*. *Bauh. hist.* 1. p. 393.
Chamæ-Genista sagittalis. *Bauh. pin.* 395.
Chamæ-Genista sagittalis pannonica. *Cam. hort.* t. 13.
Habitat in Germania, Gallia. 2
- tridentata.* 3. GENISTA ramis triquetris subarticulatis, foliis tricuspidatis.
Genistella fruticosa lusitanica latifolia. *Tournef. inst.* 646.
β. *Genista fruticosa lusitanica angustifolia.* *Tournef. inst.* 646.
Habitat in Lusitania. Læst. ♀
Præcedenti valde affinis, at majora omnia ejus articuli ramorum in folium laterale terminali, non integri sed tricuspidati.
- tinctoria.* 4. GENISTA foliis lanceolatis, ramis striatis teretibus erectis. *Hort. cliff.* 355. *Fl. succ.* 587. *Mat. med.* 347. *Roy. lugab.* 371. *Gort. gelr.* 416.
Genista tinctoria germanica. *Bauh. pin.* 395.
Genista tinctoria vulgaris. *Clus. hist.* 1. p. 101.
Habitat in Germania, Anglia. ♀
- pilosa.* 5. GENISTA foliis lanceolatis obtusis, caule tuberculato decumbente. *Hort. cliff.* 355. *Fl. succ.* 588. *Roy. lugab.* 371.
Genista ramosa, foliis hyperici. *Bauh. pin.* 395.
Chamægenista prima. *Clus. hist.* 1. p. 103.
Habitat in Pannonia, G. Narbonensi, Germania.
- * *Spinæ laterales.*
- argentea.* 6. GENISTA spinis simplicibus, ramis floriferis inermibus, foliis lanceolatis. *Hort. cliff.* 355. *Roy. lugab.* 371.
Genista minor asphaltoides. *Bauh. pin.* 395. *prodr.* 157.
Genistella *Dod. prompt.* 670.
Habitat in Angliæ ericetis humidiusculis. ♀
- germanica.* 7. GENISTA spinis compositis, ramis floriferis inermibus, foliis lanceolatis. *Roy. lugab.* 371. *Gort. gelr.* 417.
Geni-

DIADELPHIA DECANDRIA 711

Genista spinosa minor germanica. *Bauh. pin.* 395.

Genistella spinosa. *Riv. tetr.* 67.

Habitat in Germania. ♀

8. GENISTA spinis decompositis, ramis floriferis iner- *hispanica.*
mibus: foliis linearibus pilosis.

Genista spinosa minor hispanica villosissima. *Bauh. pin.*

395.

Genistella monspeliaca spinosa. *Bauh. prodr.* 157.

Genistella, montis ventosi, spinosa. *Bauh. hist.* 1. p.

400.

Habitat in Hispania, G. Narbonensi. ♀

Vix a precedenti sufficienter distincta.

9. GENISTA caule aphylo, spinis decussatis. *lusitanica.*

Genista-spartium spinosum minus. *Bauh. pin.* 394.

Scorpius secundus. *Clus. hist.* 1. p. 107. *Bauh. hist.* 1.

p. 403.

Habitat in Lusitania, Hispania. ♀

ASPALATHUS. 756

1. ASPALATHUS foliis confertis subulatis mucronatis *astroites.*
glabris, caule villosa, floribus spartis.

Genista astroites, juniperinis pungentibus foliis, æthio-
pica, floribus saturate luteis. *Pluk. mant.* 88. t. 413.

f. 3. *Raj. dendr.* 104.

Habitat in Æthiopia. ♀

2. ASPALATHUS foliis confertis subulatis mucrona- *chenopoda*
tis hispidis, floribus capitatis.

Genista africana lutea, floribus hirsutis in capitula lan-
ginosa conglobatis, foliis corruæ aculeatis subhirsu-
tis. *Herm. afr.* 11.

Chenopodium t. Chenopoda monomapatensis. *Breyn. cent.*

23. t. 11.

Habitat in Æthiopia. ♀

- 3 ASPALATHUS foliis confertis subulatis inermibus *thymifolia.*
glabris brevissimis, floribus alternis.

Genista minima æthiopica, foliis thymi confertis splen-
dentibus glabris. *Pluk. mant.* 88. t. 413. f. 1.

Habitat in Æthiopia. ♀

4. ASPALATHUS foliis confertis subulatis inermibus *ericifolia.*
hirsutis, floribus alternis, calycibus linearibus.

Genista æthiopica non spinosa, foliis ericæ villosis, flo-
ribus

Y y 4

ribus parvis spicatis luteis: calycibus longioribus immersis. *Pluk. mant.* 88. t. 413. f. 6.

Habitat in Æthiopia. ♀

- verrucosa.* 5. ASPALATHUS foliis confertis subulatis, gemmis verruca tomentosa prominentibus nudis.

Habitat in Æthiopia. ♀

- retroflexa.* 6. ASPALATHUS foliis confertis subulatis glabris minimis, ramis filiformibus reflexis, floribus solitariis terminalibus.

Habitat in Æthiopia. ♀

- eriflora.* 7. ASPALATHUS foliis confertis setaceis inermibus glabris, stipulis acutis persistentibus, floribus solitariis. calycibus ovatis.

Genista æthiopica glabra, longioribus foliis ex uno puncto plurimis, flore majore luteo. *Pluk. mant.* 88. t. 414. f. 7.

Habitat in Æthiopia. ♀

- arvensis.* 8. ASPALATHUS foliis confertis setaceis inermibus hispidis, floribus capitatis.

Genista æthiopica, flore flavo, foliolis inflexis & araneosa lanugine fimbriatis: summo ramulo circa flores glomeratis. *Pluk. mant.* 88. t. 414. f. 4. *Seb. thes.* 1. p. 38. t. 23. f. 6.

Habitat in Æthiopia. ♀

- indica.* 9. ASPALATHUS foliis quinatis sessilibus.

Genista foliis quinatis sessilibus, pedunculis axillaribus capillaribus unifloris, floribus minimis. *Fl. zeyl.* 271. *

Lotus tenuifolius maderaspatanus, siliqua singulari glabra. *Pluk. alm.* 225. t. 201. f. 2.

Dorycnium indicum, floribus singularibus rubris in pedicellis oblongis, siliquis perexiguis. *Raj. suppl.* 471.

Manelli. *Rheed. mal.* 9. p. 69. t. 37.

Habitat in India. ♀

- cretica.* 10. ASPALATHUS foliis trinibus cuneiformibus glabris: lateralibus brevioribus, stipulis obsoletis, floribus congestis.

Genista arborea cretica, foliis sempervirentibus. *Zan. hist.* 99. t. 39.

Habitat in Æthiopia. ♀

- tridentata.* 11. ASPALATHUS foliis trinibus lanceolatis glabris, stipulis tridentatis mucronatis, floribus capitatis.

Habitat in Æthiopia. ♀

DIADELPHIA DECANDRIA. 713

12. ASPALATHUS foliis trinis linearibus sericeis, sti- *argentea*,
pulis simplicibus mucronatis, floribus sparsis tomen-
tosis.

Cytisus africanus angustifolius, sericea lanagine argen-
tatus, spica lagopode. *Pluk. mant.* 63.

Habitat in Æthiopia. h

13. ASPALATHUS foliis trinis subulatis æqualibus, sti- *callosa*.
pulis subrotundis callosis, floribus spicatis glabris.

Cytisus trifolius, juniperinis foliis, floribus luteis in
spicam densiorem adactis. *Pluk. mant.* 63. t. 345. f. 4.

Habitat in Æthiopia. h. D. Royenus plurimas has A-
chyronias amicissime communicavit.

AMORPHA. 757

1. AMORPHA. *Hort. cliff.* 353. t. 19. *Hort. upf.* 208. *fruticosa*
Roy. lugdb. 393.

Barba jovis americana, pseudo-acaciæ foliis, flosculis
purpureis minimis. *Anglic. hort.* 11. t. 4.

Habitat in Carolina. h

ÆSCHYNOMENE. 758

1. ÆSCHYNOMENE caule scabro, leguminum arti- *aspera*.
culis medio scabris.

Æschynomene caule scabro, folio infimo quadripinni.
Hort. cliff. 365. *Hort. upf.* 233. *Fl. zeyl.* 298. * *Roy.*
lugdb. 384.

Mimosa non spinosa major zeylanica. *Brœyn. cens.* 52.
t. 52. *Raj. hist.* 982.

Habitat in India. c

2. ÆSCHYNOMENE caule hispido foliolis acumina- *americana*.
tis, leguminum articulis suborbiculatis. *Roy. lugdb.*
384.

Hedysarum caule hirsuto, mimosæ foliis alatis: pinnis a-
cutis minime gramineis. *Sloan. jam.* 74. *hist.* 1. p.
186. f. 3.

Habitat in Jamaica.

3. ÆSCHYNOMENE caule lævi arboreo, leguminum *arborea*.
articulis semicordatis glabris. *Roy. lugdb.* 384.

Habitat - - - h

4. ÆSCHYNOMENE caule lævi herbaceo, foliolis ob- *indica*.
tusis, leguminibus lævibus hinc obtusis rotatis. *Roy.*
lugdb. 385.

Y y 5

Neli-

714 DIADELPHIA DECANDRIA.

Nell-tali. *Rheed. mal.* 9. p. 31. t. 18.

Habitat in India.

Scriban. 5. *ÆSCHYNOMENE* leguminibus subæqualibus lævibus

Galega ægyptiaca, filiquis articulatis. *Baub. pin.* 352.

Sciban. Alp. ægypt. 81. t. 82.

Habitat in Ægypti sepibus. D. Hasselquist.

Legumina non articulata, sed lævia & aequalia sunt. Calyces campanulati, æquales, quinque-dentati, patentes.

759

GALEGA.

officinalis. 1. *GALEGA.* *Hort. cliff.* 362. *Hort. ups.* 208. *Mart. med.* 349. *Roy. lugdb.* 386.

Galega vulgaris. *Baub. pin.* 352. *Morif. hist.* 2. p. 91. f. 2. t. 7. f. 9.

Habitat in Italia, Hispania, Africa, ad margines agrorum. 2

760

CROTALARIA,

* *Foliis simplicibus.*

perfoliata. 1. *CROTALARIA* foliis cordato-ovatis perfoliatis. *Crotalaria perfoliatæ folio. Diss. elsb.* 122. t. 102. f. 122.

Habitat in Carolina.

sagittalis. 2. *CROTALARIA* foliis simplicibus lanceolatis, petiollis decurrentibus solitariis. *Hort. cliff.* 357. * *Græn. virg.* 81. *Roy. lugdb.* 374.

Crotalaria americana, caule alato, foliis pilosis, floribus in thyrso luteis. *Mart. cent.* 43. t. 43.

Crotalaria hirsuta minor americana herbacea, caule ad summum sagittato. *Herm. lugdb.* 202. t. 203. *Pluk. alm.* 122. t. 269. f. 6.

β. *Crotalaria sagittalis glabra*, longioribus foliis, americana, *Pluk. alm.* 122. t. 277. f. 2.

Habitat in Brasilia, Virginia. ☉

juncæ. 3. *CROTALARIA* foliis simplicibus lanceolatis petiolo-to-fessilibus, caule striato. *Hort. cliff.* 357. * *Hort. ups.* 210. *Roy. lugdb.* 374.

Crotalaria benghalensis, foliis genistæ hirsutis. *Pluk. alm.* 122. t. 169. f. 5.

Habitat in India. ☉

4. CRO-

4. CROTALARIA foliis simplicibus oblongis cuneiformibus retusa.
 mibus retuns. *Fl. zeyl.* 276. *
 Crotalaria asiatica, floribus luteis, folio singulari cordi-
 formi. *Herm. lugdb.* 200. t. 201.
 Tandale-Cotti. *Rheed. mal.* 9. p. 45. t. 25.
Habitat in India ☉
5. CROTALARIA foliis simplicibus ovatis sessilibus gla- *triflora.*
 bris, ramis angulatis, pedunculis ternis lateralibus uni-
 floris.
Habitat in Asia. *D. Royen.*
6. CROTALARIA foliis simplicibus ovatis villosis, pe- *villosa.*
 tiolis simplicissimis, ramis teretibus. *Hort. cliff.* 357. *
Roy. lugdb. 374.
 Crotalaria arbor africana, styracis folio molli incano.
Pluk. alm. 122. t. 185. f. 2,
 Genista arborefcens africana, styracis folio. *Herm. lugdb.*
 170. t. 271.
Habitat in Æthiopia. ☿
7. CROTALARIA foliis simplicibus ovatis, stipulis se- *verrucosa.*
 micordatis, ramis tetragonis. *Fl. zeyl.* 277. *
 Crotalaria foliis ovatis, petiolis duplici stipula auctis,
 ramis tetragonis. *Hort. cliff.* 357. *
 Crotalaria asiatica, folio singulari verrucoso, floribus cæ-
 ruleis. *Herm. lugdb.* 199. t. 199. *Raj. hist.* 1893.
 Pec-tandale-cotti. *Rheed. mal.* 9. p. 53. t. 29.
Habitat in India.
 * *foliis compositis.*
8. CROTALARIA foliis ternatis obovatis, floribus la- *latifolia.*
 teralibus subracemosis.
 Crotalaria loti folio, flore parvo variegato. *Dill. elth.*
 121. t. 102. f. 121.
 Crotalaria trifolia fruticosa, foliis glabris, flore e luteo
 viridi minore. *Sloan. jam.* 114. *hist.* 2. p. 33. t. 176.
 f. 1. 2.
Habitat in Jamaïca.
9. CROTALARIA foliis ternatis ovatis acutis, stipulis *lunatis.*
 semicordatis lunatis,
 Crotalaria foliis ternatis: foliolis sessilibus, petiolis stipu-
 la duplici auctis. *Hort. cliff.* 357. *
Habitat in Africa.
10. CROTALARIA foliis ternatis ovatis acuminatis, sti- *laburnifolia.*
 pulis nullis, leguminibus pedicellatis. *Fl. zeyl.* 273. *
 Crota

Crotalaria asiatica frutescens, foliis luteis amplis, trifolia.
Herm. lugdb. 196. t. 197. Raj. hist. 1893.
Nella-tandale-cotti. Rheed. mal. 9. p. 49. t. 27.
Habitat in Asia.

ovata.

11. *CROTALARIA* foliis ternatis ovatis acuminatis, stipulis nullis, leguminibus hirsutis.
Crotalaria foliis ternatis, petiollis nudis. Hort. cliff. 358.
** Hort. ups. 210. Roy. lugdb. 375.*
Crotalaria trifolia fruticosa, foliis rotundis incanis, floribus spicatis e viridi luteis, fructu pubescente. Sloan. jam. 141. hist. 2. p. 34. t. 179. f. 1. Raj. suppl. 466.
Habitat in Jamaica & Caribæis.

alba.

12. *CROTALARIA* foliis ternatis lanceolato-ovatis, caule lævi herbaceo, racemo terminali. *Hort. cliff. 499.**
Roy. lugdb. 375.
Anonis caroliniana perennis non spinosa, foliorum marginibus integris, floribus in thyrsio candidis. Mart. cent. 44. t. 44.
Habitat in Carolina. &

quinquefolia

13. *CROTALARIA* foliis quinatis.
Wellia-tandale-cotti. Rheed. mal. 9. p. 51. t. 28.
Habitat in India.
Foliola angusto-lanceolata, obtusa, sæpe retusa.

ONONIS.

** Floribus sessilibus.*

spinosa.

1. *ONONIS* floribus racemosis geminatis, foliis ternatis, ramis spinosis.
Ononis floribus subseffilibus solitariis lateralibus, caule spinoso. Hort. cliff. 359. Fl. suec. 623. Mat. med. 357. Roy. lugdb. 375.
Anonis spinosa, flore purpureo. Baub. pin. 389.
Anonis. Fuchj. hist. 60.
 β. *Ononis* floribus subseffilibus solitariis lateralibus, ramis inermibus. *Hort. cliff. 359. Fl. suec. 622. Roy. lugdb. 375.*
Anonis spinis carens purpurea. Baub. pin. 389.
Anonis mitior 1. Clusj. hist. 99.
Habitat in aridis Europæ. &
Hæc lævis β. est prima æstate, post aculeos α contrahit.
Loef. pruss. 17.
Folia in caule ternata; Floralia vero simplicia. Flores duo propriis pedunculis exeunt simul ex foliorum ala; Rami inferiores spina terminati.

2. O-

2. ONONIS caulibus diffusis, ramis erectis, foliis caulis repens.
nis ternatis.
Anonis maritima procumbens, foliis hirsutis pubescentibus. *Pluk. alm.* 33. *Dill. elsb.* 29. t. 25. f. 28.
Habitat in Angliæ littoribus maris. &
*Affinis admodum precedenti, differt caulibus-procumbentibus undique diffusis, & quod minor; Flores ex al-
lis solitarii (quorum non memor) speciem determina-
bunt.*
3. ONONIS floribus sessilibus lateralibus, foliis omnibus minutissima
ternatis petiolatisque, stipulis setaceis.
Ononis trifolia, calycibus ad axillas sessilibus erectis &
stipulis longas setas exerentibus. *Sauv. monsp.* 190.
Ononis floribus lateralibus sessilibus solitariis, petioli
longissimis. *Roy. lugdb.* 375.
Ononis lutea sylvestris minima. *Col. ecphr.* 1. p. 304.
t. 301
Anonis spinosa lutea minor. *Bauh. pin.* 389.
Anonis lutea montana non spinosa minima. *Barr. ic.*
1107.
Habitat in Italia, Monspeli.
4. ONONIS stipulis floralibus ovatis membranaceis inter-
mittentibus. *Roy. lugdb.* 376.
Anonis alopecuroides mitis annua purpurascens. *Dill.*
elsb. 28. t. 24. f. 27.
Anonis purpurea spicata erecta annua latifolia, siliquis
rectis latiforbibus. *Morif. hist.* 2. p. 169. f. 2. t.
17. f. 4.
Habitat in Barbados, Lusitania. ☉
5. ONONIS spicis foliosis, foliis simplicibus obtusis. *alopecuroides*
Ononis floribus spicatis. *Hort. cliff.* 358. *Roy. lugdb.* 376.
Anonis ficula alopecuroides. *Tournef. inst.* 408.
Anonis purpurea spicata alopecuroides major. *Boerh.*
lugdb. 2. p. 33.
Habitat in Sicilia, Lusitania, Hispania. ☉
6. ONONIS stipulis bracteisque acute serratis. *Roy. lugdb. variegata.*
376. *
Anonis non spinosa, flore luteo variegato, angustifolia
maritima. *Tournef. inst.* 409.
Habitat in Europæ australis maritimis.
* *Floribus pedunculatis.*
7. ONONIS pedunculis unifloris filo terminatis, foliis *Natrix.*
terna-

- ternatis. *Hort. cliff.* 358. * *Roy. lugdb.* 375. *Sauv. monsp.* 189.
Anonis viscosa spinis carens lutea major. Baub. pin. 389.
Ononis lutea. Cam epit. 445.
Natrix. Riv. tetr. 68.
β. *Anonis viscosa, spinis carens, lutea minor. Baub. pin.* 389.
Anonis viscosa minor, flore pendulo. Baub. prodr. 156.
γ. *Anonis lutea mitis oxitriphylla ad florum petiolos capreolata. Pluk. alm.* 33. t. 135. f. 5.
Habitat in G. Narbonenti, Hispania, inter segetes. &
viscosa. **8.** *ONONIS pedunculis unifloris filo terminatis, foliis simplicibus.*
Ononis pedunculis unifloris filo terminatis, foliis ternatis simplicibusque. Sauv. monsp. 189?
Anonis lutea viscosa latifolia minor, flore pallido. Barr. rar. 840. t. 1239.
Anonis viscosa, spinis carens lutea latifolia annua. Magn. monsp. 21?
Habitat Montpelii. ☉
Differt ab O. Natrice foliis simplicibus & radice annua.
ornithopodi- **9.** *ONONIS pedunculis bifloris filo terminatis. Roy. lugdb.* 376.
oides. *Anonis siliquis ornithopodii. Barb. lugdb.* 2. p. 34.
Fœnum græcum siculum, siliquis ornithopodii. Tournef. inst. 409.
Habitat in Sicilia ad Capo passaro.
tridentata. **10.** *ONONIS foliis ternatis carnosissimis sublinearibus tridentatis.*
Anonis hispanica frutescens, folio tridentato carnosissimo. Tournef. inst. 408. *Magn. monsp.* 17. t. 17. *Raj. juppl.* 466.
Habitat in Hispania. β
fruticosa. **11.** *ONONIS floribus paniculatis, pedunculis subtrifloris, stipulis vaginalibus, foliis ternatis. Hort. cliff.* 358. * *Roy. lugdb.* 376.
Anonis montana præcox purpurea frutescens. Dodart. act. 4. p. 249. t. 249.
Anonis purpurea frutescens non spinosa. Dodart. mem. 57. t. 57.
Anonis purpurea verna præcox frutescens, flore rubro amplo. Moris. hist. 2. p. 170.
Habitat in montibus Delphinatus. β

12. O-

11. **ONONIS** pedunculis axillaribus trifloris nudis, foliis *retundifolia* ternatis. *Hort. cliff.* 358. * *Roy. lugdb.* 376.
Cicer sylvestre latifolium triphyllum. Bauh. pin. 347.
Cicer sylvestre tertium. Dod. pempt. 525.
Cicer sylvestre verius. Lob. ic. 2. p. 73.
Habitat in Alpibus Helveticis.

ANTHYLLIS. 762

* *Herbaceæ.*

1. **ANTHYLLIS** herbacea, foliis quaterno-pinnatis, flo- *tetraphyllo*.
 ribus lateralibus. *Hort. ups.* 221. * *Sauv. monsp.* 237.
Anthyllis foliis quinato-pinnatis inæqualibus, floribus
lateralibus. Hort. cliff. 371. *Roy. lugdb.* 389.
Lotus pentaphyllos vesicaria. Bauh. pin. 332.
Trifolium halicacabum. Cam. hort. 171. t. 47.
Habitat in Italia, Sicilia. ☉
Folium pinnatum tetraphyllum loco pentaphylli, cum
foliolam infimum opposito careat.
2. **ANTHYLLIS** herbacea, foliis pinnatis inæqualibus, *Vulneraria*.
 capitulo duplicato.
Anthyllis foliis pinnatis, foliolis pluribus: terminali ma-
jore. Hort. cliff. 371. *Fl. succ.* 594. *Roy. lugdb.* 389.
Lotus affinis Vulneraria pratensis. Bauh. pin. 332.
Lagopodium flore luteo. Tabern. ic. 925.
 β. *Vulneraria supina, flore coccineo. Dill. elth.* 431. t.
 320. f. 413.
Habitat in pratis Europæ borealioris. &
3. **ANTHYLLIS** herbacea, foliis pinnatis æqualibus, ca. *montana*.
 pitulo terminali.
Vulneraria foliis pinnatis æqualibus: sub umbella pal-
matis. Hall. belv. 569. *Sauv. monsp.* 237.
Astragalus villosus, floribus globosis. Bauh. pin. 351.
Astragalus purpureus. Dalech. hist. 1347.
Habitat in Helvetia, G. Narbonensi, Galloprovincia. &
4. **ANTHYLLIS** herbacea, foliis pinnatis inæqualibus, *cornicina*.
 capitulis solitariis.
Habitat in Hispania. Læf. ☉
Caulis ramosus, spithameus. Folia pinnata: foliolis 5.
f. 7, ovali-lanceolatis, impari paulo majore. Pedun-
culi axillares, pilosi, longitugine foliorum. Capitula
lateralia & terminalia pedunculata ex 10 a 12 flori-
bis, rotunda, pilosa, folio ternato aut pinnato exce-
pta. Calyces subrotundi, inflati.

5. AN-

leucoides.

5. ANTHYLLIS herbacea, foliis tripartitis, calycibus prismaticis fasciculatis longitudine leguminum.

Habitat in Hispania. Læfl. ☉

Planta pilosa, procumbens. Folia caulina ternata, lateralibus basi coadunatis in unum cuneiforme, intermedia subseffili. Stipulae magnitudine & figura foliorum. Folia floralia ternata, sessilia, absque stipulis. Capitulum ex 5. s. 6. floribus, Calycibus admodum oblongis ratione plantae, angustis, prismaticis, parallelis, pilosis. Flores lutei. Legumina cylindrica, longitudine calycis, eoque vestita. Stamen decimum distinctum. Habitus ita antecedentium, ut facies vetat genere distinguere.

...*...Fruticosa.

- Barba jovis. 6. ANTHYLLIS fruticosa, foliis pinnatis aequalibus, floribus capitatis. *Hort. cliff.* 371. *Hort. ups.* 221. *Roy. lugdb.* 379. *Sæv. monsp.* 237.

Barba jovis. *Bauh. pin.* 397.

Barba jovis pulchre lucens. *Bauh. hist.* 1. p. 385.

Habitat in Helvetia. b

Cytisoides.

7. ANTHYLLIS fruticosa, foliis ternatis inæqualibus, calycibus lanatis lateralibus.

Anthyllis fruticosa, foliis ternatis: intermedio productiore. *Hort. ups.* 221.

Cytisus foliis ovatis acutis nitidis, floribus lateralibus. *Roy. lugdb.* 370.

Cytisus incanus, folio medio longiore. *Bauh. pin.* 390.

Cytisus VI. *Clus. hist.* 1. p. 96.

Spartium latifolium, parvo flore. *Barr. ic.* 1182.

Habitat in Hispania. b

hermanniae.

8. ANTHYLLIS fruticosa, foliis ternatis sessilibus, calycibus nudis.

Dorycnium foliis solitariis, floribus ad alas confertis.

Hort. cliff. 370. *

Habitat in Græcia. b

Calyces in hac vix ventricosi, at Habitus cum precedenti idem.

Erinaceæ.

9. ANTHYLLIS fruticosa spinosa, foliis simplicibus.

Genista-spartium spinosum, foliis lenticulae, floribus æ caruleo purpurascens. *Bauh. pin.* 394.

Erinacea. *Clus. hist.* 1. p. 107.

Habitat in Hispania. b

Tota

*Tota constat spinis; ad ramorum æsertum squama ovata,
Folia oblongo-ovata, tomentosa, ad flores posita sepi-
us terna.*

LUPINUS. 763

1. LUPINUS calycibus alternis inappendiculatis: labio perennis.
superiore emarginato, inferiore integro.
Lupinus calycibus alternis, radice perenni repente. *Gron.
virg. 172.*
Lupinus radice reptatrice perenni. *Roy. lugdb. 331.*
Lupinus cæruleus minor perennis virginianus repens.
Morif. bist. 2. p. 87. f. 2. t. 7. f. 6.
Habitat in Virginia. ☿
Folia ex foliolis 8, lanceolatis, obtusis, glabris. Racemi
longissimi floribus alternis, sæpius 2 ex singulo alterno
puncto, pedicellis propriis insidentibus.
2. LUPINUS calycibus alternis inappendiculatis: labio su-
periore integro, inferiore tridentato. *Hort. cliff. 499.*
Hort. ups. 209. Mat. med. 350. Roy. lugdb. 366.
Lupinus caule composito. *Hort. cliff. 359.*
Lupinus sativus, flore albo. *Bauh. pin. 347. Clus. bist.
2. p. 228.*
Lupinus flore albo. *Riv. tetr.*
Habitat - - - ☉
3. LUPINUS calycibus semiverticillatis appendiculatis: varius.
labio superiore bifido, inferiore subtridentato. *Hort.
cliff. 499. Hort. ups. 209. Roy. l. -db. 367.*
Lupinus sylvestris, flore cæruleo. *Bauh. pin. 348.*
Lupinus sylvestris, flore purpureo, semine rotundo va-
rio. *Bauh. bist. 2. p. 291.*
Habitat Meslanæ, Monspeliæ inter segetes. ☉
4. LUPINUS calycibus verticillatis appendiculatis: labio hirsutus.
superiore inferioreque integris. *Hort. cliff. 499. Hort.
ups. 209. Roy. lugdb. 367.*
Lupinus peregrinus major s. villosus cæruleus major..
Bauh. pin. 348. prodr. 148.
Lupinus exoticus hirsutissimus. *Bauh. bist. 2. p. 280.*
Habitat - - - ☉
5. LUPINUS calycibus alternis appendiculatis: labio su-
periore bipartito, inferiore integro.
Lupinus angustifolius cæruleus elatior. *Raj. bist. 908.*
Lupinus flore cæruleo minore. *Riv. tetr.*
Habitat - - - ☉

Z z

6. LU-

- lupinus.* 6. **LUPINUS** calycibus verticillatis appendiculatis: labio superiore bipartito, inferiore tridentato. *Hort. cliff.* 499. *Hort. nps.* 209. *Roy. lngdb.* 367. *Sauv. monsp.* 215.
Lupinus sylvestris, flore luteo. *Bauh. pin.* 348. *Riv. setr.*
Lupinus flore luteo, semine compresso vario. *Bauh. hist.* 2. p. 291.
Habitat in Sicilia arenosis. ☉
Hinc flores subsessiles, bracteis ovatis.

ROBINIA.

- 764
Pseudo-Acacia. 1. **ROBINIA** pedunculis racemosis, foliis impari-pinnatis. *Hort. nps.* 212.
Robinia aculeis geminatis. *Hort. cliff.* 354. *Gron. virg.* 82. *Roy. lngdb.* 372.
Acacia americana, filiquis glabris. *Raj. hist.* 1719.
Acaciae affinis virginiana spinosa, filiqua membranacea plana. *Pluk. alm.* 6. p. 73. t. 4.
Habitat in Virginia. ☿
Conf. Pseudo-Acacia hispida, floribus roseis. *Catesb. car.* 3. p. 20. t. 20.
- grandiflora.* 2. **ROBINIA** pedunculis subdivisis, foliis pinnatis, floribus foliolo majoribus.
Galega affinis malabarica arborescens, filiquis majoribus articulatis. *Raj. hist.* 1734.
Sesban affinis arbor indiae orientalis. *Breyn. prodr.* 1. p. 47. *Comm. mul.* 244.
Agaty. Rheed. mal. 1. p. 95. t. 51.
Habitat in India. ☿
Arbor crassa, inermis. Folia pinnata: Foliola obtusa. Flores inter papilionaceos maximi, magnitudine ovi galinacei. Vexillum ovatum, patens. Ala ovato-oblonga, versus apicem sursum falcata. Carina alis conformis. Stamina ad basin valde sursum flexa & reflexa. Stamina diadelphea. Stigma obtusum.
- Caragana.* 3. **ROBINIA** pedunculis simplicissimis, foliis aequalibus pinnatis. *Hort. nps.* 212.
Alpalathus arborescens, pinnis foliorum crebrioribus oblongis. *Amm. ruth.* 285.
Caragana sibirica. *Roy. lngdb.* 537.
Habitat in Sibiria. ☿

DIADELPHIA DECANDRIA. 723

4. ROBINIA pessunculis simplicissimis, foliis quater-*frutis*
natis petiolatis. *Hort. upf.* 212.

Aspalathus frutescens major latifolius, cortice aureo.
Amm. ruth. 283.

Habitat in Sibiria, Tataria. h

5. ROBINIA pedunculis simplicissimis, foliis quaterna-*pygmaea*.
tis sessilibus. *Hort. upf.* 212.

Aspalathus frutescens minor angustifolius, cortice aureo.
Amm. ruth. 282. t. 35.

Habitat in Sibiria. h

Vix hæc sufficientior, a priori, distincta.

COLUTEA. 765

1. COLUTEA arborea, foliolis obcordatis. *Hort. cliff. arborescens.*
365. *Hort. upf.* 228. *Roy. lugdb.* 374. *Sauv. monsp.* 238.

Colutea vesicaria. Bauh. pin. 396.

Colutea. Dod. pempt. 784.

β. *Colutea africana, lennæ foliis, flore sanguineo. Comm.*
rar. II. t. 11.

Habitat in Austria, G. Narbonensi, Italia præcipue ad
Vesuvium. h

2. COLUTEA fruticosa, foliolis ovato-oblongis. *Hort. frutescens.*
cliff. 366. *Hort. upf.* 228. *Roy. lugdb.* 374.

Colutea æthiopica, flore purpureo. Breyn. cent. 70. t. 29.

Habitat in Æthiopia. k. h

3. COLUTEA herbacea, foliis linearibus. *Hort. upf.* 266. *herbacea.*
Roy. lugdb. 374.

Colutea africana annua, foliolis parvis mucronatis, vesti-
culis compressis. Comm. hort. 2. p. 87. t. 44.

Habitat in africa. ☉

PHASEOLUS. 766

* *Scandentes.*

1. PHASEOLUS caule volubili, floribus racemosis ge-*volubilis*.
minis, bracteis calyce majoribus, leguminibus pendulis.

Phaseolus radice annua, caule volubili, leguminibus pen-
dulis compressis torosis. Roy. lugdb. 367. *Hort. upf.*
213. *Mat. med.* 351.

Phaseolus vulgaris. Lob. ic. 59. *Hort. cliff.* 359.

Smilax hortenensis f. Phaseolus major. Bauh. pin. 339.

Habitat in India. ☉

Z z z

2. PHA.

- coccineus.* 2. PHASEOLUS caule volubili, floribus racemosis geminis, bracteis calyce brevioribus, leguminibus pendulis.
Phaseolus puniceo flore. *Cornut. canad.* 184.
Phaseolus indicus, flore coccineo f. puniceo. *Morif. hist.* 2. p. 69.
Habitat - - - - - ☉
- acutellus.* 3. PHASEOLUS caule volubili, leguminibus acinaciformibus sublunatis lævibus.
Phaseolus benghalensis scandens, filiqua acinaciformi, semine ovato compresso striato. *Berg. viadr.* 99.
Habitat in Benghala. ☉
Foliola lateraliter exteriori latere duplo latiora. Flores minimi, virides; alis albis concavis, dein flavescentibus; carina spirali.
- tnamensis.* 4. PHASEOLUS caule volubili, florum vexillo revoluti, calycibus concolore. *Hort. cliff.* 359. *
Habitat in Africa. ☉
- farinosus.* 5. PHASEOLUS caule volubili, pedunculis subcapitatis, seminibus tetragono-cylindricis pulverulentis. *Hort. ups.* 214.
Phaseolus peregrinus, flore roseo, semine tomentoso. *Niff. act.* 1730. p. 577. t. 42.
Phaseolus indicus, hederæ folio anguloso, semine oblongo lanuginoso. *Raj. suppl.* 348.
Habitat in India.
- vexillatus.* 6. PHASEOLUS caule volubili, vexillis revolutis patulis, leguminibus linearibus strictis.
Phaseolus radice annua, leguminibus strictis erectis torosis linearibus, caule hirsuto. *Roy. lugdb.* 367.
Phaseolus flore odorato, vexillo amplo patulo. *Dill. elsb.* 313. t. 234. f. 302.
Habitat in Havana. ☉
- helveticus.* 7. PHASEOLUS caule volubili, floribus capitatis, calycibus bracteis, vexillis brevibus, alis expansis maximis.
Phaseolus radice annua, leguminibus erectis linearibus teretibus glabris. *Roy. lugdb.* 368.
Phaseolus flore helvulo, alis amplis patulis. *Dill. elsb.* 312. t. 233. f. 300.
Habitat in Carolina. ☉

8. PHASEOLUS caule volubili, floribus laxè spicatis, *alatus*,
 alis longitudine vexilli.

Phaseolus flore purpureo, alis amplis longè protensis. *Dill.*

elsh. 314. *t.* 235. *f.* 303

Habitat - - - - - ☉

9. PHASEOLUS caule volubili, vexillis earinaque spi- *Caracalla*,
 raliter convolutis.

Phaseolus radice perenni subrotunda, leguminibus fo-
 lio longioribus teretiusculis glabris. *Roy. lugdb.* 367.

Phaseolus indicus, cochleato flore. *Tournef. obs.* 93. *t.*

94. *Raj. bist.* 1890.

Habitat in India.

* *erecti.*

10. PHASEOLUS caule erecto tereti, floribus capitatis, *radiatus*,
 leguminibus cylindricis horizontalibus. *Aët. stockb.*

1742. *p.* 202. *t.* 7. *f.* 2. *Hort. upf.* 213. *Fl. zeyl.* 281.

Phaseolus zeylanicus, filiquis radiatim digestis. *Dill. elsh.*

315. *t.* 235. *f.* 304.

Habitat in China, Zeylona. ☉

11. PHASEOLUS caule erecto anguloso hispido, legu- *Mad.*
 minibus pendulis hirtis. *Hort. cliff.* 499. *Roy. lugdb.*

367. *Fl. zeyl.* 280.

Phaseolus orthocaulis, Mungo persarum, *Herm. mex.*

887.

Habitat in India. ☉

DOLICHOS.

7 6 7

* *Volubiles.*

1. DOLICHOS leguminibus ovato-acinaciformibus, se- *Lablab.*
 minibus ovatis hilo arcuato versus alteram extremita-

tem. *Roy. lugdb.* 368. *Hort. upf.* 214.

Phaseolus ægyptius, nigro semine, *Bauh. pin.* 341.

Phaseolus niger *Lablab. Alp. ægypt.* 74. *t.* 75. *Vest.*

ægypt. 27.

Habitat in Ægypto. ☉

Legumina dorso scabra. Caules ramique teretes, retror-

sum scabri. Pedunculi semiverticillati.

2. DOLICHOS leguminibus capitatis subcylindraccis: *unguiculatus*
 apice recurvo concavo. *Hort. upf.* 214.

Habitat in Barbados.

3. DOLICHOS leguminibus gladiatis dorso sulcatis, se- *eniformis.*
 minibus arillatis. *Hort. upf.* 360. *Savv. meth.* 192.

Z 2 3

Phaseo-

Phaseolus maximus, filiqua ensiformi nervis insignita & semine albo membranula incluso. *Sloan. hist.* 1. p. 177. t. 114. f. 1. 2. 4.

Habitat in Jamaica.

minimus. 4. **DOLICHOS** leguminibus racemosis compressis tetraspermis, foliis rhombæis.

Dolichos minimus, floribus luteis. *Hort. cliff.* 360. t. 21. * *Hort. ups.* 214.

Phaseolus minimus scetidus, floribus spicatis e viridi luteis, semine maculato. *Sloan. hist.* 1. p. 182. f. 115. f. 1.

Habitat in Jamaica.

scarabæoides 5. **DOLICHOS** caule volubili, foliis ovatis tomentosis, floribus solitariis, seminibus bicornibus. *Fl. zeyl.* 282. *

Phaseolus minimus hisnagaricus, foliis argenteo-villosis, filiquis torosis brevibus spadicea hirsutæ pubescentibus, fructu parvo scarabæoide nigro. *Pluk. alm.* 290. t. 52. f. 3.

Habitat in India.

erosus. 6. **DOLICHOS** caule volubili, foliis glabris multangulis dentatis.

Phaseolus nervicensis, foliis multangulis, tuberosa radice. *Pluk. alm.* 292. t. 54. f. 4.

Habitat in America. ☉

trilobus. 7. **DOLICHOS** foliolis lateralibus extrorsum gibbis: intermedio trilobo. *Hort. cliff.* 360. *

Phaseolus maderaspatanus, foliis glabris trilobis, floribus exiguis, longis petiolis ex eodem puncto geminis. *Pluk. alm.* 292. t. 214. f. 3

Habitat in India.

regularis. 8. **DOLICHOS** foliis ovatis obtusis, pedunculis multifloris, petalis æqualis magnitudinis figuræque. *Gorn. virg.* 82.

Habitat in Virginia.

lignosus. 9. **DOLICHOS** caule perenni, pedunculis capitatis, leguminibus strictis linearibus.

Dolichos caule perenni lignoso. *Hort. cliff.* 360. t. 20.

Phaseolus indicus perennis, floribus purpurascens. *Eichr. carol.* 36.

Habitat - - - - - 3

polystachios. 10. **DOLICHOS** caule perenni, spicis longissimis, pedicellis geminis, leguminibus acuminatis compressis.

Dolichos caule lignoso, pedunculis ex ala plurimis, floribus

ribus spicatis, legumine longo apice sursum acuminato. *Gron. virg.* 172.

Habitat in Virginia.

* *Erecti.*

11. DOLICHOS caule erecto flexuoso, racemis axilla-*Seja*.
ribus erectis, leguminibus pendulis hispidis subdisper-
mis. *Fl. zeyl.* 534. *Mat. med.* 363.

Phaseolus erectus, filiquis lupini, fructu pisi majoris can-
dido. *Kampf. amon.* 837. t. 838.

Habitat in India.

12. DOLICHOS caule perenni lævi, pedunculis bislo-*biformis*.
ris, leguminibus erectis. *Roy. lugdb.* 368.

Habitat in India.

PISUM.

768

1. PISUM stipulis inferne rotundaris crenatis, petiolis *ter-sationum*.
reibus, pedunculis multifloris. *Hort. ups.* 215.

Pisum stipulis crenatis. *Hort. cliff.* 369. *Mat. med.* 352.

Pisum. *Cam. epis.* 213.

β. Pisum hortense majus. *Bauh. pin.* 342.

γ. Pisum sine cortice duriore. *Bauh. pin.* 343.

δ. Pisum umbellatum. *Bauh. pin.* 344.

ε. Pisum majus quadratum. *Bauh. pin.* 342.

Habitat in Europæ agris. ☉

umbellatum
quadratum.

2. PISUM petiolis tetraphyllis, stipulis crenatis, pedun-*arvense*.
culis unithoris. *Is. W'goth.* 107. *

Pisum pulchrum, folio anguloso. *Bauh. hist.* 2. p. 297.

Morif. hist. 2. p. 47. f. 2. t. 1. f. 4.

Habitat inter Europæ segetes. ☉

3. PISUM petiolis supra planiusculis, caule angulato, sti-*maritimum*.
pulis sagittatis, pedunculis multifloris. *Fl. succ.* 608.

Is. W'goth. 190. *

Pisum stipulis integerrimis. *Hort. cliff.* 368.

Pisum marinum. *Raj. hist.* 892. *Fl. lapp.* 272.

Pisum spontaneum pene repens humile. *Morif. hist.*

2. p. 47. f. 2. t. 1. f. 5.

Habitat in Europæ borealis litoribus maris arenosis. ☉

4. PISUM petiolis decurrentibus membranaceis diphyll-*Ochrus*.
is, pedunculis unifloris. *Hort. cliff.* 370. *Hort. ups.* 215.

Ochrus folio integro capreolos emittente. *Bauh. pin.* 343.

Ervilia sylvestris. *Dod. pempt.* 522.

Habitat inter segetes Cretæ, Italiæ. ☉

Z 2 4

ORO-

769 OROBUS.

- Lathyroides.* 1. OROBUS foliis conjugatis subseffilibus, stibulis dentatis. *Hort. mps.* 220.
Lathyroides erecta, folio ovato acuminato, cæruleis vicæ floribus & siliquis, sibirica. *Amm. rath.* 151.
t. 7. f. 2.
Habitat in Sibiria. 2
- hirsutus.* 2. OROBUS foliis conjugatis petiolatis, stipulis integris. Orobus caule simplicissimo, foliolis binis ovatis. *Hort. cliff.* 366. *
 Orobus latifolius repens, flore cæruleo, foliis & siliquis hirsutis. *Boerb. lugdb.* 2. p. 46.
 Orobus sylvaticus, foliis circa caulem auriculatis. *Buxb. cent.* 3. p. 22. t. 41.
Habitat in Thracia. 2
- dentat.* 3. OROBUS foliis pinnatis ovato-oblongis, stipulis rotundato-lunatis dentatis, caule simplici.
Habitat in Sibiria. 2
Habitat Orobis verni, sed altior, crassior, major. Folia pinnata 4. f. 5. paribus, ovato-oblonga. Stipulæ latæ, subrotundo-lunares, dentatæ. Caulis simplex. Racemi axillares, secundi, floribus numerosis majoribus.
- vernus.* 4. OROBUS foliis pinnatis ovatis, stipulis semisagittatis integerrimis, caule simplici.
 Orobus caule simplicissimo, foliolis pluribus ovatis acutis. *Hort. cliff.* 366. *Fl. suec.* 595. *Roy. lugdb.* 366. *Hall. belv.* 602.
 Orobus sylvaticus purpureus vernus. *Baub. pin.* 351. *Thal. herc.* 80. t. 6.
Habitat in Europæ borealis nemoribus. 2
- inherosus.* 5. OROBUS foliis pinnatis lanceolatis, stipulis semisagittatis integerrimis, caule simplici.
 Orobus caule simplicissimo, foliolis pluribus lanceolatis integris. *Hort. cliff.* 366. *Fl. suec.* 596. *Roy. lugdb.* 366.
 Astragalus sylvaticus, foliis oblongis glabris. *Baub. pin.* 351.
 Astragalus sylvaticus. *Thal. herc.* 7. t. 1.
 Lathyrus angustifolius, radice tuberosa. *Loef. praff.* 138. t. 37.
Habitat in Europæ borealis pratis & sylvis. 2

6. ORO-

6. **OROBUS** foliis quaternopinnatis linearibus, stipulis *angustifolius* subulatis, caule simplici.

Habitat in Sibiria. 2

Habitus antecedentis, sed Folia linearia, angustissima & pinnata ex 2 oppositionibus, absque cirrho. Racemi floribus paucis luteis.

7. **OROBUS** caule ramoso, foliis sexjugis ovato-oblongis. *Hort. cliff. 366. Fl. suec. 597. Roy. lugdb. 366.*

Hall. belv. 603.

Orob. sylvaticus, vicæ foliis. Baub. pin. 352.

Orob. pannonicus 2. Claf. hist. 2. p. 230.

Habitat in Europæ borealis montosis. 2

8. **OROBUS** caule ramoso, foliis trijugis lanceolatis, *pyrenaicus*, nervosis. *Sauv. monsp. 235. †*

Orob. pyrenaicus, foliis nervosis. Tournef. inst. 235.

Orob. pyrenaicus latifolius nervosus. Pluk. phyt. 210.

f. 2.

Habitat in Pyrenæis.

LATHYRUS.

770

* *Pedunculis unifloris.*

1. **LATHYRUS** pedunculis unifloris, cirrhis aphyllis, *Aphaca*, stipulis sagittato-cordatis.

Lathyrus cirrhis aphyllis. Hort. cliff. 367. Hort. ups. 218. Roy. lugdb. 362. Dalib. parif. 217.

Vicia lutea, foliis convolvuli minoris. Baub. pin. 345.

Aphaca. Lob. ic. 2. p. 70.

Habitat in Italia, Gallia, Anglia inter segetes. ☉

Cirrhus interdum aliquis gerit Foliola conjugata, 2, lanceolata, reliquis Lathyris similima, at hoc rarissime.

2. **LATHYRUS** pedunculis unifloris, foliis simplicibus, *Nissolia*, stipulis subulatis.

Lathyrus foliis solitariis lentiformibus. Vir. cliff. 74.

Hort. cliff. 366. Roy. lugdb. 263. Dalib. parif. 218.

Lathyrus sylvestris minor. Baub. pin. 344.

Lathyrus angustifolius erectus, folio singulari, sine ca-

preotis, nissoli. Magn. hort. 112. t. 112.

Habitat in Gallia. ☉

Cirrhii nulli. Caulis erectus. Stipulæ minimæ, subulatæ.

3. **LATHYRUS** pedunculis unifloris calyce longioribus, cirrhis diphyllis simplicissimis. *amphicarpus*

Lathyrus pedunculis unifloris, cirrhis diphyllis, radicibus

Z 2 5

etiam

etiam sub terra fructificantibus. *Hort. cliff.* 367. *
Hort. xps. 216. *Roy. lugdb.* 262. *Sauv. monsp.* 192.
 Vicinæ similis supra & infra terram siliquas gerens. *Bauh.*
pin. 345.

Arachidna f. *Arachoides honorii bellii*, *Bauh. hist.* 2. p.
 323.

Habitat in Syria. ☉

Cura.

4. **LATHYRUS** pedunculis unifloris, cirrhis diphyllis, leguminibus ovatis compressis: dorso canaliculatis. *Lathyrus* pedunculis unifloris, cirrhis diphyllis, foliis lanceolato-linearibus, internodiis membranaceis. *Sauv. monsp.* 193.

Lathyrus sativus flore purpureo. *Bauh. pin.* 344.

Aracus f. *Cicera*. *Dod. pempt.* 523.

Habitat in Hispania. ☉

Medius inter antecedentem & sequentem magnitudine.
Petioli latere non membranacei. Corollæ rubrae. Legumina dorso canaliculata, at non marginibus acutis, uti antecedens & insequens. Legumen latum, ovali-oblongum.

foliatus.

5. **LATHYRUS** pedunculis unifloris, cirrhis diphyllis, leguminibus ovatis compressis dorso bimarginatis. *Hort. cliff.* 367. *Hort. xps.* 216. *Roy. lugdb.* 363. *Dalib. parif.* 216. *Sauv. meth.* 193. *

Lathyrus annuus, flore cæruleo, ochri siliqua. *Herm. lugdb.* 357.

- β. *Lathyrus sativus*, flore fructuque albo. *Bauh. pin.* 343.

Lathyrus f. *Cicercula*. *Dod. pempt.* 522.

Habitat in Hispania, Gallia. ☉

- inconspicuus* 6. **LATHYRUS** pedunculis unifloris calyce brevioribus, cirrhis diphyllis, foliolis lanceolatis.

Lathyrus orientalis, flore vix conspicuo. *Boerb. lugdb.* 2. p. 42.

Habitat in Oriente. ☉

Planta parva, spithamæ f. pedalis. Caules obsolete triangulares. Stipulæ subulato-lanceolata: dente postico acuto. Petioli brevissimi, inferiores terminati seta subulata, brevissima; superiores in cirrhum filiformem elongati. Foliola bina, lanceolata, subtus striata, acuminata. Calyx minimus & stipulis brevior ex alis petiolorum, pedunculo vix manifesto insidens. Corolla tam parva et vix calyce longior, pallida. Legomen longitudine ferme foliorum, sed latius, glabrum, strictum.

7. LA-

7. **LATHYRUS** pedunculis unifloris, cirrhis diphyllis, *setifolius*.
foliolis setaceo-linearibus.
Lathyrus pedunculis unifloris, cirrhis diphyllis: foliolis
linearibus, stipulis subulatis, caulibus nudis. *Sauv.*
monsp. 193.
Lathyrus folio tenuiore, floribus rubris. *Benth. Hist.* 2.
p. 308.
Lathyrus sylvestris major, angustissimo folio. *Bauh.*
pin. 344.
Lathyrus angustissimo f. capillaceo folio. *Bauh. prodr.*
148.
Habitat Monspelli.
8. **LATHYRUS** pedunculis unifloris cirrho terminatis, *angulatus*.
cirrhis diphyllis, foliolis linearibus.
Lathyrus pedunculis unifloris cirrhosis, foliolis linearibus,
seminibus angulatis.
Lathyrus pedunculis unifloris, cirrhis diphyllis, foliolis
capillaceis. *Sauv. monsp.* 193.
Lathyrus angustissimo folio, semine anguloso. *Tournef.*
inst. 395.
Lathyrus folio tenuiore, floribus rubris. *Buxb. cent.* 3.
p. 23. t. 42. f. 2.
Lathyrus angustifolius leptolobus, semine hexaëdro, flore
cæruleo. *Morif. bl. f.* 279
Habitat in Gallia, Hispania, Oriente. ☉
9. **LATHYRUS** pedunculis unifloris, cirrhis tetraphyllis *bithynicus*.
lis, stipulis dentatis. *Roy. lugdb.* 363.
Lathyrus pedunculis unifloris, petiolis tetraphyllis, stipulis
setaceis, caule ancipiti. *Amœn. acad.* 1. p. 166.
Chymenum bithynicum, siliqua singulari, flore minore.
Boerb. lugdb. 2. p. 43.
Habitat in Italia, Bavarie arvis. 2
Caules *angulati*. Cirrhi foliolis 2 parium, lato-lanceolatis.
Stipulæ valde dentatæ. Flores axillares, solitarii:
calyce longitudine pedunculi, vexillo violaceo, alis albidis.
Legumen erectum, compressum, ovato-oblongum villosum.
Semina sœva circiter. Media inter *Lathyros* & *vicias*.
10. **LATHYRUS** pedunculis subunifloris, cirrhis poly-*articulatus*.
phyllis: foliolis alternis. *Hort. cliff.* 368. *Hort. ups.*
217. * *Roy. lugdb.* 363.
Lathyrus, angustissimo folio, americanus variegatus.
Bauh. pin. 344.

Cly-

Clymenum hispanicum, flore vario, siliqua articulata.
Tournef. inst. 396.

β. *Lathyrus angustifolius* hemillior. *Bauh. pin.* 344.

Habitat in Bœtica. ☉

Folia sæpius tetraphylla; inferiores petioli lanceolati aphylli. Stipulæ utrinque mucronatæ.

* *Pedunculis bifloris.*

odoratus. 11. *LATHYRUS* pedunculis bifloris, cirrhis diphyllis, foliis ovato-oblongis, leguminibus hirsutis. *Hort. cliff.* 368. *Hort. upf.* 216. *Roy. lugdb.* 363.

sculus. α. *Lathyrus sculus.* *Rupp. jew.* 210.

Lathyrus distoplatyphyllos hirsutus mollis, magno & peranœno flore odoro. *Comm. hort.* 2. p. 219. t. 80.

zeylanicus. β. *Lathyrus zeylanicus*, odorato flore amœno ex albo & rubro vario. *Burm. zeyl.* 138.

Habitat α in Sicilia, β in Zeylona. ☉

hirsutus. 12. *LATHYRUS* pedunculis bifloris, cirrhis diphyllis, foliolis lineari-lanceolatis, leguminibus hirsutis, seminibus scabris. *Roy. lugdb.* 363. *Dalib. paris.* 216. *Sauv. monsp.* 113.

Lathyrus angustifolius, siliqua hirsuta. *Bauh. pin.* 344.

Lathyrus siliqua hirsuta. *Bauh. hist.* 2. p. 305.

Habitat inter Angliæ, Galliæ segetes.

tingitanus. 13. *LATHYRUS* pedunculis bifloris, cirrhis diphyllis: foliolis alternis lanceolatis. *Roy. lugdb.* 263.

Lathyrus tingitanus, filiquis orobi, flore amplo ruberrimo. *Morif. hist.* 2. p. 55.

Habitat in Mauritania. ☉

clymenum. 14. *LATHYRUS* pedunculis bifloris, cirrhis polyphyllis, stipulis dentatis. *Hort. cliff.* 368. *Hort. upf.* 217. *Roy. lugdb.* 364.

Lathyrus vicioides, vexillo rubro, labialibus petalis rostrum ambientibus cæruleis. *Morif. hist.* 2. p. 56.

Clymenum hispanicum, flore vario, siliqua plana. *Tournef. inst.* 396.

Habitat in Mauritania. ☉

* *Pedunculis multifloris.*

tuberosus. 15. *LATHYRUS* pedunculis multifloris, cirrhis diphyllis: foliolis ovalibus, internodiis nudis. *Hort. cliff.* 367. *Hort. upf.* 216. *Roy. lugdb.* 364. *Dalib. paris.* 217. *Sauv. monsp.* 194. *Gort. gelr.* 421.

La-

Lathyrus arvensis repens tuberosus. *Bauh. pin.* 344.

Apios. Fuchf. hist. 131.

Habitat inter Belgii, Genevæ, Tataricæ segetes. &

16. **LATHYRUS** pedunculis multifloris, cirrhis diphyll-^{pratenq}
lis: foliolis lanceolatis: cirrhis simplicissimis. *Hort.*
cliff. 367. *Fl. succ.* 599. *Roy. lugdb.* 364. *Dalib. pa-*
ris. 217.

Lathyrus sylvestris luteus, foliis viciz. *Bauh. pin.* 344.

Lathyrus luteus sylvestris dumetorum. Bauh. hist. 2.
p. 304. *t.* 304.

Lathyrus foliis binatis acutis, capreolo non ramoso.
Hall. belv. 597.

Habitat in Europæ pratis. &

Cirrhii quidem interdum trifidi sunt, at sæpius simpli-
cissimi.

17. **LATHYRUS** pedunculis multifloris, cirrhis di-^{sylvestris.}
phyllis: foliolis ensiformibus. *Hort. cliff.* 367. *Fl.*
succ. 598. *Dalib. paris.* 216.

Lathyrus sylvestris major. *Bauh. pin.* 433.

Lathyrus sylvestris. Clus. hist. 2. *p.* 129. *Dod. pempt.*
523.

Habitat in Europæ pratis montosis. &

18. **LATHYRUS** pedunculis multifloris, cirrhis di-^{latifolius.}
phyllis: foliolis lanceolatis, internodiis membranaceis.
Hort. cliff. 367. *Hort. ups.* 217. *Fl. succ.* 1139. *It.*
scan. 251. *

Lathyrus latifolius. Bauh. pin. 344. *Garid. prov.*
271. t. 271.

Lathyrus major latifolius, flore purpureo, speciosior.
Bauh. hist. 2. *p.* 303.

Habitat in Europæ sepibus. &

19. **LATHYRUS** pedunculis multifloris, cirrhis diphyll-^{heterophyl-}
lis tetraphyllisque: foliolis lanceolatis. *It. w-goth* 75. ^{lus.}
Lathyrus pedunculis multifloris, cirrhis diphyllis: folio-
lis gramineis trinerviis, internodiis membranaceis.
Sauv. monsp. 193.

Lathyrus major narbonensis angustifolius. *Bauh. hist.*
2. p. 304.

Habitat ad radices montium Europæ. &

20. **LATHYRUS** pedunculis multifloris, cirrhis poly-^{palustris.}
phyllis stipulis lanceolatis. *Hort. cliff.* 368. *Fl. succ.* 600.
Roy. lugab. 364. *Dalib. paris.* 217.

La-

Lathyrus foliis pinnatis, pedunculis multifloris. *Fl. lapp.* 221.

Lathyrus palustris, flore orohi nemorensis verni. *Rupp. jen.* 210. t. 1.

Vicia lathyroides f. *Lathyrus viciæformis*. *Pluk. alm.* 387. t. 71. f. 2.

Habitat in Europæ borealis pascuis paludosis. &

pissotensis. 21. **LATHYRUS** pedunculis multifloris, cirrhis polyphyllis, stipulis ovatis: basiacutis. *Hort. ups.* 217.

Habitat in Sibiria. &

Planta ut Pisum crescens. Caules acute angulati. Petioli cirrhosi, sepe hexaphyllis foliolis ovato-oblongis, acutis. Stipulæ foliis latiores, ovatae, basi acuminatae, vix angulatae. Pedunculi longitudine foliorum, multiflori. Flores violacei.

971

VICIA.

* *Pedunculis elongatis.*

viciiformis. 1. **VICIA** pedunculis multifloris, petiolis polyphyllis: foliolis ovatis.

Vicia pedunculis multifloris, petiolis polyphyllis cirrhosis, foliolis ovatis: infimis sessilibus. *Hort. cliff.* 369.*
Roy. lugdb. 365.

Pisum sylvestre perenne. *Baub. pin.* 343.

Pisum (sylvestre). *Clus. bist.* 2. p. 129.

Habitat in Pannoniæ sylvis. &

dumetorum. 2. **VICIA** pedunculis multifloris, foliolis reflexis ovatis mucronatis, stipulis subdentatis.

Vicia pedunculis multifloris, petiolis polyphyllis cirrhosis: foliolis alternis ovatis, stipulis dentatis. *Fl. succ.* 604. *Dalib. parif.* 219. *Sauv. monsp.* 234.

Vicia maxima dumetorum. *Baub. pin.* 385.

Vicia sylvatica maxima piso similis. *Baub. bist.* 2. p. 315.

Vicia sylvatica. *Riv. tetr.* 50.

Habitat in Thuringia, Gallia. &

sylvatica. 3 **VICIA** pedunculis multifloris, foliolis ovalibus, stipulis denticulatis.

Vicia pedunculis multifloris, stipulis crenatis. *Fl. lapp.* 270. *Fl. succ.* 603. *Roy. lugdb.* 365. *Sauv. monsp.* 234.

Vicia multiflora maxima perennis, tetto odore, flori-

bus albeutibus: lineis cæruleis notatis *Pluk. alm.*
387. t. 71. f. 1.

Habitat in Sueciæ, Germaniæ, Galliæ sylvis. &

4. VICIA pedunculis subsexfloris, foliolis denis ovatis *caffibita*.
acutis, stipulis integris.

Vicia pedunculis multifloris, caule fruticoso. *Hort. cliff.*
369. *Roy. lugdb.* 364.

Vicia multiflora cassubica frutescens, siliqua lentis.
Pluk. alm. 387. t. 72. f. 2.

Habitat in Germania.

5. VICIA pedunculis multifloris, floribus imbricatis, fo- *Cracca*.
liolis lanceolatis pubescentibus, stipulis integris.

Vicia pedunculis multifloris, stipulis utrinque acutis in-
tegris. *Hort. cliff.* 368. *Fl. succ.* 605. *Roy. lugdb.*
364. *Dalib. parif.* 220. *Sauv. monsp.* 234.

Vicia sylvestris spicata. *Baub. pin.* 45.

Vicia multiflora. *Baub. pin.* 343.

Cracca. Riv. tetr. 49.

Habitat in Europæ pratis, agris. &

6. VICIA pedunculis multifloris, floribus distantibus, fo- *onobrychid-*
liolis linearibus, stipulis inferne denticulatis. *des.*

Vicia pedunculis multifloris, stipulis crenatis, foliolis
linearibus. *Sauv. monsp.* 235.

Vicia onobrychidis flore. *Baub. pin.* 345. *prodr.* 149.

Habitat in Gallia. D. Sauvages.

Statura Vitiæ Craccae. Folia pinnata, cirrhosa: foliolis
sæpius 14, linearibus, angustissimis: oppositionibus di-
stantibus. Pedunculi axillares, foliis triplo longiores,
friati. Floribus 10. f. 12, subsessilibus, remotissimis,
Craccae triplo majoribus, purpureis.

7. VICIA pedunculis multifloris, foliolis oblongis, stipu- *nissoliana*
lis integris, leguminibus villosis ovato-oblongis.

Vicia leguminibus sessilibus binis ternisve subovatis nu-
tantibus, stipulis nullis. *Roy. lugdb.* 14?

Vicia orientalis, flore suave rubente, siliquis brevissimis.
Barb. lugdb. 2. p. 44?

Habitat in Oriente.

Caulis sulcatus. Cirrhi foliolis 6 paribus, oblongis, vix
retusis, acumineternatis, subvillosis. Stipulae lanceo-
late, angustissima. Pedunculi floribus 5. f. 6, mini-
mis, atropurpureis. Legumina nutantia, compressa,
ovato-oblonga, villosa, pentasperma.

- Menis.* 8. VICIA pedunculis multifloris, petiolis sulcatis subde-
decaphyllis: foliolis lanceolatis glabris.
Vicia pedunculis multifloris, petiolis polyphyllis: foliolis
lanceolatis glabris. *Hort. ups.* 219. *
Habitat in Sibiria. ♂
- benghalens.* 9. VICIA pedunculis subquinquefloris, foliolis integerri-
mis, stipulis integris, leguminibus erectiusculis.
Vicia pedunculis subquinquefloris longitudine foliorum.
Hort. cliff. 368. *Hort. ups.* 219. *Roy. lugdb.* 365.
Vicia benghalensis hirsuta & incana, siliquis pili. *Herm.*
lugdb. 623. t. 625.
Habitat in Benghala. ☉
* *Floribus axillaribus, subsessilibus.*
- Sativa.* 10. VICIA leguminibus sessilibus subbinatis erectis, foliis
retusis, stipulis notatis.
Vicia leguminibus erectis, petiolis polyphyllis: foliolis a-
cumine emarginatis, stipulis denticulatis. *Hort. cliff.*
368. *Hort. ups.* 218. *Fl. succ.* 601. *Roy. lugdb.* 365.
Dalib. paris. 218.
Vicia sativa vulgaris, semine nigro. *Bauh. pin.* 344.
Vicia sativa alba. *Bauh. pin.* 344.
Habitat inter Europæ segetes hodie. ☉
Stipulae subtus macula quasi inusta notata.
- Lathyroides.* 11. VICIA leguminibus sessilibus solitariis erectis glabris,
foliolis senis: inferioribus obcordatis,
Vicia lathyroides, purpureo-cæruleis floribus. *Herm. par.*
242. t. 242.
Vicia montana minima, flosculo purpureo. *Aman. acad.*
1. p. 167.
Habitat in Scotia, Lusatia. ☉
- Inter* 12. VICIA leguminibus sessilibus reflexis pilosis solita-
riis pentaspermis, corollæ vexillis glabris.
Vicia foliis subsessilibus obverse ovatis apice emargina-
tis, leguminibus solitariis villosis. *Dalib. pa. is.* 219.
Vicia leguminibus solitariis deorsum flexis hirsutis. *Sauv.*
monsp. 235.
Vicia sylvestris lutea, siliqua hirsuta. *Bauh. pin.* 345.
Vicia flore luteo pallido, siliquis hirsutis crassis brevibus
propendentibus. *Morij. hist.* 2. p. 62. f. 2. t. 21. f. 5
Habitat in Gallia.
Variat foliis lanceolatis & retusis, Leguminibus hirsu-
tis & subglabris. Stipulae alternatim integerrimæ &
tridentatæ.

13. VICIA leguminibus sessilibus reflexis pilosis penta-*hybrida*.
ispermis, corollæ vexillis villosis.
Habitat Monspelii, Massiliæ. ☉
Affinis *V. luteæ*. Foliola obverse ovata, retusa, parva,
pilis adpersa. Stipulæ non ad canlem fusca, sed totæ
virides. Flores flavi, supra subgrisei, pilosi. Cætera
V. luteæ.
14. VICIA leguminibus subsessilibus pendulis glabris te-*peregrina*.
traspermis, foliolis linearibus emarginatis.
Vicia peregrina angustissimis foliis, siliqua lata glabra.
Pluk. alm. 386. t. 233. f. 6.
Vicia angustifolia purpureo-violacea, siliqua lata glabra.
Magn. monsp. 276.
Habitat in Gallia. *D. Sauvages*. ☉
15. VICIA leguminibus pedicellatis subquaternis erectis, *sepium*.
foliolis ovatis integerrimis: exterioribus decrescen-
tibus.
Vicia leguminibus ascendentibus, petiolis polyphyllis,
foliolis ovatis acutis integerrimis. *Hort. cliff.* 362.
Fl. suæc. 602. *Roy. lugub.* 365. *Dalib. parif.* 219.
Sauv. monsp. 234.
Vicia sepium folio rotundiore acuto. *Bauh. pin.* 345.
Vicia sepium. *Riv. tetr.* 56.
Habitat in Europæ sepibus. &
16. VICIA leguminibus subsessilibus solitariis erectis, fo-*narbonensis*.
liolis semisubovatis, stipulis denticulatis. *Hort. upj.*
218. *Sauv. monsp.* 234
Vicia petiolis inferioribus tetraphyllis: superioribus po-
lyphyllis, foliolis ovatis cirrhiferis. *Roy. lugub.* 365.
Vicia petiolis tetraphyllis cirrhosis. *Hort. cliff.* 360.
Vicia narbonensis. *Riv. tetr.*
Faba sylvestris, fructu rotundo atro. *Bauh. pin.* 338.
Bana sylvestris. *Dod. pempt.* 516.
Habitat in Gallia, Anglia. ☉
17. VICIA caule erecto, petiolis absque cirrhis. *Hort. Faba*.
cliff. 369. *Hort. upj.* 218. *Mat. med.* 353. *Roy. lugub.*
366. *Dalib. parif.* 220.
Faba Bauh. pin. 338.
Bana s. *Phascolus*. *Dod. pempt.* 513.
6. *Faba minor* s. *equina*. *Bauh. pin.* 338.
Habitat in - - - ☉

772

CICER.

- arietinum.* 1. CICER foliolis serratis. *Hort. cliff.* 370. *Hort. ups.* 224. *Mat. med.* 359.
 Cicer sativum. *Bauh. pin.* 347.
 Cicer arietinum. *Dod. pempt.* 525.
Habitat inter Hispaniæ, Italiæ segetes. ☉

773
Lens

ERVUM.

- tetrasper-*
num 2. ERVUM seminibus compressis convexis.
 Cicer pedunculis bifloris, foliolis integerrimis, stipulis indivisis. *Hort. cliff.* 124. *Mat. med.* 360. *Sauv. monsp.* 233.
 Cicer pedunculis bifloris, seminibus compressis. *Hort. cliff.* 370. *Dalib. paris.* 346.
 Lens vulgaris. *Bauh. pin.* 346.
 Lens. *Dod. pempt.* 526.
Habitat inter Galliæ segetes. ☉
- hirsutum.* 3. ERVUM pedunculis multifloris, seminibus globosis binis.
 Cicer pedunculis multifloris, seminibus globosis. *Hort. cliff.* 370. *Fl. succ.* 607. *Roy. lugdb.* 389. *Dalib. paris.* 228. *Sauv. monsp.* 233.
 Vicia segetum cum siliquis plurimis hirsutis. *Bauh. pin.* 345.
 Cracca minor. *Tabern. ic.* 507.
Habitat in Europæ agris. ☉
- monanthos.* 4. ERVUM pedunculis unifloris.
 Vicia pedunculis unifloris, foliolis integerrimis, stipulis alternis dentatis. *Hort. ups.* 219. *
 Lens monanthos. *Herm. lugdb.* 360.
Habitat in Asia ruthenica. ☉
- Ervilia.* 5. ERVUM germinibus undato-plicatis. *Hort. ups.* 224. *Mat. med.* 361. *Sauv. monsp.* 237.
 Ervum leguminibus pendulis. *Hort. cliff.* 370.
 Orobus siliquis articulatis, flore majore. *Bauh. pin.* 346.
Habitat in Gallia, Italia. ☉

CYTI.

CYTISUS.

774

1. CYTISUS racemis simplicibus pendulis, foliis ovato-oblongis. *Hort. cliff.* 354. *Hort. ups.* 210. *Roy. lugdb.* 369. *Hall. helv.* 591.
Anagyris non foetida major alpina. Baub. pin. 391.
Anagyris non foetens minor. Baub. pin. 391.
Laburnum arbor trifolia anagyridi similis. Baub. hist. 2. p. 361.
Habitat in Helvetia, Sabaudia. b
2. CYTISUS racemis simplicibus erectis, foliolis ovato-oblongis. *Hort. cliff.* 354. *Roy. lugdb.* 369. *Sauv. monsp.* 190.
Cytisus glaber nigricans. Baub. pin. 390.
Cytisus IV. Clus. hist. 1. p. 95.
Habitat in Austria, Pannonia, Bohemia, Italia. b
3. CYTISUS racemis erectis, calycibus bractea triplici sessilifolius. auctis, foliis floralibus sessilibus.
Cytisus foliis saepius sessilibus, calycibus squamula triplici auctis. Vir. cliff. 71. *Hort. cliff.* 355. * *Roy. lugdb.* 369. *Sauv. monsp.* 191.
Cytisus glabris foliis subrotundis, pediculis brevissimis. Baub. pin. 390.
Cytisus glaber, siliqua lata. Baub. hist. 2. p. 373.
Habitat in Italia, Galloprovincia. b
4. CYTISUS racemis axillaribus erectis, foliolis sub-lanceolatis tomentosis: intermedio longius petiolato. *Fl. zeyl.* 354. * *Hort. ups.* 211.
Cytisus foliolis ovato-lanceolatis: intermedio petiolato, pedunculo ex alis multifloro. Hort. cliff. 354. *Roy. lugdb.* 369.
Cytisus folio molli incano, siliquis orebi contortis & acutis. Burm. zeyl. 86. t. 37.
Phaeocolus erectus incauus, siliquis torosis. Pluk. alm. 293. t. 213. f. 3.
Habitat in India.
5. CYTISUS pedunculis simplicibus lateralibus, calycibus hirsutis trifidis obtusis ventricosso-oblongis. *Hort. ups.* 211. *
Cytisus calycibus hirsutis sessilibus, pedunculis simplicibus brevissimis. Hort. cliff. 355. *Roy. lugdb.* 370. *Sauv. monsp.* 190.
Cytisus incanus, siliqua longiore. Baub. pin. 390.

A a a a

Cyni-

Cytisus. *Clus. hist.* 1. p. 94.

Habitat in Hispania, Sibiria, Austria, Italia. \mathfrak{h}

supinus.

6. *CYTISUS* floribus capitatis, ramis decumbentibus.

Roy. lugdb. 376. *Hort. ups.* 211.

Cytisus supinus, foliis infra & filiquis molli lanugine pubescentibus. *Bauh. pin.* 350.

Cytisus VII. species altera. *Clus. hist.* 1. p. 96.

β . *Cytisus* supinus, foliis incana lanugine inferne pubescentibus. *Bauh. pin.* 390.

Cytisus VII. species altera. *Clus. hist.* 1. p. 96.

γ . *Cytisus* incanus, folio oblongo, austriacus. *Bauh. pin.* 390.

Cytisus V. *Clus. hist.* 1. p. 195.

Habitat in Sibiria, Italia, Sicilia, Galloprovincia. \mathfrak{h}

Structura floris indicat summam cum C. hirsuto affinitatem.

æthiopicus.

7. *CYTISUS* racemis lateralibus strictis, ramis angulatis, foliolis cuneiformibus.

Cytisus æthiopicus, subrotundis incanis minoribus foliis, floribus parvis luteis. *Pluk. alm.* 128. t. 278. f. 3. *ant*

Habitat in Æthiopia. \mathfrak{h}

græcus.

8. *CYTISUS* foliis simplicibus lanceolato-linearibus, ramis angulatis.

Barba jovis, linariæ folio, flore luteo parvo. *Tournef. cor.* 44?

Habitat in Archipelagi insulis. \mathfrak{h}

monspessulanus.

9. *CYTISUS* floribus lateralibus, foliis hirsutis, caule erecto striato. *Sauv. monsp.* 191. †

Cytisus monspessulanus, medicæ folio, filiquis dense congestis & villosis. *Tournef. inj.* 648.

Habitat Monspelii.

Folia subseffilia. Foliola ovalia, tomentosa, incana. Legumina conferta, recta, villosa. Sauv.

argenteus.

10. *CYTISUS* floribus subsessilibus, foliis tomentosis, caulibus herbaceis.

Cytisus acaulis, floribus solitariis: foliolis sericeis lanceolatis. *Sauv. monsp.* 191.

Lotus fruticosus incanus filiquosus. *Bauh. pin.* 332.

Lotus asperior fruticosa narbonensis incana. *Lob. ic.* 2. p. 41.

Trifolium argenteum, floribus luteis. *Bauh. hist.* 2. p. 359.

Habitat in G. Narbonensi. \mathfrak{h}

Rami

DIADELPHIA DECANDRIA. 141

Rami versus radicem fruticantes, superne herbacei. Flores subsessiles, solitarii, duo vel tres, ex alis & apicibus ramorum. Legumina hirsuta.

11. CYTISUS foliis pinnatis.

pinnatus.

Phaseolo affinis arbor indica Coral dicta polyphyllus non spinosa, foliis mollibus. *Pluk. phys.* 104. f. 3.

Habitat in India. ♀

Folia pinnata foliolis 5, ovatis, integerrimis, glabris.

Racemus ad singulum dentem affigit flores 2. f. 3. pedicellatos.

ULEX.

775

1. ULEX foliis villosis acutis, spinis sparsis.

europæus.

Ulex folio sub singulis spinis subulato plano acuto.

Hort. cliff. 356. *Hort. nps.* 212. *Roy. lugdb.* 372.

Genista spinosa major, longioribus aculeis. *Bauh. pin.* 394.

Genista spinosa. *Dod. pempt.* 659.

β. Genista spinosa major, brevioribus aculeis. *Bauh. pin.*

394.

Habitat in Anglia, Gallia, Brabantia. ♀

2. ULEX foliis solitariis obtusis, spinis simplicibus ter- capensis.

minatibus. *Roy. lugdb.* 372.

Genista spartium bacciferum, ericæ foliis, africanum.

Pluk. alm. 166. t. 185. f. 6. *mal. Pet. gaz.* 9. t. 82.

f. 9.

Habitat in Æthiopia. ♀

ARACHIS.

776

1. ARACHIS. *Hort. cliff.* 353. *Hort. nps.* 228. *Roy. hypogæa.*

lugdb. 390.

Arachidna quadrifolia villosa, flore luteo. *Plum. gen.*

49. *Ebres. pict.* t. 5. f. 3.

Senna tetraphylla f. Absi congener hirsuta maderasp-

tensis, folliculos sub terra condens. *Pluk. alm.* 341.

t. 60. f. 2.

β. Arachiduoides americana. *Niss. act.* 1723. p. 387. t.

19.

Habitat in Brasilia, Peru. ☉

777

GLYCYRRHIZA.

1. GLYCYRRHIZA leguminibus echinatis. *Roy. lugdb. orbata.*

386.

A a a 3

Gly-

Glycyrrhiza capite echinato. Bauh. pin. 352.

Dulcis radix. Cam. epit. 423.

Habitat in Gargano Apulia, in deserto Napico Tataris. &

glabra.

2. GLYCYRRHIZA leguminibus glabris. *Hort. cliff. 490. Mat. med. 362. Roy. lugdb. 386. Sauv. monsp. 232.*

Glycyrrhiza siliquosa & germanica. Bauh. pin. 352.

Glycyrrhiza vulgaris. Dod. pempt. 341.

Habitat in Franconia, Gallia, Hispania, Italia. &

bisfuta.

3. GLYCYRRHIZA leguminibus hirsutis *Roy. lugdb. 386. Glycyrrhiza orientalis, siliquis hirsutissimis. Tournef. cor. 26.*

Habitat in Oriente.

CORONILLA.

Equestris.

1. CORONILLA fruticosa, corollarum ungvibus calyce duplo longioribus. *Roy. lugdb. 387. Hort. cliff. 363. Fl. succ. 590. It. gotl. 229.*

Colutea siliquosa f. scorpioides major. Bauh. pin. 397.

Colutea scorpioides. Cam. epit. 541.

- β. *Colutea siliquosa minor. Bauh. pin. 397.*

Habitat Geneva, Monspeli, Salerni, Vienna. &

juncos.

2. CORONILLA fruticosa, foliis quinis sublan-
ceolatis, stipulis lanceolatis.

Dorychnium luteum hispanicum carnosius. Barr. ic. 153.

Colutea caule genistæ fungoso. Bauh. hist. 1. p. 383.

Polygala major massiliotica. Bauh. pin. 349.

Habitat Massiliæ, Monspeli. &

valentina.

3. CORONILLA fruticosa, stipulis subrotundis.

Coronilla fruticosa, leguminibus teretibus, pedunculis multifloris, caule fruticoso. Hort. nps. 234. Roy. lugdb. 386. Sauv. monsp. 235.

Coronilla leguminibus teretibus, ungvibus corollæ brevibus, caule fruticoso. Hort. cliff. 363.

Coronilla leguminibus teretibus, ungvibus corollæ brevibus, cinnis ad summum enneaphyllis, foliis ovatis. Guett. stamp. 1. p. 231.

Polygala altera. Bauh. pin. 349.

Polygala valentina. Clus. hist. 1. p. 98.

Habitat in Hispania, Italia. &

4. CORONILLA fruticosa, foliolis undenis: extimo ma-
jore. *argentea*.
Colutea scorpioides odorata. *Alp. exot.* 17.
Habitat in Creta. ☉
5. CORONILLA herbacea, leguminibus falcato-gladi-
atis. *Hort. cliff.* 363. *Hort. upf.* 234. *Roy. lugdb.*
386. *Sauv. monsp.* 235.
Securidaca lutea major. *Bauh. pin.* 398.
Hedysarum primum. *Dod. pempt.* 546.
Habitat inter Hispaniæ segetes. ☉
6. CORONILLA herbacea, leguminibus erectis tereti-
bus torosis numerosis, foliis glabris. *Hort. cliff.* 363.
Hort. upf. 235. *Roy. lugdb.* 386. *Dalib. parif.* 232.
Sauv. monsp. 235.
Securidaca dumetorum major, flore vario, siliquis ar-
ticulatis. *Bauh. pin.* 349.
Securidaca 2, altera species. *Clus. hist.* 2. p. 237.
Habitat in Lusatia, Bohemia, Dania, Gallia. ☉
7. CORONILLA herbacea, leguminibus quinis erectis
teretibus articulatis. *Roy. lugdb.* 387.
Coronilla cretica herbacea, flore parvo purpurascente.
Tournef. cor. 44.
Habitat in Creta. ☉
Habet multa communia cum C. varia, at differt ab ea
præcipue umbellis ex quinque floribus, raro pluribus,
cum illa communiter viginti gerat.
8. CORONILLA caule scandente flaccido. *Roy. lugdb. scandens.*
387.
Coronilla scandens pentaphylla. *Plum. spec.* 19.
Habitat in America calidioro.

ORNITHOPUS. 779

1. ORNITHOPUS foliis pinnatis, leguminibus subar-
tuatis. *Hort. upf.* 234. *Sauv. monsp.* 236.
Ornithopus foliis pinnatis, articulis leguminum subro-
tundis lævibus. *Hort. cliff.* 364. *Roy. lugdb.* 383.
Dalib. parif. 232. *Gort. gelr.* 438.
Ornithopodium minus. *Bauh. pin.* 350.
Ornithopodium perpusillum. *Dalech. hist.* 487.
6. *Ornithopodium majus.* *Bauh. pin.* 350.
7. *Ornithopodium radice tuberculis nodosa.* *Bauh. pin.* 250.
Habitat in Angliæ, Belgii, Galliæ, Hispaniæ arenosis ☉

- compressus.* 2. ORNITHOPUS foliis pinnatis, leguminibus compressis rugosis. *Hort. cliff.* 364. *Roy. lugdb.* 383. *Sauv. monsp.* 236.
 Ornithopodio affinis hirsuta scorpioides. *Bauh. pin.* 350.
 Scorpioides leguminosa. *Dalech. hist.* 493.
Habitat in Italia, Sicilia. ☉
- scorpioides.* 3. ORNITHOPUS foliis ternatis subsessilibus appendiculatis: impari maximo. *Hort. cliff.* 364. *Hort. nps.* 234. *Roy. lugdb.* 383.
 Scorpioides portulacæ folio. *Bauh. pin.* 287.
 Scorpioides mathioli. *Dod. pempt.* 71.
Habitat in G. Narbonensi, Hispania, Italia inter segetes. ☉

HIPPOCREPIS.

- unifiliquosa.* 1. HIPPOCREPIS leguminibus sessilibus solitariis. *Hort. cliff.* 364. *Hort. nps.* 223. *Roy. lugdb.* 384. *Sauv. monsp.* 236.
 Ferrum equinum siliqua singulari. *Bauh. pin.* 349.
 Ferrum equinum vulgare. *Col. ecphr.* 1. p. 302. t. 300.
Habitat in Italia ☉
- multifligna-
sa* 2. HIPPOCREPIS leguminibus pedunculatis confertis: margine altero lobatis. *Hort. cliff.* 364. *Hort. nps.* 233. *Roy. lugdb.* 384. *Sauv. monsp.* 236.
 Ferrum equinum siliqua multiplici. *Bauh. pin.* 346.
 Ferrum equinum alterum polyceraton. *Col. ecphr.* 1. t. 300.
Habitat in Angliæ, G. Narbonensis, Hispaniæ, Italiæ cretaceis.
- comosa.* 3. HIPPOCREPIS leguminibus pedunculatis confertis: margine exteriore repandis. *Roy. lugdb.* 384. *Dalib. parif.* 232.
 Hippocrepis leguminibus pedunculatis confertis: margine exteriori lobatis. *Hort. cliff.* 364.
 Ferrum equinum germanicum, siliquis in summitate. *Bauh. pin.* 346.
 Ferrum equinum comosum, s. capitatum. *Col. ecphr.* 1. p. 302. t. 301.
Habitat in Germania, Italia, Gallia.

SCORPIURUS.

- vermiculata.* 1. SCORPIURUS pedunculis unifloris, leguminibus rectis undique squamis obtutis.

Scor-

Scorpioides filiqua crassa boeltii. *Tournef. inst.* 402.
 Scorpioides bupleuri folio, corniculis crassioribus &
 magis spongiosis instar litui contortis & in se con-
 volutis. *Morif. bist.* 2. p. 127. f. 2. t. 11. f. III.

Habitat in Europa australi. ☉

2. SCORPIURUS pedunculis bifloris, leguminibus ex-
 trorsum odruse aculeatis.

Scorpioides bupleuri folio, corniculis asperis & rugosis
 f. rigidis, striatis f. sulcatis litui instar contortis & in
 se convolutis. *Morif. bist.* 2. p. 127. f. 2. t. 11. f.
 IV.

Habitat in Europa australi. ☉

3. SCORPIURUS pedunculis subtrifloris, leguminibus *sulcata*,
 extrorsum spinis distinctis acutis.

Scorpiurus. *Hort. cliff.* 364. *Hort. nyl.* 233. *Roy. lugdb.*
 383.

Scorpioides bupleuri folio. *Baub. pin.* 287.

Scorpioides prius. *Dod. pempt.* 71.

Habitat in Europa australi. ☉

4. SCORPIURUS pedunculis subquadrifloris, legumi-
 nibus extrorsum spinis confertis acutis.

Scorpioides bupleuri folio, corniculis asperis magis in
 se contortis & convolutis. *Morif. bist.* 2. p. 127. f.
 2. t. 11. f. II.

Habitat in Europa australi. ☉

Species hasce omnes olim ex una specie ortas esse dubium
 non est, nec sufficit locus harum generationi, qui tum
 mutatus easdem redaceret; quæ itaque mixtura ha-
 rum produxerit constantes plantas? Qui has omnes
 aut conjungat aut distinguat videtur argumentis in-
 niti.

HEDYSARUM.

* *Foliis simplicibus.*

1. HEDYSARUM foliis simplicibus lanceolatis obtu-
 sis, caule fruticoso spinoso.

Genista-spartium spinosum, foliis polygoni. *Baub. pin.*
 394.

Genista-spinosa, flore rubro. *Wheel. itin.*

Agul. Rann. itin. 74. t. 74.

Habitat in Tataria, Persia, Syria, Mesopotamia. ☿

2. HEDYSARUM foliis simplicibus linearibus acutis, cau-
 le inermi

α α α 5

Or- bupleurifo-
 lium.

- Ornithopodium maderaspatanum*, bupleuri folio. *Pet. gaz.* 18. t. 11. f. 12.
Scorpioides maderaspat. graminis leucanthemi folii, filiquis nodosis. *Pluk. amaith.* 189. t. 443. f. 4.
Habitat in India.
- nummulariifolium.* 3. **HEDYSARUM** foliis simplicibus cuneiformibus. *Fl. zeyl.* 288. *
Onobrychis maderaspat. nummulariæ folio, ornithopodii filiquis. *Pet. gaz.* 41. t. 26. f. 4.
Habitat in India.
- maculatum.* 4. **HEDYSARUM** foliis simplicibus ovatis obtusis. *Hort. cliff.* 449. *Hort. ups.* 233. *Fl. zeyl.* 290. *Roy. lugdb.* 385.
Hedysarum humile, capparidis folio maculato. *Dill. elib.* 170. t. 141. f. 161.
Habitat in India.
- gangeticum.* 5. **HEDYSARUM** foliis simplicibus ovatis acutis.
Hedysarum monophyllum latifolium, siliculis plurimis spica longa digestis. *Burm. zeyl.* 113. t. 49. f. 2.
Onobrychis gangetica monophyllos, siliculis singularibus lævibus: foliatim per internodia discriminatis. *Pluk. alm.* 270. t. 50. f. 3.
Onobrychis zeylanica, folio singulari oblongo rotundo. *Raj. suppl.* 453.
Habitat in India.
- vaginale.* 6. **HEDYSARUM** foliis simplicibus cordato-oblongis, petiolis simplicibus, stipulis vaginalibus. *Fl. zeyl.* 287. *
Genista articulata repens, foliis crastiusculis planis acutis. *Burm. zeyl.* 104. t. 49. f. 1.
Habitat in India. ☉
- triquetrum.* 7. **HEDYSARUM** foliis simplicibus cordato-oblongis, petiolis alatis, ramis triquetris. *Fl. zeyl.* 286. *
Onobrychis zeylanica monophyllos, caule triangulo, petiolis foliorum alatis. *Burm. zeyl.* 176. t. 81.
Onobrychis zeylanica, aurantii folio. *Pet. ficc.* 247. *Raj. suppl.* 247.
Habitat in India.
- strobiliferum.* 8. **HEDYSARUM** foliis simplicibus, bracteis strobilorum cordatis obtusis. *Fl. zeyl.* 287. t. 3. *
Onobrychis indiæ orientalis, fagi foliis alternis, siliculis spica longa dispositis. *Raj. suppl.* 234.
Habitat in India. * *Fe-*

* *Foliis conjugatis.*

9. HEDYSARUM foliis binatis petiolatis: floralibus *diphyllum*, sessilibus. *Fl. zeyl.* 291. *

Hedysarum minus *diphyllum*. *Sloan. jam.* 73. *hist.* 1. p. 185. *Raj. suppl.* 450.

Onobrychis maderaspatana *diphyllus*, filiculis clypeatis hirsutis, minor. *Pluk. alm.* 270. t. 246. f. 2.

Nelam-mari. *Reed. mal.* 9. p. 161. t. 82. *Raj. suppl.* 404.

- β. Hedysarum bifolium: foliolis ovatis, filiculis asperis geminis inarticulatis. *Burm. zeyl.* 114. t. 50. f. 1.

Habitat in Indiis. ☉

* *foliis ternatis.*

10. HEDYSARUM foliis ternatis: bracteis strobilorum *pulchellum*, orbiculatis conjugatis *Fl. zeyl.* 292. *

Hedysarum trifoliatum frutescens, flore & fructu inter duo foliola absconditis. *Burm. zeyl.* 116. f. 52.

Onobrychis maderaspatana triphylla, filiculis eleganter foliaceis. *Raj. suppl.* 234.

Onobrychis indica triphyllus, foliis amplis mucronatis, siliquis parvis in spicam longam ex alis inter bina foliola circinnata arcte conniventia reconditis. *Pluk. amalt.* 161. t. 433. f. 7.

Habitat in India. ♀

11. HEDYSARUM foliis ternatis, pedunculis umbelli-*umbellatum*, ieris, caule fruticoso. *Fl. zeyl.* 293. *

Hedysarum trifoliatum arborecens, floribus ex alis foliorum, siliquis copiosis glabris. *Burm. zeyl.* 115. t. 51.

Habitat in India. ♀

12. HEDYSARUM foliis ternatis, caule suffruticoso, *biarticulatum*, leguminibus biarticulatis. *Fl. zeyl.* 296. *

Hedysarum triphyllum, filiculis glabris peltatis geminis inarticulatis. *Burm. zeyl.* 114. t. 50. f. 2.

Onobrychis zeylanica trifolia minor perennis. *Raj. suppl.* 457.

Habitat in India. ♀

13. HEDYSARUM foliis ternatis, floribus spicatis, leguminibus articulatis: infimo monospermo, stipulis *heterocarpis* setaceis. *Fl. zeyl.* 298. *

Hedysarum trifoliatum, filiculis inferioribus solitariis: superioribus articulatis. *Burm. zeyl.* 117. t. 53. f. 1.

Habitat in India.

14. HEDYSARUM foliis ternatis; leguminibus *mem- vifcidum* bra-

branaceis lævibus integris, caule ramisque hispids. *Fl. zeyl.* 295.

Habitat in India. 5

Legumina neutra sutura sinuata, sed omnino integerrima.

- canadense.* 15. HEDYSARUM foliis simplicibus ternatisque, floribus racemosis. *Hort. cliff.* 232.
Hedysarum foliis radicalibus simplicibus, caulinis ternatis, floribus laxè spicatis, leguminibus undulatis. *Hort. cliff.* 365. *Roy. lugdb.* 385.
Hedysarum triphyllum canadense. *Corn. canad.* 44. t. 45.
Onobrychis major perennis canadensis triphylla, siliculis articulatis asperis triangularibus. *Morif. bist.* 2. p. 130. f. 2. t. 11. f. 9.
Habitat in Virginia, Canada. 2
- canescens.* 16. HEDYSARUM foliis ternatis subtus scabris, caule hispido, floribus racemosis conjugatis. *Hort. wps.* 232.
Hedysarum foliis ternis solitariisque, caule hispido fruticoso. *Hort. cliff.* 365. *Gron. virg.* 87. *Roy. lugdb.* 385.
Hedysarum triphyllum fruticosum (lupinum), flore purpureo. *Sloan. jam.* 73. *bist.* 1. p. 185. t. 118. f. 2. *Raj. suppl.* 458.
Onobrychis americana, floribus spicatis, foliis ternis canescentibus, siliculis asperis. *Pluk. alm.* 270. t. 308. f. 7.
Habitat in Virginia, Jamaica. 2
- marilandicum.* 17. HEDYSARUM foliis ternatis: foliolis subrotundis, caule frutescente ramosissimo, leguminibus articulatis lævibus. *Gron. virg.* 174.
Hedysarum, trifoliatum, siliqua brevior. *Dill. elzb.* 174. t. 144. f. 171.
Hedysarum triphyllum marilandicum minus, siliquis compressis articulatis asperis brevioribus. *Raj. suppl.* 455.
Habitat in Carolina, Virginia.
- frutescens.* 18. HEDYSARUM foliis ternatis subovatis, caule frutescente. *Gron. virg.* 174.
Habitat in Virginia. 5
- seridiflorum* 19. HEDYSARUM foliis ternatis acutiusculis, caule erecto, racemis longissimis erectis. *Gron. virg.* 87.
Habitat in Virginia.
- hirtum.* 20. HEDYSARUM foliis ternatis ovalibus, caule fruticoso,

cofo, racemis ovatis, calycibus fructibusque hirtutis monospermis.

Trifolium fruticosum hirsutum, spicis oblongis pedunculatis. *Gron. virg.* 173. *

Habitat in Virginia. b

11. HEDYSARUM foliis ternatis ovatis, floribus gemi- *violaceum*,
natis, leguminibus nudis venosis monospermis.

Hedysarum foliis ternatis lanceolatis, leguminibus monospermis. *Gron. virg.* 87.

Habitat in Virginia.

Flores *violaceæ*, gemini, propriis pedicellis plerumque annexi; in aliis glomerati ex foliorum alis, in aliis vero filiformibus pedunculis hinc inde affixi. Legumina calyce triplo longiora, ovata, compressa, acuta, glabra, reticulata, monosperma.

22. HEDYSARUM foliis ternatis linearilanceolatis, *paniculatum*
floribus paniculatis.

Hedysarum panicula ramosa, foliolis linearilanceolatis.

Gron. virg. 86.

Onobrychis mariana triphylla, passifloræ pentaphyllæ angustiori folio & facie, siliquis dentatis asperis. *Pluk. mant.* 140. t. 432. f. 6.

Habitat in Virginia.

23. HEDYSARUM foliis ternatis, scapo florifero nu- *multiflorum*,
do, caule folioso angulato.

Hedysarum caule nudo longissimo folioso, florifero angulato *Gron. virg.* 86.

Habitat in Virginia.

24. HEDYSARUM foliis ternatis obcordatis, caulibus *repens*,
procumbentibus, racemis lateralibus.

Hedysarum caulibus procumbentibus, racemis lateralibus solitariis, petiolis pedunculo longioribus. *Gron. virg.* 86.

Habitat in Virginia.

25. HEDYSARUM foliis ternatis obcordatis, caulibus *triflorum*,
procumbentibus, pedunculis unifloris ternis. *Fl. zeyl.* 297. *

Hedysarum trifoliatum repens, acetosellæ folio glabro, caule hirsuto. *Burm. zeyl.* 119. t. 54. f. 2.

- β. Hedysarum trifoliatum repens, folio ovali glabro, caule hirsuto, siliqua plana articulata. *Burm. zeyl.* 118. t. 54. f. 1.

γ. He-

25. *Hedysarum procumbens*, trifolii-fragiferi folio. *Dill. elsb.* 172. t. 142. f. 169.
Habitat in India. ☉
- olubile.* 26. *HEDYSARUM* foliis ternatis ovato-oblongis, caule volubili. *Hort. cliff.* 499. *Roy. lugdb.* 385.
Hedysarum trifolium scandens, folio longiore splendente. Dill. elsb. 173. t. 143. f. 170.
Habitat in America septentrionali.
- * *Foliis pinnatis.*
- aipinnat.* 27. *HEDYSARUM* foliis pinnatis, leguminibus articulatis glabris pendulis, caule erecto. *Hort. nps.* 232.
Hedysarum saxatile, siliqua lævi, floribus purpureis, inodorum. Ann. rub. 116. n. 152. 153.
 β. *Onobrychis* semine clypeato lævi. *Bauh. pin.* 350. *prodr.* 149.
Habitat α. in Sibiria, β. in Helvetia. 2
Planta β differt caule humiliore, floribus majoribus paucioribus, calyce pubescente, at attributa specifica tot communia, ut vix sufficienter distinguatur.
- coronarium.* 28. *HEDYSARUM* foliis pinnatis, leguminibus articulatis aculeatis nudis rectis, caule diffuso. *Hort. cliff.* 365. *Hort. nps.* 231. *Roy. lugdb.* 385.
Onobrychis, semine clypeato aspero, major. Bauh. pin. 350.
Onobrychis altæta. Dod. pempt. 549.
Habitat in Italiæ pratis. 2
- spinosissimum* 29. *HEDYSARUM* foliis pinnatis, leguminibus articulatis aculeatis tomentosis, caule diffuso. *Hort. nps.* 231. *
Hedysarum clypeatum asperum, siliqua spinosissima, flore purpureo. Boerb. lugdb. 2. p. 51.
Habitat in Hispania. ☉
- flexuosum* 30. *HEDYSARUM* foliis pinnatis, leguminibus articulatis aculeatis flexuosis, caule diffuso.
Onobrychis major annua, siliculis articulatis asperis clypeatis undulatis junctis, flore purpureo-rubente. Moris. hist. 2. p. 130. *Raj. dist.* 929.
Habitat in Asia. ☉
- virginicum.* 31. *HEDYSARUM* foliis pinnatis, caule fruticoso, leguminibus articulatis glabris pedunculatis erectis.
Gron. virg. 174.
Habitat in Virginia.

32. HE-

32. HEDYSARUM foliis pinnatis, leguminibus mono- *Onobrychis*.
 spermis aculeatis, corollarum alis calyce brevioribus.
 Hedysarum foliis pinnatis, leguminibus subrotundis aculeatis. *Hort. cliff.* 365. *Hort. upf.* 251. *Roy. Lugdb.* 385. *Sauv. mosip.* 233.
Onobrychis folio viciæ, fructu echinato, major. Baub pin. 350.
 Caput gallinaceum belgarum. *Lob. ic.* 2. p. 81.
 β. *Onobrychis incana, foliis longioribus. Baub. pin.* 350.
prod. 149. n. *Burser. XIX.* 138.
Habitat in Sibiria, Gallia, Angliæ, Bohemiæ apricis, cretaceis. &
33. HEDYSARUM foliis pinnatis, leguminibus mono- *Caput galli*.
 spermis aculeatis, petalis æqualibus.
Onobrychis, fructu echinato, minor. Baub. pin. 350.
prod. 149. n. *Burf. XIX.* 135.
Onobrychis f. Caput Galli minus, fructu maximo insigniter echinato, Triumf. obs. 65.
Habitat in Galloprovinciæ littoribus maris. ☉
Corolla petala longitudine aequalia, calyce vix longiora.

INDIGOFERA. 783

1. INDIGOFERA leguminibus arcuatis incanis, racemosis foliis brevioribus. *Fl. zeyl.* 273. *Amœn. acad.* 1. p. 408. *Hort. upf.* 208. *Mat. med.* 348.
Indigofera foliis nudis. Hort. cliff. 487.
Anil f. Nil inodorum color. Baub. hist. 2. p. 945.
Ameri. Rheed. mal. 1. p. 101. t. 54.
Habitat in India. ♀
2. INDIGOFERA leguminibus pendulis lanatis tetragonis. *Fl. zeyl.* 272. * *Amœn. acad.* 1. p. 408.
Astragalus spicatus, filiquis pendulis hirsutis, foliis tericcis. Burm. zeyl. 37. t. 14.
Kattu-tagera. Rheed. mal. 1. p. 55. t. 30.
Habitat in India.
3. INDIGOFERA leguminibus horizontalibus teretibus, glabris foliis pinnatis ternatisque. *Fl. zeyl.* 274. *Amœn. acad.* 1. p. 408.
Colutea filiquosa glabra, ternis quinisve foliis, maderaspata, semine rubello. Pluk. alm. 113. t. 166. f. 1
Nir-pulli. Rheed. mch. 9. t. 67 *Raj. suppl.* 470.
Habitat in India. ☉

CRAC.

CRACCA.

- 784
virginiana. 1. CRACCA leguminibus retrofalcatis compressis villosis spicatis, calycibus lanatis, foliolis ovali-oblongis acuminatis. *Gen. nov.* 1090.
 Clitoria foliis pinnatis, caule decumbente. *Hort. cliff.* 498. * *Gen. virg.* 83.
 Flymus *Michx.* *gen.* 210.
 Orobus virginianus, foliis fulva lanugine incanis, foliorum nervo in spicam aheunte. *Pluk. mant.* 142.
 Cicer altragaloides virginianus hirsute pubescens, floribus amplis subrubentibus. *Pluk. alm.* 103. t. 23. f. 2.
Habitat in Virginia, Canada.
Caulis in loco natali erectus est.
- villosa*. 2. CRACCA leguminibus retrofalcatis villosis pendulis racemosis lateralibus, foliis glabris lanceolatis. *Fl. zeyl.* 199. * *Gen. nov.* 1090.
 Securidaca maderaspatensis, siliquis falcatis fulvis & villosis plurimis circum ramulos stellatim. positis. *Pluk. alm.* 399. t. 59. f. 6.
 β Coronilla zeylanica, siliquis fuscis hirsutis pilosis, flore albo. *Burm. zeyl.* 78. t. 33.
Habitat in India.
- maxima*. 3. CRACCA leguminibus strictis adscendentibus glabris, stipulis lanceolatis, foliolis oblongis glabris striatis. *Fl. zeyl.* 300. * *Gen. nov.* 1090.
Habitat in Zeylona.
- purpurea*. 4. CRACCA leguminibus strictis adscendentibus glabris racemosis terminalibus, stipulis subulatis, foliis oblongis glabris. *Fl. zeyl.* 301. * *Gen. nov.* 1090.
 Coronilla zeylanica herbacea, flore purpurascente. *Burm. zeyl.* 77. t. 32.
Habitat in Zeylona.
- undulosa*. 5. CRACCA spicis lateralibus pedunculatis, leguminibus strictis pendulis, foliolis emarginatis subtus villosis. *Fl. zeyl.* 302. * *Gen. nov.* 1090.
Habitat in Zeylona.
- sensitosa*. 6. CRACCA leguminibus binis lateralibus glabris, foliolis emarginatis subtus sericeis, caule fruticoso. *Fl. zeyl.* 303. *Gen. nov.* 1090.
Habitat in Zeylona. h

CLI-

CLITORIA. 785

1. CLITORIA foliis pinnatis. *Hort. cliff.* 360. *Hort. Ternate.* *upf.* 214. *Fl. zeyl.* 283. * *Roy. lugdb.* 369.
Flos clitoridis ternatensis. *Breyn. cent.* 76. t. 31.
Phaseolus indicus, glycyrrhizæ foliis, flore amplo cæruleo. *Comm. hort.* 1. p. 47. t. 24.
Schonga Cuspl. *Rheed. mal.* 8. p. 69. t. 38.
Habitas in India.
2. CLITORIA foliis ternatis, calycibus campanulatis *brasiliana*.
solitariis. *Hort. upf.* 215.
Clitoria foliis ternatis. *Hort. cliff.* 361. *Roy. lugdb.* 369.
Planta leguminosa brasiliana, phaseoli flore, flore purpureo maximo. *Breyn. cent.* 78. t. 32.
Habitas in Brasilia.
3. CLITORIA foliis ternatis, calycibus campanulatis *virginiana*,
geminis. *Gron. virg.* 83.
Clitorius trifolius, flore minore cæruleo. *Dill. elsb.* 90.
t. 76. f. 87.
Habitas in Virginia.
4. CLITORIA foliis ternatis, calycibus cylindricis. *mariana*,
Clitorius marianus trifolius subtus glaucus. *Pet. jec.* 243.
Habitas in America septentrionali.

786
GLYCINE.

1. GLYCINE foliis pinnatis ovato-lanceolatis. *Hort. upf. Apios.* 227.
Glycine radice tuberosa. *Hort. cliff.* 361. *Roy. lugdb.* 391. *Gron. virg.* 85.
Apios americana. *Corn. canad.* 200. t. 201. *Stiff. bot.* 29. t. 29.
Astragalus perennis spicatus americanus, scandens caulis, radice tuberosa. *Morif. hist.* 2. p. 102. f. 2.
t. 9. f. 1.
Habitas in Virginia. &
2. GLYCINE foliis pinnatis, caule perenni. *Hort. cliff. frutescens.* 361. *Roy. lugdb.* 391.
Phaseoloides frutescens caroliniana, foliis pinnatis, floribus cæruleis conglomeratis. *Hort. angl.* 55. t. 15.
Habitas in Carolina.
3. GLYCINE foliis pinnatis conjugatis: pinnis ovatis *Abrus*
oblongis obtusis. *Fl. zeyl.* 284. * *Hort. upf.* 228.
Phaseolus arborescens alatus & volubilis major *orientalis*.
B b b

lis, fructu coccineo: hilo nigro notato. *Pluk. alm.*
294. t. 214. f. 5.

Pisum Indicum minus coccineum. Bauh. pin. 213.

Abrus. Vess. Egypt. 25. *Hort. cliff.* 488.

Konni. Rheed. msk. 8. p. 71. t. 39.

Habitat in Indiis, *Ægypto.* 2, 5.

tomentosa. 4. *GLYCINE* foliis ternatis tomentosis, racemis axillaribus brevissimis, leguminibus dispermis.

Ononis caule volubili. Gron. virg. 81.

Anonis phaseoloides scandens, floribus flavis sessilibus.

Dill. elth. 30. t. 26. f. 29.

Habitat in Virginia. 2

comosa. 5. *GLYCINE* foliis ternatis hirsutis, racemis lateralibus. *Glycine* foliis ternatis. *Gron. virg.* 85.

Phaseolus inarianus scandens, floribus comosis. Pet. mus. 453.

Habitat in Virginiae madidis umbrosis.

javanica. 6. *GLYCINE* foliis ternatis, caule villoso, petiolis hirsutis, bracteis lanceolatis minimis.

Habitat in India.

Caulis volubilis, magnitudine Phaseoli vulgaris, adpersus pilis flavis retrorsum spectantibus. Folia magnitudine & figura Phaseoli, at pedicelli flavi pilis densis. Stipulae ad exortum petiolorum ovali-oblongae; at vero ad exortum petiolorum subulatae. Pedunculi foliarii, longitudine foliorum, terminati Spica ovato-oblonga, densa, ex floribus nutantibus, violaceis, interstinctis Bracteis minutissimis.

bracteata. 7. *GLYCINE* foliis ternatis nudiusculis, caule piloso, racemis pendulis, bracteis ovatis.

Glycine foliis ternatis, pedicellis bracteatis. *Gron. virg.* 173.

Habitat in Virginiae madidis, umbrosis.

Racemi nutantes. Stipulae ovatae, erectae. Caulis retrorsum griseo-pilosus. Flores Viciae Craccae dictae; Alae & Carina albae, Vexillo pallidae violaceae. Legumina compressa, semiorbiculata, Pedunculis elongatis dependentia.

bituminosa. 8. *GLYCINE* foliis ternatis, floribus racemosis, leguminibus tumidis villosis.

Phaseolus africanus hirsutus bituminosus, siliquis bullatis, flore flavo. Herm. Lugdb. 492. t. 493.

Habitat ad Cap. b. *Spei*

PHA-

PHACA. 787

1. PHACA caulescens erecta pilosa, leguminibus tereti-bericis, cymbiformibus.

Phaca leguminibus rectis. *Roy. lugdb.* 390.

Astragalus bæticus lanuginosus, radice amplissima.

Bauh. pin. 351.

Astragalus bæticus. *Clus. hist.* 2. p. 234.

Habitat in Hispania, Lusitania.

2. PHACA caulescens erecta glabra, leguminibus semi alpinis, ovatis.

Astragaloides elatior erecta, vicie foliis, floribus luteis, siliquis pendulis. *Amm. ruth.* 148.

Habitat in Lapponia, Sibiria.

3. PHACA acaulis, foliis pinnatis: pinnis digitatis. *Sibirica.*

Astragaloides incana non ramosa, floribus carneis.

Amm. ruth. 150. t. 19. f. 2.

Astragaloides hirtula minor, non ramosa, floribus purpureiscentibus. *Amm. ruth.* 249. t. 19. f. 1.

Habitat in Alpibus Sibiricis.

ASTRAGALUS. 788

* *Caulibus foliosis erectis, nec prostratis.*

1. ASTRAGALUS caulescens, spicis cylindricis sub-alopecuroides sessilibus, calycibus leguminibusque lanatis.

Astragalus capitulis oblongis sessilibus, calycibus & leguminibus lanatis. *Hort. cliff.* 361. * *Aman. acad.*

2. p. 359. * *Roy. lugdb.* 391.

Astragalus alpinus procerior alopecuroides. *Tournef. inst.* 416.

Habitat in Sibiria, Hispania.

2. ASTRAGALUS caulescens erectus, floribus glomeratis subsessilibus ex omnibus axillis foliaceis.

Astragalus orientalis maximus incanus erectus, caule ab imo ad summum florido. *Tournef. cor.* 29. itin. 2. p. 254. t. 254.

Astragalus dioica vulgo Christiana radix. *Dalech. app.* 26.

Habitat in Oriente.

Flores fere totum caulem cooperiunt.

3. ASTRAGALUS caulescens, capitulis globosis: pedunculis longissimis, foliolis emarginatis. *Hort. cliff.* 360. *Roy. lugdb.* 391.

B b b 2

Astra-

Astragalus orientalis villosissimus, capitulis rotundioribus, floribus purpureis. *Tournef. cor.* 29?
Habitat in Oriente.

pilosus.

4. *ASTRAGALUS caulescens erectus pilosus*, floribus spicatis, leguminibus subulatis pilosis.

Astragalus caulescens erectus, leguminibus erectiusculis pilosis subulatis: acumine erecto: *Hort. upf.* 226.

Astragalus villosus erectus spicatus, floribus flavescentibus. *Amm. ruth.* 166.

Cicer montanum lanuginosum erectum. *Bamb. pin.* 347. *prodr.* 148.

Habitat in Sibiria, Thuringia.

Hedysaroides 5. *ASTRAGALUS caulescens erectus*, stipulis vaginalibus.

Astragalus caule erecto ramoso, spica purpurea nitente terminato. *Hall. belv.* 567. t. 14.

Habitat in Sibiria, Helvetia. &

Facies distincta ab Astragalis. Flores *Hedysari.* Racemus ex singulis alis laxus. Stipulae vaginae instar caulem involvunt supra folium. Fructum neque ego vidi.

sulcatur.

6. *ASTRAGALUS caulescens erectus*, caule glabro sulcato.

Habitat in Sibiria. &

Planta alta, stricta, glaberrima, Caule sulcato. Folia ex oppositionibus 8. s. 9. cum impari pinnata: foliolis glabris, angusto-lanceolatis. Pedunculi erecti, stricti, foliis triplo longiores. Flores in spicam laxam sparsi, violacei, parvi, calyce brevissimo. Legumina oblonga, erecta.

golegiformis. 7. *ASTRAGALUS caulescens strictus glaber*, floribus racemosis pendulis, leguminibus triquetris utrinque inucronatis.

Astragalus caule erecto paniculato, pedunculis folio longioribus, floribus sparsis. *Hort. cliff.* 362. *Roy. lugdb.* 391.

Astragalus sylvaticus montanus procerior, albo flore, siliqua brevi glabra, semine luteo. *Amm. ruth.* 162.

Habitat in Sibiria.

Caulis bipedalis, strictissimus, glaber. Folia pinnata ex 25 ad 31 foliolis: petiolo basi tumido. Flores racemosi, penduli, flavo-albicantes. Legumina glabra, breviuscula, triquetra, acuta, intra calycem pedicellata. Semina utrinque 2.

8. A-

8. ASTRAGALUS caulescens erectiusculus, floribus ^{uliginosus.}
 spicatis, leguminibus erectiusculis nudis tumidis tere-
 ti-depressis: mucrone reflexo. *Hort. nps. 226.* *
Habitat in Sibiriae pratis subhumidis. D. Gmelin. &
Corinae floris apex violaceus.
9. ASTRAGALUS caulescens erectus laevis, peduncu- ^{carolinensis.}
 lis spicatis, leguminibus ovato-cylindricis stylo acu-
 minatis.
Astragalus floribus racemosis, pedunculis longitudine fo-
liorum, caule erecto, calycibus glabris. Roy. lugdb.
391.
Astragalus procetior non repens, flore viridi-flavescente.
Dill. elsb. 45. t. 39. f. 45.
Habitat in Carolina. &
 * *Caulibus foliosis, diffusis.*
10. ASTRAGALUS caulescens diffusus, leguminibus ^{canadensis.}
 subcylindricis mucronatis, foliolis subtus subvillosis.
Astragalus canadensis, flore viridi flavescente. Tournesf.
inst. 416. conf. Dill. elsb. 46.
Habitat in Virginia, Canada. &
Foliola subtus tenuissimis tomentosa sunt.
11. ASTRAGALUS caulescens prostratus, leguminibus *Cicer.*
 subglobosis inflatis mucronatis pilosis. *Hort. nps. 226.*
Astragalus leguminibus subglobosis, floribus spicatis, cau-
le erecto. Hort. cliff. 362. Roy. lugdb. 392.
Cicer sylvestre, foliis oblongis hispids, majus. Baub.
pin. 347.
Cicer sylvestre. Cam. epis. 205.
Habitat in Austria, Helvetia, Italia. &
Foliola oblonga, sublancoolata sunt.
12. ASTRAGALUS caulescens erecto-patulus: follo- ^{microphyllus}
 lis ovalibus, calycibus tumidiusculis, leguminibus sub-
 rotundis.
Habitat in Sibiria.
Habitus A. Ciceris dicti, at quadruplo minor. Caulis
erectus, flexuosus, pedalis, non glaber. Rami paten-
tes, breviores. Folia pinnata: foliolis 13 f. 15, ovali-
bus, obtusis, interdum emarginatis. Stipulae subulatae,
minimae. Pedunculi solitarii, foliis longiores, superne
floribus horizontalibus instructi. Calyces glabri, sub-
rotundi, parum ventricosi, pilis raris vix conspicuis
adpersi. Corollae flavae, calyce duplo longiores. Legu-
mina subrotunda, inflata, villosa.

B b b 3

13. A-

- glycyphyllos.* 13. **ASTRAGALUS** caulescens prostratus, leguminibus
 subtriquetris arcuatis, foliis ovalibus pedunculo lon-
 gioribus.
Astragalus leguminibus lunatis biventricosis, caulibus
 procumbentibus. *Hort. cliff.* 362. *Fl. suec.* 591. *Roy.*
lugdb. 392.
Astragalus luteus perennis procumbens vulgaris. *Morif.*
hist. 2. p. 107.
Glycyrrhiza sylvestris, floribus luteo-pallescentibus.
Bauh. pin. 352.
Habitat in Europæ remotis. ☉
- hamosus.* 14. **ASTRAGALUS** caulescens procumbens, legumi-
 nibus subulatis recurvatis glabris. *Hort. upf.* 226.
Sauv. monsp. 237.
Astragalus leguminibus pendulis recurvis falcatis. *Hort.*
cliff. 362. *Roy. lugdb.* 392.
Astragalus monspelianus. *Clus. hist.* 2. p. 234.
Securidaca lutea minor, corniculis recurvis. *Bauh. pin.*
 349.
Habitat Messanæ, Monspeli. ☉
- contortuplicatus.* 15. **ASTRAGALUS** caulescens procumbens, legumi-
 nibus contortuplicatis canaliculatis villosis.
Astragalus acaulis, scapo foliis brevior, leguminibus
 subulatis canaliculatis retortis hirsutis. *Hort. upf.* 227.
Astragalus repens, siliquis undulatis. *Buxb. cent.* 3. p.
 22. t. 39.
Astragalus hirsutus procumbens, viciæ foliis cordatis,
 siliquis incurvis. *Amm. ruth.* 164.
Habitat in Sibiria. ☉
Mire variat ex solo quoad magnitudinem & staturam.
- boeticus.* 16. **ASTRAGALUS** caulescens procumbens, spicis pe-
 dunculatis, leguminibus prismaticis rectis triquetris a-
 pice uncinatis. *Hort. upf.* 225.
Astragalus caule erecto, floribus spicatis, leguminibus
 oblongis acinaciformibus: apice recurvo. *Roy. lugdb.*
 391.
Astragalus annuus, foliis & siliquis longioribus glabris
 ad foliorum ortum communi pediculo innixis. *Pluk.*
alm. 60.
Astragalus luteus annuus boeticus maritimus rectus al-
 ter. *Morif. hist.* 2. p. 108. f. 2. t. 9. f. 11.
Securidaca sicula, siliquis-foliaceis. *Bocc. sic.* . . .
Raj. hist. 935.
Habitat in Sicilia, Lusitania, Hispania. ☉

17. **ASTRAGALUS** caulescens diffusus, capitulis subsessilibus lateralibus, leguminibus erectis subulatis acuminis reflexis. *Hort. cliff.* 361.

Astragalus aniuus, foliis & siliquis hirsutis plurimis in foliorum alis sessilibus. *Pluk. alm.* 60. t. 79. f. 3.

Ornithopodio affinis hirsuta fructu stellato. *Baub. pin.* 350.

Vicia sesamea spula. *Col. ephr.* 1. p. 303. t. 301.

Habitat in G. Nathonenti, Italia. ☉

18. **ASTRAGALUS** canescens procumbens, leguminibus capitatis cordatis acutis reflexis complicatis.

Astragalus siliquis cordatis acutis pendulis: lateribus conniventibus. *Hort. cliff.* 362. *Roy. Lugdb.* 392.

Astragalus hispanicus, siliqua epiglottidi similis, flore purpureo, major, *Herm. Lugdb.* 74. t. 75.

Astragaloides incana, flore purpureo lentis siliqua. *Barr. ic.* 537. f. 1.

Habitat in Hispania. ☉

19. **ASTRAGALUS** caulescens procumbens, capitulis syriacis, pedunculatis, floribus reflexis, leguminibus tomentosis.

Astragalus syriacus hirsutus. *Baub. pin.* 351?

Astragalus syriacus. *Lob. ic.* 79?

Astragalus syriacus hirsutus, magno flore. *Morif. hist.* 2. p. 96. t. 8. f. 2?

Habitat in Sibiria. 2

Planta refert figuram anthorum, sed alior est; facie proprie accedit ad *A. arenarium*, a quo differt Planta triplo majore, minus incana; Floribus numerosis capitato spicatis, nec racemosis; Floribus triplo majoribus magis reflexis. Sunt etiam Folia lineari-lanceolata. Legumina oblonga, pilosa, erecta.

20. **ASTRAGALUS** subcaulescens procumbens, floribus subracemosis erectis, foliis tomentosis.

Astragalus pedunculis folia villosa æquantibus, floribus erectis, caulibus procumbentibus. *It. scan.* 108. *

Astragalus incanus parvus nostras. *Pluk. alm.* 59. *Raj. angl.* 3. p. 326. t. 12. f. 3. mala.

Glaux montana purpurea nostras. *Raj. hist.* 939.

Habitat in Angliæ, Scaniæ arena mobili. 2

21. **ASTRAGALUS** caulescens diffusus, capitulis pedunculatis imbricatis ovatis, floribus erectis.

Ciceri sylvestri minori affinis. *Baub. pin.* 347.

B b b 4

Glaux

Glaux dioscoridis, Clus. hist. 2. p. 240. t. 241.

Habitat in Hispania.

Radix lignosa. Caules spirhami, plurimi, superne magis subvillosi. Folia pinnata: foliolis 21. f. 23, minimis, ovali-oblongis, subtus pilis albis adspersis. Pedunculi ex summis foliorum alis, foliis saepius breviores, pilosi. Capitulum trifolii facie, evatum, calycibus ante & post florescentiam apicibus nigris. Corollæ erecta, acuminata, vexillo fere dupla longiores.

Onobrychis. 22. **ASTRAGALUS** caulescens diffusus, pedunculis spicatis, vexillis flore dupla longioribus.

Onobrychis spicata, flore purpureo. Baub. pin. 350.

Onobrychis 1. Clus. hist. 2. p. 238.

Habitat in Austria. 2

alpinus. 23. **ASTRAGALUS** subcaulescens procumbens, floribus pendulis racemosis, leguminibus utrinque acutis pilosis.

Astragalus pedunculis folio longioribus, floribus laxo spicatis pendulis. Hort. cliff. 362. Fl. suec. 592. Roy. lugdb. 392.

Astragalus alpinus minimus. Fl. lapp. 267. t. 9. f. 1.

Astragalus alpinus, folio vicie. Schench. alp. 509. f. 7.

Habitat in Alpibus Lapponicis, Helveticis. 2

* *Scapo nudo, absque caule folioso.*

non recens. 24. **ASTRAGALUS** subcaulos, scapis folio longioribus, floribus laxo spicatis erectis. *Roy. lugdb. 392.*

Astragalus quidam montanus vel Onobrychis aliis. Baub. hist. 2. p. 240

Onobrychis floribus vicie majoribus cæruleo-purpureis f. foliis tragacanthæ. Baub. pin. 351.

Onobrychis IV. Clus. hist. 2. p. 240.

Habitat in Helvetia, Valtellina. 2

Differt ab A. alpino, quod caulis vix ac ne vix basiramosus: quodque flores rubicundi, erecti; nec penduli.

vegetans. 25. **ASTRAGALUS** acaulos, scapis folio longioribus laxo spicatis, calycibus leguminibusque inflatis hirsutis.

Astragalus alpinus, tragacanthæ folio, vesicarius. Tournef. inst. 417. Magn. hort. 27. t. 27. Raj. hist. 454.

Habitat in Delphinatu. 2

phryoder. 26. **ASTRAGALUS** acaulos, scapis folia æquantibus, leguminibus inflatis subglobosis nudis.

Astra-

Astragalus acaulos, leguminibus inflatis subglobosis.

*Amern. acad. 2. p. 359. **

Habitas in Sibiria.

Bractæ villosæ; Fructus Globosus, sed magis globosus.

27. **ASTRAGALUS** acaulos, scapo erecto foliis longioribus, leguminibus subulatis inflatis villosis erectis. *Hort. nralensis. nps. 226. **

Astragalus non ramosus villosus & incanus spicatus, floribus purpureo-violaceis. Amm. rnth. 167.

Habitas in Sibiria. &

28. **ASTRAGALUS** acaulis, scapis declinatis longitudine foliorum, leguminibus subulatis teretibus subarcuatis glabris. *Hort. nps. 227. Sauv. monsp. 238. monspessulanus.*

Astragalus monspessulanus. Baub. hist. 3. p. 338.

Astragalus purpureus perennis nonspeliensis. Meris. hist. 2. p. 106.

Habitas Monspeli. &

29. **ASTRAGALUS** acaulos hirsutus, scapis erectis spicatis, foliolis ovatis obtusis. *grandiflorus.*

Astragalus, tragacanthæ folio, non ramosus, floribus luteis. Amm. rnth. 168?

Habitas in Sibiria D. Gmelin. &

Folia spichamæ: foliolis 17, ovalibus, obtusis, hirsutis præsertim subtus. Scapi crassi, pilosi, terminati Spica crassiuscula. Calyces hirsuti. Corollæ magna, flavæ.

30. **ASTRAGALUS** acaulos, calycibus leguminibusque villosis, foliolis acutis. *It. æl. 50. Fl. succ. 593. * campestris.*

Astragalus acaulos, foliis peracutis, calyce & fructu villosis. Hall. belv. 567. t. 13.

Astragalus perennis supinus, foliis & siliquis hispidis, flore luteo. Buxb. Hall. 32.

Habitas in Oelandia, Germania, Helvetia. &

31. **ASTRAGALUS** subacaulos, scapis subbifloris, leguminibus hamatis subulatis bicaripatis. *trimestris.*

Phaca leguminibus arcuatis. Roy. lugdb. 390.

Habitas in Ægypto. ☉

Planta tenera. Caules vix ulli aut breves. Scapi e radice s. pedunculi e caule, pauciflori subbiflori. Flores flavi. Legumen subulatum, in hamum flexum, compressum, externe duobus angulis compressis carinatum.

* *Caule lignoso.*

tragacanthoides.

32. ASTRAGALUS subacaulos, floribus radicalibus numerosis subsessilibus.

Astragalus acaulos, flore luteo. *Tournef. cor.* 28. *Bauh. sent.* 3. p. 21. t. 38. f. 2

Astragalus repens, barbæ jovis foliis, acaulos ferme polyanthos. *Amma. ruth.* 170.

Habitat in Sibiria, Armeria. 2

Caulis aut Scapus nullus, sed radicis ramis supraterram diffusis, & depressis, vestiti Foliis pinnatis, parvis, villosis. Flores aggregati ad radicem, sessiles. Calyces hirsuti dentibus nigricantibus. Corollæ flavæ. Legumina subrotunda, lævia.

Tragacantha

33. ASTRAGALUS caudice arhorescente, petiolis spiniscentibus.

Astragalus aculeatus fruticosus massiliensis. *Pluk. alm.* 60.

Tragacantha. Bauh. pin. 388. *Cam. epit.* 446. *Hort. cliff.* 361. *Mat. med.* 348.

3. Tragacanthæ affinis lanuginosa, f. Poterium. *Bauh. pin.* 388.

Tragacantha altera, Poterium forte. Clus. hist. 108.

Habitat in littore Massiliensi, Æthiæ, Olympo. b

BISERRULA.

789
Pelecinus.

1. BISERRULA. *Hort. cliff.* 361.

Astragalus purpureus annuus peregrinus, siliquis utrinque serræ similibus. *Morif. hist.* 2. p. 107. f. 2. t. 9. f. 6.

Securidaca siliquis planis utrinque dentatis. *Bauh. pin.* 349.

Securidaca peregrina. *Clus. hist.* 2. p. 238.

Habitat in Sicilia, Hispania, Galloprovincia. ☉

PSORALEA.

790
pinnata.

1. PSORALEA foliis pinnatis, floribus axillaribus. *Hort. upf.* 225. *

Psoralea foliis pinnatis: foliolis linearibus acuminatis. *Roy. lugdb.* 373.

Dorychnium caule fruticoso, foliis pinnatis: foliolis linearibus. *Hort. cliff.* 370.

Spartium africanum. *Riv. tetr.* 5.

Genistæ affinis arbor africana monospermos, flore caeruleo, foliis pinnatis. *Herm. lugdb.* 272. t. 273.

Ha-

- Habitat in Æthiopia. b*
 Pedunculi axillares, solitarii, eglames, hirsuti.
3. PSORALEA foliis ternatis, foliolis cuneiformibus re-aculatis.
 curvo-mucronatis.
 Psoralea foliis ternis: foliolis cuneiformibus triquetris a-
 culatis. *Roy. Ingd.* 373.
 Genita Spartium africanum trifolium. floribus cæruleis,
 foliis minimis in spinulam deinentibus. *Raj. denar.*
 104.
Habitat in Æthiopia. b
 Pedunculi axillares, plures, glumosi.
4. PSORALEA foliis inferioribus ternatis; superioribus tenuifolia.
 simplicibus lanceolato-subulatis.
Habitat in Æthiopia. b
 Pedunculi axillares, solitarii, glumosi.
5. PSORALEA foliis omnibus ternatis: pedunculis spi- bituminosa.
 catis folio longioribus. *Hort. upf.* 225.
 Psoralea trifolia. *B. Juss. act.* 1744. p. 381.
 Trifolium capitulis subrotundis, calycibus ventricosis,
 caule arborescente. *Hort. cliff.* 373.
 Trifolium bitumen redolens. *Bauh. pin.* 327.
 Trifolium bituminosum. *Dod. pempt.* 560.
 6. Trifolium bitumen redolens angustifolium. *Boerb.*
Ingd. 2. p. 32.
 Trifolium bituminosum inodorum. *Bast. evst. ast. t.* 11.
f. 2.
Habitat in Sicilia, Italia, Narbonæ collibus mariti-
mis. b
Varietas 6. differt foliis radicalibus numerois; Radi-
ce bicenni; Caule minore, angustiore, magis herbaceo.
5. PSORALEA foliis ternatis: foliolis subrotundis an-americana.
 gulatis, spicis axillaribus pedunculatis.
 Trifolium americanum. *Lob. sc. 2. p.* 31. *Dod. pempt.*
 567. *Bauh. pin.* 327.
Habitat in America?
 Canles diffusi, herbacei, punctis glandulosis adspersi.
 Folia ternata: intermedio majore, petiolato, subrotun-
 da, obtusissima, obsolete dentata aut angulata, adspersa
 punctis glandulosis, ut in congeneribus. Spicæ axil-
 lares, oblongæ, pedunculis longitudine foliorum. *Lob.*
 & *Dod. figuræ bene sunt. Hoc ære mihi non visum*
Hortis Europæis.

6. PSO-

764 DIADELPHIA DECANDRIA.

- Corylifolia.* 6. PSORALEA foliis simplicibus ovatis. *Hort. ups.* 225.
 Psoralea folio integro subrotundo. *Juss. act.* 1744. p. 382.
 Dorycnium foliis simplicibus ovatis. *Roy. lugdb.* 389.
 Loto affinis coryli folio. *Dodart. act.* 4. p. 289. t. 289.
Pluk. alm. 227. t. 96. f. 5.
Habitat in India. ☉
- pentaphylla.* 7. PSORALEA foliis digitato-quinatis: foliolis inaequalibus. *Hort. ups.* 225.
 Psoralea pentaphylla, radice crassa. *B. Juss. act.* 1744. p. 381. t. 17.
Habitat in Mexico vel Malabarica. 2
- Dalea.* 8. PSORALEA foliis pinnatis, spici terminalibus.
 Dalea. *Hort. cliff.* 363. t. 22. *
Habitat in America. ☉
 Foliola subius punctis glandulosis adpersa. Spica densa
 ut in Plantagine, alba aut cerulea.

EBENUS.

- 791
cretica. 1. EBENUS.
 Ebenus cretica. *Alp. exot.* 279. t. 278. *Pon. ital.* 128.
 Trifolium spici ovatis villosis, caule fruticoso. *Roy. lugdb.* 380.
 Anthyllis fruticosa, foliolis ternatis ac quinatis lanceolatis tomentosis. *Saur. meth.* 237.
 Cytisus incanus creticus. *Bauh. pin.* 390.
 Barba jovis cytisi folio, flore rubello. *Barr. rar.* 1389. t. 377. & 913.
 Loto affinis alata, folio & facie pentaphylloides fruticosi, floribus in spicam longiorem positus. *Pluk. alm.* 227. t. 67. f. 5.
Habitat in Creta.
 Bractea inter flores in spica distinguunt genus abunde a Trifoliis.

TRIFOLIUM.

- 792
 * Meliloti leguminibus nudis polyspermis.
M. carulea. 1. TRIFOLIUM spici oblongis, leguminibus seminudis mucronatis, caule erecto. *Hort. cliff.* 375. *Roy. lugdb.* 381.
 Lotus hortensis odorata. *Bauh. pin.* 330.
 Lotus sylvestris. *Cam. epit.* 724.
Habitat in Bohemia, Lybia.

2. TRI-

2. **TRIFOLIUM** leguminibus racemosis nudis mono-*Fl. indica*
spermis, caule erecto. *Hort. ups.* 223.
Trifolium erectum, caule teretiusculo, pedunculis com-
presso angulatis, pericarpis racemosis rotundis rugo-
sis nudiusculis monospermis. *Fl. zeyl.* 552. *
3. **Melilotus** futea indiae orientalis erecta, folliculis rotun-
dis parvis. *Pluk. alm.* 246. t. 45. f. 4.
β. **Melilotus** humilis erecta, floribus exiguis odoratis al-
bis. *Pluk. alm.* 347. t. 45. f. 5.
Habitat in India, Africa. ☉
3. **TRIFOLIUM** leguminibus racemosis nudis disper-*M. polonica*
mis lanceolatis, caule erecto.
Habitat in Polonia. ☉
Differt a T. Meliloti officinali Caule omnino tereti; Fo-
liis minoribus, acuminatis, superne acute serratis.
Racemo floribus rarissimis longiusque pedicellatis.
Pedunculis teretibus nec sulcatis. Florum vexillis rep-
licatis, nec alis extrorsum marginibus longitudina-
liter conniventibus. Leguminibus dispermis, parum
rugosis, lanceolato-acuminatis, longioribus. Statura &
odor conveniunt.
4. **TRIFOLIUM** leguminibus racemosis nudis disper-*M. officinalis*
mis, caule erecto. *Hort. cliff.* 376. *Hort. ups.* 223.
Fl. suec. 619. *Mat. med.* 355. *Ray. lugab.* 223.
Delib. parif. 225.
Trifolium odoratum f. Melilotus. *Dod. pempt.* 567.
Melilotus officinarum germaniae. *Bauh. pin.* 331.
γ. **Melilotus** officinarum germaniae, flore albo. *Tournef.*
inst. 407.
Lotus sylvestris, flore albo. *Tabern. hist.* 893.
Habitat in Europae campestribus. ☉. δ
5. **TRIFOLIUM** leguminibus racemosis nudis disper-*M. italica*
mis rugosis obtusis, caule erecto, foliolis integris.
Melilotus italica, folliculis rotundis. *Bauh. pin.* 331.
Melilotus italica. *Cam. hort.* 99. t. 29.
Habitat in Italia. ☉
6. **TRIFOLIUM** leguminibus racemosis nudis disper-*M. cretica*
mis membranaceis, caule erectiusculo.
Trifolium peltatum creticum. *Bauh. pin.* 359. *prodr.*
142. t. 142.
Habitat in Creta. ☉
Habitus Meliloti italicae, foliis crassis, vix serratis car-
neosis, inodoris, obtusis.

7. TRI-

M. cornutus
lata.

7. TRIFOLIUM floribus spicatis, leguminibus nudis mucronatis falcatis declinatis. *Hort. cliff.* 376. * *Roy lugdb.* 381. *Gort. gelr.* 435.

Melilotus, corniculis reflexis, major. *Bamb. pin.* 331.

Melilotus lutea major, corniculis reflexis ex eodem centro ortis. *Morif. hist.* 2. p. 162. f. 2. t. 16. f. 11.

Trifolium corniculatum 2. *Dod. pempt.* 573.

Habitat in Gargano, Apuliæ.

M. ornitho-
podioides.

8. TRIFOLIUM leguminibus nudis octospermis subternis calyce duplo longioribus, caulibus declinatis.

Medicago leguminibus ternatis erectis recurvis descendibus, pedunculo communi. *Hort. cliff.* 376.

Trifolium filiquosum, loto affine, filiquis ornithopodii. *Pluk. alm.* 375. t. 68. f. 1.

Fœna græcum humile repens, ornithopodii filiquis brevibus erectis. *Raj. angl.* 3. p. 331. t. 14. f. 1.

Habitat in Anglia, Gallia. ☉

* *Lotoidea leguminibus teetis polyspermis.*

Lupinaster.

9. TRIFOLIUM capitulis dimidiatis, foliis quinatis sessilibus, leguminibus polyspermis. *Hort. ups.* 225.

Lupinaster. *Buxb. act.* 2. p. 345. t. 20.

Lupinaster floribus purpureis & albis, filiquis minoribus.

Amm. rusb. 143. 144.

Habitat in Sibiria. 2

reflexum.

10. TRIFOLIUM capitulis fructiferis reflexis, leguminibus trispermis.

Trifolium leguminibus polyspermis, foliolis obverse ovatis denticulatis, floribus tetrapetalis, capitulis fructiferis reflexis. *Gron. virg.* 84.

Trifolium montano simile virginianum, floribus amœne purpureis amplioribus & magis patulis summo caule glomeratis per maturitatem reflexis. *Pluk. mant.* 485.

Habitat in Virginia.

lyridum.

11. TRIFOLIUM capitulis umbellaribus, leguminibus tetraspermis, caule adscendente.

Trifolialtrum pratense corymbiferum. *Mich. gen.* 28. t. 25. f. 2. 6.

3. Trifolium caule fistuloso erecto, capitulis subglobosis, foliolis obverse ovatis serratis. *Dalib. parif.* 226.

Trifolium orientale altissimum, caule fistuloso, flore albo. *Faill. parif.* 195. t. 22. f. 5.

Tri-

Trifolium flore albo. Riv. tetr.

Habitat in Europæ Cæltis.

22. TRIFOLIUM capitulis umbellaribus, leguminibus *repens*, tetraspermis, caule repente.

Trifolium capitulis subrotundis, flosculis pedunculatis, leguminibus tetraspermis, caule procumbente. Hort. cliff. 375. Fl. succ. 612. Mat. med. 354. Roy. lugdb. 373. Gron. virg. 85. Dalib. parif. 224.

Trifolium pratense album. Bauh. pin. 327.

Trifolium repens. Riv. tetr. 17. Fl. lapp. 274.

Habitat in Europæ pascuis. &

23. TRIFOLIUM capitulis umbellato-globosis imbricatis, vexillis deflexis persistentibus, leguminibus tetraspermis. *Roy. lugdb. 379.*

Lagopus americanus, floribus majoribus comosis. Pet. mus. 214.

Habitat in America.

24. TRIFOLIUM scapo nudo, leguminibus disperinis alpinum, pendulis, foliis lineari-lanceolatis.

Trifolium scapo nudo simplicissimo, foliis lineari-lanceolatis. Hort. cliff. 499. Sauv. monsp. 185.

Trifolium alpinum, flore magno, radice dulci. Bauh. pin. 328. prodr. 143.

Trifolium angustifolium alpinum. Pen. bald. 340. t. 340.

Anonis acaulis, floribus longissimis, foliis non serratis. Hall. belv. 588.

Habitat in Alpibus Italicis, Helveticis, Pyrenæis, Baldo. &

* *Lagopoda calycibus villosis.*

25. TRIFOLIUM capitulis villosis quinquefloris, involucri centrali reflexo rigido fructum involvente. *Wm.*

*Hort. cliff. 374. * Hort. ups. 222. Roy. lugdb. 379.*

Dalib. parif. 224.

Trifolium blesense. Dodart. mem. 4. p. 313.

Trifolium album tricoccum subterraneum gastonium reticulatum. Morif. bist. 2. p. 132. f. 2. t. 14. f. 5.

Trifolium pratense supinum catholiceps. Barr. ic. 881.

Habitat in Gallia, Italia. ☉

26. TRIFOLIUM capitulis villosis globosis, calycibus *globosum*, superioribus flosculo destitutis. *Hort. cliff. 374. **

Roy. lugdb. 379.

Tri-

Trifolium globosum repens. Baub. pin. 329. prodr. 143.

Trifolium glomerulis perforatæ cherleri. Baub. hist. 2.

p. 277.

Lagopus minor supinus, molli & compresso capite. Barr. 16. 859.

β. *Trifolium capitulis villosis globosis, calycibus omnibus fertilibus tomento connexis. Roy. lugdb. 379.*

Habitat Montspeli.

rubens. 17. **TRIFOLIUM** spicis villosis longis, corollis monopetalis, caule erecto, foliis ferrulatis. *Hort. cliff. 375.*

Roy. lugdb. 380. Saur. monsp. 184.

Trifolium montanum, spica longissima rubente. Baub. pin. 328.

Lagopus major alter. Dod. pempt. 578.

Habitat in Italia, G. Narbonenti, Helvetia. ☉

pratense. 18. **TRIFOLIUM** spicis subvillosis, cinctis stipulis oppositis membranaceis, corollis monopetalis.

Trifolium spicis villosis, caule diffuso, foliolis integerrimis. Hort. cliff. 375. Fl. suec. 612. Roy. lugdb. 580. Dalib. parif. 222.

Trifolium pratense, flore monopetalo. T. Fl. lapp. 273.

Trifolium pratense purpureum. Baub. pin. 327.

Trifolium pratense. Cam. epit. 582.

Habitat in Europæ graminosis. &

Spica sessilis inter Folia duo, opposita, sessilia, quorum Stipulae membranaceae, dilatatae, calycem quasi communem constituunt.

squarrosum. 19. **TRIFOLIUM** spicis subpilosis, calycum infimo dente longissimo reflexo, caule herbaceo.

Trifolium purpureum lagopoides hirsutum annuum rotundifolium, spica dilute rubente. Morif. hist. 2. p. 140. f. 2. t. 13. f. 1.

Trifolium angustifolium hispanicum, spica dilute rubente. Baub. pin. 329.

Habitat in Hispania. ☉

Habitus Trifolii pratensis. Folia subpilosa, lanceolata aut ovalia, emarginata. Spicae terminales, ovatae; calyces parum villosi, 5-dentati: dentibus nudis, setaceis 4 erectis; 5:0 seu infimo reliquis triplo majore, reflexo, unde Spica squarrosa & uncinata valde existit.

lappaceum. 20. **TRIFOLIUM** spicis globosis, calycibus setis rigidis obvallatis, caule erecto.

Tri-

Trifolium globosum f. capitulo lagopi rotundiore. *Bauh.*

pin. 329. *prodr.* 143

Trifolium capitulo glomerato rigido. *Bauh. hist.* 2.

p. 377.

Habitat Monspeli.

11. TRIFOLIUM spicis villosis oblongis obtusis aphyll- *incarnatum.*

lis, foliolis subrotundis. *Roy. lugdb.* 380. *Hort. ups.*

222. *Dalib. parif.* 225.

Trifolium spica rotunda rubra. *Bauh. pin.* 328.

Trifolium album incarnatum spicatum f. *Lagopus* maximum. *Bauh. hist.* 1. *p.* 376.

Lagopus maximum, folio & facie trifolii pratensis.

Lob. ic. 39.

Trifolium alopecurum latifolium, spica longa. *Barr.*

ic. 697.

Habitat in Italia. ☉

12. TRIFOLIUM spicis villosis conico-oblongis, den- *angustifoli-*

tibus calycinis setaceis subæqualibus, foliolis linearibus. *Hort. cliff.* 375. *Hort. ups.* 222. *Roy. lugdb.*

379.

Trifolium montanum angustissimum spicatum. *Bauh.*

pin. 328.

Trifolium alopecurum angustifolium elatius. *Barr. ic.*

698.

Habitat in G. Narbonensi, Italia. ☉

13. TRIFOLIUM spicis villosis ovalibus, dentibus ca- *arvense.*

lycinis setaceis æqualibus *Hort. cliff.* 375. *Fl. suec.*

616. *Roy. lugdb.* 379. *Dalib. parif.* 225.

Trifolium arvense humile spicatum f. *Lagopus.* *Bauh.*

pin. 328.

Lagopus. *Fuchs. hist.* 494. *Cam. epit.* 724.

Habitat in Europa, America septentrionali. ☉

14. TRIFOLIUM spicis pilosis, calycibus patentibus, *stellatum.*

caule diffuso, foliolis cuneiformibus.

Trifolium spicis ovatis pilosis, denticulis calycinis lanceolatis patulis. *Hort. cliff.* 375. * *Roy. lugdb.* 738.

Sav. monsp. 184.

Trifolium stellatum. *Bauh. pin.* 329. *prodr.* 143.

Lagopus minor erectus, capite globoso stellato. *Barr.*

ic. 860. & 755

Habitat in Sicilia, Italia, G. Narbonensi. ☉

15. TRIFOLIUM spicis ovatis, calycibus patulis: la- *clypeatum.*

C c c

cinia

cinia infima maxima lanceolata, foliolis ovatis. *Hort. cliff.* 373. * *Roy. lugdb.* 377.

Trifolium clypeatum argenteum. Alp. exot. 307. t. 306.

Habitat in Oriente. ☉

scabrum. 26. **TRIFOLIUM** capitulis sessilibus ovatis, calycibus villosis inæqualibus.

Trifolium spicis sessilibus lateralibus ovatis scabris. Hort. cliff. 373. * *Roy. lugdb.* 377. *Hall. botv.* 583. *Dalib. parif.* 225. *Sauv. monsp.* 183.

Trifolium capitulo oblongo aspero. Bauh. pin. 329. *prodr.* 140.

Trifolium ejus caules ex geniculis glomerulos oblongos proferunt. Bauh. hist. 2. p. 378.

Trifolium flosculis albis in glomerulis asperis cauliculis adnatis. Vaill. parif. 196. t. 33. f. 1.

Habitat in Anglia, Gallia, Italia. ☉

glomerotum. 27. **TRIFOLIUM** capitulis sessilibus globosis, calycibus striatis patulis æqualibus.

Trifolium capitulis sessilibus lateralibus subglobosis.

Hort. cliff. 373. * *Roy. lugdb.* 377. *Sauv. meth.* 183.

Trifolium cum glomerulis, ad caulium nodos rotundis. Raj. angl. 3. p. 329.

Trifolium arvense supinum verticillatum. Barr. ic. 882.

Habitat in Angliâ.

striatum. 28. **TRIFOLIUM** capitulis sessilibus sublateralibus ovatis, calycibus striatis rotundatis.

Trifolium capitulis sessilibus lateralibus ovatis, calycibus subrotundis villosis striatis. Fl. suec. 614. *Dalib. parif.* 225.

Trifolium parvum hirsutum, flore parvo dilute purpureo in glomerulis mollioribus oblongis, semine magno. Vaill. parif. 196. t. 33. f. 2.

Habitat in Germania, Gallia, Hispania.

montanum. 29. **TRIFOLIUM** capitulis terminalibus, corollarum vexillis subulatis, caule erecto, foliolis lanceolatis ferratis. *Fl. suec.* 611. *Dalib. parif.* 223.

Trifolium montanum album. Bauh. pin. 328.

Trifolium pratense album. Fuchs. hist. 818.

Habitat in pratis siccis Europæ septentrionalis. &

fruticans. 30. **TRIFOLIUM** spicis subvillosis: lacinia calycum infima maxima, caule fruticoso, foliis sessilibus.

Trifolium spicis subrotundis: lacinia calycum infima

ma-

maxima, foliis fere sessilibus. *Hort. cliff.* 373. * *Roy. lugdb.* 378.

Trifolium africanum fruticans, flore purpurascens.
Comm. hort. 2. p. 211. t. 106.

Habitat in Æthiopia. b

31. TRIFOLIUM acaule, pedunculis unifloris, floribus ~~unifloris~~.
foliisque commixtis.

Spica trifolia. *Alp. exot.* 169. t. 168.

Melilotus cretica humillima humifusa, flore albo magno. *Tournef. cor.* 28.

Habitat in Creta.

* *Vesicaria calycibus inflatis ventricosis.*

32. TRIFOLIUM spicis ovatis, calycibus inflatis glabris quinquedentatis, involucris univertalibus pentaphyllis. *Hort. cliff.* 373. * *Roy. lugdb.* 378.

Trifolium capitulo spumoso lævi. *Bauh. pin.* 329. *prodr.* 140.

Trifolium caule nudo, glomerulis glabris. *Bauh. hist.* 2. p. 379.

Habitat in Gallia, Italia, Apulia.

33. TRIFOLIUM spicis subovatis, calycibus inflatis ~~resupinatum~~ dorso gibbis, caulibus prostratis. *Hort. cliff.* 373. * *Roy. lugdb.* 378. *Gort. gelr.* 429.

Trifolium pratense folliculatum. *Bauh. pin.* 329.

Trifolium pratense folliculatum, flore inverso. *Barr. rar.* 824. t. 872.

Trifolium folliculatum f. *vesicarium minus purpureum.*
Bauh. hist. 2. p. 379.

Habitat in Anglia, Belgio.

Corollæ *resupinata in hoc sunt, ut vexillum respiciat peripheriam, carina centrum.*

34. TRIFOLIUM capitulis sessilibus globosis hirsutis, tomentosum. calycibus inflatis reflexis. *Sauv. monsp.* 175. *

Trifolium fragiferum tomentosum. *Magn. monf.* 265. t. 264.

Trifolium glomerulis tomentosis per caulium longitudinem. *Bauh. hist.* 2. p. 379.

Habitat in G. Narbonensi, Hispania, Lusitania, D. Sauvages. Læsing. 2

Caulis nudique decumbentes, non repentes. Folia obovata denticulata, obtusa, glabra, stipulis acuminatis; Pedunculi axillares, foliis non longiores. Capitula parva. Caly-

C c c 2

cis

cis dentes 3 superiores elongati, inferiores 2 parvi. Calyx crassissimo tomento involutus, ut vasa minime conspiciantur. Accedit propius *T. folliculato*, quam fragifero. Involucrum universale dodecaphyllum. Corollæ fere resupinatae.

- fragiferum.* 35. **TRIFOLIUM** capitulis subrotundis, calycibus infatis bidentatis reflexis, caulibus repentibus. *Hort. cliff.* 373. * *Fl. suec.* 613. *Roy. lugdb.* 378. *Dalib. paris.* 224. *Gert. gelr.* 428.

Trifolium fragiferum friscium. Baub. pin. 329. *Clas. cur. app.* 38. t. 39.

Trifolium fragiferum. Vaill. paris. 195. t. 22. f. 2.

Trifolium caule nudo, glomerulis glabris. Baub. hist. 2 p. 379. *fig. majore.*

Habitat in Suecia, Gallia. Anglia. &

Pedunculi in hoc foliis longiores, & Canles radices agunt. Calyces parum pubescentes.

* *Lupulina vexillis corollæ inflexis.*

- agrarium.* 36. **TRIFOLIUM** spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus nudis, caule erecto. *Fl. suec.* 617. *Hort. cliff.* 374. *Roy. lugdb.* 378. *Dalib. paris.* 226.

Trifolium pratense luteum, capitulo lupuli f agrarium. Baub. pin. 328. *Vaill. paris.* 196. t. 22. f. 3.

Trifolium agrarium. Dod. pempt. 576.

Habitat in Europæ pratis. ☉

Calyx ante florescentiam vix manifeste villosus f. nunc alterove pilo ad apices instructus. Corollæ flavae nec spadicæ.

- montanum.* 37. **TRIFOLIUM** spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus pilosis, caule erecto. *Trifolium montanum lupulinum. Baub. pin.* 328. *prodr.* 140.

Trifolium pratense, flore rufescente. Vaill. paris. 196.

Lupulinum montanum, capitulis spadicæis. Rapp. sen. 207.

Lotus montanus aureus, amplo lupuli capitulo, annuus. Barr. ic. 1024.

Habitat in Europæ pratis siccis. ☉

Differt a precedenti vexillis ferrugineis, calycibus præcipue ante florescentiam manifeste pilosissimis.

- procumbens.* 38. **TRIFOLIUM** spicis ovalibus, imbricatis, vexillis deflexis persistentibus, caulibus procumbentibus. *Fl. suec.* 618. *Is. goth.* 257.

Tri-

DIADELPHIA DECANDRIA. 773

Trifolium pratense luteo-croceum. *Vaill. parif.* 196.
Trifolium lupulinum alterum minus. *Raj. angl.* 3. p. 330. t. 14. f. 3.
Habitat in Europæ campestribus. 2

39. TRIFOLIUM spicis subimbricatis, vexillis deflexis *filiforme*.
 persistenteribus, calycibus pedicellatis, caulibus procum-
 bentibus.

Trifolium lupulinum minimum. *Morif. hist.* 2. p. 142.
Raj. angl. 3. p. 331. t. 14. f. 4.
Habitat in Angliâ. ☉

*Caulis filiformes, procumbentes. Pedunculi filiformes, fo-
 liis longiores, sustinentes Flores 3 ad 5, in satis sæpe
 12. ad 15, deflexos, Calycibus fructum ferentibus
 insidentibus manifestis & distinctis pedicellis.*

40. TRIFOLIUM spicis bifloris sessilibus, involucris hi- *biflorum*.
 spidis infundibuliformibus, foliis lanceolatis.

Trifolium caule piloso, foliolis subserratis, floribus late-
 ralibus subsolitariis, leguminibus ovatis. *Gron. virg.* 84.
Anonis mariana lutea, foliis angustioribus. *Pet. sic.* 84.
Habitat in Virginia, Canada.

*Caulis erectiusculus. Folia ternata, lanceolata: Petioli
 breves: Stipulis vaginalibus, dentibus duobus acumi-
 natis longitudine vaginæ. Capitula axillaria, infundi-
 buliformia, sessilia, foliolis pluribus obvoluta leviter
 trifidis: laciniâ mediâ longiore. Flores duo, alternatim
 prodeuntes.*

LOTUS. 793

- * *Leguminibus rarioribus nec capitulum constituentibus.*

1. LOTUS leguminibus solitariis membranaceo-quadrangulis, bracteis lanceolatis. *It. æl.* 143. *Fl. succ.* 610.
Sauv. monsp. 188.

Lotus tetragonolobus maritimus, flore luteo. *Bart. æst.*
 2. p. 346.

Lotus siliquosa lutea, siliquis pinnatis strictioribus & lon-
 gioribus. *Raj. hist.* 967.

Habitat in Europæ maritimis. 2

2. LOTUS leguminibus solitariis membranaceo-quadrangulis, bracteis ovatis. *Hort. ups.* 220.

*Lotus leguminibus subsolitariis: angulis quatuor mem-
 branaceis. Hort. cliff.* 372. *Roy. lugdb.* 388.

Lotus ruber, siliqua angulosa. *Bamb. pin.* 332.

C c c 3

Lo-

Lotus pulcherrima tetragonolobus. *Cam. hort.* 91. t. 26.
Habitat in Siciliae collibus. ☉

- conjungata.* 3. LOTUS leguminibus conjugatis membranaceo-quadrangulis, bracteis oblongo-ovatis.
 Lotus luteus, siliqua angulosa. *Boerb. lugdb.* 2. p. 37.
Habitat Monspeli. ☉
Convenit cum 2 habitu, aetate, foliis, hirsutiae, leguminibus; differt vero corollis duplo minoribus, leguminibus duobus, eorumque alis integris minus crispis, foliorum basilicalem non amplexante, sed ab opposito latere dehiscente.
- erecta.* 4. LOTUS leguminibus folitariis rectis teretibus terminalibus, caule erecto. *Sauv. monsp.* 189. †
 Lotus pratensis siliquosa lutea. *Bauh. pin.* 332. *minor*
et mollior. Tournesf. inst. 403.
 Lotus tritolia pratensis siliquosa monspeliensium. *Lob. ic.* 2. p. 42.
Habitat Monspelij.
- edulis.* 5. LOTUS leguminibus subfolitariis gibbis incurvis. *Hort. cliff.* 370. *Roy. lugdb.* 388.
 Lotus pentaphyllos, siliqua cornuta. *Bauh. pin.* 332
 Lotus edulis cretica. *Raj. hist.* 967.
 Lotus oligoceros cretica lutea, siliquis binis curvis pro-pendentibus. *Morif. hist.* 2. p. 176, f. 2 t. 18 f. 5.
Habitat in Italia, Sicilia, Creta.
- peregrinus.* 6. LOTUS leguminibus subbinatis linearibus compressis nutantibus, *Hort. cliff.* 372. *Roy. lugdb.* 388. *Sauv. monsp.* 188.
 Lotus, siliquis geminis, peregrina *Boerb lugdb.* 2. p. 38
Habitat in Europa australi.
- angustissimus* 7. LOTUS leguminibus subbinatis linearibus strictis erectis, caule erecto, pedunculis alternis,
 Lotus leguminibus saepius ternatis linearibus strictis erectis. *Hort. cliff.* 372. *Raj. lugdb.* 388. *Sauv. monsp.* 188.
 Lotus pentaphyllos minor hirsutus, siliqua angustissima. *Bauh. pin.* 332.
 Lotus corniculata, siliquis singularibus s. binis, tenuis. *Bauh. hist.* 2. p. 356.
Habitat in G. Narbonensi.

8. LOTUS leguminibus subquinatis arcuatis compressis, *ornithopodi-*
caulibus diffusis. *Hort. cliff.* 372. *Hort. upf.* 220. *oides.*
Roy. lugdb. 388. *Sauv. monsp.* 188.
Lotus filiquis ornithopodif. *Bauh. pin.* 332.
Lotus peculiaris filiquosa. *Cam. hort.* 91. t. 25.
Habitat in Sicilia. ☉

9. LOTUS leguminibus fabternatis, caule herbaceo ere- *jacobaeus.*
cto, foliolis linearibus. *Hort. cliff.* 372. *Roy. lugdb.*
388.
Lotus angustifolia, flore luteo purpurascente, insulae f.
jacobaei. *Comm. hort.* 2. p. 165. t. 83.
Habitat in Insula S. Jacobi. ☉

* *Pedunculis multifloris in capitulum.*

10. LOTUS capitulis dimidiatis, caule fruticoso, foliis *cretica.*
nitidis. *Hort. cliff.* 372. *Roy. lugdb.* 387.
Lotus argentea cretica. *Pluk. alm.* 226. t. 43. f. 1.
Lotus polyceratos fruticosa cretica argentea, filiquis lon-
gissimis propendentibus rectis. *Morif. hist.* 2. p. 177.
Habitat in Syria, Creta.

11. LOTUS capitulis hirsutis, caule erecto hirtio, legu- *hirsuta.*
minibus ovatis. *Hort. upf.* 220. *Sauv. monsp.* 188.
Lotus caule fruticoso, florum capitulis depresso, caly-
cibus lanigeris. *Hort. cliff.* 372. *Roy. lugdb.* 387.
Lotus pentaphyllos filiquosus villosus. *Bauh. pin.* 372.
Lotus polyceratos frutescens incana alba, filiquis erectis
crassioribus & brevioribus rectis. *Morif. hist.* 2. p.
177. f. 2. t. 18. f. 14.
Habitat in G. Narbonensi, Italia. &

12. LOTUS capitulis subglobosis, caule erecto, legumi- *recta.*
nibus rectis glabris. *Hort. upf.* 221.
Lotus caule fruticoso, florum capitulis globosis. *Hort.*
cliff. 372. *Roy. lugdb.* 387. *Sauv. monsp.* 188.
Lotus filiquosus glaber, flore rotundo. *Bauh. pin.* 332.
Lotus polyceratos frutescens hirsuta alba major latifo-
lia, filiquis tenuibus curtis rectis. *Morif. hist.* 2. p.
177. f. 2. t. 18. f. 13.
Habitat in G. Narbonensi, Sicilia, Calabria. &

13. LOTUS capitulis depresso, caulibus decumbentibus, *corniculata.*
leguminibus cylindricis.
Lotus caule herbaceo, florum capitulo depresso, legu-
minibus decumbentibus teretibus. *Hort. cliff.* 372.
Fl. spec. 609. *Roy. lugdb.* 388. *Dalib. parif.* 221.
C c c 4 Lotus

- Lotus f. Melilotus pentaphyllos minor glabra. *Bauh. pin.* 332.
 Trifolium corniculatum. *Dod. pempt.* 573.
 β. Lotus pentaphyllos, flore majore luteo splendente. *Bauh. pin.* 222.
 renuifolia. γ. Lotus pentaphyllos frutescens, tenuissimis glabris foliis. *Bauh. pin.* 332.
 Trifolium corniculatum frutescens, tenuissimis foliis. *Bauh. prodr.* 144. *Burf. XVIII.* 73.
Habitat in Europa. 2
Hujus forte varietas etiam γ. est, cujus Caules duplo longiores & angustiores, Folia linearia, & Legumina angustiora. Videtur in umbra nata; Barserus C. Bambizo hanc attulit.
 γ. 14. LOTUS capitulis dimidiatis, caule diffuso ramosissimo, foliis tomentosis. *Roy. lugdb.* 387.
 Lotus siliquosa maritima lutea, cythi facie. *Barr. rar.* 71. t. 1031.
Habitat in maritimis Europæ australis.
 Dorycnium. 15. LOTUS capitulis aphyllis, foliis sessilibus quinatis. *Dorycnium. foliis digitatis sessilibus. Hort. cliff.* 371.
Roy. lugdb. 389. *Sauv. monsp.* 214.
 Dorycnium monspeliensium. *Lob. ic.* 51.
 Trifolium album angustifolium, floribus veluti in capitulum congestis. *Bauh. pin.* 329.
Habitat in Hispania, G. Narbonensi, Austria. 2

TRIGONELLA.

294. γ. 1. TRIGONELLA leguminibus pedunculatis congestis pendulis linearibus rectis, foliolis sublanceolatis. *Trigonella leguminibus pedunculatis. Vir. cliff.* 76. *Hort. cliff.* 378. * *Roy. lugdb.* 382.
 Melilotus lupina angustifolia, medicæ facie, siliqua compressa. *Amm. ruth.* 119.
 Lotus montana humilior trifolia ad caulem alata lutea, siliqua modo recta, modo medicaginis instar falcata nigra. *Amm. ruth.* 119.
Habitat in Sibiria.
 Foliola lanceolata, obtusissima, superne acute serrata, dentata. Stipulæ integerrime. Legumina linearia.
 platycarpes. 2. TRIGONELLA leguminibus pedunculatis congestis pendulis ovalibus compressis, caule diffuso, foliolis subrotundis. *Hort. cliff.* 229.

Meli-

Melilotus supina latifolia, filiqua lata membranacea compressa. *Amm. rnb.* 158. *J. F. goett.* 1. p. 213. t. 13.

Melilotus filiqua membranacea compressa. *Alt. petrop.* 8. p. 209. t. 12.

Habitat in Sibiria. ♂

Foliola subovata, longitudinaliter acute serrata. Stipulae sapius dentatae. Legumina ovalia.

3. **TRIGONELLA** leguminibus pedunculatis congestis polyceratis. erectis rectis parallelis linearibus.

Medicago leguminibus confertis sessilibus erectis. *Roy. lugdb.* 382. *Sauv. monsp.* 187.

Medicago leguminibus confertis longis rectis parallelis, pedunculo communi. *Hort. cliff.* 376. *

Fœnum græcum sylvestre alterum polyceration. *Bauh. pin.* 348.

Fœnum græcum sylvestre alterum. *Dod. pempt.* 547.

Habitat in Hispania.

4. **TRIGONELLA** leguminibus pedunculatis congestis spinosis, declinatis subfalcatis compressis, pedunculo communi spinoso foliis longiore.

Medicago leguminibus sapius ternatis sessilibus arcuatis declinatis, spina ex alis. *Hort. cliff.* 377.

Fœnum græcum sylvestre polyceration creticum majus. *Breyn. cent.* 79. t. 33. f. 1.

Habitat in Creta. ☉

Stipulae dentatae; Pedunculus fit spina rigida ad cuius basin saepe Legumina 5, venosa, glabra.

5. **TRIGONELLA** leguminibus pedunculatis congestis monspeliaca, declinatis subfalcatis, pedunculo communi inermifoliis longiore.

Medicago leguminibus confertis sessilibus arcuatis inclinat. *Hort. cliff.* 377. *Roy. lugdb.* 382. *Dalib. parif.* 231. *Sauv. monsp.* 187.

Fœnum græcum sylvestre polyceratum minus monspeliacum. *Breyn. cent.* 80. t. 33. f. 2.

Habitat Montspeli. ☉

Stipulae simplices. Pedunculus apice desinit in acumen molle, sub quo Legumina 10 ad 12.

6. **TRIGONELLA** leguminibus sessilibus strictis erectiunculis subfalcatis acuminatis. *Fœnum græcum.* *Hort. upf.* 229. *Mat. cun. med.* 356.

C c c 5

Tri-

Trigonella leguminibus sessilibus. *Vir. cliff* 76. *Roy. lugdb.* 383. *Dalib. parif.* 228. *Sauv. monsp.* 187.

Medicago leguminibus subsolitariis sessilibus erectis reflexo-falcatis acuminatis. *Hort. cliff.* 376.

Fœnum græcum sativum. *Baub. pin.* 348.

Fœnum græcum. *Fuchf. hist.* 798. *Cam. epit.* 199.

β. *Fœnum* græcum sylvestre. *Baub. pin.* 348.

Habitat Monspelii. ☉

indica.

7. *TRIGONELLA* leguminibus sessilibus subsolitariis subsolatis, foliolis integerrimis, caule diffuso. *Fl. zeyl.* 235. *

Lotus maderaspatenus villosus, ornithopodii siliqua singulari. *Pluk. alm.* 226. t. 200. f. 7.

Habitat in India.

MEDICAGO.

1. *MEDICAGO* leguminibus lunatis margine integerrimis, caule arboreo. *Hort. cliff.* 376. *Hort. ups.* 230. *Roy. lugdb.* 382.

Cytisus incanus, siliquis falcatis. *Baub. pin.* 389.

Cytisus marantæ. *Lob. ic.* 2. p. 46.

Habitat in Rhodo, Neapoli. ☿

virginica.

2. *MEDICAGO* caule erecto ramosissimo, floribus fasciculatis terminalibus. *Gron. virg.* 86.

Lotus affinis trifoliata frutescens glabra. *Pluk. mant.* 120.

Habitat in Virginia.

radiata.

3. *MEDICAGO* leguminibus reniformibus margine dentatis, foliis ternatis. *Hort. cliff.* 377. * *Hort. ups.* 230. *Roy. lugdb.* 381. *Berg. viaar.* 95.

Medicago annua, siliqua falcata striata. *Morif. hist.* 2. p. 158. f. 2. t. 16. f. 3.

Trifolium siliqua falcata. *Baub. pin.* 330.

Lunaria radiata italorum. *Lob. ic.* 2. p. 38.

Habitat in Italia ☉

circinnata.

4. *MEDICAGO* leguminibus reniformibus margine dentatis, foliis pinnatis. *Hort. cliff.* 377. * *Hort. ups.* 230. *Roy. lugdb.* 382.

Lotus affinis siliquis hirsutis circinnatis. *Baub. pin.* 333.

Habitat in Hispania, Italia. ☉

sativa.

5. *MEDICAGO* pedunculis racemosis, leguminibus contortis, caule erecto glabro. *Hort. cliff.* 377. *Roy. lugdb.* 281.

Medica legitima. *Clasf. hist.* 2. p. 242.

Medica sativa *Morif. hist.* 2. p. 158. f. 2. t. 16. f. 2.

Fœ-

Fœnum burgundicum. Lob. ic. 2. p. 38.

Habitat in Hispaniâ, Galliâ apricis. 2

6. **MEDICAGO** pedunculis racemosis, leguminibus lufalcatis, caule prostrato. *Fl. suec. 620. Dalib. parif. 229.*

Trifolium sylvestre luteum, filiqua cornuta. Bauh. pin. 330.

Medica flavo flore. Clus. hist. 2. p. 243.

Habita: in Europæ pratis apricis, siccis. 2

7. **MEDICAGO** spicis ovalibus, leguminibus reniformi lupulina. bus monospermis, caule procumbente *Fl. suec. 621. Dalib. parif. 281.*

Trifolium leguminibus spicatis reniformibus nudis monospermis, caule procumbente. Hort. cliff. 375. Roy. lugdb. 380.

Trifolium pratense luteum, capitulo brevior. Bauh. pin. 328.

Trifolium pratense luteum. Fuchf. hist. 819.

Habitat in Europæ pratis. 3

8. **MEDICAGO** pedunculis racemosis, leguminibus marina. cochleatis spinosis, caule procumbente tomentoso. *Hort. cliff. 378. * Roy. lugdb. 381. Sauv. monsp. 186.*

Trifolium cochleatum maritimum tomentosum. Bauh. pin. 329.

Medica marina. Clus. hist. 2. p. 243.

Habitat in maris Mediterranei littoribus.

9. **MEDICAGO** leguminibus cochleatis, stipulis dentatis, caule diffusio. *Hort. cliff. 378. Hort. ups. 230. Roy. lugdb. 381. Dalib. parif. 230.*

Trifolium cochleatum, fructu nigro hispido. Bauh. pin. 329.

- a. **Medicago** leguminibus solitariis cochleatis compressis orbicularis, planis, stipulis ciliatis, caule diffusio. *Sauv. monsp. 186.*

Medica major dicarpos capsula compressa orbiculata nigra plana oris crispis. Morif. hist. 2. p. 152. f. 2. t. 15. f. 1.

Medica orbiculata. Bauh. hist. 2. p. 384.

Trifolium cochleatum f. scutellatum, folio latiore, fructu minuto ab uso. Bauh. pin. 329. prodr. 140.

- β. **Medica** cochleata major dicarpos, capsula rotunda scutellata. globo-

- globosa scutellata. *Morif. hist. 2. p. 152. f. 2. t. 15. f. 4.*
Medica scutellata. Bauh. hist. 3. p. 384.
Trifolium cochleatum, fructu latiore. Bauh. pin. 329.
- tornata. 7. *Medica tornata major & minor lenis. Park. theatr. 1116.*
- turbinata. 8. *Medicago fructu turbinato. Sauv. monsp. 187.*
Medica dicarpos, fructus capsula turbinata rugosa. Morif. hist. 2. p. 153.
Medica magna turbinata. Bauh. hist. 2. p. 385.
Trifolium cochleatum turbinatum, fructu compresso oblongo. Bauh. pin. 329.
- intertexta. 8. *Medica cochleata spinosa major dicarpos, capsula f. spinis longioribus sursum & deorsum tendentibus. Morif. hist. 1. p. 153. f. 2. t. 15. f. 8. 9. 7.*
- arabica. 8. *Medica cochleata minor polycarpos annua, capsula majore alba, folio cordato macula fusca. Morif. hist. 2. p. 144. f. 2. t. 15. f. 17.*
Medica arabica. Cam. hort. 97. t. 27.
Trifolium cochleatum, folio maculato cordato. Bauh. pin. 329.
- coronata. 7. *Medicago fructu minimo, coronulam utrinque dentatam referente. Sauv. monsp. 187.*
Medicago cochleata polycarpos annua, capsula minima coronata. Morif. hist. 2. p. 155, f. 25. t. 15. f. 21.
Medica coronata cherleri parva. Bauh. hist. 2. p. 386.
Trifolium folio obtuso, folliculis coronatis. Bauh. pin. 329. prodr. 141. n. 13.
- ciliaris. 9. *Medicago leguminibus congestis globosis hispidis, stipulis ciliatis. Sauv. monsp. 186.*
Medica annua, fructu ciliari. Tournef. inst. 410.
Trifolium fructu cochleato ciliari. Magn. monsp. 269.
- hirsuta. 8. *Medicago leguminibus solitariis globosis brevissime echinatis, stipulis ferratis. Sauv. monsp. 186.*
Medica echinata hirsuta. Bauh. hist. 2. p. 386.
Trifolium echinatum arvense. Bauh. pin. 329.
- rigidula. 10. *Medicago triphylla, leguminibus cochleatis spinosis, foliolis inferioribus cuneiformibus retusis, superioribus subrotundis. Dalib. parisi. 231.*
Medica hirsuta, echinnis rigidioribus. Bauh. hist. 2. p. 385.
Trifolium fructu compresso spinis horrido. Bauh. pin. 329.
- minima. 2. *Medicago leguminibus cochleatis aculeis uncinatis alternis externe donatis, stipulis integris. Guett. stamp. 249. Dalib. parisi. 230.*
 Medi-

DIADELPHIA DECANDRIA. 781

- Medica echinata minima. Bauh. hist. 2. p. 386.*
Trifolium echinatum, fructu minore. Bauh. pin. 330.
Medica cochleata dicarpos capsula spinosa rotunda mi- muricata.
nore. Moris. hist. 2. p. 153. f. 2. t. 15. f. 11.
Medicago fructu echinato, foliis linearibus dentatis. laciniata.
Sauv. monsp. 187.
Trifolium fructu echinato, foliis eleganter dissectis.
Magn. monsp. 271. t. 270.
Trifolium cochleatum spinosum syriacum, foliis laci-
niatis. Breyn. cent. 81. t. 34.

Clas-

POLYADELPHIA.

PENTANDRIA.

794

THEOBROMA.

Cacao.

1. THEOBROMA foliis integerrimis. *Hort. cliff.* 397. *Mat. med.* 364.
Cacao. Clus. exot. 55. *Sloan. jam.* 134. *bist.* 2. p. 15. t. 160. *Mer. surin.* 26. t. 26. *Geoffr. mat.* 409.
Catesb. car. 3. p. 6. t. 6.
 Arbor cacavifera americana. *Pluk. alm.* 40. t. 268. f. 3.
 Amygdalis similis guatimalensis. *Bauh. pin.* 442.
Habitat in America meridionali, Antillis. b

Guazuma.

2. THEOBROMA foliis serratis. *Hort. cliff.* 379. *Roy. lugdb.* 47.
 Guazuina arbor ulmifolia, fructu ex purpura nigro. *Plum. gen.* 36.
 Cenchramidea jamaicensis ulmifolia, fructu ovali integro verrucoso. *Pluk. alm.* 92. t. 77. f. 2.
 Alni fructu morifolia arbor, flore pentapetalo flavo. *Sloan. jam.* 135. *bist.* 2. p. 13. *Raj. dendr.* 11.
Habitat in Jamaicae campesribus. b

ICOSANDRIA.

797

CITRUS.

Medica.

1. CITRUS petiolis linearibus. *Hort. cliff.* 379. *Hort. upf.* 236. *Mat. med.* 366. *Roy. lugdb.* 266.

Malus Medica. *Bauh. pin.* 435.

Limon.

- β. Limon vulgaris. *Ferr. besp.* 193

Malus Limonia acida. *Bauh. pin.* 436.*Habitat in Asia: Media, Assyria, Persia. b**Varietates hujus hodie numerosissima una cum sequenti mixtae, de quibus Hesperides consulenda.*

Aurantium.

2. CITRUS petiolis alatis. *Hort. cliff.* 379. *Hort. upf.* 236. *Fl. zeyl.* 304. *Mat. med.* 367. *Roy. lugdb.* 266.

Malus Aurantia major. *Bauh. pin.* 346.Aurantia mala. *Cam. epist.* 150.

β. Ma-

- β. *Malus Aurantia*, cortice eduli. *Bauh. pin.* 346. *sinensis.*
 γ. *Malus aurantia* fructu rotundo maximo pallecente ca-
 put humanum excedente. *Sloan. jam.* 212. *bist.* 1.
p. 41. *t.* 12. *f.* 2. 3.
Pumpelmus. Meist. ikon. 84.
Habitat in India. δ

POLYANDRIA.

HYPERICUM.

798

* *Pentagyna.*

1. **HYPERICUM** floribus pentagynis, caule fruticoso, *balearicum.*
 foliis ramisque cicatrisatis.
Hypericum floribus pentagynis, folio & ramis verrucos-
 tis. *Hort. cliff.* 380. *Roy. lugdb.* 473.
Hypericum f. *Afcyrum* frutesceus, magno flore. *Magn.*
char. 260.
Myrto-Cistus pentaxi. *Clus. bist.* 1. *p.* 68.
Habitat in Majorca. δ
2. **HYPERICUM** floribus pentagynis, caule fruticoso, *kalmianum.*
 foliis lineari-lanceolatis.
Habitat in Virginia. *Kalm.* δ
Frutex. Rami tetragoni. Folia *Rosmarini* aut *Laven-*
dulae, glabra. Flores in *Corymbo* dichotomo, termina-
 li, ex tribus aut septem floribus. Capsula ovata, lon-
 gitudine calycis. Calyx lanceolatus, obtusiusculus. Sty-
 li 5, tenuissimi.
3. **HYPERICUM** floribus pentagynis, foliis lanceolatis *Lafiantha.*
 ferratis. *Hort. cliff.* 380.
Alcea floridana quinquecapsularis, laurinis foliis leviter
 crenatis, seminibus coniterarum instar alatis. *Pluk.*
amaltb. 7. *t.* 352. *f.* 3. *Gatesb. car.* 1. *p.* 44. *t.* 44.
Amm. acl. 7. *t.* 352. *f.* 3.
Habitat in Carolina, Surinamo. γ. *Bartsch.*
4. **HYPERICUM** floribus pentagynis, caule tetragono *Afcyrum.*
 herbaceo simplici, foliis lævibus integerrimis. *Hort.*
usp. 236.
Hypericum floribus pentagynis, foliis ovato-oblongis
 glabris integerrimis. *Hort. cliff.* 380. *Roy. lugdb.* 473.
Androsæmum flore & theca quinquecapsulari omnium
 maximis. *Meris. bist.* 2. *p.* 472.

Andro-

784 POLYADELPHIA POLYANDRIA.

Androsæmum constantinopolitanum, flore maximo.
Wheel. itin. 172. *Raj. hist.* 1017.
Habitat in Oriente, Sibiria, Canada. &

* *Trigyna*

Androsæ-
mum.

5. *HYPERICUM* floribus trigynis, fructu baccato, caule fruticoso ancipiti. *Hort. ups.* 237.

Hypericum floribus trigynis, fructu baccato, foliis ovalis pedunculo longioribus. *Hort. cliff.* 380. *Roy. lugdb.* 374. *Dalib. paris.* 235.

Androsæmum maximum frutescens. *Bauh. pin.* 280.

Androsæmum. Dod. pempt. 78.

Habitat in Angliæ, Ilvæ sepibus. &

Olympicum. 6. *HYPERICUM* floribus trigynis, calycibus acutis, staminibus corolla brevioribus, caule fruticoso. *Hort. cliff.* 380. *Roy. lugdb.* 374.

Hypericum montis olympi. Wheel. itin. 222. *Dill. elsb.* 182. t. 151. f. 183.

Ascyrum magno flore. Bauh. pin. 280. *prodr.* 130. *Burf. XVI.* 25.

Habitat in Pyrenæis, Olympo. &

canariense.

7. *HYPERICUM* floribus trigynis, calycibus obtusis, staminibus corolla brevioribus, caule fruticoso. *Hort. cliff.* 381. *Roy. lugdb.* 374.

Hypericum frutescens canariense multiflorum. Comm. hort. 2. p. 135. t. 68.

Hypericum s. Androsæmum magnum canariense ramosum, copiosis floribus, fruticosum. Pluk. alm. 189. t. 305. f. 1.

Habitat in Canariis. &

bifretidum.

8. *HYPERICUM* floribus trigynis, staminibus corolla longioribus caule fruticoso ancipiti. *Hort. cliff.* 331. *Hort. ups.* 237. *Roy. lugdb.* 374.

Hypericum foetidum frutescens minus. Dill. elsb. 182.

Androsæmum foetidum, capitulis longissimis filamentis donatis. Bauh. pin. 280.

Tragium. Gluf. hist. 2. p. 205.

Habitat in Sicilia, Calabria, Creta. &

egypticum.

9. *HYPERICUM* floribus trigynis, nectariis petalorum lanceolatis, caulibus suffruticosis compressis.

Habitat in Ægypto. D. B. Fussians. &

Frutex spitanæus, ramosissimus, brachiatus: ramulis compressis. Folia (Polygoni) ovata, minima, acuminata, aenea,

avenia, sessilia, articulis ramorum longiora. Flores saepius bini in ramulis. Squamula nectarifera oblonga, intra unguis cujusvis petali. Facies plantae Arenariae.

10. HYPERICUM floribus trigynis, stipulis reflexis *o-orientale*, vatis crenatis.

Hypericum orientale, parmicæ foliis. Tournesf. cor. 18. itin. 2. p. 220. t. 220.

Habitat in Oriente.

11. HYPERICUM floribus trigynis, foliis linearibus im *ericoides*, bricatis.

Hypericum ericoides minimum, foliis cinereis. Pluk. alm. 189. t. 93. f. 5.

Hypericum tomentosum lusitanicum minimum. Tournesf. inst. 256.

Habitat in Lusitania, Hispania.

12. HYPERICUM floribus trigynis, foliis lineari-lanceolatis, caule quadrangulo, pericarpis coloratis.

Habitat in Canada. Kalm.

Caules herbacci, teretes, ramosi, glabri. Folia linearia f. subanceolata, integerrima, glabra. Panicula dichotoma, floribus pedicellatis. Flores parvi admodum, magnitudine Spergulae. Pericarpium calyce duplo longius, conicum, rubrum.

13. HYPERICUM floribus trigynis, caule quadrato quadrangulo herbaceo. *Hort. cliff. 380. Fl. succ. 624. Roy. lugdb. lum. 473. Dalib. paris. 234.*

Hypericum Ascyron dictum, caule quadrangulo. Benth. bist. 3. p. 382.

Ascyron. Dod. pempt. 78.

Habitat in Europæ pratis. 2

14. HYPERICUM floribus trigynis, caule ancipiti, foliis obtusis pellucido-punctatis. *Hort. cliff. 380. Fl. succ. 625. Mat. med. 368. Roy. lugdb. 474. Hull. belv. 630. Dalib. paris. 233.*

Hypericum vulgare Benth. pin. 279.

Hypericon. Dod. pempt. 76.

Habitat in Europæ pratis. 2

15. HYPERICUM floribus trigynis, caulibus ancipiti-bus prostratis. †

Hypericum foliis ovatis, caulibus supinis. Guett. Raimp. 2. p. 174. Dalib. paris. 234.

D d d

Hyper-

Hypericum humilifolium foliis perforatis punctis in margine nigris. *Hali. helv.* 360.

Hypericum minus supinum f. *supinum* glabrum. *Bauh. pin.* 279.

Hypericum supinum III. *minimum*. *Clus. hist.* 2. p. 181.

Habitat in Europa australiore. &

Affinis minimum antecedentis.

hirsutum. 16. *HYPERICUM* floribus trigynis: calycibus serrato-glandulosis, caule erecto, foliis ovatis.

Hypericum foliis lanceolatis, calycinis laciniis linearibus. *Guett. stamp.* 176.

Hypericum floribus trigynis, calycum ferraturis capitatis, caule tereti. *Hort. cliff.* 380. *Fl. suec.* 626. *Koy. lugdb.* 474. *Dalib. parif.* 234.

Ascyron f. *Hypericum* bifolium glabrum non perforatum. *Bauh. pin.* 280.

Androsæmun campoclarense. *Col. ecphr.* 1. p. 73. t. 74. *It. wgoth.* 213.

β. *Hypericum* foliis ovato-oblongis brevipediculo donatis, calycinis laciniis oblongo-acutis. *Guett. stamp.* 178.

Androsæmun Ascyron dictum, caule rotundo hirsuto. *Morif. hist.* 2. p. 271. f. 5. t. 6. f. 11.

Androsæmun hirsutum. *Bauh. pin.* 280. *caute, Habitat in Europæ collibus & montibus. &*

tomentosum. 17. *HYPERICUM* floribus trigynis: calycibus serrato-glandulosis, foliis subovatis hirsutis.

Hypericum supinum tomentosum minus & monspeliacum. *Bauh. pin.* 279. *Sauv. monsp.* 149.

Hypericum supinum tomentosum alterum. *Clus. hist.* 2. p. 181.

β. *Hypericum* supinum tomentosum majus & hispanicum. *Bauh. pin.* 279.

Hypericum supinum tomentosum hispanicum. *Clus. hist.* 2. p. 181.

Habitat in G. Narbonensi, Hispania. &

pulchrum. 18. *HYPERICUM* floribus trigynis: calycibus serrato-glandulosis, foliis cordatis glabris. †

Hypericum foliorum basi latissima, foliis calycis brevioribus obtusis. *Hall. belv.* 361

Hypericum foliis cordatis connatis, laciniis calycinis ovatis obtusis. *Guett. stamp.* 2. p. 175.

Hypericum minus erectum. *Bauh. pin.* 279.

PCLYADELPHIA POLYANDRIA. 787

Hypericum minus glabrum erectum pulchrum. Moris.

hist. 2. p. 470.

Hypericum pulchrum tragi. Bauh. hist. 3. p. 383.

Habitat in Europa australiori. &

19. **HYPERICUM** floribus trigynis: calycibus serrato-^{nummulari-}
glandulosis, foliis cordato-orbiculatis glabris. ^{vinum,}

Hypericum nummulariæ folio. Bauh. pin. 279. prodr.
130.

Hypericum fere orbiculato folio, floribus amplis pallide
luteis, petalis in ambitu crenatis. Pluk. alm. 188. t.
93. f. 4.

Androsæmum supinum saxatile, nummulariæ folio aly-
sa parte rubente. Bocc. mus. 2 p. 134. t. 91.

Habitat in Pyrenæis. &

20. **HYPERICUM** floribus trigynis: calycibus serrato-*Coris*,
glandulosis, foliis subverticillatis.

Hypericum f. Coris legitima ericæ similis: Moris. hist.
2. p. 468. f. 5. t. 6. f. 4.

Hypericoides Coris quorundam. Bauh. hist. 3. p. 384.

Coris lutea. Bauh. pin. 280.

Habitat in Europa australi.

Stipulæ utrinque solitariæ f. ad singulum folium duæ,
magnitudine & figura foliorum, hinc, quum folia sint
opposita, quasi sena articulos cingunt, quorum quatuor
sunt stipulæ.

* Digyna.

21. **HYPERICUM** floribus digynis, foliis ovatis sessilibus. ^{mutidum.}
bns. Gron. virg. 88.

Habitat in Virginia, Canada.

22. **HYPERICUM** floribus digynis, foliis linearibus. ^{fetofum.}
Gron. virg. 88.

Hypericum virginianum parvum fruticosum, ramulis e-
quiseti. Pluk. alm. 189.

Habitat in Virgiuia.

ASCYRUM. 789

1. **ASCYRUM** foliis ovatis, caule tereti, panicula dicto- *Cruz andrea*
toma.

Hypericoides ex terra mariana, floribus exiguis luteis.

Pluk. mant. 104. Raj. suppl. 496.

Habitat in Virginia.

D d d 2

2. A-

783 POLYADELPHIA POLYANDRIA.

- Hypericoides* 2. ASCYRUM foliis ovatis, caule compresso.
 Ascyrum foliis ovatis. *Hort. cliff.* 494. *Gron. virg.* 88.
Hypericoides frutescens erecta, flore luteo. Plum. gen.
 51.
Hypericum pumilum sempervirens, caule compresso
ligneo ad bina latera alato, flore luteo tetrapetalo,
Pluk. mant. 104. *Raj. suppl.* 495.
Habitat in virginia.
- villosum.* 3. ASCYRUM foliis hirsutis, caule stricto.
Hypericum virginianum, frutescens pilosissimum. Pluk,
alm. 189. *t.* 245. *f.* 6.
Habitat in Virginia. †

Clas-

Classis XIX.

SYNGENESIA.

POLYGAMIA ÆQUALIS.

TRAGOPOGON. § 00

1. TRAGOPOGON calycibus corollæ radium æquantibus, foliis integris strictis. *pratenſe*.
 Tragopogon calycibus florem superantibus. *Hort. cliff.* 382. *Fl. succ.* 648. *Mat. med.* 372. *Roy. lugdb.* 119.
Dalib. parisi. 242.
 Tragopogon pratensis luteum majus. *Bauh. pin.* 274.
 Tragopogon. *Fuchs. hist.* 827.
Habitat in Europæ pratis apricis. ♂
2. TRAGOPOGON calycibus corollæ radio brevioribus, foliis integris subundulatis. *orientale*.
 Tragopogon orientale angustifolium, flore maximo luteo. *Tournef. cor.* 36.
 Barba hirci. *Cam. epit.* 312. *figura*.
Habitat in Oriente. ♂
3. TRAGOPOGON calycibus corollæ radiis longioribus, foliis integris strictis, pedunculis superne incrassatis. *porrifolium*.
Hort. nps. 243.
 Tragopogon purpureo-cæruleum, porri folio, quod Arteni vulgo. *Bauh. pin.* 274. *Morif. hist.* 3. p. 8. f. 2. 9. f. 5.
 Tragopogon alterum f. Barba hirci. *Dalcch. hist.* 1079.
Habitat - - - ♂
4. TRAGOPOGON calycibus corollæ radio longioribus, foliis integris, seminibus levibus: disci plumosis; radii setaceis. *hybridum*.
Hort. nps. 243. *
 Tragopogon gramineis foliis hirsutis. *Bauh. pin.* 275.
 Tragopogon alterum, gramineo folio, flore suave rubente. *Col. esphr.* 1. p. 232. t. 231.
Habitat in Italia. ☉
5. TRAGOPOGON foliis radicalibus lyratis rotundatis, caulinis indivisis.
 Tragopogon foliis lanceolatis sessilibus amplexicaulibus dentatis, caule ramoso. *Gron. virg.* 91?
 D d d 3. *Habitat*

790 SYNGENESIA: POLYGAMIA ÆQUALIS.

Habitat in Virginia, Canada. Kalm.

Folia radicalia plurima, lyrata: lobis rotundatis, glabris. Caules subnudi, erecti, striati. Folio uno alterove lanceolato, integro amplexicali, Ramis paucis. Rami terminantur Pedunculis tribus, ad quorum exortum folia 2, lanceolata, opposita: altero minore. Calyx in 12 folia ad basin divisus, æqualia. corolla breviora. Flores saturate lutei. Habitus Arnice gerberia. Pappus plumosus non est, nec stipitatus, sed simplex. Statura Hyoseridis; Calyx Tragopogonis. Semina Hieracii.

anicampii. 6. TRAGOPOGON calycibus corolla brevioribus infermibus, foliis lyrato-sinuatis. *Hort. upf.* 244. *Hort. cliff.* 382. * *Sauv. monsp.* 295.

Chondrilla foliis cichorei tomentosis. *Bauh. pin.* 130.

Chondrilla prior. *Clas. hist.* 2. p. 143.

Habitat in Hispania. Z

microides. 7. TRAGOPOGON calycibus corolla brevioribus aculeatis, foliis pinnato-hastatis. *Hort. cliff.* 382.

Picris calycibus simplicibus aculeatis, foliis incrinibus hastato-sinuatis. *Hort. upf.* 241. *Sauv. monsp.* 295.

Sonchus asper laciniatus creticus. *Bauh. pin.* 124. *prodr.* 60.

Hieracium majus, folio sonchi, semine incurvo. *Bauh. pin.* 127.

Chondrilla creticæ nomine missa semine crispo. *Bauh. hist.* 2. p. 1022.

Habitat in Creta, Montpelii. ☉

asperum. 8. TRAGOPOGON calycibus corolla brevioribus hispidis, foliis integris; caulinis oblongis. *Sauv. monsp.* 82.

Sonchus asper, subrotundo folio, major & minor. *Bauh. pin.* 124. *prodr.* 60.

Habitat Montpelii.

SCORZONERA.

humilis.

1. SCORZONERA caule subnudo unifloro, foliis nervosis planis. *Hort. cliff.* 382. *Fl. succ.* 647. *Mat. med.* 371.

Scorzonera humilis latifolia nervosa. *Bauh. pin.* 275.

Scorzonera humilis latifolia pannonica. *Clas. hist.* 2. p. 138.

Habitat in Europæ septentrionalioris pratis apricis. Z

2. SCOR-

SYNGENESIA: POLYGAMIA ÆQUALIS. 797

2. SCORZONERA caule ramoso, foliis amplexicauli-*hispanica*,
bus integris serrulatis. *Hort. cliff.* 383. *Hort. npsf.*
242. *Gmel. fib.* 2. p. 6.
Scorzonera latifolia sinuata. *Bauh. pin.* 275.
Scorzonera major hispanica. 1. *Clus. hist.* 2. p. 137.
Habitat in Hispania, Sibiria. 2
3. SCORZONERA foliis lineariformibus integris *graminifolia*
carinatis.
Scorzonera lusitanica, gramineo folio, flore pallide-lu-
teo. T. *Buxb. cent.* 2. p. 26. t. 21.
Scorzonera caule ramoso, foliis linearibus acuminatis
carinatis, calycibus acutis. *Gmel. fibir.* 2. p. 4.
Habitat in Lusitania, Sibiria.
4. SCORZONERA foliis lineariformibus integris pla-*purpurea*
nis, pedunculis cylindricis.
Scorzonera caule ramoso tereti, foliis linearibus acumi-
natis, calycibus obtusiusculis. *Gmel. fib.* 2. p. 7. t. 2.
Scorzonera angustifolia subcærulea. *Bauh. pin.* 275.
Scorzonera sylvestris, foliis angustis, flore cæruleo.
Mentz. pag. 1. f. 1.
Habitat in Marchia Brandeburgica, Sibiria.
Folia alterna, plana, remota. Pedunculi filiformes.
5. SCORZONERA foliis subulatis integris, pedunculo *angustifolia*
incrassato, caule basi villosa.
Scorzonera angustifolia prima. *Bauh. pin.* 275.
Scorzonera humilis angustifolia pannonica III. *Clus.*
hist. 2. p. 137 t. 138.
Tragopogon pinifolium hispanicum. *Barr. ic.* 496.
Habitat in Hispania. *Læsing.*
Caulis vix spithameus, basi ramosus, villosus. Folia con-
serta, linearia, subulata, longitudine fere caulis. Pe-
dunculi terminales, versus florem crassiores ipso caule.
Flos magnus. Corolla lutea, subtus purpurascens.
6. SCORZONERA foliis linearibus dentatis. *Ray. Inqdb. laciniata.*
120. *Hort. npsf.* 242. *Dalib. parif.* 242.
Scorzonera foliis laciniatis. *Tournef. inst.* 477. *Sauv.*
monsp. 692.
Tragopogon laciniatum luteum. *Bauh. pin.* 274. *Raj.*
hist. 251.
Habitat in Germania, Gallia. 3
7. SCORZONERA foliis omnibus antrorsum dentato-*tingitana*,
sinuatis amplexicaulibus.

D d d 4

Scorzo-

792 SYNGENESIA: POLYGAMIA ÆQUALIS.

Scorzonera foliis omnibus extrorsum dentatis amplexicaulibus, squamis calycinis margine membrenaceis.

Hort. ups. 242.

Sonchus floribus solitariis, foliorum laciniis extrorsum flexis. *Hort. cliff.* 385.

Sonchus tingitanus papaveris folio. *T. Raj. suppl.* 137.

Chondrilla tingitana, floribus luteis papaveris hortensis folio. *Herm. lugdb.* 657. t. 659.

Habitat in Tingide. ☉

Picroides. 8. **SCORZONERA** foliis superioribus amplexicaulibus integerrimis, inferioribus lyratis.

Crepis foliis radicalibus ovato-oblongis denticulatis, caulinis lanceolatis amplexicaulibus, pedunculis unidioris squamosis. *Roy. lugdb.* 127.

Sonchus lævis angustifolius. *Bauh. pin.* 124?

Crepis dalechampia. *Daleb. hist.* 562. *

Habitat - - - ☉

Semina incurva & crenulata sunt.

5802

PICRIS.

Echioides. 1. **PICRIS** perianthiis involucreo maximo cinctis. *Hort. ups.* 241.

Picris calycibus simplicibus perianthio maximo obvallatis. *Hort. cliff.* 387. *Roy. lugdb.* 128. *Dalib. paris.* 240.

Hieracium echioides, capitulis cardui benedicti. *Bauh. pin.* 128.

Buglossum echioides luteum hieracio cognatum. *Loeb. ic.* 577.

β. *Hieracium* capitulis cardui benedicti. *Herm. par.* 185. t. 185.

Habitat in Angliæ, Galliæ sylvis caduis, aggeribus. ☉

Hieracioides. 2. **PICRIS** perianthiis imbricatis. *Hort. cliff.* 387. *Fl. suec.* 611. *Roy. lugdb.* 128. *Dalib. paris.* 240. *Gmel. sibir.* 2. p. 39.

Cichorium pratense luteum hirsutie asperum. *Bauh. pin.* 126.

Hieracium asperum, majore flore, in agrorum limitibus. *Bauh. hist.* 2 p. 1029.

Habitat in Germaniæ, Angliæ, Belgii, Galliæ versuris agrorum. ☿

pyrenaica. 3. **PICRIS** perianthiis laxis, caule piloso, foliis dentato-sinuatis.

Hic-

SYNGENESIA: POLYGAMIA ÆQUALIS. 793

Hieracium pyrenaicum, blattariæ folio, minus pilosum.

Herm. parad. 184. t. 184.

Habitat in Pyrenæis. 2

4. *PICRIS* foliis pinnatifidis: lobis alternis rotundatis. † *asplenoides*.
 Hein in herbaria hispidosa asplenoides, succo sulphureo *Vaill. act.* 1721. p. 206.
Habitat in Ægypto.

SONCHUS.

803

1. *SONCHUS* pedunculis hispidis, floribus racemosis, *canadensis*.
 foliis lyrato-hastatis.
Chondrilla foliis pinnato hastatis denticulatis. *Gron. virg.* 89.?
Habitat in Canada. Kalm.
 Folia *S. floridani*, *dentata*, *subtus glauca*. Pedunculi *longi, terminales, adspersi pilis glutiniferis*. Racemi *breves*. Flores *magnitudine sequentis, caerulei*.
2. *SONCHUS* pedunculis calycibusque hispidis subum-
 bellatis, foliis basi sagittatis.
Sonchus pedunculis calycibusque hispidis, floribus distanti-corymbosis. *Hort. ups.* 244.
Sonchus caule erecto, foliis pinnato-hastatis apiceque hastatis, floribus congestis villosis. *Hort. cliff.* 385.
Rev. lugdb. 129. *Dalib. paris.* 244.
Sonchus asper arborescens. *Bauh. pin.* 124.
Sonchus lævior austriacus 5 *altissimus*. *Clus. hist.* 2. p. 147.
Habitat in Belgii, Angliæ, Galliæ, Hungariæ pratis paludosis. 2
Facies a S. arvensi diversissima; Structura tamen ita eadem, ac si S. palustris ex S. arvensi olim ortus fuisset.
3. *SONCHUS* pedunculis calycibusque hispidis subum-
 bellatis, foliis basi cordatis.
Sonchus pedunculis calycibusque hispidis, floribus congestis. *Fl. succ.* 642. *Dalib. paris.* 243.
Sonchus foliis lanceolato-oblongis dentatis, floribus congestis hispidis. *Hort. cliff.* 384.
Hieracium majus, folio sonchi, 1. *Hieracium sonchites*.
Bauh. pin. 126.
Hieracium majus. Fuchs. hist. 319.
Habitat in Europæ agris argillosis. 2

D d d 5

4. SON-

794 SYNGENESIA: POLYGAMIA ÆQUALIS.

- oleraceus.* 4. **SONCHUS** pedunculis tomentosis, calycibus glabris.
Fl. succ. 643. *Dalib. parif.* 243.
 Sonchus annuus ramosus diffusus, foliis laciniatis. *Fl. lapp.* 289. *Hort. cliff.* 384.
 Sonchus caule ramossissimo diffuso, foliis summis amplexicaulibus, pedunculis inæqualibus. *Roy. lugdb.* 129.
- lævis.* α. Sonchus lævis laciniatus latifolius. *Bauh. pin.* 124.
 Sonchus lævis. *Cam. epit.* 279.
 β. Sonchus lævis minor, paucioribus laciniis. *Bauh. pin.* 124.
 Sonchus lævis latifolius. *Tabern. ic.* 190.
- asper.* γ. Sonchus asper laciniatus & non laciniatus. *Bauh. pin.* 124.
 Sonchus asper laciniatus latifolius & angustifolius. *Loef. pruss.* 257. t. 77. 78.
 δ. Sonchus asper non laciniatus. *Tournef. inst.* 474.
 Sonchus aspera. *Fuchsf. hist.* 674.
Habitat in Europæ cultis. ☉
- tenuerrimus.* 5. **SONCHUS** pedunculis tomentosis, calycibus pilosis
 Sonchus lævis in plurimas & tenuissimas lacinias divisus. *Bauh. pin.* 124. *prodr.* 61.
 Chondrilla lutea. *Bauh. hist.* 2. p. 120.
Habitat Monspeli, Florentiæ. ☉
 Caulis *ramosissimus: pilis sparsis, glutiniferis.* Pedicelli albo tomento vestiti. Calyces cylindrici: pilis patentibus adspersi. Folia glabra, tenera, pinnatifida, inermia.
- alpinus.* 6. **SONCHUS** pedunculis squamosis, floribus racemosis.
Fl. succ. 644.
 Sonchus caule erecto, foliis pinnato-hastatis: apice cordato-triangulis, floribus racemosis. *Hort. cliff.* 385. *
 Sonchus lapponum altissimus, floribus cæruleis. *Fl. lapp.* 290.
 Sonchus lævis laciniatus cæruleus f. Sonchus alpinus cæruleus. *Bauh. pin.* 124.
 Sonchus cæruleus. *Cam. epit.* 284.
Habitat ad latera Alpium Lapponiæ, Helvetiæ, Austriæ. ☉
- floridanus.* 7. **SONCHUS** pedunculis subsquamosis, foliis lyrato-hastatis.
 Sonchus caule erecto, foliis pinnato-dentatis: radicalibus.

SYNGENESIA: POLYGAMIA ÆQUALIS. 795.

bus latissimis obtusis, floribus racemosis. *Roy. lugdb.* 129.

Sonchus annuus canadensis altissimus laciniatus, flore cæruleo. *Vail. alt.* 1721. p. 198.

Lactuca altissima, folio sonchi laciniato, flore parvo cæruleo. *Boerb. lugdb.* 1. p. 81.

Habitat in Virginia, Canada.

8. *SONCHUS* pedunculis squamatis, foliis lanceolatis *fibriosis*, indivisis sessilibus. *Hort. upf.* 244.

Sonchus foliis lanceolatis sessilibus plerumque denticulatis, floribus corymbosis, caulibus glabris *Gmel. fibr.* 2. p. 11. t. 3.

Lactuca salicis folio, flore cæruleo. *Amn. rurb.* 211. 212.

Habitat in Sibiria, Suecia septentrionali, Finlandia. 2

LACTUCA. 804

1. *LACTUCA* foliis pinnato-sinuatis denticulatis acutis: *quercina*, subtus lævibus, caule glabro. *St. gest.* 289. * *Fl. suec.* 646. *

Lactuca foliis quernis. *Raj. hist.* 221.

Habitat in Insula Carolina Balthici. 2

2. *LACTUCA* foliis rotundatis, caule corymboso. *fativa.*
Lactuca sativa. *Baub. pin.* 122. *Hort. cliff.* 384. *Hort.* upf. 242. *Mat. med.* 370.

β. *Lactuca capitata.* *Baub. pin.* 123. *Morif. hist.* 3. p. 57. *capitata*, f. 7. t. 2. f. 2.

γ. *Lactuca crispa.* *Baub. pin.* 123. *Dod. pempt.* 614. *crispa.*
Habitat - - - - - ☉

Cultura mangorico forte producta ex proximis.

3. *LACTUCA* foliis carina aculeatis dentatis. *viresc.*

Lactuca caule foliisque aculeatis. *Hort. cliff.* 384. * *Roy. lugdb.* 121. *Dalib. parif.* 236. *Sauv. monsp.* 295.

Lactuca sylvestris, odore viroso. *Baub. pin.* 123. *Morif. hist.* 3. p. 58. f. 7. t. 2. f. 16.

β. *Lactuca sylvestris annua, costa spinosa, folio integro, colore cæcio.* *Morif. hist.* 3. p. 58.

γ. *Lactuca sylvestris italica, costa spinosa, sanguineis maculis adspersa.* *Herm. parad.* 191. t. 194.

δ. *Lactuca sylvestris, costa spinosa.* *Baub. pin.* 123.

Lactuca sylvestris laciniata. *Morif. hist.* 3. p. 58. f. 7. t. 2. f. 17.

Habitat in Europæ australioris aggeribus, sepibus.

Folia.

796 SYNGENESIA: POLYGAMIA ÆQUALIS.

Folia prima f. radicalia in a. integra, in d. sinuata
Raj. angl. 3. p. 161.

canadensis. 4. LACTUCA foliis lanceolato-ensiformibus dentatis
inermibus.

Lactuca canadensis altissima angustifolia, flore pallido
luteo. *Tournef. inst.* 474.

Sonchus sylvestris, folio laciniato glauco; costa non spinosa. *Raj. suppl.* 137.

Habitat in Canada. *Kalm.*

Differt a sativa caule altiore; Foliis quadruplo longioribus, angustioribusque; Racemo terminali longo composito, nec corymboso.

saligna

5. LACTUCA foliis hastato-linearibus sessilibus carinatis
aculeatis *Amoen. acad.* 1. p. 150.

Lactuca foliis linearibus hastatis: inferioribus dentatis
Roy. lugd. 121.

Lactuca foliis imis pinnatifidis, cæteris ensiformibus basi
hastatis, dorso serratis. *Sauv. monsp.* 692.

Lactuca foliis radicalibus linearibus dentato pinnatis: laciniis
sursum dentatis: caulinis alato-lanceolatis, floralibus linearibus sessilibus. *Gnett. stamp.* 374. *Dalib. parif.* 236.

Lactuca sylvestris altissima, foliis inferioribus dentatis
superioribus acuminatis integris. *Raj. suppl.* 135.

Lactuca sylvestris altera, angusto saligno folio: costa
albicante. *Rupp. jen.* 167. *Hall. jen.* 207. t. 4. optim.

Lactuca sylvestris, angusto-laciniato folio. *Magn. monsp.* 298. *

Endivia minor lactucina spinosa. *Barr. ic.* 136.

Habitat in Gallia, Lipsiæ.

perennis.

6. LACTUCA foliis linearibus dentato-pinnatis: laciniis
sursum dentatis. *Hort. cliff.* 384. *Roy. lugdb.* 121.

Dalib. parif. 236.

Chondrilla cærulea altera, cichorii sylvestris folio. *Banb. sin.* 130.

Chondrilla. *Dod. pempt.* 637.

Apat. *Dalech. hist.* 561.

Habitat in Germania, Helvetia, Gallia. 2

CHONDRILLA.

juncea.

1. CHONDRILLA. † *Hort. cliff.* 383. *Roy. lugdb.* 120. *Hall. belv.* 755. *Gmel. sibir.* 2. p. 8.

Chondr.

SYNGENESIA: POLYGAMIA ÆQUALIS. 797

- Chondrilla foliis radicalibus pinnatifidis, caulinis lanceolato linearibus. *Wach. utr.* 223. *Bœhm. lips.* 89.
 Chondrilla juncea viscosa arvensis. *Bauh. pin.* 30.
 Chondrilla juncea. *Tabern. hist.* 487.
 Chondrilla viminea. *Bauh. hist.* 2. p. 1021. *
Habitat in Germania, Helvetia, Gallia *ad agrorum margines.*

PRENANTHES. 806

- . **PRENANTHES** foliis linearibus integerrimis. *tenuifolia.*
Prenanthes angustifolius, flore purpureo. *Vaill. act.* 1721. p. 193.
 Chondrilla angustissimo, longissimo integroque folio. *Juss. act.* 1709.
Habitat in Alpibus Europæ australis.
 . **PRENANTHES** foliorum ramentis cauli adnatis. *viminea.*
Prenanthes foliorum laciniis extrorsum dentatis, ramentis foliaceis cauli ramisque adnatis. *Vir. cliff.* 78. *Roy. lugdb.* 120 *Sauv. monsp.* 692.
 Chondrilla viscosa, caule foliis obducto. *Bauh. pin.* 130.
Habitat in Gallia, Lusitania.
 . **PRENANTHES** flosculis quinis, foliis lanceolatis purpureis, denticulatis. *Hort. cliff.* 383.
Lactuca montana purpureo-cærulea minor. *Bauh. pin.* 123.
Sonchus lævior pannonicus 4, purpureo flore. *Clus. hist.* 2. p. 147.
 . *Lactuca montana*, purpureo-cærulea, major. *Bauh. pin.* 123.
Sonchus montanus, purpureus tetrapetalos. *Col. ecphr.* 1. p. 245.
Habitat in Germaniæ, Helvetiæ, Italiæ nemoribus montanis.
 . **PRENANTHES** flosculis quinis, foliis lyrato-hastatis muralis. *tis.* *Hort. cliff.* 383. *Fl. suec.* 645. *Roy. lugdb.* 120. *Hall. helv.* 755. *Dalib. parif.* 235.
Sonchus lævis laciniatus muralis, parvis floribus. *Bauh. pin.* 124.
Sonchus lævis vulgator 2. *Clus. hist.* 2. p. 136.
Habitat in Europæ nemoribus umbratis.
 . **PRENANTHES** erecta; flosculis quinis, foliis trilobis. *altissima.*
Prenanthes canadensis altissima, foliis variis, flore luteolo. *Vaill. act.* 1721. p. 194.

Son-

798 SYNGENESIA: POLYGAMIA ÆQUALIS.

Sonchus elatus f. *dendroides virginianus*, ari in modum
ariculatis foliis, ramolissimus, floribus luteis parvis pen-
tapetalis. *Pluk. alm.* 355. t. 317. f. 2.
Habitat in Virginia, Canada.

6. *PRENANTHES* flosculis plurimis, floribus nutanti-
bus subumbellatis, foliis hastato-angulatis. *Hort. cliff.*
383. *Gron. virg.* 89.

Prenanthes nov' anglicanus, chenopodii foliis, radice
bulbosa, sanguineo caule, floribus racemosis candidis-
simis. *Pluk. amaltb.* 195.

β. *Prenanthes autumnalis*, flore dilute purpureo deorsum
nutante, spicatum ad caulem disposito, foliis scabris
incisilis, caule singulari. *Gron. virg.* 89.

Habitat in Carolina, Virginia, Pennsylvania.

altissima. 7. *PRENANTHES repens*, foliis trilobis. *Amoen. acad.*
2. p. 360. f. 23. *

Habitat in Sibiria.

LEONTODON.

207
Taraxacum. 1. *LEONTODON* calyce inferne reflexo. *Fl. lapp.* 280.
Fl. succ. 627. *Mat. med.* 368. *Hort. cliff.* 386. *Gron.*
virg. 176. *Roy. lugdb.* 122. *Dalib. parif.* 240. *Gmel.*
fibir. 2. p. 14.

Dens leonis. latiore folio. *Bauh. pin.* 126.

Hedynois. *Fuchs. hist.* 680.

β. Dens leonis angustiore folio. *Bauh. pin.* 126.

Habitat in Europæ pascuis. &

bulbosum. 2. *LEONTODON* foliis oblongo-ovatis subdentatis gla-
bris, calycibus lævibus, pedunculis hirtis.

Taraxacum calyce toto erecto subhirtuto, radice bulbo-
sa. *Sauv. monsp.* 297.

Chondrilla bulbosa. *Bauh. pin.* 130.

Chondrilla pulilla marina lutea bulbosa. *Lob. ic.* 230.

Habitat Monspelii, inque Italia.

Dandelion. 3. *LEONTODON* foliis ensiformibus integris, calyce
erecto simplici. *Gron. virg.* 90. *

Habitat in Virginia.

Pedunculi versus flores hispidi, & glutinosi evadunt.

Folia radicalia ensiformia, longa, glabra, integerrima.

autumnale. 4. *LEONTODON* calycibus erectis, pedunculis squa-
mosis, foliis glabris dentatis integerrimis. *Fl. succ.* 629.

Crepis foliis longis dentatis linearibus, caule declinato
nudo. *Hort. cliff.* 386.

Hiera

INGENESIA: POLYGAMIA ÆQUALIS. 799

Hieracium, chondrillæ folio glabro, radice succisa, majus & minus. *Bauh. pin.* 127.

Hieracium minus. *Luchs. hist.* 320.

Habitat in Europæ pratis, pascuis, autumno. 2

5. LEONTODON foliis dentatis hirsutis, calyce erecto *tuberosum*, acuto.

Taraxaconoides chondrillæ folio hirsuto, asphodeli radice. *Vaill. art.* 1721. p. 170.

Dens Leonis asphodeli bulbulis. *Bauh. pin.* 126.

Chondrilla altera dioscoridis montpelientium. *Lob. ic.* 232.

Habitat in Hetruriæ, Galloprovinciæ, Narbonæ pratis. 2

6. LEONTODON calyce toto erecto hispidum, foliis *hispidum*, hispidis dentatis integerrimis. *Hort. cliff.* 386. *Fl. succ.*

628. *Roy. lugdb.* 122. *Dalib. parif.* 241.

Picris calyce erecto, foliis hispidis dentatis: dentibus integerrimis. *Sauv. monsp.* 388.

Hieracium asperum, folio magno dentis leonis. *Bauh. pin.* 127.

Hieracium, caule aphylo, hirsutum. *Bauh. hist.* 2. p. 1037.

β. Hieracium, dentis leonis folio, hirsutie asperum minus laciniatum. *Bauh. pin.* 127. *prodr.* 63.

γ. Hieracium, dentis leonis folio, hirsutie asperum magis laciniatum. *Bauh. pin.* 127.

Hieracium parvum hirtum, caule aphylo, crispum ubi ficcatum. *Bauh. hist.* 1038.

Habitat in Europæ pratis.

Varietates, quamvis habitu diversas ejusdem esse speciei ex pilis apice stellatis judicari.

HIERACIUM. 808

* *Scapo nudo, unifloro.*

1. HIERACIUM foliis integris subdenticulatis lanceo-*incanum*, latis scabris, scapo unifloro.

Hieracium montanum angustifolium nonnihil incanum. *Bauh. pin.* 129.

Hieracium VI montanum. *Clus. hist.* 2. p. 141. *Bauh. hist.* 3. p. 1038.

Habitat in Alpibus Danubio imminentibus, Helvetiæ, Tauro.

Stolones nulli. Folia vix manifeste denticulata.

2. HIE-

- Hedypnoidis* 2. **HIERACIUM** foliis dentatis viscidis hirsutis subasperis, scapo nudo unifloro, calyce hirt. *Sauv. monsp.* 298. †
Hieracium montanum, dentis leonis folio, f. laciniatum, lanuginoso folio. *Bauh. pin.* 468.
Hieracium folio Hedypnoidis. *Clus. hist.* 2. p. 142.
Habitat in Lotharingia, Monspelii.
- alpinum*. 3. **HIERACIUM** foliis oblongis integris dentatis, caule subnudo unifloro, calyce piloso.
Hieracium caule unifloro, calyce villosa. *Fl. lapp.* 283.
Fl. jucc. 632.
Hieracium alpinum pumilum, folio lanuginoso. *Bauh. pin.* 129.
Hieracium villosum alpinum, flore magno singulari.
Raj. angl. 3. p. 169. t. 6. f. 2. *
Pilosella monoclonos non repens alpina minor lanuginosa, amplo flore. *Morsf. hist.* 3. p. 78. f. 7. t. 7. f. 5.
Habitat in Alpibus Lapponicis, Britannicis.
- venosum*. 4. **HIERACIUM** foliis cuneiformibus hirtis, scapo nudo crassissimo erecto. *Gron. virg.* 89.
Hieracium species 2. *Cold. noveb.* 175.
Hieracium marianum perelegans, lapathi venis sanguineis inscriptis foliis, flore parvo flavescente. *Pluk. mant.* 102.
Hieracium fruticosum latifolium, foliis punctis & venis sanguineis notatis. *Bauist. virg.*
Habitat in Virginia.
- Pilosella*. 5. **HIERACIUM** foliis integerrimis ovatis, caule repente, scapo unifloro. *Hort. cliff.* 388. *Fl. jucc.* 633.
Nat. med. 369. *Roy. lugdb.* 122. *Hall. betu.* 742.
Dalib. parif. 238.
Pilosella major repens hirsuta. *Bauh. pin.* 262.
Pilosella major. *Cam. epit.* 709. *Fuchf. hist.* 605.
Habitat in Europæ pascuis aridis. &
- * *Scapo nudo multifloro.*
- rubrum*. 6. **HIERACIUM** foliis integris ovato-oblongis, stolonibus repentibus, scapo nudo multifloro. *Fl. jucc.* 634. *Dalib. parif.* 237. *Sauv. monsp.* 14.
Pilosella major repens minus hirsuta. *Bauh. pin.* 262.
Habitat in Suecia. &
- Auricula*. 7. **HIERACIUM** foliis integerrimis lanceolatis, scapo nudo multifloro. *Fl. lapp.* 282. *Fl. jucc.* 635. *
Hiera-

SYNGENESIA: POLYGAMIA ÆQUALIS. 801

Hieracium foliis integerrimis, caule repente, scapo nudo multifloro. *Hort. cliff.* 388. *

Pilosella major erecta altera. *Bauh. pin.* 262.

Pilosella repens minor, caule pedali, polyanthos, foliis angustis oblongis. *Raj. suppl.* 147.

Habitat in Europæ pratis aridis, juxta agros. &

8. *HIERACIUM* foliis ovaus subdentatis, scapo nudo *præmorsum*, racemoso, floribus superioribus primoribus. *Fl. suec.* 636. *

Hieracium foliis ovatis tomentosis, caule nudo floribus spicatis luteolis terminato. *Hall. helv.* 744. *Gmel. fib.* 2. p. 32. t. 13. f. 2.

Hieracium pratense latifolium non sinuatum majus. *Bauh. pin.* 129.

Hieracium latifolium, præmorsa radice, caule singulari, pilosellæ major affine. *Bauh. hist.* 2. p. 1035.

Pilosella major erecta. *Bauh. pin.* 262. *Raj. hist.* 241.

Intybus aphyllocalis major. *Thal. barc.* 66.

Habitat in Helvetia, Hærcynia, Upiandia. &

9. *HIERACIUM* foliis integris, caule subnudo simpli-aurantiacum
cissimo piloso corymbifero. *Hort. cliff.* 388. *Hort. ups.* 238.

Hieracium hortense, floribus atro-purpurascens. *Bauh. pin.* 128. *prodr.* 65.

Hieracium alpinum non laciniatum, flore fusco. *Bauh. pin.* 128. *prodr.* 65.

Hieracium germanicum 1. *Col. ecphr.* 2. p. 28. t. 30.

Pilosella polyclonos repens major syriaca, flore amplo aurantiaco. *Morif. hist.* 3. p. 78. f. 7. t. 8. f. 7.

Habitat in Syria, Helvetia.

10. *HIERACIUM* foliis caulinis elongato-dentatis glaberrimis; radicalibus lanceolatis integris, caule ramoso. *det.*

Hieracium alpinum pumilum, chondrillæ folio. *Bauh. pin.* 129. *prodr.* 64. *Burf. VI.* 87.

Habitat in Sneberg Austria inferioris.

Caulis *spithameus*, *levis*. Folia radicalia lanceolata, petiolata, integra, glabra. Caulina 5 ad 6, alterna glabra, longa: dentibus longis, recurvatis, facie Leon-
rod. autumnalis. Pedunculi ex alis superioribus foliorum, altitudine caulis, foliarii, nudi, uniflori: Bractæ una aut altera setacea, apice subvillosa. Calyces erecti, imbricati foliolis linearibus. Corolla pallide flava.

E e e

11. HI-

- Gronovii.* 11. **HIERACIUM** caule paniculato subnudo, foliis radicalibus obovatis integerrimis pilosis.
Hieracium foliis radicalibus obverse ovatis pubescentibus: caulinis ovatis amplexicaulibus, floribus paniculatis, caule erecto. *Gron. virg.* 90.
Habitat in Virginia, Pennsylvania. 2
Radix præmorsa. Folia radicalia obovata, obtusa, integerrima, tenuitate membranacea, supra pilis raris adspersa, subtus subviolacea. Caulis pedalis, filiformis, uno alterove folio lanceolato, paniculatus, panicula inæquali.
- Gmelini.* 12. **HIERACIUM** caule paniculato nudo, foliis radicalibus ovatis serratis glabris.
Hieracium foliis ex sinuato-dentatis, caulibus supra ramosis subnudis, pedunculis hirtutis unifloris. *Gmel. fib.* 2. p. 23. t. 8. f. 2.
Habitat in Sibiria. 2
 * *Caule folioso.*
- paniculatum* 13. **HIERACIUM** caule erecto, foliis alternis lanceolatis nudis dentatis, panicula capillari.
Habitat in Canada. *Kalm.*
Caulis pedalis, erectus, teres, inferne albo-lanatus, superne glaber. Folia lanceolata, latiuscula, basi attenuata, acuminata, remote dentata, tenera, glabra: inferiora subtus, præsertim ad carinam pilosa. Panicula terminalis varie ramosa: Pedunculis tenuissimis, fere capillaribus, divaricatis. Bractæ ad ramificationes brevissimæ, setaceæ. Flores parvi.
- porrifolium.* 14. **HIERACIUM** caule composito ramosissimo, foliis lanceolato-linearibus integerrimis. *Roy. lugdb.* 125. *
Hieracium phalangii foliis. *Vaill. act.* 1721. p. 184
Hieracium montanum, asphodeli foliis acuminatis. *Bocc. mus.* 147. t. 106. *Raj. suppl.* 142.
Habitat in Alpibus Austriæ, italiæ.
Folia radicalia subulata, fere Statices. Caulis pedalis, Ramis longis, raris, parum subdivisis. Folia in caule minora & tantum ad exortum ramorum.
- maritimum.* 15. **HIERACIUM** caule ramoso, foliis radicalibus ovatis dentatis; caulino minori. *Hort. cliff.* 388. *Fl. succ.* 637. *Roy. lugdb.* 123. *Dalib. parisi.* 237. *Gmel. fibir.* 2. p. 125.
Hieracium caule ramoso, foliis ovatis dentatis. *Fl. lapp.* 284.

Hiera-

SYNGENESIA: POLYGAMIA ÆQUALIS. 803

Hieracium murorum, folio pilosissimo. *Bauh. pin.* 129. pilosissimum.
Corchorus. Dalech. hist. 565.

β. *Hieracium murorum laciniatum minus pilosum. Bauh. sylvaticum*
pin. 129. *Is. wgoth.* 257.

Pulmonaria gallica femina. Tabern. hist. 504.

γ. *Hieracium macrocaulon hirsutum, folio longiore. Raj.*
angl. 3. p. 169.

Habitat in Europæ apricis datis.

16. **HIERACIUM** caule paniculato, foliis amplexicaulibus dentatis glabris, calycibus hispidis. *Fl. lapp.* 285. *paludosum.*
Fl. succ. 638.

Hieracium cichorei folio. Is. Wgoth. 32.

Hieracium montanum latifolium glabrum minus. Bauh.
pin. 129. *Morif. hist.* 3. p. 69. f. 7. t. 5. f. 47.

Habitat in Europæ borealis nemoribus paludosis.

Calyces pilis nigris exstantibus hirsuti sunt. Varietas forte sola antecedentis, quamvis ex loco evasit facie diversissima.

17. **HIERACIUM** caule multifloro, foliis lyratis glabris, calycibus pedunculisque hispidis. *lyratum.*

Hieracium caule ramoso, foliis semiamplexicaulibus teneris oblongis plerumque petiolatis: petiolis infimorum dentatis. Gmel. sib. 2. p. 24. t. 9.

Habitat in Sibiria. D. Gmelin.

Affinis maxime H. paludoso. Caulis levis. Calyces pubescentes, & quod in precedenti non obtinet, Pedunculi etiam pubescentes. Folia magis lyrata, petiolata, angulis obtusioribus.

18. **HIERACIUM** foliis radicalibus obovatis denticulatis, caulibus oblongis semiamplexicaulibus. *cerinthoides.*
124. Hort. nup. 238.

Hieracium pyrenaicum, foliis cerinthes, latifolium. Tournef. inst. 472.

Habitat in Pyrenæis. 2

19. **HIERACIUM** foliis amplexicaulibus cordatis subdentatis, pedunculis unifloris hirsutis, caule ramoso. *amplexicaule.*
Hort. cliff. 387. * *Roy. lugdb.* 123.

Hieracium pyrenaicum longifolium amplexicaule. Tournef. inst. 472.

β. *Hieracium pyrenaicum rotundifolium amplexicaule. Tournef. inst.* 472.

Habitat in Pyrenæis.

804 SYNGENESIA: POLYGAMIA ÆQUALIS.

- pyrenaicum*. 20. **HIERACIUM** foliis obovato-lanceolatis retrorsum dentatis, caule simplici, floribus terminalibus confertis. *Amæn. acad.* 1. p. 151. *
Habitat in Pyrenæis. Burf. VI. 74.
- Blattarioides* 21. **HIERACIUM** foliis lanceolatis amplexicaulibus dentatis, floribus solitariis, calycibus laxis. *Hort. cliff.* 387. * *Roy. lugdb.* 123. *Amæn. acad.* 1. p. 151.
Hieracium pyrenaicum, blattariæ folio, minus hirsutum. Tournef. inst. 472.
Habitat in Pyrenæis.
- glutinosum*. 22. **HIERACIUM** foliis lanceolatis lyrato-dentatis scabriusculis, floribus umbellatis.
Hieracium dentis leonis folio, floribus parvis. Bauh. pin. 127. *prodr.* 63.
Habitat in G. Narbonensi. ☉
Habitus Crepis testorum. Folia summa lanceolata, integerrima. Caulis striatus. Folia mollia, glutinosa.
- halmii*. 23. **HIERACIUM** caule erecto multifloro, foliis lanceolatis dentatis, pedunculis tomentosis.
Habitat in Pennsylvania. Kalm.
Caulis erectus, H. sabauda angustior, lavis. Folia lanceolata, alterna, subsessilia, acuminata, parva, nulla, argutius extrorsum dentata, quam ulla alia hujus generis. Pedunculi ad apicem caulis alterni, saepius simplices & uniflori, tomentoso-albidi. Bracteis varioribus, linearibus. Flores parvi, terminales, erecti.
- sprengeria-num*. 24. **HIERACIUM** caule ramoso folioso, foliis semiamplexicaulibus oblongis repandis hispidis. *Roy. lugdb.* 123.
Hieracium pulchrum ramosum sprengertianum. Bauh. hist. 2. p. 1026. *Morif. hist.* 3. p. 65. f. 7. t. 5. f. 17.
Habitat in Lusitania.
- sabaudum*. 25. **HIERACIUM** caule erecto multifloro, foliis ovato-lanceolatis dentatis semiamplexicaulibus. *Roy. lugdb.* 124. *Hort. upf.* 238.
Hieracium fruticosum latifolium hirsutum. Bauh. pin. 129.
Hieracii sabaudi varietas. 1. Bauh. hist. 2. p. 1030.
Habitat in Germania. 2.
- umbellatum*. 26. **HIERACIUM** foliis linearibus subdentatis sparsis, floribus subumbellatis. *Fl. lapp.* 287. *Fl. suec.* 639. *Hort.*

SYNGENESIA: POLYGAMIA ÆQUALIS. 805

Hort. cliff. 387. *Roy. lugdb.* 123. *Dalib. parif.* 237.

Gmel. fib. 2. p. 25.

Hieracium fruticosum angustifolium majus. Baub. pin.
129.

Hieracium sabaudum. Dalech. hist. 570.

Habitat in Europæ pascuis siccis. &

CREPIS. 809

1. CREPIS foliis ovatis integris villosis petiolatis, caule pygmao procumbente. †

Hieracium alpinum incanum saxatile, brunellæ foliis integris. Bocc. mus. 2. p. 33. t. 24. *Raj. suppl.* 142.

Habitat in Alpibus Italiæ ad Nursiam prope lacum pilati.

2. CREPIS foliis lanceolatis dentato-sinuatis hispida: seminibus nudiculis, scapo unifloro.

Hieracium, dentis leonis folio, hirsutie asperum magis & minus laciniatum. Baub. pin. 127. *prodr.* 63. t. 633. n. 1.

Habitat in G. Narbonensi, Hispania.

3. CREPIS foliis pinnatifidis crenatis, scapo nudo paucifloro.

Hieracium ficulum, bursæ pastoris folio. Bocc. mus. 2. p. 147. t. 106. & 112. *Raj. suppl.* 142.

Hieracium minus panormitanum, cichorei folio. Cyp. catb. 95. *suppl.* 3.

Habitat in Sicilia. &

4. CREPIS involucris calyce longioribus: squamis seminibus sparsis. *Hort. ups.* 239.

Hieracium calycibus fetis longissimis distantibus patentibus obvallatis. Hort. cliff. 387. *Sauv. monsp.* 83.

Hieracium proliferum falcatum. Baub. pin. 128.

Hieracium calyce barbato. Col. ecphr. 2. p. 28. t. 27. f. 1.

Habitat in Monspelii, Vesuvii, Siciliæ, Messanæ arenosis maritimis. ☉

5. CREPIS involucris ovatis concavis obtusis patentibus.

Hieracium montanum rapifolium. Baub. pin. 129. *prodr.* 749. *Raj. hist.* 237. *Morif. hist.* 3. p. 65. *secundum.* *Burf.* VI: 70.

- β. Cichorium pratense hirsutum vesicarium. *Baub. pin.* 126.

E e e 3

Cicho-

806 SYNGENESIA: POLYGAMIA ÆQUALIS.

Cichorium sylvestre vesicarium pratense. Col. ecphr. 1. p. 238. t. 237.

Habitat a in Helvetia, B in Apulia.

Plantæ a Radix crassa. Folia lyrata, angulata, dentata, scabra. Caulis brevis. Rami longi, nudi, bifidi s. trifidi, uniflori, pedunculorum Bractæe ovate, solitariae, concavae, integrae, pallidae. Flores lutei. Calyces singuli cincti Involucro longitudine calycis, ex foliolis 5. s. 6, patentibus, ovatis, obtusis, concavis, membranaceis, pallidis. Planta B foliis a priori manifeste differt, flores vero conjungunt utramque.

rubra,

6 CREPIS foliis amplexicaulibus lanceolatis, omnibus dentatis; radicalibus sinuatis. *Vir. cliff. 79. Hort. mps. 239. Sawv. monsp. 295.*

Hieracium foliis caulinis sinuatis, calycibus ante florescentiam nutantibus. Hort. cliff. 385.

Hieracium apulum, flore suaverubente. Col. ecphr. 1. p. 244.

Hieracium dentis leonis folio, flore suaverubente. Bauh. pin. 127.

Chondrilla purpurascens foetida. Bauh. pin. 130. prodr. 68. t. 68.

Habitat in Apulia. ☉

Pappus stipitatus ut in C. alpina.

alpina,

7. CREPIS foliis amplexicaulibus oblongis acuminatis: inferioribus superne, summis inferne denticulatis. *Hort. mps. 238.*

Hieracioides annua, endiviæ folio, capite magno. Vaill. act. 1721. p. 246.

Hieracium alpinum, scorzonerae folio. Tournef. inst. 472.

Leontodon calyce toto erecto: inferiore squamis siccis, foliis amplexicaulibus. Gmel. sibir. 2. p. 16. t. 5.

Habitat in Alpibus Italiae. ☉

pudica,

8. CREPIS foliis sagittatis dentatis, pedunculis superne turbinato-inflatis. *Guett. stamp. 378. Dalib. parif. 238. n. 1.*

Chondrilla, hieracii folio, annua Tournef. inst. 238.

Hieracium pulchrum. Bauh. hist. 2. p. 1025.

Hieracium montanum hirsutum minus. C. B. Raj. hist. 234.

Hieracioides annua glutinosa, floribus parvis. Vall. act. 1721.

Habitat in Gallia. ☉

9. CRE-

SYNGENESIA: POLYGAMIA ÆQUALIS. 807

9. CREPIS foliis ovatis denticulatis, petiolis alatis am *sibirica.*
plexicaulibus dentatis. *Hort. upf.* 239. *
Hieracium caule ramoso, foliis infirmis: infirmis petio-
latis, reliquis ex ovato lanceolatis: omnibus sinuosis, pe-
tiolorum instar dentatis. *Gmel. fib.* 2. p. 26. t. 24.
Habitat in Sibiria. &
Pappus *sessilis ut in C. hiemi & C. tectorum.*
10. CREPIS foliis lanceolatis sagittatis sessilibus lævibus: *tectorum*
inferioribus dentatis. *Fl. suec.* 640. *Dalib. parif.* 238.
Gmel. fib. 2. p. 29.
Hieracium foliis radicalibus pinnato-dentatis, caulinis
lanceolatis subdenticulatis. *Fl. lapp.* 286. *Hort. cliff.*
387
Hieracium majus erectum angustifolium, caule lævi.
Bauh. pin. 127.
Hieracium majus dioscoridis. *Tavern. ic.* 180.
Habitat in Europæ aridis, tectis. ☉
Calyx *calyculatus, sulcatus, adpersus longitudinaliter*
pilis glutiniferis.
11. CREPIS foliis pinnato-hastatis scabris: basi superne *biennis.*
dentatis, calycibus muricatis.
Crepis scanensis. *It. scan.* 171. *
Hieracium, chondrillæ folio, glabrum. *Bauh. pin.* 127.
Hieracium foliis & facie chondrillæ. *Lob. ic.* 239.
Hieracium, chondrillæ folio, hirsutum. *Bauh. pin.* 127.
Hieracium V. *Tubern. hist.* 492.
Hieracioides foliis variis glabris, caule ramosissimo. *Hall.*
helv. 751.
Cichorium pratense luteum lævius. *Bauh. pin.* 126.
Habitat in Scanæ & Europæ australioris apricis. &
Calyx *calyculatus, sulcatus, muricatus longitudinaliter*
aculeis mollibus.
12. CREPIS foliis hastato-lyratis, caule piloso, *hirtum.*
Hieracioides foliis variis asperis, caule altissimo folioso
unifloro. *Hall. helv.* 751.
Hieracium, chondrillæ folio, hirsutum. *Bauh. pin.* 127.
Hieracium erucærolium hirsutum. *Bauh. hist.* 2. p. 1025.
Morif. hist. 3. p. 65. f. 7. t. 5. f. 14.
Habitat in Helvetia, Gallia. ☉
- 13 CREPIS foliis pinnatis angulatis, petiolis dentatis. *fatida.*
Roy. lugdb. 126. *Dalib. parif.* 239.
Hieracium luteum, cichorei sylvestris folio, amygdalas
amarus oleus. *Morif. hist.* 3. p. 63. f. 7. t. 4. f. 4.
E c e 4 Sene

808 SYNGENESIA: POLYGAMIA ÆQUALIS.

Senecio hirsutus. Baub. pin. 131.
Erigeron tertium. Dod. pempt. 641.
Habitat in Gallia.

810

ANDRYALA.

integrifolia. 1. **ANDRYALA** foliis integris. *Gaett. stamp. 2. p. 384.*
Hort. upf. 240. Sauv. monsp. 83.
Sonchus villosus luteus major. Baub. pin. 124.
Sonchus lanatus. Dalech. hist. 1116.
Hieracium villosum. Rai. hist. 231.
Habitat in Gallia, Sicilia, &

finata. 2. **ANDRYALA** foliis dentato-hastatis.
Andryala foliis inferioribus dentatis, summis integris.
Hort. cliff. 387. Sauv. monsp. 692.
Eriophorus foliis inferioribus ad costam usque laciniatis.
Vaill. act. 1721. p. 212.
Sonchus villosus luteus minor. Baub. pin. 124. prodr. 61.
Habitat Monspelii, inque Sicilia.

HYOSERIS.

* *Caule nudo.*

foetida. 1. **HYOSERIS** scapis simplicissimis unifloris, foliis pinnatifidis, seminibus nudis.
Taraxacnastrum, dentis leonis folio, radice foetida.
Vaill. act. 1721. p. 180.
Leontodontoides alpinus glaber, erysimi folio, radice crassa foetida. Mich. gen. 31. t. 28.
Dens Leonis tenuissimo folio. Baub. pin. 126. prodr. 62.
Hieracium foetidum. 111. Col. ecphr. 2. p. 29. t. 31.
Habitat in Alpibus Italiae, Austriae superioris.
Folia pinnatifida, glabra lobis latere postico subdentatis. Scapi simplices, nudi. Calyx simplex, basi auctus. Semina nuda.

radiata. 2. **HYOSERIS** scapis unifloris nudis, foliis glabris lyrate-hastatis angulatis.
Hyoseris foliis glabris hastato-pinnatis. Roy. lugdb. 127.
Sauv. monsp. 300.
Taraxacnastrum dentis leonis folio ad summitatem radiato. Vaill. paris. 121. p. 180.
Dens leonis minor, foliis radiatis. Baub. pin. 129. Pluk alm. 130. t. 37. f. 2.
Habitat in Hispania, G. Narbonensi. &

Pr-

SYNGENESIA: POLYGAMIA ÆQUALIS. 809

Pedunculus & rachis foliorum farinacea. Lobi foliorum imbricatim reflexi utrinque dentati, laeves. Semina membranacea: Pappo simplici, sessili.

3. HYOSERIS scapo simplicissimo unifloro: foliis scabris, seminibus papposis. *Hort. cliff.* 386 *Roy. lugdb.* 127.

Taraxacnastrum, erysimi folio, flore minimo. *Vaill. act.* 1721. p. 235.

Dens leonis minimus. foliis hirsutis, calycis segmentis a flore delapso erectis & semina complexis. *Raj. suppl.* 147.

Habitat in Sicilia.

4. HYOSERIS scapis unifloris, foliis lanceolatis lyratis virginica, glabris. *Gron. virg.* 60. *

Habitat in Virginia.

Folia radicalia prima ovata, tum lanceolata, demum lyrata, acuta, glabra. Scapi nudi, uniflori, foliis triplo longiores. Calyces simplicissimi, nec imbricati, ex decem foliolis lanceolatis. Corolla saturate lutea. Semina tetragona, coronata margine membranaceo integro & insuper setis 3. f. 4. longis.

5. HYOSERIS caule diviso nudo. *Hort. cliff.* 386. *Fl. minima.* *succ.* 630. *Roy. lugdb.* 127. *Sauv. monsp.* 14.

Hyoseris angustifolia. *Tabern. ic.* 180.

Hieracium minimum. *Cluj. hist.* 2. p. 142.

Habitat in Europæ arvis apricis.

* *Eque folioso.*

6. HYOSERIS fructibus subglobosis glabris, caule ramoso. *Hedypnois*.

Lapsana fructibus subglobosis glabris, seminibus disci calyculato-papposis. *Hort. upf.* 246. * *Sauv. monsp.* 83.

Lapsana calycibus fructus capitatis dehiscentibus, pedunculis incrassatis, floribus nutantibus. *Hort. cliff.* 387. *Roy. lugdb.* 131.

Rhagadioloides calthæ folio, calyce glabro. *Vaill. act.* 1721. p. 202.

Hedypnois annua. *Tournef. inst.* 478.

Habitat in Europa australi. ☉

7. HYOSERIS fructibus globosis pilosis, caule ramoso. *Rhagadioloides*.

Lapsana fructibus globosis pilosis, seminibus disci calyculato-papposis. *Hort. upf.* 246. *

Habitat in Europa australi. ☉

E e e 5

8. HYO-

810 SYNGENESIA: POLYGAMIA ÆQUALIS.

- erecta.* 8. **HYOSERIS** fructibus ovaris scabris, caule ramoso.
Lappana fructibus ovaris scabris, seminibus disci calycu-
lato-papposis. *Hort. nps.* 246. *
Habitat in Creta.

812

HYPOCHÆRIS.

- erecta.* 1. **HYPOCHÆRIS** foliis dentatis, caule ramoso folio-
so, calycibus hispidis. †
Hypochæris hieracii barbati folio, calyce hispido. *Vaill.*
act. 1721. p. 215.
Hieracium alpinum, dentis leonis folio acuto piloso pro-
scisso, capitulis pungenti lanugine oblitis *Cap. cath.*
95. *Raj. suppl.* 146.
Habitat in Sicilia.
Calycis squamæ asperæ spinulis pungentibus.
- Achyrophor.* 2. **HYPOCHÆRIS** calycibus æqualibus hispidis. *Hort.*
cliff. 385. * *Hort. nps.* 240. *
Achyrophorus fere glaber, beilidis glauco dentatoque
folio. *Vaill. act.* 1721. p. 278.
Hieracium ramosum, floribus amplis, calycibus valde
hirsutis, foliis oblongis obtusis: dentibus majoribus
inæqualibus incis. *Raj. suppl.* 144.
Habitat in Creta. ☉
- pontana.* 3. **HYPOCHÆRIS** foliis lanceolatis dentatis, caule
simplici folioso unifloro.
Hypochæris non ramosa, endiviæ folio prælongo, mo-
nanthos. *Vaill. act.* 1721. p. 215.
Hieracium latifolium pontanum præaltum glabrum, en-
diviæ folio. *Bucc. mus.* 2. p. 148. t. 113. *Raj. suppl.*
142.
Habitat in Sabaudia montibus.
- maculata.* 4. **HYPOCHÆRIS** caule subnudo: ramo solitario, fo-
liis ovato-oblongis integris dentatis. *Hort. cliff.* 385.
Fl. suec. 631. *Roy. lugdb.* 125. *Dalib. paris.* 239.
Hypochæris hirsuta, endiviæ folio, magno flore. *Vaill.*
act. 1721. p. 385. *Fl. lapp.* 288.
Hieracium alpinum latifolium hirsutie incanum, flore
magno. *Bauh. pin.* 128.
Hieracium latifolium 1. *Clus. bist.* 2. p. 139.
Hieracium alpinum, foliis dentatis, flore magno. *Bauh.*
pin. 128. *prodr.* 65. *Hall. helv.* 760. t. 24.
Habitat in Europæ frigidioris pratis asperis. 2

5. HY-

SYNGENESIA: POLYGAMIA ÆQUALIS. 811

5. **HYPOCHÆRIS** glabra, calycibus oblongis imbricatis, caulibus nudis, foliis dentato sinuatis. *Hort. cliff.* 386. n. 4. *Hort. nps.* 240. * *Roy. lugdb.* 126. *Dalib. parif.* 239. *Sauv. monsp.* 300.

Hypochæris chondrillæ folio, parvo flore. Vaill. a3.
1721. p. 214.

Hieracium minus, dentis leonis folio oblongo glabro.
Bauh pin. 127.

Hyoferis altera. Tabern. hist. 490.

Habitat in Dania, Germania, Belgio. ☉

6. **HYPOCHÆRIS** foliis dentato-sinuatis scabris, caule radicato, ramoso nudo, pedunculis squamosis. *Hort. cliff.* 386. *Roy. lugdb.* 125. *Dalib. parif.* 239. *It. scan.* 225. * *Sauv. monsp.* 299.

Hieracium, dentis leonis folio obtuso, majus. Bauh.
pin. 127. *Morif. bist.* 3. p. 66. f. 7. t. 4. f. 27.

Hieracium longius radiculatum Lob. hist. 120. *Raj. bist.*
230.

Hieracium tertium. Dod. pempt. 639.

Habitat in Europæ cultioris pascuis.

LAPSANA. 813

1. **LAPSANA** calycibus fructus angulatis, pedunculis tenuibus ramotissimis. *Hort. cliff.* 384. *Fl. fusc.* 649. *Roy. lugdb.* 130. *Hort. nps.* 246. *Dalib. parif.* 244. *Gmel. fib.* 2. p. 40.

Lamplana. Dod. pempt. 675.

Soncho affinis Lapsana domestica. Bauh. pin. 124.

Habitat in Europæ cultis.

2. **LAPSANA** calycibus fructus torulosis depressis ob- *Zacinthe,*
tusis sessilibus.

Lapsana calycibus fructus subrotundis angulatis, pedunculis incrassatis, floribus erectis. Roy. lugdb. 131
Hort. cliff. 390.

Chondrilla verrucaria, foliis cichorei viridibus. Bauh.
pin. 130.

Cichorium verrucatum, zacintha. Clus. bist. 2. p. 144.

Habitat in Italia. ☉

*Toruli calycis fructiferi octo, gibbi, profunde distincti
nodis rufescentibus, exteriora vero calycis foliola se-
mera & mollia persistant.*

3. **LAPSANA** calycibus fructus undique patentibus radiatis subulatis, foliis lanceolatis indivisis. *Hort. nps.*
245. * *Sauv. monsp.* 8. *Hie-*

812 SYNGENESIA. POLYGAMIA ÆQUALIS.

Hieracium filiqua falcata. Bauh. pin. 128.
Hieracium stellatum. Bauh. hist. 2. p. 1014.
Habitat Mouspelii, Bononiæ. ☉

- Rhagadiolus* 4. **LAPSANA** calycibus fructus undique patentibus: radiois subulatis, foliis lyratis. *Hort. upf. 245. **
Hieracium falcatum alterum. Raj. hist. 256.
Rhagadiolus lapsanæ foliis. Tournesf. cor. 36.
Rhagadiolus edulis hieraciis affinis. Bauh. hist. 2. p. 1014.
Habitat in Oriente.

- Chondrilloides.* 5. **LAPSANA** calycibus pyramidatis, seminibus papposis, foliis oblongis subdentatis scabris. *Hort. upf. 245. * Sauv. monsp. 83.*
Hieracium montanum alterum leptomacrocaulon. Col. cephr. 1. p. 248. t. 249.
Hieracium annuum montanum fruticosius, caule canaliculato. Morisf. hist. 3. p. 68. f. 7. t. 5. f. 37.
Habitat in Gallia, Italia. ☉

- capillaris.* 6. **LAPSANA** seminibus papposis, foliis radicalibus lanceolatis obtusis subdentatis; caulinis hastatis, caule filiformi.
Hieracium minus glabrum, foliis eleganter virentibus. Bauh. pin. 127. prodr. 63.
Habitat in Helvetiæ, Italiæ agris.
Folia radicalia lanceolata, obtusa, glabra, subdentata; caulina lanceolata, hastata, subamplexicaulia, integerrima, acuta, parva. Caules erecti, filiformes, subdichotomi, spithamei. Flores parvi, lutei. Calyces admodum parce imbricati.

CATANANCHE.

- 814
cærulea. 1. **CATANANCHE** squamis calycinis inferioribus ovatis. *Hort. cliff. 390. Roy. lugdb. 122. Sauv. monsp. 308.*
Catananche cærulea, semistofculorum ordine simplici. Vaill. act. 1721. p. 215.
Chondrilla cærulea, cyani capitulo. Bauh. pin. 130.
Chondrillæ species tertia. Dod. pempt. 638.
Habitat in G. Narbonensi. ☉
- lutea.* 2. **CATANANCHE** squamis calycinis inferioribus lanceolatis. *Hort. cliff. 390. Hort. upf. 247. Roy. lugdb. 122. Sauv. monsp. 308.*
Catananche lutea, longo nervoso dentatoque folio. Vaill. act. 1721. p. 216.
Chon-

SYNGENESIA: POLYGAMIA ÆQUALIS. 813

Chondrilla cyanoides lutea, coronopi folio non diviso.

Bocc. musf. 2. p. 21. t. 7. *Barr. sc.* 1135.

Stoebe plantaginifolia. *Alp. exot.* 287. t. 286.

Habitat in Creta. ☉

3. CATANANCHE foliis incisif.

græca.

Catananche lutea, foliis varie incisif. *Vaill. act.* 1721.

p. 216.

Scorzonera græca saxatilis & maritima, foliis varie laciniatis. *Tournef. cor.* 36. *itin.* 1. p. 223. t. 223.

Habitat in Græciæ maritimis

CICHORIUM. 815

1. CICHORIUM caule simplici, foliis dentato-sinuatis. *Intybus.*

Fl. suec. 650. *Dalib. parif.* 244. *Mat. med.* 373.

Cichorium caule simplici. *Hort. cliff.* 389.

Cichorium sylvestre f. *officinarum.* *Bauh. pin.* 126.

Intybus sylvestris. *Cam. epit.* 285. *Fuchsf. hist.* 979.

Habitat in Europa ad margines agrorum viarumque. 2

2. CICHORIUM caule simplici, foliis integris c. sinatis. *Endivia.*

Hort. cliff. 389. *Hort. upf.* 247. *Mat. med.* 374.

Cichorium latifolium f. *Endivia vulgaris.* *Bauh. pin.*

125.

Intybum sativum. *Dod. pempt.* 634.

β. Endivia crispa. *Bauh. pin.* 125.

Habitat - - - ☉

3. CICHORIUM caule dichotomo spinoso. *Hort. cliff. spinosum.*

388.

Cichorium spinosum. *Bauh. pin.* 126. *prodr.* 62. t. 62.

Chondrillæ genus elegans, cæruleo flore. *Clusf. hist.* 2.

p. 145.

Habitat in Cretæ, Siciliæ collibus arenosis maritimis. 3

SCOLYMUS. 816

1. SCOLYMUS foliis margine attenuatis.

maculatus.

Scolymus annuus. *Hort. upf.* 244.

Scolymus. *Hort. cliff.* 388.

Scolymus Chrysanthemus annuus. *Dodart. mem.* 4. p.

303.

Scolymus theophrasti narbonensis. *Clusf. hist.* 2. p. 153.

Habitat in G. Narbonensi, Italia. ☉

2. SCOLYMUS foliis margine incrassatis.

hispanicus.

Scolymus vivax. *Sauv. monsp.* 294.

Scolymus Chrysanthemos. *Bauh. pin.* 384.

Sco-

814 SYNGENESIA: POLYGAMIA ÆQUALIS.

Scolymus theophrasti hispanicus. *Clus. hist.* 2. p. 153.
Carduus Chrysanthemos. *Dod. pempt.* 725.
Habitat in Italia, Sicilia, G. Narbonensi. 2

817
scabror.

ELEPHANTOPUS.

1. **ELEPHANTOPUS** foliis oblongis scabris. *Hort. cliff.* 390. *Hort. upf.* 147. *Gron. virg.* 176. *Roy. lugdb.* 131.
Elephantopus conyzæ folio. *Dill. elth.* 126. t. 106. f. 126.

Bidens frutescens, foliis oblongis utrinque acuminatis venosis & lanuginosis. *Brevn. ic.* 32. t. 32.

Echinophoræ indicæ affinis, semine & floribus in capitulis lævibus in caulium cymis. *Pluk. alm.* 132. t. 388. f. 6.

Anaschovadi. *Reed. mal.* 10. p. 13. t. 7.

Habitat in Indiis. 2

- tomentosus.* 2. **ELEPHANTOPUS** foliis ovatis tomentosis. *Gron. virg.* 90. *

Habitat in Virginia.

Affinis forte nimis præcedenti.

818

GUNDELIA.

- Tournefortii* 1. **GUNDELIA.**

Gundelia orientalis, acanthiaculeati folio, capite glabro. *Tournef. cor.* 51. *Itin.*

Eryngium syriacum, foliis chamæleontis longis & spinosis. *Morif. hist.* 3. p. 167.

Silybum diofcoridis f. *Hacub alcardeg serapionis*. *Rauv. itin.* 74. t. 74.

Hacub f. *Silybum* quibusdam. *Bauh. hist.* 3. p. 84.

- β. *Gundelia orientalis*, acanthi aculeati folio, floribus intense purpureis, capite araneosa lanugine oblitto. *Tournef. cor.* 51.

Habitat in Armenia, Syria. 2

819
sphærocephalus.

ECHINOPS.

1. **ECHINOPS** calyculis unifloris, caule multifloro, foliis spinosis supra nudis.

Echinops floribus capitatis, calycibus unifloris. *Hort. cliff.* 390. *Hort. upf.* 248. *Roy. lugdb.* 144. *Dalib. parif.* 245.

Echi-

SYNGENESIA: POLYGAMIA ÆQUALIS. 815

Echinopus major. Bauh. hist. 3. p. 69.

Carduus sphærocephalus latifolius vulgaris. Bauh. pin. 381.

Chalcejos. Dalech. hist. 1480.

Habitat in Italia. 2

2. ECHINOPS calyculis unifloris, caule unicapitato. *Rure.*

*Echinops caule subunifloro. Hort. ups. 248. **

Echinops caule subunifloro, foliis duplicato-pinnatifidis: foliolis latiusculis vicinis. Gmel. fib. 2. p. 100.

Carduus sphærocephalus cæruleus minor. Bauh. pin. 381.

Ritro floribus cæruleis. Lob. ic. 2. p. 8.

Crocodylium monspeliensium. Dalech. hist. 1476.

β. *Echinops caule subunifloro, foliis duplicato-pinnatifidis: foliolis linearibus remotis. Gmel. fib. 2. p. 102. t. 46.*

Habitat in Gallia, Sibiria.

3. ECHINOPS calycibus fasciculatis unifloris: lateralibus *strigosus.*

sterilibus, foliis pinnatifidis supra strigosis.

Carduus tomentosus, capitulo majore. Bauh. pin. 328.

Scabiosa carduifolia annua. Herm. parad. 224. t. 224.

Spina alba. Lob. ic. 2. p. 9.

Habitat in Hispania. Læff. ☉

*Folia pinnatifida: lanceola laciniis lanceolatis, spinamini-
mima terminatis, lateralibus inermibus, supra pilis ri-
gidis strigosa, subtus tomentosa. Calyce non in capitulum
globosum aucti, sed fasciculati, lateralibus. f.
inferioribus minoribus sterilibus.*

4. ECHINOPS floribus fasciculatis, calycibus multiflo- *corymbosus.*

ris. Hort. cliff. 391. Roy. Lugdb. 144.

*Carduus Chamæleon dictus, capitulis pluribus minori-
bus cæruleis corymbatim dispositis. Moris. hist. 3.*

p. 159. f. 7. t. 33. f. 17.

Chamæleon niger umbellatus, flore cæruleo-hyacinthino.

Bauh. pin. 380.

Chamæleon niger. Dalech. hist. 1454. Dod. pempt. 727.

*Habitat in Apulia, Hellesponto, Lemno, Tracia, in
campis. 2*

5. ECHINOPS capitulis lateralibus sessilibus.

fruticosus.

*Echinops frutescens, ad nodos floridus, foliis linearibus subtus
argenteo. Plum. spec. 10.*

Habitat in America meridionali. 3

ARCTI-

816 SYNGENESIA: POLYGAMIA ÆQUALIS.

820

ARCTIUM.

Lappa.

1. ARCTIUM foliis cordatis inermibus petiolatis.
Arctium. Hort. cliff. 391. Fl. succ. 651. Mat. med. 375.
Roy. lugdb. 144. Dalib. parif. 245.
Lappa major f. Arctium diofcoridis. Bauh. pin. 198.
Bardana f. Lappa major. Dod. pempt. 58.
Personata. Cam. epis. 887.
- β. Lappa major montana, capitulis tomentosis. *B. pin. 198.*
Habitat in Europæ cultis ruderatis. δ

Personata.

2. ARCTIUM foliis decurrentibus ciliato-spinosis: radicalibus pinnatis: caulinis oblongo-ovatis.
Carduus mollis, foliis imis laciniatis, superioribus integris, squamis calycinis reflexis. Hall. belu. 678. t. 19.
Carduus mollis latifolius, lappæ capitulis. Bauh. pin. 377. prodr. 155. Bursf. XXI. 13.
Carduus capitulis in vertice congestis lanigeris: squamis reflexis. Gmel. fib. 2. p. 62. t. 24.
Habitat in Alpibus Helvetiæ, Genève, Taurero Austriæ.
Folia caulina ovato-oblonga, subtus tomentosa, serrata serraturis spinulosis, nec non spinulis minimis interjectis. Folia hæc, basi quamvis attenuata, decurrunt per caulem tenuissimis marginibus spinosis. Pedunculi tomentosi & marginibus spinulosis armati. Habitus antecedentis elucet in textura foliorum, panicula, calycis figura, magnitudine.

821
nitoria.

SERRATULA.

præalta.

1. SERRATULA foliis pinnatifidis: pinna terminali maxima. *Hort. cliff. 391. Hort. npsf. 249. Fl. succ. 660. Roy. lugdb. 143. Dalib. parif. 246.*
Serratula. Bauh. pin. 235. Dod. pempt. 24.
- β. Serratula præalta centauroides montana italica. *Bocc. mus. 2. p. 45. t. 37.*
Carduus inermis, foliis glabris pinnatis: lacinia externa maxima, capitulis squalidis. Gmel. fib. 2. p. 49. t. 20.
Habitat in Europæ borealioris pratis β. in Italia, Sibiria. 2
Planta β. quadruplo omnibus partibus major, at structura omnibus partibus cum communi eadem.

alpina.

2. SERRATULA calycibus subhirsutis ovatis.
Serratula foliis petiolatis: radicalibus ovato-oblongis dentatis; caulinis integerrimis. Fl. succ. 661.
Serratula foliis ovato-lanceolatis: radicalibus serratis, caule thyrsifloro. Fl. lapp. 291. Hort. npsf. 391.

Cirs-

SYNGENESIA: POLYGAMIA ÆQUALIS. 817

Cirsium inerme, foliis ex ovato-lanceolatis denticulatis infra lanugine candidis. *Gmel. fib. 2. p. 67 t. 26.*

Carduo-Cirsium minus britannicum, floribus congestis. *Pluk. alm. 83. t. 134. f. 3.*

β. Cirsium humile montanum, cynoglossi folio, polyanthemum. *Morif. bist. 3. p. 148. Dill. elzb. 82. t. folia.*
70. f. 81. *vide. Gmel. fib. 2. p. 76. t. 32.*

γ. Cirsium polyanthemum, molli hastato folio. *Morif. bist. lapatifolia.*
3. p. 148. f. 7. t. 29. f. 1. t. 22.

Cirsium foliis triangularibus lunate dentatis. *Hall. helv. 683.*

Carduus mollis, lapathi foliis. *Bauh. pin. 377.*

Carduus mollior 2. *Clus. bist. 2. p. 151.*

δ. Cirsium inerme, foliis linearibus utrinque viridibus, calycibus hirsutis. *Gmel. fib. 2. p. 78. t. 33.*

Habitat in Alpibus Lapponiæ, Austriæ, Helvætiæ, Arvoniz, Sibiriz. &

Planta floribus distinctissima, foliorum figura diversissima; sed quæ causa produxerit has constantes varietates? certe locus non suffecerit.

3. SERRATULA foliis linearilanceolatis alternis subsalicifolia. tus incanis sessilibus integerrimis.

Cirsium inerme erectum, foliis ex linearilanceolatis: infra candidis. *Gmel. fib. 2. p. 63. t. 27.*

Habitat in Sibiriz apricis, siccis.

Caulis angulatus, superne ramosus, ut in congeneribus.

Folia salicina, integerrima, subtus albida, basi vix ac ne vix decurrentia. Flores rubri; Calyces facie S. alpina, tres vel quatuor in singulo ramo.

4. SERRATULA foliis lanceolatis subtus villosis submultiflora. decurrentibus integerrimis, caule corymbofo, calycibus cylindricis.

Serratula elatior, floribus umbellatis parvis. *Gerb. tan. 294.*

Cirsium inerme, caulibus adscendentibus, foliis linearibus infra cinereis. *Gmel. fib. 2. p. 71. t. 28? sed flores majores.*

Habitat in Sibiria. D. Gmelin.

Caulis angulatus, corymbosus, ramis itidem corymbosis, ut terminetur densissima sylva florum, fere infinitorum. Folia saligna, subtus albo villo vestita. Calyces cylindrici squamis glabris, acutis, purpurascentibus; similis præcedenti, sed folia basi parum decurrentia, subtus villosa, & Calyces copiosiores, argutiores, glabri magis & latius colorati.

F f f

5. SER-

518 SYNGENESIA: POLYGAMIA ÆQUALIS.

- noveboracensis.* 5. **SERRATULA** foliis lanceolato-oblongis serratis pendulis. *Hort. cliff.* 392. *Roy. lugdb.* 143.
Serratula noveboracensis maxima, foliis longis serratis. *Dill. elsb.* 255. t. 263. f. 342.
Serratula noveboracensis altissima, foliis doræ mollibus subincanis. *Morif. bist.* 3. p. 133. *Raj. suppl.* 208.
Centaurium medium noveboracense luteum, solidaginis folio integro tenuiter crenato. *Pluk. alm.* 93. t. 109. f. 3.
Habitat in Noveboraco, Virginia, Carolina, Canada. Kamtschatka. 2
- pratensis.* 6. **SERRATULA** foliis lanceolato-oblongis serratis patentibus subtus hirsutis.
Serratula virginiana, persicæ folio subtus incano. *Dill. elsb.* 356. t. 264. f. 343.
Serratula præalta, angusto plantaginis aut persicæ folio. *Bocc. mus.* 2. p. 45. t. 32.
Eupatoria virginiana, serratulæ noveboracensis latioribus foliis. *Pluk. alm.* 141. t. 280. f. 6.
Habitat in Carolina, Virginia, Pennsylvania.
- glauca.* 7. **SERRATULA** foliis ovato-oblongis acuminatis serratis, floribus corymbosis, calycibus subrotundis.
Gron. virg. 92.
Serratula marilandica, foliis glaucis cirsii instar denticulatis. *Dill. elsb.* 354. t. 262. f. 341.
Centaurium medium marianum, folio integro cirsii nostratis more spinulis fimbriato. *Pluk. mant.* 40.
Habitat in Marilandia, Virginia, Carolina. 2
- squarrosa.* 8. **SERRATULA** foliis linearibus, calycibus squarrosis sessilibus acuminatis. *Hort. cliff.* 392. * *Gron. virg.* 92. *Roy. lugdb.* 143.
Cirsium tuberosum, capitulis squarrosis. *Dill. elsb.* 83. t. 71. f. 82.
Stœbe virginiana tuberosa latifolia, capitulis sessilibus: squamis foliaceis acutis donatis. *Morif. bist.* 3. p. 137. f. 7. t. 27. f. 10. *bona.*
Habitat in Virginia. 2
- ramosa.* 9. **SERRATULA** foliis lanceolatis integerrimis, calycibus squarrosis pedunculatis obtusis lateralibus.
Cirsium non ramosum, foliis lateralibus, flores ferens pauciores majores. *Gron. virg.* 92.
Jacea altera non ramosa, tuberosa radice, foliis latioribus, flores ferens pauciores majores. *Banist. virg.* 1929. *Pluk. mant.* 105. *Eupa-*

SYNGENESIA. POLYGAMIA ÆQUALIS. 319

Eupatorio affinis americana bulbosa, floribus scariosis, capitulis contextis. *Pluk. alm.* 142. t. 177. f. 4.

Habitat in Virginia.

Diversissima a precedente. Calycibus squarrosis, obtusis. Pedunculis lateralibus, elongatis, unifloris, foliolis, aliquot instructis. Foliis lato-lanceolatis, minime nervosis, sessilibus, nec decurrentibus.

10. SERRATULA foliis linearibus, floribus sessilibus *spicata*. lateralibus spicatis, caule simplici.

Serratula foliis linearibus, floribus solitariis sessilibus. *Gron. virg.* 92.

Cirsium tuberosum, lactucæ capitulis spicatis. *Dill. elth.* 85. t. 72. f. 83.

Jacea non ramosa, tuberosa radice, floribus plurimum rigidis perangustis. *Banist. virg.* 1927.

Jacea angustifolia, tuberosa radice, virginiana. *Pluk. alm.* 190. t. 424. f. 6.

Stœbe virginiana tuberosa angustifolia, capitulis sessilibus. *Morif. bist.* 3. p. 137. f. 7. t. 27. f. 10.

Habitat in America septentrionali. &

11. SERRATULA foliis linearibus sparsis revolutis, *ChamaPent-* caule fruticoso. *ce.*

Centaurea calycibus inermibus: squamis lanceolatis, foliis linearibus confertis integerrimis. *Hort. cliff.* 421.

Roy. lugdb. 138.

Jacea fruticans, pini folio. *Bauh. pin.* 271. *Pluk. alm.* 190. t. 94. f. 3.

Chamaepeuce. *Alp. exot.* 77. t. 76.

Stœbe capitata, chamaepinus fruticosa cretica. *Pon. ital.* 75.

Stœbe cretica fruticans, pinæ angustis foliis. *Morif. bist.* 3. p. 133. f. 7. t. 26. f. 8.

Habitat in Creta. &

12. SERRATULA foliis lanceolatis, squamis calycinis *amaræ* apice membranaceis obtulis patulis coloratis, floribus terminalibus.

Cirsium inermis, foliis scabris: inferioribus ex sinuato dentatis, squamis calycum subrotundis membranaceis.

Gmel. fib. 2. p. 72. t. 29.

Habitat in Sibiria.

Habitus *S. alpinae*. Folia lanceolata, nonnihil decurrentia, superiora integerrima, margine præsertim scabra. Caucis angulati, Rami erecti, singuli terminati pedun-

F f f 2

cu-

820 SYNGENESIA: POLYGAMIA ÆQUALIS

culis floribusque sæpius tribus. Flores magnitudine Cyani rubri, Pappus albus; Calyces squarrosi squamis membranaceis, mollibus ut Centaureæ nonnullæ, coloratis; pedunculi duo s. tres ramos terminant.

centaureoides 13. SERRATULA foliis pinnatifidis acutis glabris inermibus, squamis calycinis mucronatis: interioribus membranaceis.

Cirsium inerme, foliis scabris: inferioribus pinnatifidis Gmel. fib. 2. p. 80. t. 35.

Habitat in Sibiria. &

Herba pedalis. Rami alterni (in culta), stricti, fere glabri, uniflori. Folia utrinque glabra, venosa, acuta, inermia, lanceolata: dentibus lanceolatis. Calyx ovato-oblongus, imbricatus squamis mucrone inermi, quarum flosculis propiores majores exaridæ. Corollæ purpureæ absque flosculis marginalibus. Habitus Centaureæ moschata, sed caret flosculis neutris.

arvensis. 14. SERRATULA foliis dentatis spinosis. Fl. succ. 662 Dalib. parif. 246.

Carduus radice repente, foliis lanceolatis dentatis margine aculeatis. Fl. lapp. 293. Roy. lugdb. 133.

Carduus vincarum repens, fonchi folio. Bauh. pin. 387. prodr. 156.

Carduus in avena proveniens. Bauh. pin. 377.

Ceanothus theophrasti. Col. ceph. 1. p. 45. t. 45.

Habitat in Europæ cultis agris. &

822

CARDUUS.

* *Folius decurrentibus.*

Acarna. 1. CARDUUS foliis lanceolatis dentatis ciliatis decurrentibus: spinis marginalibus duplicibus.

Carduus caule alato spinoso, foliis spinosis: spinis duplicibus. Roy. lugdb. 133.

Acarna major, caule folioso. Bauh. pin. 379.

Chamæleon salmaticensis. Clus. hist. 2. p. 155.

β. Acarna humilis, caule folioso. Bauh. pin. 379.

Picnomon cretæ saloniensis. Dalech. hist. 1456.

Habitat in Hispania.

lencographeus. 2. CARDUUS foliis dentatis spinosis decurrentibus, pedunculis nudis longissimis unifloris, calycibus spinosis inclinatis.

Carduus foliis sinuatis aculeatis, caule crispo, pedunculis unifloris. Roy. lugdb. 134.

Cirs-

SYNGENESIA: POLYGAMIA ÆQUALIS. 824

Cirsium maculis argenteis notatum. Tournef. inst. 448.

Carduus leucographus f. *mariz* hirsutus, capitulo minore. *Morif. biesf. 244.*

Habitat in Campania.

3. **CARDUUS** foliis decurrentibus hastato-pinnatis spinosis: laciniis divaricatis, calycibus ovatis spinosis villosis, caule piloso. *lanceolatus.*

Carduus foliis lanceolatis decurrentibus: denticulis superficieque spinosis. Hort. cliff. 392. Fl. suec. 654. Roy. lugdb. 132. Dalib. parif. 247.

Carduus lanceolatus latifolius. Bauh. pin. 385.

Carduus lanceolatus f. *sylvestris dodonæi. Bauh. hist. 3. p. 58.*

Habitat in Europæ cultis ruderatis.

Calycis intima squama inermes, setacea, exaride.

4. **CARDUUS** foliis spinosis semidecurrentibus, floribus cernuis: squamis calycinis superne patentibus. *nutans.*
Fl. suec. 655. Guett. stamp. 2. p. 321. Dalib. parif. 248. Gmel. sib. 2. p. 42.

Carduus foliis sinuatis decurrentibus margine spinosis, floribus solitariis nutantibus. Hort. cliff. 393.

*Carduus nutans. Bauh. hist. 3. p. 56. It. goth. 293. **

Carduus spinosissimus latifolius sphærocephalus vulgaris. Bauh. pin. 385.

Habitat in Europa ad pagos. ♂

5. **CARDUUS** foliis sinuatis decurrentibus margine spinosis, calycibus pedunculatis solitariis erectis villosis. *acanthoides*

Carduus foliis laciniatis: subtus tomentosis ex margine spinosis, capitulis sessilibus congestis aculeatis. Hall. herb. 677. Dalib. parif. 248.

*Carduus acanthoides. Bauh. hist. 3. p. 59. It. scan. 297. **

Carduus spinosissimus, capitulis minoribus. Raj. angl. 3. p. 194.

Carduus polyacanthos, capitulis longioribus & tenuioribus, foliis albicantibus. Morif. hist. 3. p. 153.

Habitat in Europæ ruderatis. ☉

Differt a Carduo nutante calycibus erectis, a C. crispo calycibus solitariis villosis.

6. **CARDUUS** foliis sinuatis decurrentibus: margine spinosis, flosculis aggregatis terminalibus nudis. *crispus.*
Hort. cliff. 393. Fl. suec. 658. Roy. lugdb. 133. Dalib. parif. 248.

822 SYNGENESIA: POLYGAMIA ÆQUALIS.

Carduus spinosissimus angustifolius vulgaris. Loef. pruss
43. t. 5.

Carduus caule crispo. Bamb. hist. 3. p. 9.

Habitat in Europæ septentrionalioris agris, cultis. ☉

palustris. 7. *CARDUUS* foliis dentatis decurrentibus: margine spinosis, floribus racemosis erectis, pedunculis inermibus. *Carduus foliis linearibus dentatis spinosissimis, caule crispo spinoso, floribus erectis. Roy. lugab. 134. Fl. suec. 659. Dalib. paris. 249.*

Carduus palustris. Bamb. pin. 377. prodr. 156.

Carduus spinosissimus erectus angustifolius palustris.

Morif. hist. 3. p. 153. f. 7. t. 32. f. 13.

Carduus polycephalus angustifolius. Dalech. hist. 1743.

Habitat in Europæ pratis subpaludosis.

difsectus. 8. *CARDUUS* foliis lanceolatis decurrentibus denticulis inermibus, calyce spinoso. *Hort. cliff. 392.* Dalib. paris. 249.*

Cirsium anglicum. Lob. ic. 583. Dalech. hist. 584. figura tenuis.

Cirsium majus, singulari capitulo magno, f. incanum varie difsectum. Bamb. pin. 377.

Habitat in Anglia, Gallia. &

cyanoides. 9. *CARDUUS* foliis pinnatifidis linearibus integerrimis inermibus petiolaris decurrentibus: subtus tomentosus. *monoclonos.* *Carduus caule simplici, foliis pinnatifidis rarius denticulatis: infra candidis, squamis calycum lanceolatis adscendentibus. Gmel. sib. 2. p. 42. t. 15.*

polyclonos. β. *Carduus inermis, caule ramosissimo, foliis inferioribus pinnatis; superioribus linearibus integerrimis; infra candidis, squamis calycum lanceolatis adscendentibus. Gmel. sib. 2. p. 44. t. 16.*

Habitat in Tataria.

Folia subtus albo tomento, absque serraturis aut setis, α fere Sorbi; β angustissima & linearia. Calycum squamæ lineares, acuminatæ, subvillosæ: in α magnitudine Scabiosæ, in β. Cyani. Habitus α. Jacea moschata J. B. 3. p. 34.

monspeffulans. 10. *CARDUUS* foliis lanceolatis subrepandis glabris decurrentibus inæqualiter ciliatis, pedunculis alternis, calycibus inermibus.

Carduus foliis lanceolatis glabris integris ciliatis decurrentibus, calycibus inermibus. sauv. monsp. 79.

Car.

SYNGENESIA · POLYGAMIA ÆQUALIS. 813

Carduus Cirsium monspelianum, folio longo glabro.

Bauh. hist. 3. p. 44.

Cirsium foliis non hirsutis, floribus compactis. *Bauh. pin.* 377.

Cirsion. *Dod. pempt.* 377.

Habitas Monspeli.

Habitus C. Serratuloides. Folia lanceolata, utrinque glabra, glauca, integra; radicalia vero repanda, margine spinis setaceis, inæqualibus ciliata. Capitata nonnulla in summitate caulis pedunculis brevibus.

* *Foliis amplexicaulibus.*

11. **CARDUUS** foliis lanceolatis sessilibus integerrimis: *casabonæ*, margine spinis ternatis. *Hort. cliff.* 393.

Acarua major, caule non folioso. *Bauh. pin.* 379.

Acarua theophrasti angvillaræ. *Lob.* 486. f. 1. *Dalech. hist.* 1484.

Polyacanthus casabonæ, *Acaruæ* similis. *Bauh. hist.* 3. p. 92.

Habitas - - -

12. **CARDUUS** foliis integris subtus tomentosis, spinis *stellatis*, ramosis lateralibus. *Roy. lugdb.* 133.

Carduus humilis aculeatus, *ptarmicæ austriacæ* foliis.

Triumf. obs. 96. t. 96.

Carduus stellatus leucoji lutei foliis. *Dedar. mem.* 4. p. 261.

Habitas - - - ☉

13. **CARDUUS** foliis amplexicaulibus hastato-pinnatifidis *marianus*, spinosis, calycibus aphyllis: spinis canaliculatis duplicato-spinosis.

Carduus squamis calycinis margine apiceque spinosis.

Hort. cliff. 393. * *Hort. upf.* 250. *Roy. lugdb.* 133.

Dalib. paris. 247. *Mat. med.* 316. *Sauv. monsp.* 291.

Carduus albis maculis notatus vulgaris. *Bauh. pin.* 281

Carduus mariæ. *Dalech. hist.* 1475.

Habitas in Angliæ, Galliæ, Italiæ *aggeribus ruderatis.* ☉

14. **CARDUUS** foliis amplexicaulibus angulato-spinosis *syriacus*, tis, floribus solitariis subsessilibus obvallatis foliolis subquinis. *Hort. cliff.* 393. * *Hort. upf.* 250. *

Carduus albis maculis notatus exoticus. *Bauh. pin.* 381.

Carduus luteus syriacus. *Cam. hort.* 35. t. 10.

Habitas in Syria, Creta, Hispania. ☉

15. **CARDUUS** foliis sessilibus bisariam pinnatifidis, *la-crisophorus* ciniis alternis erectis, calycibus globosis villosis. *Hort. upf.* 249. * *Sauv. monsp.* 290.

F f f 4

Car-

824 SYNGENESIA: POLYGAMIA ÆQUALIS.

- Carduus foliis sinuatis decurrentibus: denticulis super-
ficieque spinosis, calycibus lanigeris. Roy. lugdb. 133.
Dalib. parif. 247.*
- Carduus capite rotundo tomentoso. Baub. pin. 382.*
- Carduus tomentosus corona fratrum herbariorum. Lob.
ic. 2. p. 9.*
- spurius.* β. *Carduus foliis semidecurrentibus duplicato pinnatifidis:
laciniis alternis erectis, calycibus ovatis. Hort. upf.
249. not.*
- Carduus tomentosus pyrenaicus, floribus purpureis glo-
meratis. Tournef. inst. 441?*
- Habitat in Anglia, Gallia, Hispania, Lusitania. ♂*
- altissimus.* 16. **CARDUUS** foliis pinnatifidis sinuatis serratis sessi-
libus inermibus, caule ramulissimo, calycibus villosis
subferratis.
- Cirsium altissimum, laciniato folio subtus tomentoso.
Dill. elth. 81. t. 69. f. 80.*
- Habitat in Carolina. ♀*
- virginianus.* 17. **CARDUUS** caule folioso unifloro, foliis lanceola-
tis spinulosis: subtus tomentosis, caule inermi. *Gron.
virg. 92.*
- Cirsium minus virginianum, singulari capitulo, caule
folioso. Morif. hist. 3. p. 150. Raj. suppl. 197.*
- Habitat in Virginia.*
- tuberosus* 18. **CARDUUS** foliis pinnatifidis spinosis sessilibus, cau-
le inermi, capitulis solitariis.
- Carduus foliis semipinnatis: pinnis semibilobis, capitu-
lis inermibus singularibus. Sauv. monsp. 292.*
- Carduus pratensis, asphodeli radice, latifolius. Baub.
pin. 377. Morif. hist. 3. p. 7. t. 29. f. 27. 28.*
- Carduus bulbosus monspeliensium. Lob. ic. 10.*
- β. *Carduus pratensis, asphodeli radice, foliis tenuiter inci-
sis. Baub. pin. 377.*
- Jacea aculeata s. tuberosa. Tabern. ic. 154.*
- Habitat Mospelii, Liptiæ, inque Bohemia, Austria, Hel-
vetia.*
- heterophyl-
lus.* 19. **CARDUUS** caule subunifloro, calyce inermi acuto,
foliis amplexicaulibus lanceolatis ciliatis integris laci-
niatisque. *Hort. cliff. 392. * Fl. suec. 657. * Roy.
lugdb. 134. Gmel. sib. 2. p. 51.*
- Carduus calyce inermi, foliis lanceolatis margine cilia-
tis. Fl. lapp. 192.*

Cri-

SYNGENESIA: POLYGAMIA ÆQUALIS. 825

Cirsium majus, singulari capitulo magno, f. incanum varie dissectum. *Bauh. pin.* 377?

Habitat in Europæ frigidioris partibus depressis. &

Caulis sesquipedalis f. brevior. Foliis 5 ad 10, nondentatis, ciliatis. Radice repente.

10. **CARDUUS** foliis lanceolatis dentatis amplexicanli-*belentoides*, bus: spinulis inæqualibus ciliatis, caule inermi. *Hort. cliff.* 392. *Hort. upf.* 250. *

Cirsium foliis longe lanceolatis serratis: subtus tomentosis. *Hall. herb.* 683. t. 21.

Cirsium singulari capitulo squamato, vel incanum alterum. *Bauh. pin.* 377.

Cirsium anglicum, radice hellebori nigri modo fibrosa, folio longo. *Bauh. hist.* 3. p. 45?

Habitat in Anglia, Sibiria. &

Antecedenti *simillimus*, duplo f. homine altior, radice vix repente, foliis omnibus indivisis, subtus albis, at non ut illi niveis. Caule toto folioso; Foliis 40 a 50, semiamplexicaulibus (auriculis rotundis, reflexis, adnatis), dentatis, ciliatis spinis inermibus.

11. **CARDUUS** foliis lanceolatis integris subamplexicaulibus: serraturis spinoso-setaceis, pedunculis unifloris. *Hort. upf.* 248. (sub *Serratula*) * *Sauv. monsp.* 79.

Cardus calycibus inermibus acutis, foliis utrinque viridibus firmis integerrimis inæqualiter ciliatis. *Gmel. fib.* 2. p. 52. t. 22. & 23. f. 1.

Cirsium singularibus capitulis parvis. *Bauh. pin.* 377?

Cirsium 3 montanum 1. *Clus. hist.* 2. p. 149.

Habitat in Sibiria, Helvetia, Montspeli. &

12. **CARDUUS** foliis lanceolatis amplexicaulibus, serraturis spinoso-setaceis, capitulis triphyllis.

Habitat in Sibiria. &

Habitus & altitudo C. Serratuloides, ut facile pro varietate sumeretur, a quo differt Foliis duplo latioribus, magis dentatis, vix ac ne vix subtus tomentosis; Capitulis foliolis 3, rarius duobus obvallatis, nec unico, ut ille. Foliis basi pungentibus. Floribus calyce cylindrico, nec globofo. Colore florum albo ucc purpureo, at Pappoplumoso convenit.

13. **CARDUUS** foliis lanceolatis integris inermibus dentatis glabris, capitulis aphyllis inermi spinosis.

F f f 5

Habi-

826 SYNGENESIA: POLYGAMIA ÆQUALIS

Habitat in Hispania. L'hsing.

Caulis *levis*. Folia *lanceolata*, *basi attenuata*, *argute dentata*, *inermia*. Calyces *cylindrico-ovati*, *terminati squamis aciformibus*, *mitibus*, *patulis*. Corolla *albo-flavescentis*.

Y23

CNICUS.

benedictus. 1. CNICUS caule diffusio, foliis dentato-sinuatis. *Hort. cliff.* 395. *Hort. upf.* 250. *Mat. med.* 379.

Carduus lylveltris hirsutus s. *Carduus benedictus*. *Baub. pin.* 378.

Carduus benedictus. *Cam. epit.* 562.

Habitat in Chio, Lemno, Hispania ad versuras agrorum. ☉

oleaceus. 2. CNICUS caule erecto, foliis inferioribus laciniatis; superioribus integris concavis. *Hort. cliff.* 394. *Fl. succ.* 711.

Carduus pratensis lausfolius. *Baub. pin.* 376.

Carduus pratensis. *Lab. ic.* 2. p. 11.

Habitat in Europæ septentrionalioris pratis subnemore-sis. ☿

Calyces *laterales sessiles*, *sapius bini*, *intermedio subpedunculato*.

spinosissimus. 3. CNICUS foliis amplexicaulibus sinuato-pinnatis spinosis, caule simplici, floribus sessilibus.

Carlina polycephalos alba. *Baub. pin.* 380

Cirsium alpinum spinosissimum, floribus ochroleucis inter flavescentia folia congestis. *Hall. helv.* 679. t. 20.

Habitat in Alpibus Helvetiæ, Austriæ.

Centauroides 4. CNICUS foliis pinnatifidis, squamis calycinis membranaceis.

Carduus inermis, foliis pinnatifidis serratis inermibus, squamis calycinis membranaceis acuminato-lanceolatis. *Hort. cliff.* 392.

Centaurium majus, foliis cinaræ cornuti. *Morif. hist.* 3. p. 131. f. 7. t. 25. f. 2.

Habitat in Pyrenæis. ☿

perennis. 5. CNICUS foliis cordatis, petiolis crispis spinosis amplexicaulibus, floribus cernuis. *Hort. upf.* 251. *

Carduus foliis ex cordato-lanceolatis margine serratis & spinosis, squamis calycum membranaceis laceris spinosis, capitulis nutantibus. *Gmel. sib.* 2. p. 47. t. 19.

Habitat in Sibiria. ☿

ONON

SYNGENESIA: POLYGAMIA ÆQUALIS. 827

ONOPORDUM. 824

1. ONOPORDUM calycibus squarrosis, foliis ovato-*acanthium*.
oblongis sinuatis.

Onopordum foliis decurrentibus margine spinosis. *Hort. cliff.* 393. *Fl. suec.* 653. *Roy. lugdb.* 134. *Dalib. paris.* 246.

Spina alba tomentosa latifolia sylvestris. *Bauh. pin.* 382.
Loef. pruss. 261 t. 82.

Acanthium. *Dod. pempt.* 721.

Habitat in Europæ ruderatis, cultis. 3

2. ONOPORDUM calycibus squarrosis, foliis linearibus pinnatifidis.

Onopordum foliis decurrentibus ligulatis pinnatifidis.
Sauv. montp. 270.

Spina tomentosa altera spinosior. *Bauh. pin.* 382.

Carduus, quibusdam dictus Acanthium illyricum. *Bauh. hist.* 3. p. 55.

Acanthium illyricum. *Lob. ic.* 1.

Habitat in Europa australi.

Folia vix caule latiora. *Sauv.*

3. ONOPORDUM calycibus imbricatis *Hort. upf.* 249. *arabicum*,
Carduus tomentosus, acanthi folio, altissimus lusitanicus.

Tournef. inst. 441.

Carduus tomentosus, Acanthium dictus, arabicus. *Pluk. alm.* 85. t. 154. f. 5.

Acanthium altissimum lusitanicum. *Morif. hist.* 3. p. 153.

Habitat in Lusitania, G. Narbonensi. 3

CYNARA. 825

1. CYNARA foliis subspinosis pinnatis indivisisque, ca-*Scolymus*.
lycinis squamis ovatis.

Cynara foliis pinnatis: laciniis ferratis. *Hort. cliff.* 394.
Hort. upf. 251. *Roy. lugdb.* 135.

Cynara sylvestris latifolia. *Bauh. pin.* 384.

- β Cynara hortensis aculeata. *Bauh. pin.* 383.

Scolymus diofcoridis. Clus. hist. 2. p. 153.

- γ Cynara hortensis, foliis non aculeatis. *Bauh. pin.* 383.

Habitat in G. Narbonensis, Italiæ, Siciliæ agris. 2

2. CYNARA foliis spinosis: omnibus pinnatifidis, caly-*Cardunculus*
ciis squamis ovatis.

Cina-

528 SYNGENESIA. POLYGAMIA. ÆQUALIS.

Cinara foliis pinnatis: foliolis oblongis indivisis, caule spinoso. *Sauv. meth.* 239.

Cinara spinosa, cujus pediculi esitantur. *Bauh. pin.* 383.

Scolymus aculeatus. *Tabern. hist.* 1075.

Habitat in Creta. &

Hybridam f. natam ex precedenti semine asseverat J. B.

Flos saturate cæruleus corollis & pistillis, at stamina grisea.

humilis.

3. *CYNARA* foliis spinosis pinnatifidis subtus tomentosis, calyceibus squamis subulatis.

Cinara sylvestris bætica. *Clus. enr.* 35.

Carduus tingitanus, flore magno cæruleo, folio atrætylidis, divisura subtus incano, spinis durioribus horrido. *Pluk. alm.* 85. t. 81. f. 2.

Carduus humilis tingitanus cæruleus magno strobilo tenuius lacinia. *Morif. hist.* 3. p. 158.

Habitat in agro Tingitano in Bætica. &

Semina, in hac specie minore, reliquis longe majora & singularia.

826

CARLINA.

acaulis.

1. *CARLINA* caule unifloro flore brevior. *Hort. cliff.* 395. *Hort. upf.* 252. *Mat. med.* 378. *Roy. lugdb.* 135. *Sauv. monsp.* 293.

Carlina acaulos, in magno flore albo. *Bauh. pin.* 380.

Chamæleon albus. *Clus. hist.* 2. p. 155. *Cam. epit.* 428.

Habitat in Italiæ, Germaniæ montibus. &

lanata.

2. *CARLINA* caule trifloro dichotomo: intermedio sessili. *Sauv. monsp.* 293. †

Acarna flore purpureo rubente patulo. *Bauh. pin.* 379.

Acanthoides parva apula. *Col. ecphr.* 1. p. 29. t. 27. f. 2.

Habitat in Italia, G. Narbonensi. ☉

Calyx squamis lateribus inermibus constat.

umbellatis.

3. *CARLINA* caule multifloro subdiviso, floribus sessilibus. *Roy. lugdb.* 135.

Acarna capitulis parvis luteis in umbella. *Bauh. pin.* 379.

Acarna apula umbellata. *Colum. ecphr.* 27.

Habitat in Italia. ☉

corymbosa.

4. *CARLINA* caule multifloro corymboso, floribus terminali-

SYNGENESIA: POLYGAMIA ÆQUALIS. 829

minalibus. *Hort. cliff.* 395. *Fl. suec.* 652. *Roy. lugdb.* 135. *Gmel. fib.* 2. p. 82.

Carlina sylvestris vulgaris. Clus. hist. 2. p. 155.

Cnicus sylvestris spinosior. Bauh. pin. 378.

Atractylis mitior. Fuchf. hist. 121.

Habitat in Europæ montosis, aridis, sabulosis.

5. **CARLINA** floribus sessilibus lateralibus paucissimis. *racemosa. Sauv. meth.* 293.

Carlina sylvestris minor hispanica. Clus. hist. 2. p. 157

Acarna flore luteo patulo. Bauh. pin. 379.

Habitat in Hispaniæ desertis. ☉

6. **CARLINA** caule multifloro, foliis decurrentibus. *pyrenaica. Carduus tomentosus pyrenaicus spinosissimus, flore albo. Bursf. XXI.* 43.

Habitat in Pyrenæis versus Hispaniam copiose.

Habitus Onopordi, magnitudo antecedentis. Tota planta villis lanata. Folia pinnata, lobis rotundatis, spinisque validis armatis, decurrentia per caulem totaliter spinis aequè validis. Calyces copiosi in summo canale: squamis lanceolatis, longitudine floris, mucronatis: interioribus longioribus, membranaceis, lanceolatis, albis.

ATRACTYLIS. 827

1. **ATRACTYLIS** foliis oblongo-ovatis denticulatis spinosis, calycibus patentibus, caule fruticoso. *fruticosa. Hort. cliff.* 295. *Roy. lugdb.* 137.

Carthamus africanus frutescens, folio ilicis, flore aureo. B. Walt. hort. 13. t. 7.

Crocodiloides atractylidis folio, flore sulphureo coronato. Vaill. act. 1718. p. 207.

Habitat in Æthiopia. ♀

2. **ATRACTYLIS** flore acaule. *gummifera.*

Cnicus, carlinæ folio, acaulos gummiifer aculeatus.

Tournef. cor. 33.

Carlina acaulos gummiifera. Bauh. pin. 380.

Carduus pinca theophrasti. Alp. exot. 126. t. 124. 125.

Chameleo albus dioscoridis. Colum. ecphr. 1. p. 1. t. 12.

Habitat in Creta, Italia.

3. **ATRACTYLIS** foliis dentato-sinuatis flore radiato: obvallato involuero patente, caule herbaceo.

Cnicus aculeatus purpureus humilior. Tournef. inst. 451.

Cro-

830 SYNGENESIA: POLYGAMIA ÆQUALIS.

Crocodiloides atractylidis folio, flore purpureo coronato. *Vaill. act.* 1713. p. 172.

Habitat Madritii in collibus. *Læf. VII:* 35.

Habitus *A. cancellata*, sed folia sinuata. Flos radiatus. Calycis squamæ mucronatæ. Involucrum ex foliis veris patentibus.

candelata. 4. ATRACTYLIS involucris cancellatis ventricosissimis linearibus dentatis, calycibus ovatis, floribus flosculosis.

Atractylis foliis linearibus dentatis, calycibus conniventibus. *Hort. cliff.* 395. *Koy. lugdb.* 137.

Acarna capitulis globosis. *Bauh. pin.* 379.

Eryngium parvum palmare, foliis serratis. *Morif. hist.* 3. p. 166. f. 7. t. 36. f. 16.

Carduus parvus. *Bauh. hist.* 3. p. 93. *Ruj. hist.* 316.

Carduus minimus. *Alp. exot.* 254.

Habitat in Hispania, Sicilia, Cretæ agris. ☉

Receptaculum rectum paleis coalitis. Pappus plumosus, basi quasi monophyllos sub florescentia, corollulis longior. *Læf.*

828

CARTHAMUS.

stultiorius. 1. CARTHAMUS foliis ovatis integris serrato-aeuleatis. *Hort. cliff.* 394. *Hort. upf.* 251. *Mat. med.* 377.

Cnicus sativus f. Carthamus officinarum. *Bauh. pin.* 378.

Cnicus vulgaris. *Clus. hist.* 2. p. 152.

Habitat in Ægypto. ☉

lanatus. 2. CARTHAMUS caule piloso: superne lanato, foliis inferioribus pinnatifidis, summis amplexicaulibus dentatis. *Hort. upf.* 251. *Sauv. monsp.* 292.

Carthamus foliis amplexicaulibus acuminato-dentatis.

Hort. cliff. 394. *Dalib. parif.* 250.

Atractylis lutea. *Bauh. pin.* 379.

Atractylis theophrasti & dioscoridis sacco sanguineo.

Col. ecphr. 1. p. 19. t. 23.

Atractylis. *Dod. pempt.* 736.

Habitat in Gallia, Italia, Creta. ☉

cæruleus. 3. CARTHAMUS foliis lanceolatis acuminato-serratis. *Hort. cliff.* 135. *Koy. lugdb.* 136.

Cnicus perennis cæruleus tingitanus. *Herm. lugdb.* 162. t. 163.

Carduus cæruleus erectus tingitanus, cnici facie, foliis magis integris. *Morif. hist.* 3. p. 159. β. Cui-

SYNGENESIA: POLYGAMIA ÆQUALIS. 831

β. Cnicus cæruleus asperior. *Bauh. pin.* 378.

Cnicus alter clusii, cæruleo flore. *Clus. hist.* 2. p. 152.

Carduus cæruleus erectus, cnici foliis dissectioribus. *Morif. hist.* 3 p. 159. f. 7. t. 34 f. 19.

Habitat in agro Tingitano & Hispalensi inter segetes. 2

4. CARTHAMUS foliis inermibus: radicalibus dentatis, caulinis pinnatis. *Guss. stamp.* 2. p. 330. *Dalib. parif.* 250.

Carthamoides cærulea humilis & mitior. *Vaill. act.* 1718.

Carduncellus montis lupi minor species. *Morif. hist.* 3. p. 159.

Habitat circa Parisios.

5. CARTHAMUS foliis caulinis linearibus pinnatis longitudine plantæ. *Carduncellus.*

Carthamus foliis caule longioribus ligulatis pinnatifidis.

Savv. monsp. 292. *

Carthamus foliis radicalibus superne ferratis, inferne dentatis; caulinis dentato-pinnatis. *Roy. lugdb.* 136.

Cnicus cæruleus humilis montis lupi. *Herm. lugdb.* 164.

Carduncellus montis lupi. *Lob. ic.* 2. p. 20.

Eryngium montanum minimum, capitulo magno. *Bauh. pin.* 386.

Habitat Monspeli.

6. CARTHAMUS foliis ensiformibus sinuato-dentatis. *arborescens.*

Roy. lugdb. 136.

Carthamoides lutea altissima & foetidissima. *Vaill. act.* 1718. p. 772.

Cnicus hispanicus arborescens foetidissimus. *Tournef. inst.* 451.

Habitat in Hispania. 5

STOEBE.

829

1. STOEBE. *Hort. cliff.* 390. *

Coryza africana frutescens, foliis ericæ hamatis & in-ethiopica. canis. *Tournef. inst.* 455.

Eupatorioides capensis capitatus. *Pet. gaz.* 13. t. 8. f. 1?

Helichrysoides juniperi creberrimis aduncis foliis, floribus in ramulorum cymis. *Vaill. act.* 1719. p. 393.

Habitat in Æthiopia. 5

BIDENS.

850

1. BIDENS corollis flosculosis, calycibus subfoliosis, se- tripartitis minibus erectis, foliis trifidis.

Bi.

832 SYNGENESIA: POLYGAMIA ÆQUALIS.

Bidens corona seminum retrorsum aculeata, seminibus erectis. *Hort. cliff.* 399. *Fl. suec.* 663. *Roy. lugdb.* 157.

Bidens foliis tripartito-diviſis. *Tournef. inst.* 462. *Fl. lapp.* 294.

Cannabina aquatica, folio tripartitum diviſo. *Banb. pin.* 321.

Hepatorium aquatile. *Dod. pempt.* 595.

Habitat in Europæ inundatis. ☉

cernua.

2. *BIDENS* seminibus erectis: foliis lanceolatis amplexicaulibus, floribus cernuis. *Fl. suec.* 664.

Cannabina aquatica, folio non diviſo. *Banb. pin.* 321

Euphorbium cannabinum chrysanthemum. *Tabern. ic.* 117.

Habitat in Europa ad fontes & fassas ☉

Conferatur *Coreopsis bidentis* ejusque affinitas.

nodiflora.

3. *BIDENS* foliis oblongis integerrimis, caule dichotomo, floribus solitariis sessilibus.

Bidens nodiflora, bruellæ folio. *Dill. elth.* 52. t. 44. f. 52.

Habitat in Benghala.

frondosa.

4. *BIDENS* foliis pinnatis serratis, seminibus erecto-distantibus, calycibus frondosis, corollis radiatis.

Bidens canadensis latifolia, flore luteo. *Tournef. inst.* 362.

Chrysanthemum cannabinum bidens virginianum, caule erecto firmo subrubente. *Morif. bist.* 3. p. 17. f. 6 t. 5. f. 21.

Habitat in America septentrionali. ☉

Folia ternata aut ex 5 pinnata, supra glabra, striata, serrata. Calyces frondosi; rudius in corolla brevissimus, saepe mutilatus.

pilosa.

5. *BIDENS* foliis pinnatis subpilosis, corollis flosculosis, calycibus simplicibus, seminibus divergentibus.

Bidens corona seminum retrorsum aculeata, seminibus undique patentibus. *Hort. cliff.* 399 * *Hort. ups.* 255.

Bidens latifolia hirsutior, semine angustiore radiato. *Dill. elth.* 51. t. 43. f. 51.

Habitat in America. ☉

bipinnata.

6. *BIDENS* foliis bipinnatis incisis, calyce involucrato, corollis semiradiatis, seminibus divergentibus.

Chrysanthemum aquaticum, foliis multifidis cicutæ nonnihil similibus, virginianum. *Herm. lugdb.* 416.

Chrysanthemum americanum, coridis indi folio. *Herm. par.* 123. t. 123.

Chrys-

SYNGENESIA. POLYGAMIA ÆQUALIS. 833

- Chrysanthemum cannabinum* bidens virginianum, cicurariæ foliis, flosculis conniventibus. *Morif. hist.* 3. p. 17.
- Habitat in Virginia.* ☉
- Confer* *Chrysanthemum chinense*, foliis plurifariam divisis halicacabi pereigrini æmulis. *Pluk. phyt.* 22. f. 4.
- Chrysanthemum curassavicum*, foliis multifidis latioribus, seminibus longissimis. *Morif. hist.* 3. p. 17.
- Hybridas esse aliquas inter Bidentes conicas, frondosas & bipinnatas, suadet habitus & semina divergentia. Flores interdum carent radio, interdum flosculo uno alterove radiante & sterili instructi sunt.*
- . *BIDENS* foliis simplicibus serratis petiolatis, floribus *niveis*. globosis, pedunculis elongatis, seminibus lævibus.
- Bidens* foliis cordatis serratis petiolatis, floribus globosis, pedunculis filiformibus. *Roy. lugdb.* 158.
- Ceratocephalus* foliis cordatis s. triangularibus, flore albo. *Vaill. act.* 1720. p. 327. *vide. Dill. elsh.* t. 47. f. 55. n. 3.
- . *Bidens* scabra, flore niveo, folio trilobato. *Dill. elsh.* 55. t. 47. f. 55.
- . *Bidens* scabra, flore niveo, folio panduræformi. *Dill. elsh.* 54. t. 46. f. 54.
- Habitat in Carolina.* ☿
- . *BIDENS* foliis oblongis integris: inferne alternis, superne oppositis, floribus verticillatis. *Hort. cliff.* 399. * *verticillatis.*
- Bidens americana* procumbens, polygoni folio subtus incano. *Houft. mss.*
- Habitat in Vera Cruce*
- . *BIDENS* foliis ovatis serratis petiolatis, caule frutico-*fruticosa.*
- fo. Hort. cliff.* 399. *
- Habitat in America* ☿
- . *BIDENS* foliis oppositis ovatis acuminatis integerri-*scandens.*
- mis, caule scandente fruticoso, floribus oppositis paniculatis. *Hort. cliff.* 399. *
- Habitat in Vera Cruce.*
- . *BIDENS* foliis ovatis serratis: inferioribus oppositis, *bullata.*
- superioribus ternatis: intermedio majore.
- Habitat in America.* ☉
- Caulis erectus. Folia subovata, serrata, crassa, villosa,*
G g g
sere

200

34 SYNGENESIA: POLYGAMIA ÆQUALIS

fere bullata: inferiora simplicia, superiora ternata, subseffilia: foliolis lateralibus minimis. Flores axillares, solitarii, subseffiles, lutei; Calyces basi cincti foliolis ovato-oblongis.

CACALIA.

* *Frutescentes.*

- papillaris.* 1. CACALIA caule fruticoso, obvallato spinis petiolaribus truncatis.
Kleinia caule carnosio, petiolis truncatis obvallato. *Hort. cliff.* 395.
Cacalianthemum caudice papillari. *Dill. elth.* 63. t. 55. f. 63.
Habitat in Æthiopia. h

- Anteuphorbium.* 2. CACALIA caule fruticoso, foliis ovato-oblongis, petiolis basi linea triplici deductis.
Kleinia foliis carnosiss planis ovato-oblongis. *Hort. cliff.* 395.
Anteuphorbium. Bauh. pin. 387. *Dod. pempt.* 378. *Morif. hist.* 3. p. 345. f. 7. t. 37. f. 10. *Dill. elth.* 63. t. 55. f. 2. 3
Habitat in Æthiopia. h

- Kleinia* 3. CACALIA caule fruticoso composito, foliis lanceolatis planis, petiolorum cicatricibus obsoletis.
Kleinia foliis lanceolatis planis, caule lævi ventricosio. *Hort. cliff.* 395.
Cacalianthemum folio nerii glauco. *Dill. elth.* 61. t. 54. f. 2.
Frutex indiae orientalis, lavendulae folio. *Bauh. pin.* 401.
Nec Cacalia, nec Cacaliastrum, an Tithymaloides frutescens nerii folio. *Klein. monogr.*
Habitat in Canariis, forse etjam in India. h

- Ficoides,* 4. CACALIA caule fruticoso, foliis compressis carnosiss.
Kleinia foliis carnosiss lanceolatis compressis, caule tereti. *Hort. cliff.* 395.
Senecio africanus arborefcens, ficoidis folio & facie. *Comm. rar.* 40. t. 40.
Habitat in Æthiopia. h

* *Herbaceae.*

- Porophyllum* 5. CACALIA caule herbaceo indiviso, foliis ellipticis subcrenatis.

SYNGENESIA: POLYGAMIA ÆQUALIS. 835

Porophyllum foliis ellipticis. *Hort. cliff.* 494. *

Tagetes foliis integris perforatis. *Plum. spec.* 10.

Senecio indicus, atriplicis folio glabro. *Morif. hist.* 3. p. 106. f. 7. t. 17. f. 7.

Chrysanthemum americanum frutescens, balsaminæ foliis nigris maculis punctatis. *Pluk. alm.* 100. t. 161. f. 1.

Habitat in America. ☉

6. CACALIA caule herbaceo, foliis lyratis dentatis. *Sonchifolia.*

Kleinia caule herbaceo, foliis lyratis. *Fl. zeyl.* 305. *

Chondrilla zeylanica minor marina, folio sinapios. *Burm. zeyl.* 61.

Tagolina luzonum, flore purpureo. *Pet. gaz.* t. 80. f. 13. *Camell. luz.* 3. n. 8.

Habitat in Zeylona, China. ☉.

7. CACALIA caule herbaceo, foliis trilobis acuminatis *basata* ferratis.

Senecio foliis ex dentato-trilobis acuminatis ferratis *Gmel. fib.* 2. p. 136. t. 66.

Habitat in Sibiria. D. *Gmelin.*

Caulis erectus, altus, purpureus. Folia petiolata, triloba, lobis acuminatis: intermedio duplo majore, serrata inter lobos, basi rotundata, subius parum villosa. Flores in panicula parva, pedunculis incanis subtomentosis, absque bracteis setaceis præterquam ad subdivisiones. Facies accedit ad suaveolentem.

8. CACALIA caule herbaceo, foliis hastato-sagittatis *suaevoleus*. denticulatis, petiolis superne dilatatis. *Hort. ups.* 254. *

Habitat in Virginia, Canada. 2

Planta humana altitudinis. Folia sagittato-triangularia f. deltato-hastata, margine serrata; pedunculi etiam supra ramificationes bracteis setaceis adspersi, glabri nec incani.

9. CACALIA caule herbaceo, foliis subcordatis dentato-sinuatis, calycibus quinquefloris.

Cacalia virginiana glabra, foliis deltoidibus sinuatis subtus glaucis. *Morif. hist.* 3. p. 94. f. 7. t. 15. f. 7

Porophyllum foliis deltoidibus angulatis. *Gron. virg.* 94

Nardus americana procerior, foliis cæsiis. *Pluk. alm.* 251. t. 101. f. 2

Habitat in Virginia, Canada. 2

Magnitudo & statura antecedentis. Folia inferiora cordata, superiora latiora quam longa, sinuata, subtus glau.

G g g 2

glauca, laevia. Differt a precedente non modo figura foliorum & glauco colore, sed & floribus quintuplo minoribus, tantum ex quinqueflosculis compositis, cum antecedentes gaudeant pluribus.

alpina.

10. CACALIA foliis reniformibus acutis denticulatis.

Cacalia. Hall. belv. 705.

Tussilago caule ramoso. Roy. lugdb. 159. Sauv. monsp. 113.

11. *Cacalia foliis crassis hirsutis. Barb. pin. 197. Moris. hist. 3. p. 94. f. 7. t. 12. f. 1.*

Cacalia incano folio. Clus. hist. 2. p. 115.

12. *Cacalia foliis cutaneis acutioribus & glabris. Barb. pin. 198. Moris. hist. 3. p. 94. f. 7. t. 12. f. 6.*

Cacalia glabro folio. Clus. hist. 2. p. 115.

Cacalia. Lob. ic. 592.

Habitat in Alpibus Helvetiæ, Austriæ. &

Hanc speciem genere cum antecedentibus convenire docuit antopsis, hinc genere conjungenda; Cacalia cum Kleinii.

832

EUPATORIUM.

* *Calycibus quadrifloris.*

- hyssopifolia.* 1. EUPATORIUM foliis lanceolato-linearibus trinerviis integerrimis.

Eupatorium virginianum, folio angusto, floribus albis.

Dill. el. 141. t. 115. f. 140.

Eupatorium virginianum, flore albo, hyssopi foliorum æmulum. Moris. hist. 3. p. 98.

Eupatoria hirsuta, hyssopi foliorum æmula, virginiana.

Pluk. alm. 141. t. 88. f. 2. Raj. suppl. 189.

Habitat in Virginia. &

scandens.

2. EUPATORIUM caule volubili, foliis cordatis dentatis acutis. *Hort. cliff. 396. Hort. upf. 253. Roy.*

lugdb. 156. Gron. virg. 93. Cold. noveb. 182.

Clematidis novum genus, cucumeris folio, virginianum.

Pluk. alm. 109. t. 163. f. 3.

Habitat in aquosis virginiae. &

*consonia-
num.*

3. EUPATORIUM caule volubili, foliis ovatis integerrimis. *Hort. cliff. 396. **

Conyza americana scandens, foliis subrotundis mucronatis, floribus spicatis singulis quatuor flosculis constantibus. Amm. herb. 451.

Habitat in Vera Cruce.

• *Caly-*

SYNGENESIA: POLYGAMIA ÆQUALIS. 837

* *Calycibus quinquefloris.*

4. EUPATORIUM foliis ovato-hastatis petiolatis den- *zeylanicum.*
tatis. *Fl. zeyl.* 306. *
Eupatorium zeylanicum, foliis dentatis ad pediculos bre-
ves auriculatis. *Morif, hist.* 3. p. 99.
Cacalia foliis auriculatis ferratis supra nigricantibus, in-
fra lanugine alba obductis. *Burm. zeyl.* 52. t. 21.
Jacobca, senecionis flore, zeylanica odorata. *Raj. suppl.*
188.
Habitat in Zeylona.
5. EUPATORIUM foliis sessilibus amplexicaulibus dis- *sessilifolium*
tinctis lanceolatis. *Hort. upf.* 254. *
Eupatorium virginianum, flore albo, foliis menthæ an-
gustioribus sessilibus minutim dentatis. *Morif, hist.* 3. p.
98. *Raj. suppl.* 188.
Habitat in Virginia. &
6. EUPATORIUM foliis ovatis petiolatis ferratis. *chinense,*
Habitat in China. *Osbeck.*
Caulis *seres, brachiatus, nudiusculus.* Folia *opposita,*
ovata, acuminata, grosse serrata, petiolata, glabra,
sabtus pallidiora. Corymbi *terminales, flosculis quinis.*
7. EUPATORIUM foliis sessilibus distinctis subrotun- *rotundifo-*
do-cordatis. *linum,*
Eupatoria *valerianoides virginienfis, triflaginis folio abs-*
que pediculis. *Pluk. alm.* 141. t. 88. f. 4. *Raj. suppl.*
189.
Cacalia foliis rotundioribus ad caulem sessilibus. *Morif,*
hist. 3. p. 94.
Habitat in Virginia, Canada.
8. EUPATORIUM foliis lanceolatis nervosis: inferior- *altissimum.*
ribus extimo subserratis, caule suffruticoso. *Hort. upf.*
253.
Eupatorium virginianum, longissimis & angustissimis
foliis. *Morif. hist.* 3. p. 97. *Raj. suppl.* 187.
Habitat in Pennsylvania. &
9. EUPATORIUM foliis ternis.
Eupatorium caule erecto, foliis ovato-lanceolatis serratis *trifoliatum,*
petiolatis ternatis. *Gron. virg.* 178.
Eupatorium cannabinum, foliis in caule ad genicula ter-
nis, marilandicum. *Raj. suppl.* 189.
Habitat in Virginia.

838 SYNGENESIA: POLYGAMIA ÆQUALIS.

- cannabinum* 10. EUPATORIUM foliis digitatis. *Hort. cliff.* 396. *Fl. succ.* 665. *Mat. med.* 380. *Roy. lugdb.* 155.
Eupatorium cannabinum. *Bauh. pin.* 320.
Eupatorium adulterinum. *Fuchs. hist.* 265.
Habitat in Europa ad aquas.

* *Calycibus octofloris.*

- purpureum*. 11. EUPATORIUM foliis subverticillatis lanceolato-ovatis serratis petiolatis rugosis.
Eupatorium foliis verticillatis. *Cold. noveb.* 180.
*Eupatorium foliis ovato-lanceolatis obtuse serratis nup-
 tiolos desinentibus.* *Gron. virg.* 93.
Eupatorium enulæ folio. *Corn. canad.* 72. t. 72.
*Eupatorium canadense elatius, longioribus foliis rugosis
 integris & caulibus ferrugineis.* *Morif. hist.* 3. p. 97.
 f. 7. t. 13. f. 4.
 β. *Eupatorium foliis lanceolato-ovatis serratis petiolatis,
 caule erecto.* *Hort. cliff.* 396. *Roy. lugdb.* 155.
*Eupatorium novæ angliz, urticæ foliis, floribus purpu-
 rascentibus, caule maculato.* *Herm. par.* 158. t. 158.
Morif. hist. 3. p. 97. f. 7. t. 18. f. 3. *Raj. suppl.* 187.
Habitat in America septentrionali. 2
*Caulis teres, erectus, viridis, punctis linearibus longitu-
 dinalibus purpurascens. Folia terna, quaterna,
 f. sena, lato-lanceolata, lanceolato-ovata, serrata, rugo-
 sa, scabrinuscula, petiolata, utrinque viridia. Corym-
 bus terminalis. Calyces florum incarnati. Flosculi
 octo, Corollis albidis, Antheris purpureis, stylis lon-
 gissimis.*

* *Calycibus multifloris.*

- æstivum*. 12. EUPATORIUM foliis cordato-ovatis obtuse serratis petiolatis, calycibus multifloris.
Eupatorium foliis cordatis serratis petiolatis. *Gron. virg.* 94.
Eupatorium scordoniæ folio, flore cæruleo. *Dill. elib.* 140. t. 114. f. 139.
*Eupatorium marianum, scrophulariæ foliis, capitulis
 globosis colore cœlestino.* *Pluk. mant.* 71. t. 39.
 f. 4.
Habitat in Carolina, Virginia.

* *Calycibus quindecim flosculis.*

- perfoliatum*. 13. EUPATORIUM foliis connatis tomentosis. *Hort. cliff.* 396. *Hort. ups.* 253. *Roy. lugdb.* 156. *Gron. virg.* 177. *Cold. noveb.* 181.

Eup

SYNGENESIA: POLY GAMIA ÆQUALIS. 839

Eupatorium virginianum, salviæ foliis longissimis acuminatis, perfoliatum. *Pluk. alm.* 140. t. 87. f. 6. *bon. Raj. suppl.* 189.

Eupatorium virginianum, mucronatis rugosis & longissimis foliis, perfoliatum. *Morif. hist.* 3. p. 97.

Habitat in Virginix aquosis. 2

* *Calycibus multifloris: viginti circiter flosculis:*

14. **EUPATORIUM** foliis ovatis obtuse serratis petio-*aromaticum.*
latis trimerviis, calycibus simplicibus.

Eupatorium caule erecto ramoso, foliis ovatis obtuse serratis petiolatis. *Gron. virg.* 177.

Eupatoria valerianoides, flore niveo, teucris foliis cum pediculis, virginiana. *Pluk. alm.* 141. t. 88. f. 3. *Morif. hist.* 3. p. 98.

Habitat in Virginia. 2

Caulis teres, quadripedalis, strictus, ramis brachiatum, facie *Scutellaria*. Folia ovato-oblonga, obtuse serrata, trinervia, subrugosa, petiolata. Racemi terminales. Flores calyce duplo longiores, toti nivei flosculis 18 ad 28, stylis vix flosculo longioribus. Differt a reliquis imprimis calycibus non imbricatis sed ferme aequalibus & stylis non flore longioribus, uti *Ageratum*; sed huic pappus seminum verus.

AGERATUM. 835

1. **AGERATUM** foliis ovatis, caule piloso.

Ageratum. *Hort. cliff.* 396. *Hort. upf.* 253.

Eupatorium humile africanum, senecionis facie, folio lamii. *Herm. par.* 161, t. 161.

Habitat in America. ☉

2. **AGERATUM** foliis ovatis crenatis obtusis, caule ciliato, glabro.

Centaureum ciliare minus bisnagaricum, origani foliis amplioribus, floribus in umbellis. *Pluk. alm.* 93.

81. f. 4.

Habitat in Bisnagar.

3. **AGERATUM** foliis ovato-cordatis rugosis, floralibus alternis, caule glabro.

Eupatorium foliis ovatis serratis petiolatis, caule glabro. *Hort. upf.* 254.

Eupatorium caule erecto, foliis cordatis serratis *Hort. cliff.* 396. *Gron. virg.* 93.

Eupatorium scrophulariæ foliis glabris, flore albo *Morif. hist.* 3. p. 98 f. 7. t. 18. f. 11. Va.

840 SYNGENESIA: POLYGAMIA ÆQUALIS.

Valeriana urticae folio, flore albo. *Corn. canad.* 20. t. 21.
Habitat in Canada, Virginia. &

434

STÆHELINA.

gnaphaloides 1. STÆHELINA foliis tomentosis, squamis calycinis lanceolatis.

Gnaphalium foliis linearibus, floribus terminalibus, calycibus cylindraceis: squamis reflexis. *Roy. lugdb.* 151.
Jacea æthiopica, *stoechados citrinæ* majoribus tomentosis foliis, capitulorum spinis & squamulis ex aureo colore nitentibus. *Pluk. alm.* 193. t. 302. f. 3.
Habitat in Æthiopia.

centauroides 2. STÆHELINA foliis subtrigonis, squamis calycinis crenatis.

Santolina foliis linearibus, flore solitario terminali, squamis calycinis crenatis. *Hort. cliff.* 398. *
Habitat in Æthiopia.

3. STÆHELINA foliis linearibus denticulatis, squamis calycinis lanceolatis.

Gnaphalium caule fruticoso, foliis linearibus denticulatis, capitulis oblongis. *Roy. lugdb.* 151.
Stæchadi citrinæ affinis, capitulis longioribus. *Banh. hist.* 3. p. 156.

Elichrysum sylvestre, flore oblongo. *Banh. pin.* 265.
Jacea capitata, *rosmarini* folio. *Tournef. inst.* 444.
Habitat in Hispania, G. Narbonensi. b

Singularis planta quasi media inter Serratulas, Gnaphalia & Stæbelinas. Fructificatio in viva planta debet attentè examinari & genus certius determinari.

835

CHRYSOCOMA.

* *Frutescentes.*

Coma aurea. 1. CHRYSOCOMA fruticosa, foliis linearibus dorso decurrentibus. *Hort. cliff.* 397. *Hort. ups.* 252. *Roy. lugdb.* 146.

Coma aurea africana fruticans, foliis linearibus angustis, major. *Com. hort.* 2. p. 89. t. 45.

Gonyza æthiopica, flore bullato aureo, pinastri brevioribus foliis læte viridibus. *Pluk. alm.* 400. t. 327. f. 2.
Habitat in Æthiopia. b

crana.

2. CHRYSOCOMA subfruticosa, foliis linearibus subtus pilosis, floribus ante florescentiam cernuis. *Hort. cliff.* 397. * *Roy. lugdb.* 146.

Coma.

SYNGENESIA: POLYGAMIA ÆQUALIS. 341

Coma aurea, foliis linearis angustioribus, minor. *Comm. amst.* 2. p. 89. t. 45. f. 1.
Habitat in Æthiopia. 5

3. CHRYSOCOMA suffruticosa, foliis linearibus recurvis scabris ciliatis, floribus erectis.
 Conyza africana tenuifolia subfrutescens, flore aureo.
Dill. elsb. 104. t. 88. f. 13.
Habitat in Æthiopia. 5

* Herbaceæ.

4. CHRYSOCOMA herbacea, foliis linearibus glabris, Linosyris calycibus laxis.

Chrysocoma calycibus laxis. *Hort. cliff.* 396. *Roy. lugdb.* 145.

Chrysocome dioscoridis & plinii. *Col. ecphr.* 1. p. 81. t. 82.

Linosyris nuperorum. *Lob. hist.* 223.

Osyris austriaca. *Clus. hist.* 1. p. 325.

Linaria, folioso capitulo luteo, major. *Bauh. pin.* 213.

Habitat in Europa temperatiore. 2

5. CHRYSOCOMA herbacea paniculata, foliis lanceo-bistornatis trinerviis punctatis nudis.

Aster calycibus oblongis laxis, foliis ex lineari-lanceolatis integerrimis trinerviis: infra scrobiculis excavatis. *Gmel. fib.* 2. p. 189. t. 82. f. 1.

Conyza lini foliis asperis rigidis & nervosis, floribus luteis umbellatis. *Amm. ruth.* 192.

Habitat in Sibiria.

Flos variat cum & sine radio.

5. CHRYSOCOMA herbacea, foliis linearibus lanceo-graminifoliis latis glabris, floribus corymbosis.

Habitat in Canada. *Kalm.*

Habitus *Achilleæ* *Parmicæ* dictæ. Caulis levis, erectus, herbaceus, parum angulatus. Folia alterna, remotæ, lineari-lanceolata, acuminata, glabra, integerrima, margine scabro subcarinata, obsolete trinervia. Corymbus foliatus, ex floribus numerosissimis, sæpe 3 f. 4 simul in singulo pedicello, subsessilibus, magnitudine *Achilleæ*. Calyces semi-ovati, imbricati, glabri. Pappus simplex.

7. CHRYSOCOMA herbacea, foliis lanceolatis villosis, villosis calycibus arctis.

G g g s

Aster

842 SYNGENESIA: POLYGAMIA ÆQUALIS.

Aster incanus, calycibus laxis oblongis, foliolis ex lineari-lanceolatis, floribus nudis umbellatis. *Gmel. sib.* 2. p. 192. t. 82. f. 2.

Conyza tomentosa & *incana*, floribus luteis umbellatis, echii scorpioidis folio. *Amm. ruth.* 190.

Habitat in Sibiria, Tataria.

836

TARCHONANTHUS.

camphoratus 1. **TARCHONANTHUS.** *Hort. cliff.* 398. *Roy. lugdb.* 152.

Elichryso affinis arbor africana, flore purpureo-violaceo, folio salviae, odore rosmarini. *Herm. lugdb.* 228. t. 229.

Pluk. alm. 134. t. 174. f. 1. *Morif. hist.* 3. p. 90

Habitat in Æthiopia. b

837

SANTOLINA.

chamaCy- 1. **SANTOLINA** pedunculis unifloris, foliis quadrifariam
pariflor. dentatis. *Hort. cliff.* 397. *Hort. ups.* 252. *Mat. med.* 383. *Roj. lugdb.* 146.

Abrotanum femina, foliis teretibus. *Bauh. pin.* 136.

Abrotanum femina vulgare. *Clus. hist.* 1. p. 341.

Habitat in Europa australi. b

rosmarinifolia. 2. **SANTOLINA** pedunculis unifloris, foliis linearibus. *Hort. cliff.* 397. *Roy. lugdb.* 145.

Abrotanum femina, foliis rosmarini, majus. *Bauh. pin.*

137. *Morif. hist.* 3. p. 12. f. 6. t. 3. f. 22.

Abrotanum femina 4. *Clus. hist.* 1. p. 342.

β *Abrotanum* femina, foliis rosmarini, minus. *Bauh. pin.* 137.

γ *Abrotanum* femina viridis. *Bauh. pin.* 137.

δ *Abrotanum* femina, flore majore. *Bauh. pin.* 137. &

Habitat in Hispania.

crenata. 3. **SANTOLINA** flore solitario terminali, squamis calycinis crenatis, foliis linearibus. *Hort. cliff.* 398. *

Habitat in Æthiopia.

stans 4. **SANTOLINA** corymbis simplicibus coarctatis, foliis pinnatifidis dentatis.

Elichrysum inodorum glabrum, coronopi folio, annuum magnoli. *Herm. lugdb.* 228. t. 227.

Helichrysum inodorum glabrum, coronopi folio, annuum. *Magn. monsp.* 307. *

Chrysanthemum corymbiferum. *Triumph. obs.* 85. t. 86.

Bel-

SYNGENESIA POLYGAMIA ÆQUALIS. 843

Bellis polyclonos annua africana, coronopi folio. floribus nudis compactis. *Morif. hist.* 3. p. 30.

Habitat in Africa ☉

5. SANTOLINA corymbis compositis fastigiatis, foliis *dentatis*. inferioribus linearibus dentatis, superioribus ovatis serratis. *Hort. cliff.* 398. *Roy. lugdb.* 147.

Coma aurea africana frutescens, foliis inferioribus incis, superioribus dentatis. *Comm. rar.* 41. t. 41.

Habitat in Æthiopia.

6. SANTOLINA corymbis simplicibus fastigiatis, foliis *trifurcata*. is trilobis cuneiformibus.

Santolina corymbo simplici terminali, foliis trifidis.

Hort. cliff. 397. *Roy. lugdb.* 146

Coma aurea africana fruticans, foliis glaucis & in extremitate trifidis. *Comm. hort.* 2. p. 97. t. 49.

Habitat in Æthiopia.

7. SANTOLINA corymbis simplicibus fastigiatis, foliis *crithmifolia*. semitrifidis linearibus.

Santolina corymbo simplici, foliis trifidis. *Hort. ups.* 252.

Coma aurea africana fruticans, foliis crithmi marini.

Comm. hort. 2. p. 99. t. 50.

Jacobææ æthiopica, foliis abrotani trifidis, summo caule capitulis parvis glomeratis. *Pluk. alm.* 194. t. 302. f. 7.

Habitat in Æthiopia.

Præcedenti nimis affinis.

TANACETUM. 888

* *Herbacea.*

1. TANACETUM foliis pinnatis: pinnis linearibus *re-crithmifolium*. motis integerrimis.

Santolina corymbo composito terminali, foliis linearibus dentatis: dentibus longitudine folii. *Hort. cliff.* 397. *Roy. lugdb.* 146.

Elichrysium africanum frutescens, foliis crithmi marini.

Comm. hort. 2. p. 113. t. 57.

Ageratum capense, crithmi folio, capitulis parvis. *Pet. gaz.* t. 34. f. 1.

Habitat in Æthiopia.

2. TANACETUM foliis pinnato-multifidis: laciniis linearibus subdivisis acutis. *Hort. cliff.* 398. *Roy. lugdb.* *sum.*

844 SYNGENESIA: POLYGAMIA ÆQUALIS.

Abrotanum africanum fruticans multiflorum, foliis tanacetii decuplo-minoribus. *Comm. hort.* 2. p. 199. t. 100.

Habitat in Æthiopia. 5

* *Fruticosa.*

frutescens. 3. TANACETUM foliis pinnatifidis: laciniis lanceolatis obtusiusculis integerrimis.

Tanacetum foliis pinnatis integerrimis. *Hort. cliff.* 398.

Roy. lugdb. 154.

Tanacetum africanum arborescens, foliis lavendulae multifido folio. *Comm. hort.* 2. p. 201. t. 101.

Habitat in Æthiopia. 5

Sibiricum. 4. TANACETUM foliis pinnatis: laciniis lineari-filiformibus, corymbis glabris, caule herbaceo.

Tanacetum foliis pinnatis multifidis: laciniis linearibus trifidis. *Gmel. sib.* 2. p. 134. t. 65. f. 2.

Habitat in Sibiria.

Folia linearia, pinnata: pinnis linearibus, filiformibus, saepe bifidis trifidisve, integerrimis. Corymbus terminalis & axillares pauciflori floribus luteis.

incanum. 5. TANACETUM foliis bipinnatis tomentosis, corymbo ovato composito.

Artemisia caule simplici, corymbo composito ovato terminali. *Hort. cliff.* 404. *

Absinthium orientale incanum tenuifolium, floribus luteis in capitulum congestis & sursum spectantibus.

Tournef. cor. 34.

Habitat in Oriente. 2

annuum. 6. TANACETUM foliis bipinnatifidis linearibus acutis, corymbis tomentosis. *Roy. lugdb.* 155.

Tanacetum annuum villosum, absinthii austriaci folio.

Vaill. alt. 1719. p. 281.

Absinthium corymbiferum annuum. *Tournef. inst.* 458.

Elichrysum foliis abrotani. *Bauh. pin.* 264.

Elichryson. *Clas. hist.* 1. p. 325. t. 326.

Habitat in Hispania, Hetruria. 6

Rami longiores quam umbella primaria.

vulgare. 7. TANACETUM foliis bipinnatis incisiss serratis. *Hort. cliff.* 398. *Fl. succ.* 666. *Mat. med.* 381. *Roy. lugdb.* 155. *Gmel. sib.* 2. p. 133. t. 65. f. 1.

Tanacetum foliis pinnatis planis: pinnis serratis. *Fl. lapp* 295.

Ta-

SYNGENESIA: POLYGAM. SUPERFLUA. 845

- Tanacetum vulgare luteum. *Bauh. pin.* 132.
 Athanasia f. Tanacetum. *Dalech. hist.* 955.
 Artemisia inonoelonos. *Fuchf. hist.* 46.
 β. Tanacetum foliis crispis. *Bauh. pin.* 132. crispum.
 Tanacetum crispum. *Dod. pempt.* 36.
Habitat in Europæ aggeribus. 2
8. TANACETUM foliis ovatis integris serratis. *Hort. Balsamit. cliff.* 398. *Hort. ups.* 255. *Mat. med.* 382. *Roy. lugdb.* 154.
 Mentha hortensis corymbifera. *Bauh. pin.* 226.
 Costus hortensis. *Dalech. hist.* 678.
Habitat in Hetruria, Narbona. 2

POLYGAMIA-SUPERFLUA.

ARTEMISIA. § 39

* Fruticosa erecta.

1. ARTEMISIA foliis linearibus confertis minimis di-*arbiopica*,
 visis, caule fruticoso tomentoso.
 Abrotanoides africanum, foliis cinereis muscosis, ca-
 pitulis florum globosis magnis. *Roy. suppl.* 232.
 Absinthium africanum arborescens, folio vermicula-
 tæ incano. *Tournesf. inst.* 358.
Habitat in Æthiopia. 3
 Caulis fruticosus, paniculatus, albus. Folia Erica ma-
 gnitudine, minima, coacervata. Flores in racemis
 subrotundis, magnitudine A. Absinthii.
2. ARTEMISIA foliis ramosissimis setaceis, caule ere-*Abrotanum*.
 cto suffruticoso. *Hort. cliff.* 403. *Hort. ups.* 257.
Mat. med. 385.
 Abrotanum mas angustifolium majus. *Bauh. pin.* 136.
 Abrotanum mas. *Dod. pempt.* 21.
*Habitat in Syriæ, Galatiæ, Cappadociæ montibus a-
 pricis. 3*
- * Procumbentes ante florescentiam.
3. ARTEMISIA foliis caulinis linearibus pinnato-mul-*Santonium*.
 tificis, ramis indivisis, spicis secundis reflexis.
 Artemisia fruticosa incana ramosissima, corymbis sessi-
 libus spicatis subrotundis, foliis superioribus lineari-
 bus brevissimis obtusiusculis. *Gmel. sib.* 2. p. 115. t.
 51.
 Semen sanctum. *Lob. ic.* 756.

Habi-

Habitat in Tataria, Persia. &c.

Habitus A. campestris, sed erectus. Caulis paniculatus, subcanus. Folia inferiora pinnato-multifida, linearia, canescentia. Rami virgati, Racemis s. Spicis alternis, secundis, recurvatis, Pedicellis adspersis foliis linearibus, obtusiusculis, recurvis. Flores solitarii, cylindrici, in Planta fructificante omnes Canes erecti, tum amifere incanum colorem. Folia in ramis minima, linearia, indivisa.

- campestris.* 4. ARTEMISIA foliis multifidis linearibus, caulibus procumbentibus virgatis. *Hort. cliff.* 403. *Fl. jucc.* 668. *Roy. lugdb.* 154. *Gmel. sib.* 2. p. 117.

Abrotanum campestre. *Bauh. pin.* 136.

Ambrosia altera. *Cam. epit.* 597.

Habitat in Europæ campis apricis, aridis. &

- palustris.* 5. ARTEMISIA foliis linearibus pinnatis integerrimis, floribus glomeratis subsessilibus.
Artemisia herbacea, foliis multifidis linearibus, corymbis in spica interrupta glomeratis sessilibus. *Gmel. sib.* 2. p. 119. t. 55.

Absinthium capillaceis foliis, flosculis aureis densissime congestis. *Amm. rub.* 197.

Habitat in Sibiria.

Caules procumbentes, adscendentes. Folia Coronopi, linearia, pinnatifida: segmentis 5. s. 7. Flores glomerati, subsessiles, flavi.

- crabrofolia.* 6. ARTEMISIA foliis compositis divaricatis linearibus carnosis glabris, caule adscendente paniculato.

Abrotanum hispanicum maritimum, folio crasso splendente & rigido. *Tournef. infl.* 459.

Absinthium inodorum, foliis crithmi lucidis obscure virentibus. *Morif. hist.* 3. p. 11. *Raj. suppl.* 232.

Habitat in Lusitaniæ litoribus arenosis. *Læssing.*

Habitus A. campestris, glaber. Canles semipedales, s. pedates, adscendentes, Paniculae florum terminati. Folia pinnatifida trifidaque, linearia, glabra, carnosae. Flores A. campestris.

- maritima.* 7. ARTEMISIA foliis multipartitis tomentosis, racemis cernuis, flosculis femineis ternis.

Artemisia foliis compositis multifidis tomentosis, ramis floriferis nutantibus. *lt. xl.* 112. " *Fl. jucc.* 671.

Arte-

SYNGENESIA: POLYGAM. SUPERFLUA. 847

Artemisia foliis compositis multifidis, racemis simplicibus alternis. floribus erectis sessilibus. Roy lugdb. 113. Sauv. monsp. 264.

Abinthium seriphium belgicum. Baub. pin. 139. Baub. hist. 3. p. 178.

β. *Abinthium seriphium germanicum. Baub. pin. 139. Habitat in Europæ septentrionalioris littoribus maris. &*

ARTEMISIA foliis pinnatis, caulibus adscendentibus *rupifris*, hirsutis, floribus globosis cernuis: receptaculo papposo. *it. gosl. 285. t. 285. * Fl. succ. 669. **

Artemisia foliis pinnatis pilosis, ramis adscendentibus, floribus oblongis suberectis. Gmel. lib. 2. p. 111. t. 49. t. 1.

Abinthium pumilum repens & supinum. C. B. Pinck. alm. 3. t. 37. f. 1.

Abinthium pumilum palmeum minus, argenteo sericeoque folio Bocc. mus. 2. p. 81. t. 71

Abinthium alpinum incanum. Baub. pin. 139. prodr. 71²

β. *Abinthium viride, foliis multifidis linearibus. Gmel. lib. 2. p. 131. t. 64. f. 2.*

Habitat in Sibiria, Oelandiæ rupibus calcareis.

Differt C. B. (ex Bursero) planta ramis longioribus, aliisque forte ex loco notis; Sibirica magis fruticosa est. Planta in rupibus alba sericea. In horto major, atro-viridis, nuda.

** Erecta herbaceæ, foliis compositis.*

γ. **ARTEMISIA** foliis multipartitis subtus tomentosis, *pontica*, floribus subrotundis nutantibus: receptaculo nudo. *Hort. upf. 257. Hort. cliff. 404. Mat. med. 387.*

Abinthium ponticum tenuifolium incanum. Baub. pin. 138.

Abinthium tenuifolium. Dod. pempt. 24.

Habitat in Hungaria interiore, Pannonia, Thracia, Myfia. &

10. **ARTEMISIA** foliis triplicato-pinnatis utrinque glabris, *annua*, floribus subglobosis nutantibus: receptaculo glabro conico *Hort. upf. 257.*

Artemisia, radice annua, viridis foliis pinnatis: pinnis pinnatifidis, calycibus subrotundis fere erectis ex alis copiosissimis. Gmel. lib. 2. p. 125.

Abinthium, tanacetii folio, odoratissimum. Amm. raph. 142. t. 193. f. 13. imperfecte.

β Ab-

848 SYNGENESIA: POLYGAM. SUPERFLUA.

β. Absinthium, folio tenui multifido, camphoram spirans.

Boerb. lugdb. 1. p. 126.

Habitat in Sibiria montosis. ☉

tanacetifolia 11. ARTEMISIA foliis bipinnatis subtus tomentosis nitidis, pinnis transversis, racemis simplicibus.

Artemisia radice perenni, foliis pinnatis pinnatifidis incisis serratis, calycibus subrotundis viridibus nutantibus. *Gmel. fib. 2. p. 123. t. 58.*

Habitat in Sibiria. &

Absinthium. 12. ARTEMISIA foliis compositis multifidis, floribus subglobois pendulis: receptaculo villosa. *Hort. cliff 404. Fl. suec. 670, Mat. med. 386. Roy. lugdb. 153*

Absinthium incanum, foliis compositis latiuscule multifidis, floribus subglobois pendulis. *Gmel. fib. 2. p. 129. t. 63.*

Absinthium ponticum f. romanum, officinarum f. discoridis. *Bauh. pin. 138.*

Absinthium. *Cam. epit. 452.*

Habitat in Europæ ruderatis aridis. &

vulgaris. 13 ARTEMISIA foliis pinnatifidis planis incisis: subtus tomentosis, racemis simplicibus, floribus ovatis: radio quinquefloro.

Artemisia foliis pinnatifidis planis laciniatis, floribus erectis. *Hort. cliff. 404. Fl. suec. 667. Mat. med. 384.*

Roy. lugdb. 153. Gmel. fib. 2. p. 108.

Artemisia vulgaris major. *Bauh. pin. 137. Fl. lapp. 379.*

Artemisia vulgaris. *Bauh. hist. 3. p. 184.*

Habitat in Europæ cultis, ruderatis. &

** Foliis simplicibus.*

integrifolia. 14. ARTEMISIA foliis lanceolatis subtus tomentosis integerrimis dentatisque.

Artemisia foliis planis lanceolato-linearibus: inferioribus sæpe ex pinnato-dentatis. *Gmel. fib. 2. p. 109. t. 48. f. 1.*

Habitat in Sibiria.

Habitus absolute Artemisia vulgaris; at Folia lanceolata, rigidiuscula, glabra, subtus cano tomentosa, margine integerrima, rarius uno alterove dente erecto notata. Racemi axillares, breves, terminales, glomerati. Flores rotundiusculi: flosculi feminei sæpius quinque.

carulescens. 15. ARTEMISIA foliis lanceolatis integris: lateralibus sæpe palmatis. *Hort. cliff. 403.*

Ab-

SYNGENESIA: POLYGAM. SUPERFLUA. 849

Absinthium maritimum, lavendulæ folio. *Bauh. pin.* 139.

Morij. hist. 3. p. 7. f. 6. t. 1. f. 5.

Absinthium angustifolium. *Dod. pempt.* 26.

β. *Absinthium maritimum*, foliis superioribus in aliquot laciniis divis. *Bauh. pin.* 139.

Abrotanum latifolium rarius, artemisiæ folio. *Col. ecphr.*

2. p. 75. t. 76.

Habitat in Europæ australis littoribus maris.

Folia utrinque villosa. Flores cylindrici, mutantes.

16. ARTEMISIA foliis lanceolatis glabris integerrimis. *Dracunculus*

Hort. cliff. 403. *Hort. upf.* 256. *Gmel. sib.* 2. p. 126.

1. 59. & 60. f. 1.

Abrotanum lini folio acriori & odorato. *Tournef. inst.*

459.

Dracunculus hortensis. *Bauh pin.* 93.

Draco herba. *Dod. pempt.* 709.

Habitat in Sibiria, Tataria. 2

17. ARTEMISIA foliis simplicibus tomentosis obtusis chinensis.

lanceolatis: inferioribus cuneiformibus trilobis.

Artemisia foliis radicalibus a caulinis diversis. *Gmel. sib.*

2. p. 61. f. 1. 2.

Habitat in China, Lagerström. Sibiria

Folia tomentosa, pollicem transversum longa: inferiora cuneiformia, triloba; superiora lanceolata, obtusa. Flores in summitate conferti, in Racemum sæpius ovatum, simplicem, subrotundum, Calyce laxo, tomentoso.

18. ARTEMISIA foliis simplicibus lyrato-sinuatis cau- *maderaspata-*

libus procumbentibus, floribus pedunculatis solitariis globo- *tana.*

globosis oppositifoliis.

Absinthium maderaspatanum. *Pluk. phys.* 357. f. 3.

Absinthium minus odoratum gangeticum, floribus chamæmeli solitariis e foliorum alis. *Pluk. alm.* 2. t. 1.

f. 2. mal.

Habitat in India. 2

Herba procumbens, terræ adpressa, Ramis alternis, teretibus, flexuosis, striatis, pubescentibus. Folia pinnato f. densato-sinuata, lyrata, extrorsum latiora, mollia. Pedunculi solitarii, folio oppositi, uniflori, nudi, striati, villosi. Flores subglobosi disco convexo, magni, lutei. Flosculi marginales feminei trifidi, stylo bifido tenuiore; discoides hermaphroditi: stylo simplici.

19. ARTEMISIA foliis cuneiformibus repandis, caule *minima.*

procumbente, floribus axillaribus sessilibus.

H h h

Habi-

850 SYNGENESIA: POLYGAM. SUPERFLUA.

Habitat in China. Lagerström. ☉

Planta omnium minima, Folia glabra. Flores ex 6. f. 7. flosculis, minimi, sessiles, axillares, solitarii.

840

GNAPHALIUM.

* *Argyrocomæ calycibus albis.*

- dicium.* 1. GNAPHALIUM caule simplicissimo, corymbo simplici terminali, sarmentis procumbentibus. *Hort. cliff. 400. Fl. succ. 672. Mat. med. 388. Roy. lugdb. 147. Gmel. fib. 2. p. 105.*
Gnaphalium caule simplicissimo, floribus coloratis terminato. Fl. lapp. 672.
Gnaphalium montanum, flore rotundiore. Banb. pin. 263.
Pilosella minor. Dod. pempt. 68. f. exterior.
- femina.* 2. GNAPHALIUM montanum longiore & follo & flore. *Banb. pin. 263.*
Pilosella minor. Dod. pempt. 68. f. interiori
Habitat in Europæ apricis aridis. &
- plantagini-folium.* 2. GNAPHALIUM caule simplicissimo, foliis radicalibus ovatis maximis, sarmentis procumbentibus.
Gnaphalium stolonibus reptatricibus longissimis, foliis ovatis, caule capitato. Gron. virg. 95.
Gnaphalium, plantaginis folio, virginianum. Pluk. alm. 171. t. 348. f. 9.
Habitat in Virginia. &
Habitus omnino precedentis; sed Folia radicalia pollice majora, ovalia. Vidi solam feminam; an precedentis sola varietas?
- margaritaceum.* GNAPHALIUM foliis lineari-lanceolatis acuminatis alternis, caule superne ramoso, corymbis fastigiatis.
Hort. cliff. 401. Hort. apf. 255. Gmel. fib. 2. p. 107.
Gnaphalium latifolium americanum. Banb. pin. 263.
Gnaphalium americanum. Clus. hist. 1. p. 327.
Habitat in America septentrionali, kamtschatca. &
- grandiflorum.* 4. GNAPHALIUM foliis amplexicaulis ovatis nervosis utrinque lanuginosis.
Gnaphalium tomentosum, foliis inferioribus subrotundis; superioribus acuminatis. Burm. afr. 213. t. 76. f. 1.
Elichrysium africanum lanuginosum latifolium, calyce floris argenteo & amplissimo. Old. afr. 27.
Habitat in Æthiopia.

5. GNA.

SYNGENESIA: POLYGAM. SUPERFLUA. 851

5. **GNAPHALIUM** foliis amplexicaulibus integerrimis *foetidum*, acutis subtus tomentosis, caule ramoso. *Hort. cliff.* 402. *Hort. upf.* 256.
Gnaphalium atricaum latifolium foetidum, capitulo argenteo. *Comm. hort.* 2. p. 111. f. 56.
Conyza africana graveolens, capitulis argenteis. *Pluk. alm.* 117. t. 243. f. 1. *Morif. bist.* 3. p. 115 f. 7. s. 20. f. 32.
Habitat in Æthiopia. ☉
6. **GNAPHALIUM** foliis amplexicaulibus lanceolatis, *heliantemifolium*, corymbis compositis, caule ramosissimo.
Gnaphalium africanum, floribus minimis albicantibus. *Volk. herb.* 194. s. 94.
Habitat Æthiopia. 2
7. **GNAPHALIUM** foliis semiamplexicaulibus ensiformibus repandis obtusis utrinque pubescentibus, floribus conglomeratis. *Roy. lugdb.* 149.
Elichrysum sylvestre latifolium, capitulis conglobatis. *Bauh. pin.* 264.
Gnaphalium majus, lato oblongo folio. *Bauh. pin.* 263. *Pluk. alm.* 171. t. 31. f. 16.
Gnaphalium ad stœchadem citrinam accedens. *Bauh. bist.* 3. p. 160.
Habitat in Helvetia, G. Narbonensi, Hispania, Lusitania. ☉
Planta valde lanata. Calyces flavo-albicantes, molles, squamis ovato-lanceolatis.
8. **GNAPHALIUM** foliis lanceolatis, caule tomento- *obtusifolium*, so ramoso, floribus terminalibus glomeratis conicis.
Gnaphalium foliis lanceolatis, caule tomentoso, corymbis supradecompositis, floribus sessilibus confertis. *Gren. virg.* 95.
Elichrysum obtusifolium, capitulis argenteis conglobatis. *Dill. el.* 130. t. 108. f. 131.
Helichrysum Chrysocoma Gnaphaloides virginiana annua, foliis obtusioribus, capitulis argenteis conglobatis. *Morif. bist.* 3. p. 88. f. 7. s. 10. f. 19.
Habitat in Virginia, Pennsylvania. ☉
Caulis tomentosus, pilosus, ramosus, erectus. Folia lanceolata, recurvata, nudiuscula. Flores terminales, conglobati, inaequales, subsessiles. Calyces conici, albi acuti. Corollæ flava.

852 SYNGENESIA: POLYGAM. SUPERFLUA

- indalatum*. 9. **GNAPHALIUM** foliis decurrentibus acutis undatis
subtus tomentosis, caule ramoso. *Hort. cliff.* 402.
Elichrysum graveolens acutifolium, caule alato. *Dill.*
chb. 130. t. 108. f. 130.
Habitat in Africa. ☉
- indicum*. 10. **GNAPHALIUM** foliis lanceolatis alternis, caule
herbaceo ramosissimo diffuso, corymbis inæqualibus.
Fl. zeyl. 307. *
Gnaphalium, latiore folio, minus ramosum, capitulis pal-
lescentibus. *Pluk. alm.* 172. t. 187. f. 5.
Habitat in India.
- ulvum*. 11. **GNAPHALIUM** foliis filiformibus tomentosis sub-
imbricatis, caule ramosissimo fruticoso.
Gnaphalium incanum, folio lineari, caule acumbente.
Burm afr. 215. t. 77. f. 1.
Elichrysum æthiopicum, tenuissimis stœchadis citrinæ
foliis confertis, ramosissimum, comis argenteis.
Pluk. mant. 67. *Raj. suppl.* 171.
Habitat in Æthiopia.
- maritimum*. 12. **GNAPHALIUM** fruticosum, calycibus coronatis,
foliis revolutis obliquis mucronatis.
Gnaphalium caule fruticoso, foliis lanceolatis contor-
tis fasciculatis, calycibus cylindricis longis. *Hort.*
cliff. 403.
Gnaphalium fruticosum, foliis lanceolatis congestis, sto-
rum calyce tubuloso. *Burm. afr.* 221. t. 79. f. 1.
Argyrocome capitis b. spei, thymi foliis. *Pet. gaz.* t. 7.
f. 3.
Xeranthemum africanum frutescens, folio ericæ s. ju-
niperi angustiore brevior acuto, floribus congloba-
tis exiguis albis. *Raj. suppl.* 182.
Xeranthemum stœchadis citrinæ foliis rigidis pungen-
tibus, flore albo duriusculo. *Pluk. mant.* 190.
Gnaphaloides æthiopicum frutescens, stœchadis citrinæ
foliis spinosis. *Pluk. alm.* 92. *Raj. suppl.* 193.
Eupatorioides capitis b. spei, fatnrejæ foliis rigidis.
Pet. cent. 208. *Raj. suppl.* 239.
Lychnis coronaria monomotapensis globulifera, foliis
ex albidis genistæ aculeatæ. *Pluk. amatib.* 134. t. 406.
f. 6.
13. *Gnaphalium* fruticosum, foliis rarioribus, capitatum.
Burm. afr. 223. t. 79. f. 2.
Xeranthemum africanum, foliis brevioribus canescen-
ti-

SYNGENESIA: POLYGAM. SUPERFLUA. 853

tibus, floribus minimis conglobatis albis. *Raj. suppl.* 182.

Eupatorioides capensis capitatus. *Pet. gaz.* t. 8. f. 1. *Raj. suppl.* 239.

Frutex æthiopicus cineraceus, foliis nepetæ aculeatis, floribus albis coronatus. *Pluk. phyt.* 410. f. 2.

7. Gnaphalium frutescens, foliolis lanceolatis æqualibus, umbellatum. *Burm. afr.* 223. t. 79. f. 3.

Elichrysum æthiopicum frutescens, coridis foliis incanis, capitulis parvis glomeratis. *Pluk. alm.* 72.

Habitat in Æthiopia. 5

* *Chrysocomæ calycibus luteis.*

13. GNAPHALIUM foliis confertis angusto-lanceola *orientale.* tis, caule fruticoso; corymbo composito. *Hort. cliff.* 402. *Hort. upf.* 256.

Elichrysum orientale. *Baub. pin.* 264. *prodr.* 123.

Elichrysum africanum frutescens, angustis & longioribus foliis incanis. *Comm. hort.* 2. p. 109. t. 55.

Helichrysum frutescens latifolium, flore corymbifero toto aureo. *Morif. hist.* 3. p. 86. f. 7. t. 10. f. ult.

Habitat in Africa. 2-5.

14. GNAPHALIUM foliis linearibus, caule fruticoso *Stæchas.* ramofo, corymbo composito. *Hort. cliff.* 401. *Hort. upf.* 256

Elichrysum f. *Stæchas citrina angustifolia.* *Baub. pin.* 264.

Helichrysum f. *Chrysocome angustifolia vulgaris.* *Morif. hist.* 3. p. 401. f. 7. t. 11. f. 7.

Chrysocome f. *Stæchas citrina minor.* *Barr. ic.* 410. 409. 278.

Stæchas citrina. *Dod. pempt.* 268.

Habitat in Germania, Gallia, Hispania 5

15. GNAPHALIUM foliis linearibus alternis, caule frutescente, floribus paniculatis subternis terminalibus sessilibus. *Hort. cliff.* 401. *

Elichrysum sylvestre angustifolium, capitulis conglobatis. *Baub. pin.* 264.

Stæchadi citrinæ affinis, capitulis parvis raris. *Baub. hist.* 3. p. 156. *Raj. hist.* 283.

Stæchas citrina spuria, longioribus foliis. *Barr. ic.* 368. 277.

Ageratum aliud quorundam. *Dalech. hist.* 778.

Habitat in G. Narbonensi. 5

H h h 3

16. GNA-

854 SYNGENESIA: POLYGAM. SUPERFLUA.

- teretifolium.* 16. **GNAPHALIUM** caule ramoso fruticoso, corymbis ramosis, foliis confertis teretiusculis. *Hort. cliff.* 401. *
- Grnaphallum frutescens*, foliis tenuissimis teretibus, ramis creberrimis. *Burm. afr.* 217. t. 77. f. 3.
- Millefolium æthiopicum*, ericæ foliis, ineanum, flore specioso. *Pluk. alm.* 251. t. 308. f. 2.
- Habitat in Æthiopia.* 5
- petiolatum.* 17. **GNAPHALIUM** caule fruticoso, foliis ovatis integerrimis petiolatis, floribus confertis terminalibus. *Hort. cliff.* 402. *
- Grnaphallum sericeum tomentosum*, foliis orbiculatis subtus incanis. *Burm. afr.* 214. t. 76. f. 2.
- Habitat in Æthiopia.* 5
- Annatum.* 18. **GNAPHALIUM** foliis cuneiformibus dentatis, caule fruticoso, corymbo simplici. *Roy. lugdb.* 151. *
- Coma aurea incana*, foliis obtusis tridentatis, capitulis oblongis. *Burm. afric.* 185. t. 68. f. 3.
- Habitat in Æthiopia.* 5
- arenarium.* 19. **GNAPHALIUM** foliis lanceolatis, inferioribus obtusis, caule herbaceo simplicissimo, corymbo composito. *Fl. suec.* 674. *Mat. med.* 389. *Gmel. fib.* 2 p. 107.
- Elichrysum* f. *Stoechas citrina latifolia*. *Baub. pin.* 264. *It. æl.* 160.
- Stoechas citrina germanica*, latiore folio. *Baub. hist.* 3 p. 153. *Raj. hist.* 281.
- Habitat in Europæ campis arenosis.* ☉
- ratilens.* 20. **GNAPHALIUM** foliis lineari-lanceolatis, caule inferne ramoso, corymbo decomposito terminali. *Hort. cliff.* 401. *
- Elichrysum africanum*, folio oblongo angusto, flore rubello postea aureo. *B. Dill. elstb.* 127. t. 107. f. 127.
- Habitat in Africa.* 2
- igneus.* 21. **GNAPHALIUM** foliis linearibus tomentosis integerrimis sessilibus, corymbis alternis conglobatis, floribus globosis. *Roy. lugdb.* 149.
- Elichrysum flore suaverubentre*. *Baerb. lugdb.* 1. p. 120.
- Habitat* - - -
- guthnium.* 22. **GNAPHALIUM** foliis lanceolatis nudis, caule erecto simplicissimo, floribus spicatis lateralibusque sessilibus. *Roy. lugdb.* 148. *Gron. virg.* 178.

Ell-

SYNGENESIA: POLYGAM. SUPERFLUA. 855

Elichrysium spicatum obtusifolium, basi angustiore. *Dill. elsb.* 131. t. 109. f. 132.

Gnaphalium spicatum majus non ramosum erectum virginianum, foliis obtusioribus. *Morif. hist.* 3. p. 92.

Habitat in Carolina, Virginia, Pennsylvania. ☉

23. GNAPHALIUM foliis lanceolatis semianplexicaulibus, caule inferne ramoso terminali. *Hort. cliff.* 401. *

Elichrysium africanum, folio longo subtus cano supra viridi, flore luteo. *B Dill. elsb.* 128. t. 107. f. 128.

Habitat in Africa. 2

24. GNAPHALIUM foliis lanceolatis tomentosis, caule ramoso, squamis calycinis reflexis. 200

Gnaphalium incanum angustifolium, calycis, squamis ferrugineis reflexis. *Burm. afr.* 226. t. 80. f. 2.

Habitat in Æthiopia.

25. GNAPHALIUM foliis oblongis tomentosis sessilibus, capitulis foliosis villosissimis. *Roy. lugdb.* 150.

Gnaphalium, latiore folio, æthiopicum, flore roseo, calyculis spinosis. *Pluk. alm.* 171. t. 323. f. 1.

Habitat in Æthiopia.

26. GNAPHALIUM foliis amplexicaulibus spatulatis, ramis patentibus, corymbis aggregatis. *Hort. cliff.* 402.

n. 15. *

Elichrysium africanum, folio oblongo tomentoso caulem amplexente, flore luteo. *Boerb. lugdb.* 1. p. 121.

Habitat in Æthiopia.

27. GNAPHALIUM foliis decurrentibus obtusis mucronatis utrinque tomentosis planis. *odoratissimum,*

Elichrysium latifolium villosum, alato caule, odoratissimum. *Pluk. alm.* 134. t. 173. f. 6.

Habitat - - -

* *Filagine calycibus cinereis.*

28. GNAPHALIUM caule simplicissimo, capitulo terminali, bracteis hirsutissimis radiato. *Leontopodium.*

Gnaphalium alpinum, magno flore, folio oblongo. *Baub. pin.* 264.

Gnaphalium alpinum pulchrum. *Baub. hist.* 3. p. 161.

Gnaphalium alpinum. *Clas. hist.* 1. p. 328.

Leontopodium. *Dod. pempt.* 68.

- β. *Gnaphalium* magno flore, folio brevi. *Baub. pin.* 264.

H h h 4

Habi-

Habitat in Alpibus Helvetiæ, Valesiæ, Corinchi, Austriæ, Sibiria. &

- alpinum.* - 29. GNAPHALIUM caule simplicissimo, capitulo terminali aphylo, floribus oblongis. *Fl. lapp.* 301. *Fl. suec.* 673.

Gnaphalium latiori folio, caule non ramoso, spica nuda fusca terminato. *Hall. helv.* 701.

Gnaphalium alpinum minus. *Baub. pin.* 264.

Filago alpina minor erecta. *Schenck. alp.* 133.

Habitat in Alpibus Lapponiæ, Helvetiæ. &

Caulis simplicissimus, digitis brevior, foliis 3 s. 4 lanceolatis. Folia radicalia lanceolato-cuneiformia, supra viridia lævia, subtus tomento striata albida, ut margo etiam a latere superiore albus conspiciatur. Ad radicem sarmenta ut G. dioico. Flores terminales, paniculi, conferti, absque foliis Calyx cylindricus, basi ventricosus, squamis apice membranaceis, lanceolatis, acutis, erectis. Flosculi seminei, in planta a me culta, absque staminibus. an G. dioici varietas obsexum, sarmenta, caulem nudum.

- uliginosum.* 30. GNAPHALIUM caule ramoso diffuso, floribus confertis terminalibus. *Fl. lapp.* 300. *Fl. suec.* 676.

Hort. cliff. 402. *Roy. lugdb.* 142. *Gmel. sib.* 2. p. 105.

Gnaphalium longifolium humile ramosum, capitulis nigris. *Raj. hist.* 295. *angl.* 3. p. 16.

Gnaphalium annuum serotinum, capitulis nigricantibus, in humidis gaudens. *Morif. hist.* 3. p. 92.

Habitat in Europæ paludibus, ubi aque stagnans. ☉

- sylvaticum.* 31. GNAPHALIUM caule simplicissimo, floribus sparsis. *Fl. lapp.* 298. *Fl. suec.* 675. *Hort. cliff.* 402. *Roy. lugdb.* 148. *Gmel. sib.* 2. p. 106.

Gnaphalium minus, angusto oblongo folio, alterum. *Baub. pin.* 263.

Gnaphalium rectum. *Baub. hist.* 3. p. 160.

Habitat in Europæ sylvis arenosis. ♂

- arvense.* 32. GNAPHALIUM paniculatum hirsutum. *Fl. suec.* 679.

Gnaphalium caule erecto ramosissimo, glomerulis florum ad alas longe testibus. *Hall. helv.* 702.

Filago incana tomentosa erecta. *Vaill. act.* 1719 p. 391.

Habitat in Europæ campis aridis, arvisque sabulosis. ☉

SYNGENESIA: POLYGAM. SUPERFLUA. 857

33. GNAPHALIUM caule erecto diviso, floribus py-*ramidalibus*.
 ramidatis lateralibus axillaribusque *Fl. succ.* 678.
 Gnaphalium minus repens. *Bauh. pin.* 262.
 Gnaphalium minimum. *Lob. ic.* 481. *Bauh. hist.* 3. p.
 155. *Raj. hist.* 296.
Habitat in Europæ collibus sabulosis. ☉
34. GNAPHALIUM caule erecto dichotomo, flori-*gallicum*.
 bus pyramidatis axillaribus.
 Gnaphalium vulgare medium. *Morif. hist.* 3. f. 7. t.
 11. f. 14. *Pet. herb.* t. 18. f. 12.
 Gnaphalium minimum alterum nostras, stœchadis citri-
 næ foliis tenuissimis. *Pluk. alm.* 172. t. 298 f. 2.
Habitat in Anglia, Gallia.
Differt a G. germanico Floribus in capitulis paucis; a G.
montano, Foliis subsomentosis, tamen glabris nec bir-
sutis.
35. GNAPHALIUM caule erecto dichotomo, capitu-*germanicum*.
 lis globosis axillaribus terminalibusque.
 Gnaphalium caule erecto dichotomo, floribus in alis
 sessilibus. *Fl. lapp.* 299. *Fl. succ.* 677. *Roy. lugdb.* 148.
 Gnaphalium vulgare majus. *Bauh. pin.* 262.
 Gnaphalium decimum. *Pet. herb.* 18. f. 10.
Habitat in Germania ☉
36. GNAPHALIUM foliis alternis linearibus acutis *saxatile*.
 planis, pedunculis longissimis unifloris. *Hort. cliff.*
 401.
 Helichrysum capitulis singularibus brevibus. *Morif. hist.*
 3. p. 87. f. 7. t. 10. f. 16.
 Helichrysum saxatile, singulari capitulo amplo, angusto
 stœchadis folio. *Bocc. mus.* 142. t. 104.
 Elchryso sylvestri, flore oblongo, similis. *Bauh. pin.*
 265. *prodr.* 123. t. 123.
Habitat in Italia. Sicilia.

XERANTHEMUM. 841

- * *Receptaculo paleaceo, semin. corona quinqueseta.*
1. XERANTHEMUM foliis lanceolatis patentibus. *annuum*.
 Xeranthemum receptaculis paleaceis, seminum pappo
 quinqueseto. *Hort. cliff.* 400.
 Xeranthemum, oleæ folio, capitulis simplicibus incanis,
 non foetens, flore majore violaceo. *Morif. hist.* 3. p.
 43. f. 6. t. 2. f. 2.
 Jacea oleæ folio, capitulis simplicibus. *Bauh. pin.* 272.
 H h h ; Ptar-

858 SYNGENESIA: POLYGAM. SUPERFLUA

- Ptarmica austriaca. Clus. hist. 2. p. 11.*
- inapertum.** β . *Xeranthemum*, oleæ folio, capitulis simplicibus, incanum foetens, flore purpurascante minore. *Morif. hist. 3. p. 43. f. 6. s. 12. f. 1.*
Xeranthemum capitulis inapertis. Hall. belv. 709. t. 23.
Jaceæ oleæ folio, minore flore. Bauh. pin. 272.
- orientale.** γ . *Xeranthemum orientale*, fructu maximo. *Tournef. cor. 38.*
Jaceæ oleæ folio, capitulis compactis. Bauh. pin. 272.
Habitat α . in Austria, β . in Italia, Helvetia G. Narbonensi. ☉
- crucifolium.** 2. *XERANTHEMUM* foliis pinnatifidis. *Gmel. sib. 2. p. 108. t. 47. f. 1. †*
Habitat ad Tanain prope oppidum Cavilnense.
** Receptaculo nudo, seminum corona pilosa.*
- vestitum.** 3. *XERANTHEMUM* fruticosum erectum, foliis lanceolato-linearibus, ramus unifloris foliosis.
Xeranthemum frutescens lanuginosum, foliis longis mollibus, flore argenteo amplissimo. Burm. afr. 177. t. 66. f. 1.
Xeranthemum speciosum capense, flore albo maximo. Pet. gaz. 1. t. 1. f. 9.
Chrysocome f. Argyrocome africana cricoides, flore albo. Seb. thes. 2. p. 45. t. 43. f. 5.
Habitat in Æthiopia. ☿
- speciosissimum.** 4. *XERANTHEMUM* fruticosum erectum, foliis lanceolatis, ramis unifloris subnudis.
Xeranthemum tomentosum latifolium, flore maximo. Burm. afr. 178. t. 66. f. 2.
Elichrysum africanum arborescens, foliis incanis latio-ribus. Boerb. lugdb. 1. p. 121.
Elichrysum africanum lanuginosum latifolium, calyce floris argenteo & amplissimo. Breyn. ic. 27. t. 16. f. 1.
Chrysocome f. Argyrocome gnaphaloides africana, amplissimis floribus. Seb. thes. 2. p. 45. t. 43. f. 6.
- proliferum.** 5. *XERANTHEMUM* caule prolifero, foliis subrotundis imbricatis obsoleteis.
Elichrysum abrotanti feminae foliis. Breyn. ic. 28. t. 17. f. 1.
Habitat in Æthiopia.
** Receptaculo nudo, seminum coronis plumosiss.*
- retortum.** 6. *XERANTHEMUM* caulibus frutescentibus provolutis, foliis tomentosis recurvatis. **Xc**

SYNGENESIA: POLYGAM. SUPERFLUA. 859

Xeranthemum receptaculis nudis, seminum pappo plumoso. *Hort. cliff.* 400. *

Xeranthemoides procumbens, polii folio. *Dill. elsb.* 423. t. 322. f. 415.

Habitat in Æthiopia. 5

7. *HERANTHEMUM* caulibus frutescentibus provo-*elliatum*.
lutis, foliis ciliatis hirsutis.

Aster africanus frutescens, splendentibus parvis & reflexis foliis. *Comm. hort.* 2. p. 55. t. 28.

Habitat in Æthiopia. 6

8. *XERANTHEMUM* squamis calycinis mucronato-*spinosum*.
spinosis.

Xeranthemum spinosum æthiopicum, foliolis simplicibus, capite spinoso. *Burm. afr.* 182. t. 73. f. 3.

Xeranthemum spinosum æthiopicum capitulis compatis purpurascens. *Breyn. prodr.* 2. p. 105.

Habitat in Æthiopia.

9. *XERANTHEMUM* ramis unifloris imbricatis, foliis *sesamoides*.
obsoletis.

Xeranthemum ramosum, foliis squamosis linearibus, floribus argenteis. *Burm. afr.* 181. t. 76. f. 2.

β. *Xeranthemum* *sesamoides*, flore albo, ericæ foliis caulii tomentoso adlatis; ad radicem vero stœchadis citrinæ longioribus & solutis. *Pluk. amalth.* 213. t. 449. f. 5.

Chrysocome f. *Argyrocome* africana ericoides, capitulis b. spei. *Seb. thes.* 2. t. 43. f. 5.

Elichrysium africanum lanuginosum, angustissimo folio, calyce floris argenteo & amplissimo. *Breyn. ic.* 27. t. 16. f. 2.

Habitat in Æthiopia.

10. *XERANTHEMUM* floribus paniculatis, foliis li-*paniculatum*.
neari-lanceolatis.

Xeranthemum incanum, foliis oblongo-acutis, capitulis plurimis argenteis. *Burm. afr.* 180. t. 77. f. 1.

Habitat in Æthiopia

CARPESIMUM. 842

1. *CARPESIMUM* floribus terminalibus.

Carpesium. *Act. nps.* 1741. p. 81. *Hort. nps.* 257.

Aster atticus, foliis circa florem mollibus. *Barb. pin.* 266.

Cornutum.

Aster

Aster cernuus. Col. ecphr. 1. p. 251. t. 252.

Chrysanthemum conyzoides cernuum, foliis circa florem mollibus. Moris. hist. 3. p. 18. f. 6. t. f. 27.

Habitat in Italia. x

obovatoideis 2. **CARPESIMUM** floribus lateralibus.

Habitat in China Osbeck.

Caules ramosi, duriusculi. Folia alterna, lato-lanceolata, in petiolo attenuata, obsolete crenata, digitum longa Flores sparsi, saepe solitarii ad axillas foliorum, longitudinaliter, secundum ramum, sessiles aut brevissimis pedunculis unijloris insidentes, magnitudine unius corylli, nutantes. Calyces umbilicati, cincti foliolis parvis. Flosculi numerosi. Semina oblonga, nuda, glabra, balsamo quasi illita.

BACCHARIS.

ovifolia.

1. **BACCHARIS** foliis lanceolatis longitudinaliter dentato-serratis. *Hort. cliff. 404. Roy. lugdb. 152*

Conyza frutescens, foliis angustioribus nervosis. Few. peruv. 750. t. 37.

Eupatorium africanum, agerato affinis peruvianæ, floribus albis. Pluk. alm. 400. t. 328. f. 2.

Pseudo-Helichrysum frutescens peruvianum, foliis longis serratis. Moris. hist. 3. p. 90.

Habitat in Virginia, Peru. h

tenuifolia.

2. **BACCHARIS** foliis ovato-lanceolatis serratis, caule suffruticoso. *Hort. cliff. 404 **

Conyza africana tenuifolia subfrutescens, flore aureo. B. Dill. elth. 104. t. 88. f. 103.

Habitat in Africa. h

seriifolia.

3. **BACCHARIS** foliis lanceolatis superne uno alterove denticulo serratis. *Hort. cliff. 404. * Roy. lugdb. 152.*

Arbuscula foliis nerii. Boerb. lugdb. 2. p. 263.

Habitat in Æthiopia. h

halimifolia.

4. **BACCHARIS** foliis obovatis superne emarginato crenatis. *Hort. cliff. 405. Gron. virg. 96. Roy. lugdb. 153.*

Senecio virginianus arborecens, triplicis folio. Raj. hist. 1799. Herm. parad. 225.

Pseudo-Helichrysum virginianum frutescens, halimi latioris foliis glaucis. Moris. hist. 3. p. 90. f. 7. t. 10. f. 4.

Eli-

SYNGENESIA: POLYGAM. SUPERFLUA. 86:

- Elichryso affinis virginiana frutescens, foliis chenopodii glaucis. *Pluk. alm.* 134. t. 27. f. 2.
 Argyracome virginiana, atriplicis folio. *Pet. gaz.* t. 7. f. 4.
Habitat in Virginia. 5
5. BACCHARIS foliis obovatis denticulatis. *indica.*
Habitat in India. Braad.
 Rami striis elevatis. Folia obovata, glabra, medio subferrato-dentata, magnitudine extimi pollicis. Corymbus magnus, terminalis: pedunculis angulatis bracteis aliquot subulatis. Calyces cylindrici, laeves. Rami ex alis foliorum superiorum, corymbu longiores.
6. BACCHARIS foliis lanceolatis serrato dentatis, co-fatida, rymbis foliosis. *Gron. virg.* 96.
 Conyza americana frutescens foetidissima. *Dill. elsh.* 106. t. 89. f. 105?
Habitat in Virginia. 2

CONYZA. 8 C 1 4

1. CONYZA foliis lanceolatis acutis, caule annuo corym-squarrosa, boso. *Hort. cliff.* 405. *Hort. upf.* 257.* *Roy. lugdb* 157.
 Conyza major vulgaris. *Bauh. pin.* 265.
 Conyza. *Cum. epit.* 612.
Habitat in Germania, Belgii, Angliæ, Galliæ siccis. 3
2. CONYZA foliis lineari-lanceolatis integerrimis, co-limisolia, rollis radiatis.
 Aster americanus albus, mezerei arabum exasperatis foliis, florum petalis reflexis. *Pluk. alm.* 56. t. 79. f. 2.
Habitat in America septentrionali.
3. CONYZA foliis lato-lanceolatis subserratis, corollis Asteroider. radiatis.
 Aster marilandicus, pharmicæ capitulis, flore albo, baccharidis monspeliensium folio & effigie. *Pluk. mant.* 29.
Habitat in America septentrionali. 2
 Habitus Asteris; Corymbus subfastigiatus. Flores præcedentis radiati.
4. CONYZA foliis ovato-oblongis amplexicaulibus. *bifrons.*
Hort. cliff. 405.
4. Conyza pyrenaica, foliis primulæ veris. *Herm. parad.* radiata.
 127. t. 127.

β. Eu-

862 SYNGENESIA: POLYGAM. SUPERFLUA.

- paniculosa.* 3. *Eupatoria coryzoides maxima canadensis*, foliis caulem amplexantibus. *Pluk. alm.* 141. t. 87. f. 4.
Habitat in Pyrenæis, Canada
- lobata.* 5. *CONYZA* foliis inferioribus trifidis, superioribus ovato-lanceolatis obsolete serratis, floribus corymbosis. *Hort. cliff.* 405.
Conyza arborescens lutea, folio trifido. *Plum. spec.* 9.
Habitat in Vera Cruce. 5
- bifolcata.* 6. *CONYZA* foliis ovalibus dentatis, pedunculis diphyllis: bracteis oppositis.
Eupatoria conyzoides, integro jacobææ folio molli & inano. *Pluk alm.* 140. t. 177 f. 1.
Habitat in India.
- tortuosa* 7. *CONYZA* caule tortuoso fruticoso, foliis ovato-oblongis integerrimis, racemis reflexis. *Hort. cliff.* 405. *
Conyza inagdagascariensis fruticosa, tortuoso caule, corni folio. *Vaill. alt.* 1719. p. 396.
Habitat in Magdagascar, Vera Cruce. 5
- candida.* 8. *CONYZA* foliis ovatis tomentosis, floribus confertis, pedunculis lateralibus terminalibusque. *Hort. cliff.* 405. *Roy. lugdb.* 157.
Conyza cretica fruticosa, folio molli candidissimo tomentoso. *Tournef. cor.* 33.
Jacobæa cretica incana, integro limonii folio. *Barr. rar.* t. 217.
Habitat in Creta.
- cinerea.* 9. *CONYZA* foliis oblongis, floribus paniculatis.
Senecioides. *Fl. zeyl.* 419.
Senecio indica foliis ternis crenatis. *Burm. zeyl.* 211. t. 96. f. 1.
Senecio indicus, atriplicis folio. *Morif. hist.* 3. p. 106. f. 7, t. 17. f. 7.
Eupatorium, angustiore folio, bermudense, floribus flavescens. *Plum. alm.* 141. t. 243. f. 3.
Habitat in India. ☉
Variant Folia figura lanceolata. f. ovata, crenata f. serrata.
- chinensis.* 10. *CONYZA* foliis lanceolato-ovatis serratis subtus tomentosis, floribus terminalibus congestis.
Habitat in China. *Toren.*
Raro plures quam tres flores simul congesti.

11. CO-

SYNGENESIA: POLYGAM. SUPERFLUA. 863

11. CONYZA foliis ovalibus integerrimis scabris subtus *hirsuta*, hirsutis.

Habitat in China.

Caulis *subvillosus*. Folia *oblonga f. ovalia, integerrima, scabra, subtus valde hirsuta pallida*. Flores *in racemis conferti*. Calyces *Asteris*.

ERIGERON. 845

1. ERIGERON pedunculis unifloris lateralibus, calyci-*viscosum*, bus squarrosis. *Hort. nps. 258. **

Aster foliis serratis, pedunculis simplicibus lateralibus unifloris longitudine folii, foliatis. *Hort. cliff. 409.*

Conyza mas theophrasti, major dioscoridis. *Bauh. pin. 265.*

Conyza major. *Dod. pempt. 51.*

Habitat in Narbonensi, Hispania, Italia.

Folia *adpersa pilis, interjectis glandulis viscidis.*

2. ERIGERON caule paniculato, floribus subsolitariis *carolinianum*, terminalibus, foliis linearibus integerrimis.

Virga aurea carolinensis, linariae montepessulanæ foliis.

Dill. elth. 412. t. 306. f. 393.

Habitat in Carolina.

3. ERIGERON caule floribusque paniculatis. *Hort. cliff. canadense. 407. Hort. nps. 258. Gron. virg. 96. Gort. gelr. 475.*

Conyza annua acris alba elatior, linariae foliis. *Morif. hist. 3. p. 115. f. 11. t. 20. f. 29. Bocc. sic. 85. t. 46.*

Habitat in Canada, Virginia, nunc in Europa australi. ☉

4. ERIGERON foliis basi revolutis.

bonariense.

Erigeron foliis inferioribus dentato-laciniatis: superioribus integris. *Hort. cliff. 407. Hort. nps. 258.*

Senecio bonariensis purpurascens, foliis imis coronopii. *Dill. elth. 314. t. 257. f. 334.*

Habitat in America australi. ☉

5. ERIGERON caule multifloro, foliis lanceolatis sub-*philadelphicum*, ferratis; caulinis semiamplexicaulibus.

Habitat in Canada. Kalm. 2

Caulis *pedalis, parum ramosus*. Folia *lanceolata, scabra, una alterave acuta serratura*; Caulina *semiamplexicaulia, breviora*. Flores *sape 7 a 8, radio albo aut purpurascens; pappo albo.*

6. ERIGERON pedunculis alternis unifloris. *Hort. cliff. aro. 407. Fl. suec. 691. Roy. lugdb. 165. Erig.*

864 SYNGENESIA: POLYGAM. SUPERFLUA.

Erigeron vulgare. Fl. lapp. 308.

Conyza cœrulea actris. Banb. pin. 265.

Amellus montanus æquicolorum. Col. cephr. 2. p. 25. t. 26.

Habitat in Europæ apricis, siccis. &

- alpinum.* 7. **ERIGERON** caule subbifloro, calyce subhirsuto.
Conyza cœrulea alpina major. Banb. pin. 265. prodr. 124.
Asteri montano purpureo similis f. globulariæ. Banb. hist. 2. p. 107.
Habitat in Alpibus Helvetiæ. &
Pappus ruffo-ferrugineus est.
- uniflorum.* 8. **ERIGERON** caule unifloro, calyce piloso. *Hort. cliff. 407. Fl. succ. 692.*
Aster caule unifloro, foliis integerrimis, calyce villoso subrotundo. Fl. lapp. 307. t. 9. f. 3.
Aster caule unifloro, calyce tomentoso. Hall. belv. 724.
Habitat in Alpibus Lapponiæ, Helvetiæ. &
- gramineum.* 9. **ERIGERON** caule unifloro, foliis linearibus ciliatis scabris.
Aster caule unifloro longitudine foliorum linearium. Gmel. fib. 2. p. 174. t. 76. f. 2.
Aster acaulos albus, foliis gramineis. Amm. rub. 215.
Habitat in Sibiria. D. Gmelin. &
Minima planta, obvallata foliis emarcidis, aridis.
- ficulum.* 10. **ERIGERON** squamis calycinis inferioribus laxis florem superantibus. *Hort. cliff. 407.*
Conyza sicula annua lutea, foliis atro-virentibus, caule rubente. Bocc. sic. 62. t. 31. f. 4. Moris. hist. 3. p. 115. f. 7. t. 20. f. 28.
Habitat in Siciliæ paludosis. ☉
- campoboratum.* 11. **ERIGERON** foliis lanceolato-ovatis villosis: serraturis apice cartilagineis. *Hort. ups. 259*
Baccharis foliis ovato-lanceolatis serratis, caule herbaeo. Gron. virg. 97.
Habitat in Virginia. ☉
- tuberosum.* 12. **ERIGERON** foliis lanceolatis, ramis unifloris, caule suffruticoso.
Erigeron caule subunifloro, foliis lanceolato-linearibus rigidis, radice tuberosa. Sauv. monsp. 55.
a. Chondrilla bulbosa syriaca, foliis angustioribus. Banb. pin. 130.

SYNGENESIA: POLYGAM. SUPERFLUA. 865

Conyza marina. Moris. hist. 3. p. 114. f. 7. t. 19. f. 20. bona.

β. *Chondrilla bulbosa syriaca, foliis latioribus. Bauh. pin. 130.*

Conyza tuberosa lutea, foliis angustis & rigidis. Magn. monsp. 77. Moris. hist. 3. p. 114. f. 7. t. 20. f. 15.

γ. *Aster conyzoides gesneri. Moris. hist. 3. p. 118. f. 7. t. 22. f. 7.*

Habitat in Galloprovincia, G. Narbonensi, Syria.

Caulis brevis, lignosus; Rami simplices, uniflori. Folia sparsa, lanceolata, subpetiolata, integerrima. Flores lutei, Asteris facie, terminales, sessiles. Pappus griseus. Media inter Asters & Erigera.

TUSSILAGO.

846

1. *TUSSILAGO* scapo unifloro, foliis lyrato-ovatis. *Anandria.*

Tussilago scapo unifloro, calyce clauso. *Hort. ups. 259.*

*t. 3 f. 1. * Gmel. fib. 2. p. 143. t. 68. f. 1.*

*Anandria. Amoen. acad. 1. p. t. 11. **

β. *Tussilago* scapo unifloro, calyce subaperto. *Gmel. fib.*

1. p. 143. t. 67. f. 2.

Habitat in Sibiria. &

Una eademque planta, quæ in solo libero calycem clausum flore occultato gerit, in olla loco calidiori florem præcociorem radiatum expandit, ut variis annis observavimus.

2. *TUSSILAGO* scapo subnudo unifloro, foliis cordato orbiculatis crenatis. *Hort. cliff. 411. Roy. lugdb. 159.*

Tussilago alpina rotundifolia glabra. Bauh. pin. 197.

Tussilago alpina 2. Clus. hist. 2. p. 113.

β. *Tussilago alpina rotundifolia canescens. Bauh. pin. 197.*

Tussilago alpina 1, flore evanido. Clus. hist. 2. p. 113.

Habitat in Alpibus Helvetiæ, Austriæ, Bohemiæ. &

3. *TUSSILAGO* scapo imbricato unifloro, foliis subcor-datis angulatis denticulatis. *Hort. cliff. 411. Fl. suec. 680. Mat. med. 391. Roy. lugdb. 159. Dalib. paris. 256. Gmel. fib. 2. p. 140.*

Tussilago vulgaris. Bauh. pin. 197.

Tussilago Clus. hist. 140. Cam. epit. 590. 591.

Habitat in Europæ argillosis subtus humidis. &

4. *TUSSILAGO* thyrsifloro fastigiato, floribus radiatis. *Fl. frigida. lapp. 303. Fl. suec. 682.*

l i i

Tus.

866 SYNGENESIA: POLYGAM. SUPERFLUA

Tussilago scapo imbricato, floribus spicatis radiatis, foliis infra incanis acutis. *Gmel. fib.* 2. p. 150. & 70.

Petasites scapo paucifloro, foliis subtus tomentosis albis-
simis. *Hall. belv.* 707.

Cacalia tomentosa. *Bauh. pin.* 198. *prodr.* 102. *Scheuch.*
alp. 130. t. 18. f. 1.

Habitas in Alpium Lapponiae, Helvetiae, Sibiriae con-
vallibus. &

albo.

5. *TUSSILAGO* thyrsis fastigiato, flosculis femineis nu-
dis paucis. *Hort. cliff.* 411. *Fl. suec.* 681. *Roy. lugdb.*
159.

Petasites minor. *Bauh. pin.* 197.

Petasites flore albo. *Cam. epit.* 593.

Habitas in Europa. &

hybrida.

6. *TUSSILAGO* thyrsis oblongo, flosculis femineis nu-
dis plurimis. *Hort. cliff.* 411. *Roy. lugdb.* 159. *Gort.*
gcl. 478.

Petasites major, floribus pedunculis longis insidentibus.
Dill. elieb. 309. t. 230. f. 297.

Petasites in medio majoribus flosculis, reliquis minori-
bus. *Buxb. bal.* 258.

Habitas in Germania, Hollandia. &

Petasites.

7. *TUSSILAGO* thyrsis ovato, flosculis omnibus her-
maphroditis. *Hort. cliff.* 411. *Fl. suec.* 685. *Mat. med.*
390. *Roy. lugdb.* 159. *Dalib. paris.* 256.

Tussilago major. *Cam. epit.* 592.

Petasites major & vulgaris. *Bauh. pin.* 197.

Habitas in Europa temperatiore. &

847 SENECIO.

* *Floribus flosculosis.*

bicarinifolius 1. *SENECIO* corollis nudis, foliis amplexicantibus la-
ceris, caule herbaceo erecto. *Hort. wps.* 261.

Senecio foliis lanceolatis amplexicaulibus levibus acute
sinuatis denticulatis, caule herbaceo. *Hort. cliff.* 406.
Roy. lugdb. 163.

Senecio americanus altissimus, blattariae s. hieracii folio.
Herm. par. 226. t. 226.

Habitas in America septentrionali. ☉

divaricatus. 2. *SENECIO* corollis nudis, foliis lanceolatis. dentatis
suavis ramulis floriferis divaricatis.

Habitas in China. Osbeck.

Can-

SYNGENESIA: POLYGAM. SUPERFLUA. 867

Caulis erectus, angulatus, angustus, sesquipedem altus, Folia alterna, remota, lanceolata, petiolata, serrato-dentata, tomentoso-scabra. Ramorum tantum rudimenta ex alis foliorum. Ex supremis alis foliorum aliquot ramuli cum floribus, basi admodum divaricati. Flores terminales, aliquot, pedunculati, magnitudine precedentis, Calycibus basi squamis aliquot laxis subulatis adspersis.

3. SENECIO corollis nudis, scapo subnudo longissimo. *Pseudo-China* Roy. *ludb.* 164.

Senecio maderaspatanus, rari folio, floribus maximis, cujus radix a nonnullis China dicitur. *Dill. els.* 345. t. 258. f. 335.

Habitat in India. *

4. SENECIO corollis nudis, foliis pinnato-sinuatis amplexicaulibus, floribus sparsis. *Roy. ludb.* 165. *Hall. belv.* 733. *Fl. suec.* 690. *Dalib. parif.* 257.

Senecio foliis pinnatifidis denticulatis: laciniis æqualibus patentissimis: rachi lineari. *Hort. cliff.* 406.

Senecio minor vulgaris. *Bauh. pin.* 131. *Fl. lupp.* 296.

Erigeron minus. *Dod. pempt.* 641.

Habitat in Europæ cultis, ruderalis, succulentis. ☉

* Floribus radiatis: radio revolutis.

5. SENECIO corollis revolutis, foliis sessilibus sinuatis, trifloris. calycibus conicis: squamis minimis intactis. *Hort. upf.* 261.

Senecio foliis pinnatifidis: laciniis inæqualibus erectis. *Hort. cliff.* 406.

Senecio foliis sinuatis petiolatis, pedunculis folio longioribus, florum radiis disco brevibus. *Roy. ludb.* 165.

Jacobæa ægyptica, senecionis folio, semisifosculis vix conspicuis. *Vaill. æt.* 1720. p. 295.

Habitat in Ægypto. ☉

6. SENECIO corollis revolutis, foliis amplexicaulibus ægyptiis. sinuatis, squamis calycinis brevioribus integris spha-celatis. *Hort. upf.* 261. *

Jacobæa ægyptica, senecionis folio, flore minore. *Vaill. æt.* 1720. p. 298.

Habitat in Ægypto. ☉

7. SENECIO corollis revolutis, foliis amplexicaulibus lividis. lanceolatis dentatis, squamis calycinis brevissimis intactis. *Hort. upf.* 261. *

l i i a

Senec

868 SYNGENESIA: POLYGAM SUPERFLUA.

Senecio foliis sinuatis: basi latioribus amplexicaulibus, floribus corymbosis, radiis disco brevioribus. Roy. lugdb. 165?

Habitat in Hispania. ☉

- erilebus.* 8. **SENECIO** corollis revolutis, foliis amplexicaulibus, squamis calycinis imbricatis laceris. *Hort. ups. 261. **
Habitat in Hispania. ☉

- viscosus.* 9. **SENECIO** corollis revolutis, foliis pinnatifidis viscidis, squamis calycinis laxis longitudine perianthii.
Senecio foliis pinnatifidis sessilibus: laciniis angulosis: superioribus latioribus, florum radiis disco brevioribus. Roy. lugdb. 142. it. Wgoth. 142. Dalib. parif. 258.
Senecio hirsutus viscidus graveolens. Dill. elth. 347. t. 258. f. 336.
Senecio incanus pingvis. Banb. pin. 131.
Habitat in Europæ pagis, urbibus. ☉

- lytuatus.* 10. **SENECIO** corollis revolutis, foliis pinnatifidis denticulatis, caule corymboso erecto.
Senecio foliis pinnatifidis denticulatis, florum radiis revolutis linearibus. It. Wgth. 326. Fl. sac. 689. Dalib. parif. 258.
Senecio minor, latiore folio, f. montanus. Banb. pin. 131. Dill. elth. 258. t. 258. f. 337.
Jacobæa, senecionis folio incano, perennis. Raj. hist. 285. Hall. jen. 177. l. 4.
Habitat in Europæ borealis sylvis ceduis. ☉

- glaucus.* 11. **SENECIO** corollis revolutis, foliis amplexicaulibus, linearibus pinnato-dentatis: laciniis linearibus integris.
Roy. lugdb. 164.
Jacobæa ægyptia annua, coronopi folio glauco. Vaill. act. 1720. p. 297.
Jacobæa ægyptia, folio glauco coronopi. Boerb. lugdb. 1. p. 99.
Habitat in Ægypto. ☉

- hastatus.* * *Floribus radiatis: radio patente, foliis pinnatifidis.*
12. **SENECIO** corollis radiantibus, petiolis amplexicaulibus, pedunculis folio triplo longioribus, foliis pinnato-sinuatis. *Roy. lugdb. 164.*
Jacobæa asra perennis viscosa lutea, asplenii foliis. Vaill. act. 1720. p. 298.
Habitat in Africa.

SYNGENESIA: POLYGAM. SUPERFLUA. 869

13. **SENECIO** corollis radiantibus, foliis pinnatifidis *æ-elegans*.
qualibus patentissimis: rachi inferne angustata. *Hort.*
cliff. 406. *Hort. upf.* 260.
Jacobæa africana frutescens, flore amplo purpureo ele-
gantissimo, senecionis folio. *Volck. norib.* 225. t.
225.
Aster africanus frutescens, foliis senecionis crassioribus.
Comm. hort. 2 p. 61. t. 31.
Habitat in Æthiopia.
- 14 **SENECIO** corollis radiantibus planis calyce longio- *squalidus*.
ribus integris, foliis pinnatifidis: laciniis linearibus di-
stantibus. *Hort. upf.* 260.
Habitat in Europa Australi. ☉
15. **SENECIO** corollis radiantibus, foliis pinnatifidis den-
tatis subhirtis, caule erecto.
Senecio scanorensis. *it. scan.* 225.
Jacobæa altissima, foliis crucæ artemisiæve similibus &
æmulis. *Rupp. gen.* 164.
Jacobæa incana repens. *Barr. rar.* 1075. t. 153.
Habitat in aggeribus Europæ temperatæ.
16. **SENECIO** corollis radiantibus, foliis utrinque to- *incanus*.
mentosis semipinnatis: laciniis subdentatis, corymbo
subrotundo. *Hall. belv.* 731.
Chrysanthemum alpinum incanum, foliis laciniatis.
Bauh. pin. 133.
Chrysanthemum alpinum 1. *Clus. hist.* 1. p. 333.
Jacobæa alpina incana minor. *Barr. ic.* 262.
Habitat in Alpibus Helvetiæ, Austriæ, Pyrenæo-
rum.
17. **SENECIO** corollis radiantibus, foliis pinnato-multifi- *abrotanifoli-*
dis linearibus. † *us.*
Senecio foliis decompositis. *Guet. stamp.* 360.
Chrysanthemum alpinum. foliis abrotani multifidis.
Bauh. pin. 134.
Chrysanthemum alpinum II. *Clus. hist.* 1. p. 333. t.
334.
Habitat in Alpibus Stiriæ, Pyrenæicis.
18. **SENECIO** corollis radiantibus, foliis bipinnatis li- *canadensis*,
nearibus.
Habitat in Canada. Kalm.
Caulis erectus, levis. Folia bipinnata, linearia, glabra:
summa simpliciter pinnata. Corymbus terminalis,
1113 *com-*

870 SYNGENESTA: POLYGAM. SUPERFLUA

compositus, fastigiatus, flavus. Calyces rufescentes.

- Jacoba.* 19. **SENECIO** corollis radiantibus, foliis pinnato-lyratis: laciniis lacinulatis, caule erecto. *Hort. cliff.* 406. *Fl. suec.* 688. *Roy. lagdb.* 174. *Dalib. parisi.* 257. *Jacobæa vulgaris laciniata. Baub. pin.* 131. *Herba f. Jacobi. Fuchs. hist.* 742. *Habitat in Europæ pascuis. &*
- virg.* 20. **SENECIO** corollis radiantibus, foliis crenatis: infimis cordatis petiolatis; superioribus pinnatifidis lyratis. *Gron. virg.* 98. *Jacobæa virginiana, foliis imis alliaræ glabris; caulescentibus barbæræ. Moris. hist.* 3. p. 110. *Raj. suppl.* 180. *Habitat in Virginia, Canada. &* *Antecedenti nimis affinis.*
- * *Floribus radiatis: radio patente, foliis indivisis.*
- paludosus.* 21. **SENECIO** corollis radiantibus, foliis ensiformibus acute serratis subtus suovillosis, caule stricto. *Senecio foliis integris serratis. Grev. stamp.* 2. p. 559. *It. scan.* 396 * *Solidago caule simplici, corymbo terminali, foliis subtus tomentosis, flosculis femineis sedecim. Leeb. scan.* 68. *Fl. suec.* 686. *Jacobæa foliis longis integris & mucronatis. Moris. hist.* 3. p. 110. *f. 7. s. 7. f. 22.* *Conyza palustris serratifolia. Baub. pin.* 266. *β. Solidago palustris altissima incana & tomentosa, foliis longioribus serratis. Amm. ruth.* 220. *Gmel. sib.* 2. p. 162. *Habitat in Europæ paludibus maritimis. &* *Variat foliis utrinque tomentosis.*
- nemorensis.* 22. **SENECIO** corollis radiantibus octonis, foliis lanceolatis biserratis subtus villosis, caule ramoso. *Senecio foliis ovato-lanceolatis subtus hirsutis, floribus umbellatis. Hall. belv.* 734. *Itin.* 23. *Jacobæa nemorensis, latiore rigidior & hirsuto folio. Rupp. jen.* 142. *Solidago foliis lanceolatis serratis, floralibus ad pedunculum minimis. Gmel. sib.* 2. p. 165. *Habitat in Germaniæ, Sibiræ nemoribus. &*

22. SE-

SYNGENESIA: POLYGAM. SUPERFLUA. 87r

23. SENECEO corollis radiantibus, floribus corymbosis, *sarracenicus*.
foliis lanceolatis serratis glabriusculis. *Hort. ups.* 266.

Senecio foliis lanceolatis serratis, floralibus pedunculos
æquantibus. *Vir. cliff.* 84. *Roy. lugdb.* 163. *Gort.*
gelr. 481.

Solidago caule simplici, corymbo terminali: pedunculis
partialibus alternis nudis longitudine folii. *Hort. cliff.*
410.

Solidago sarracenicæ. *Fuchf. hist.* 728.

Virga aurea angustifolia serrata. *Baub. pin.* 268.

Habitat in Helvetiæ montanis, nemorosis. &

Dignoscitur facie, latis foliis, magnitudine, radice ma-
xime reptatrice.

24. SENECEO corollis radiantibus, foliis obovatis car-*balimisfolius*.
natis subdentatis, caule fruticoso.

Solidago foliis obverse ovatis carnosissimis crenatis, caule
fruticoso, corymbo ramoso. *Hort. cliff.* 411.

Doria africana arborescens, foliis crassis & succulen-
tis atriplicem referentibus. *B. Dill. elth.* 124. t. 104.
f. 124.

Habitat in Æthiopia. &

25. SENECEO corollis radiantibus, foliis serrato-subspi-*byzantium*.
natis.

Senecio foliis crenato-dentatis: inferioribus pinnato-ly-
ratis: superioribus pinnatifidis. *Roy. lugdb.* 530.

Habitat Byzantii. &

Caulis quadripedalis, teres, striatus, viridis, adpersus
pilis albis. Rami in inferiore parte caulis, patentissi-
mi. Folia lanceolata, sessilia, amplexicaulia, revoluta,
serrata dentibus acutis, sinibus obtusis, acuminibus
spinoso-inermibus; supra viridia, nitida, scabriuscu-
la; subtus venosa, villosa, viridia. Ramorum rudi-
menta ex singulis alis foliorum. Scapi terminales, te-
retes, caule longe angustiores, filiformes, longissimi,
leviores, striati, inferne adpersi foliis linearibus, in-
divisis, non dentatis, sed serrato-spinosis; subnudi.
Corymbi terminales, alterni, subdivisi. Calyces api-
ce sphacelati. Corollæ luteæ: Radio octifloro, ovato,
vix tridentato. Facies Serratula arvensis.

26. SENECEO corollis radiantibus, foliis sagittatis am-*ilicifolius*.
plexicaulibus dentatis, caule fruticoso. *Vir. cliff.* 84.
Hort. cliff. 406. *Roy. lugdb.* 164.

872 SYNGENESIA: POLYGAM. SUPERFLUA.

Jacobææ africana frutescens, foliis incisfis & subtus cinereis, Comm. rar. 42. t. 42.

Habitat in Æthiopia. b.

- rigidus.* 26 **SENECIO** corollis radiantibus, foliis cordato-oblongis amplexicaulibus scabris acuminatis serratis, caule fruticoso. *Hort. cliff. 406. Roy. lugdb. 164.*
Jacobææ africana frutescens, foliis rigidis & hirsutis. Comm. hort. 2. p. 149. t. 75.
Jacobææ africana ramosissima, foliis senecionis pinguis rigidis & lucentibus subtus incanis. Raj. supp. 179.
Habitat in Æthiopia. b

848

ASTER

* *Integrifolii, pedunculis nudis.*

- fruticosus.* 1. **ASTER** foliis linearibus fasciculatis punctatis, pedunculis unifloris nudis, caule fruticoso rugoso. *Hort. cliff. 409. Roy. lugdb. 163.*
Aster africanus frutescens, foliis angustis & plerumque conjunctis. Comm. hort. 2. p. 53. t. 27
 β. *Aster maritimus fruticosus, hysopi foliis confertis, flore albo. Pluk. mant. 29. t. 340. f.*
Habitat ad Cap. b. Spei in saxosis. b

- alpinus.* 2. **ASTER** foliis lanceolatis hirtis: radicalibus obtusis, caule simplicissimo unifloro.
Aster montanus cæruleus, magno flore, foliis oblongis. Baub. pin. 267?
Aster VII austriacus V. Clus. hist. 2. p. 15?
 β. *Aster atticus alpinus alter. Baub. pin. 267.*
*Aster atticus cæruleus alpinus. Baub. pin. 905.**
 γ. *Aster hirsutus austriacus cæruleus, magno flore, foliis subrotundis. Baub. pin. 267. prodr. 124.*
Habitat in Austria, Valleis, Helvetia, Pyrenæis. &

- sibiricus.* 3. **ASTER** foliis lanceolatis venosis scabris extimoseratis, caulibus striatis, pedunculis tomentosis.
Aster foliis ovatis oblongis supra serratis, caulibus striatis, pedunculis unifloris umbellatis. Gmel. sib. 2. p. 186. t. 80. f. 10.
Habitat in Sibiria.

- Tripolium.* 4. **ASTER** foliis lanceolatis integerrimis carnosiss glabris, ramis inæquatis, floribus corymbosis. *Roy. lugdb. 168. Fl. succ. 697. Gmel. sib. 2. p. 187. t. 80. f. 2.*
Aster maritimus Tripolium dictus. Raj. bist. 270. Hort. cliff. 408.

Tri-

SYNGENESIA: POLYGAM. SUPERFLUA. 873

Tripolium majus cæruleum. Bauh. pin. 267.

Tripolium. Dod. pempt. 379.

Habitat in Europæ littoribus maritimis & ad Sibiria læcus salsos. &

5. **ASTER** foliis lanceolatis obtusis scabris trinervis integris, pedunculis nudiusculis corymbosis, squamis calycinis obtusis. *Amellus.*

Aster foliis lanceolatis scabris semiamplexicaulibus subterratis, calycibus laxis: squamis lanceolatis. Hort. cliff. 407. Roy. lugdb. 168. Gmel. fib. 2. p. 185.

Aster atticus cæruleus vulgaris. Bauh. pin. 267.

Aster atticus. Dod. pempt. 266.

Aster VIII. italorum fuchii. Clas. hist. 2. p. 16.

Habitat in Europæ australis asperis collibus. &

6. **ASTER** ramis divaricatis, foliis ovatis ferratis, floribus integerrimis. *divaricatus.*

Aster caule infirmo, foliis ovatis acuminatis integerrimis, pedunculis unifloris nudis, calycibus simplicibus. Gron. virg. 99.

Aster americanus latifolius albus, caule ad summum brachiato. Pluk. alm. 56. t. 79. f. 1.

Habitat in Virginia. &

** Integrisfolii, pedunculis squamosis.*

7. **ASTER** foliis linearibus integerrimis, caule paniculato. *damosus.*

Hort. cliff. 408. Roy. lugdb. 167. Gron. virg. 99.

Aster novæ anglia, linearis foliis, chamæmeli floribus.

Herm. par. 95. t. 55.

Aster americanus multiflorus, flore albo bellidis: disco luteo. Pluk. alm. 56. t. 78. f. 6.

Habitat in America septentrionali. &

Caulis paniculato-ramosissimus, ramis filiformibus. Folia caulina angusta lanceolata. Ramorum linearia. Pedunculi filiformes, striati, uniflori, adpersi foliolis angustissimis. Flores parvi Calyce erecto, imbricato, laxo; Radius copiosus albus; Discus flavus, radio parvior.

8. **ASTER** foliis sublinearibus integerrimis, pedunculis tenuifoliis, foliolotis.

Aster americanus, belvedere foliis, floribus ex cæruleo albicantibus, spicis longis. Pluk. alm. 56. t. 78. f. 5.

Habitat in America septentrionali. &

Caules tenuis, angulati, laeves. Folia alterna, linearisub-

sublaucolata, integerrima, minus scabra. Caulis parum ramosus. Flores terminales, solitarii, Pedunculis foliolis subulatis, minimis adspersis.

linariifolius. 9. ASTER foliis linearibus integerrimis mucronatis scabris carinatis, pedunculis foliolosis.

After americanus frutescens, saturejæ foliis scabris, floribus amplis saturate violaceis. *Pluk. alm. 56. t. 14. f. 7.*

After marilandicus, rosmarini foliis angustioribus, in caule crebris inordinatis, floribus in summitate paucis. *Raj. suppl. 175.*

Habitat in America septentrionali. 2

Caules purpurascens. Folia linearia, viridia, valde scabra, acuta, sparsa. Pedunculi alterni, foliolis minimis adspersi. Flores terminales, solitarii.

linifolius. 10. ASTER foliis linearibus acutis integerrimis, caule corymbofo ramosissimo. *Hort. cliff. 408. Gron. virg. 99. Roy. lugdb. 167.*

After tripolii flore, angustissimo & tenuissimo folio. *Morif. hist. 3. p. 121.*

Habitat in America septentrionali. 2

sedifolius. 11. ASTER foliis lanceolato-linearibus subcarnosis integerrimis planis, floribus corymbosis fastigiatis, pedunculis foliolosis.

After caule erecto umbellifero, venis simplicibus, foliis ligulatis. *Sauv. monsp. 54.*

After tripolii flore. *Bauh. pin. 267.*

After minor narbonensium, tripolii flore, linearis folio.

Lob. ic 349. bene. Pluk. alm. 56. t. 271. f. 3?

Habitat in Hungaria interamni, Hispania, Mouspelii.

concolor. 12. ASTER caule simplicissimo, foliis ovatis sessilibus integerrimis, racemo terminali. *Gron. virg. 178. **

After purpureus non ramosus, foliis parvis rigidis, floribus concoloribus seriatim ad caulem dispositis. *Clayt. ibid.*

Habitat in Virginia. 2

rigidus. 13. ASTER floribus terminalibus solitariis, foliis linearibus alternis. *Gron. virg. 98.*

After foliis parvis rigidis crebris, caule non ramoso ligneo infirmo, flore unico specioso: barbulis purpureis longis, flosculis in disco ferrugineis, calyce squamoso tumido rotundo, *Clayt. ibid.*

Habitat in Virginia. 2

14. A-

SYNGENESIA: POLYGAM. SUPERFLUA. 875

14. ASTER foliis cordato-lanceolatis undulatis, flori-^{undulatis.}bus racemosis adscendentibus. *Hort. cliff.* 408. *Roy. lugdb.* 167.
 Aster novæ anglæ purpureus, virgæ aureæ facie & foliis undulatis. *Herm. par.* 96. t. 96.
 Aster virginianus comosus, foliis latioribus & flosculis minimis cæruleis. *Morif. hist.* 3. p. 120.
Habitat in America septentrionali. 2
15. ASTER foliis lanceolatis alternis integerrimis semi-^{nova anglia}amplexicaulibus, floribus terminalibus. *Hort. cliff.* 408. *Gron. virg.* 100. *Roy. lugdb.* 166.
 Aster novæ anglæ altissimus hirsutus, floribus amplis purpuro-violaceis. *Herm. par.* 98. t. 98.
 Aster novæ anglæ altissimus hirsutus, floribus omnium maximis purpuro-violaceis. *Tournef. inst.* 482.
Habitat in Nova Anglia. 2
 Caulis fuscus. Pedunculi imbricati foliolis. Corollæ radii cæruleus.
16. ASTER caule paniculato, pedunculis racemosis: pe-^{ericoides.}dicellis foliosis: foliolis linearibus integerrimis. *Gron. virg.* 100.
 Aster caule paniculato, floribus racemosis, calycibus patulis oblongis, foliis subulato-linearibus. *Roy. lugdb.* 168.
 Aster ericoides dumosus *Dill. els.* 40. t. 36. f. 40.
Habitat in America septentrionali. 2
- * *Serratifolii, pedunculis lævibus.*
17. ASTER foliis cordatis serratis petiolatis, caule pa-^{cordifolius.}niculato. *Hort. cliff.* 408. *Roy. lugdb.* 167.
 Aster latifolius glaber humilis ramosissimus, flore parvo cæruleo, foliis ad basin cordatis. *Morif. hist.* 3. p. 121. f. 7. t. 22. f. 34.
 Aster latifolius autumnalis. *Corn. canad.* 64. t. 65.
Habitat in America & Asia septentrionali. 2
18. ASTER foliis semiamplexicaulibus lanceolatis ser-^{punicus.}ratis scabris, pedunculis alternis subunitis, calycibus discum superantibus. *Hort. cliff.* 408. *Roy. lugdb.* 166.
 Aster americanus latifolius, punicis caulibus. *Herm. lugdb.* 649. t. 651. *Morif. hist.* 3. p. 120.
Habitat in America septentrionali. 2
19. ASTER foliis lanceolatis: lateribus inferiorum cre-^{arctatus.}natis, radice annua, caule corymboso, pedunculis nud

876 SYNGENESIA: POLYGAM. SUPERFLUA.

nudis. *Hort. cliff.* 409. *Hort. upf.* 262. * *Roy. lugdb.* 169.

Aster annuus ramosus albus latifolius canadensis. *Morif. hist.* 3. p. 122.

Bellis ramosa umbellifera. *Corn. canad.* 193. t. 194.

Habitat in Canada. ☉

vernus.

20. ASTER caule subnudo filiformi subramoso, pedunculis nudis, foliolis radicalibus lanceolatis integerrimis obtusis. *Gron. virg.* 99.

Aster palustris, foliis bellidis non serratis, barbulis tenuibus albis, disco luteo, caule viridi hirsuto erecto. *Clayt. ibid.*

Habitat in Virginia. &

* *Serratifolii, pedunculis squamosis.*

indicus.

21. ASTER foliis ovato-oblongis serratis. floralibus ovali-lanceolatis integerrimis, ramulis unifloris.

Aster conyzoides ind. orient. ramulior, caulibus sparsis. *Pluk. alm.* 57. t. 149. f. 4.

Habitat in China.

lævis.

22. ASTER foliis lanceolatis sessilibus lævibus subserratis, pedunculis foliolosis subdivisis.

Habitat in America septentrionali. *Kalm.* &

Caulis erectus, lævis, angulatus, angustus. Folia alterna, remota, medio parum serrata, sessilia, lato-lanceolata, utrinque glaberrima. Panicula minus ramosa: Pedunculis foliolis subulatis, minimis adspersis. Radius corollarum caruleus.

mutabilis.

23. ASTER foliis lanceolatis, calycibus imbricatis basi iquarrolis.

Aster cæruleus americanus non fruticosus serotinus angustifolius humilis, flore amplo floribundus. *Pluk. alm.* 56. t. 326. f. 1.

Habitat in America septentrionali. &

Differt a *A. serotino* Calycibus magis arcte imbricatis & Corollæ radiis saturate purpureis; Discus florum primum luteus, dein purpureus.

tradescenti.

24. ASTER foliis lanceolato-linearibus medio serratis, pedunculis foliosis, caule racemoso, calycibus erectis. *Hort. cliff.* 408. *Hort. upf.* 262. * *Roy. lugdb.* 167.

Aster virginianus ramosissimus serotinus, parvis floribus albis tradescenti. *Morif. hist.* 3. p. 121.

Habitat in Virginia. &

25. A-

SYNGENESIA: POLYGAM. SUPERFLUA. 877

25. ASTER foliis lanceolatis subserratis sessilibus, cau- *novi belgii.*
le paniculato, ramulis unifloris solitariis, calycibus
squarrosis. *Hort. cliff.* 408. *Hort. nps.* 262. *

Aster novæ belgiæ latifolius umbellatus, floribus diftute
violaceis. *Herm. lugdb.* 66. t. 67.

Habitat in Virginia, Pennsylvania. *h*

26. ASTER caule corymboso, foliis lanceolatis reflexis, *grandiflorus*
floribus solitariis, calycibus patulis. *Roy. lugdb.* 168.

Aster foliis lanceolatis semiamplexicaulibus crenatis sca-
bris, ramis unifloris foliolis. *Gron. virg.* 99.

Aster grandiflorus asper: squamis reflexis. *Dill. elsb.*
41. t. 36. f. 41.

Aster virginianus pyramidatus, buglossi foliis asperis, ca-
lyctis squamulis foliaceis. *Mart. cent.* 19. t. 19.

Habitat in America septentrionali. *z*

27. ASTER floribus ovatis, disco radiis longiore. *miser.*

Aster ericoides, meliloti agricæ umbone. *Dill. elsb.* 40.
t. 35. f. 39.

Habitat in America septentrionali. *z*

Habitus proximorum. Caulis sesquipetalis, crassus,
viridis, minus paniculatus, quam in reliquis. Folia
caulis parum serrata, mutantis. Ramorum lanceolata,
integerrima. Calyces cum disco corolla ovati, imbrici-
cati foliolis erectiusculis, longitudine radii. Radius
albus, brevissimus & miser. Discus parvus, valde
convexus, pallidus, stylis flavis.

28. ASTER caule fruticoso, ramis unifloris, foliis ovato- *reflexus.*
lanceolatis recurvis serrato-ciliatis.

Aster africanus frutescens, splendentibus parvis reflexis
foliis. *Comm. hort.* 2. p. 55. t. 28.

Habitat in Æthiopia. *h*

29. ASTER foliis ovatis angulatis dentatis petiolatis, ca- *chinensis.*
lycibus terminalibus patentibus foliosis. *Hort. cliff.*

407. *Hort. nps.* 262 *Roy. lugdb.* 169.

Aster, chenopodii folio, annuus, flore ingenti specioso.

Dill. elsb. 38. t. 34. f. 38.

Habitat in China? *o*

* *Pinnatifolii.*

30. ASTER foliis pinnatis. *Hort. cliff.* 407. *

Aster americanus, foliis pinnatis & serratis, floribus au-
rantis. *Hauft. nss. c. fig. Amm. herb.* 466.

Habitat in Vera Cruce.

SO.

849

SOLIDAGO.

- sempervirens* 1. **SOLIDAGO** foliis lanceolatis subcarneosis glaberrimis margine scabriusculis, paniculacorymbosa.
Solidago panicula corymbosa, racemis reflexis, floribus adscendentibus, foliis glaberrimis. *Gron. virg.* 97.
Solidago maxima. *Corn. canad.* 168.
Virga aurea noveboracensis glabra, caulibus rubentibus, foliis angustis glabris. *Herm. flor.* 26.
Virga aurea canadensis, foliis carnosissimis non serratis: latioribus s. angustioribus. *Morif. hist.* 3. p. 124. f. 7. t. 23. f. 15.
Virga aurea s. Solidago procerior americana, caule multiplici. *Pluk. alm.* 389. t. 235. f. 5.
Habitat in Noveboraco, Canada. 2
Notabilis Caule rubro, homine altiore, Foliis glaberrimis, subcarneosis, margine parum scabris, tota hyeme persistentibus; tempore florendi nimis fero, ut apud nos hyems sapissime flores suffocet.
- canadensis.* 2. **SOLIDAGO** paniculato-corymbosa, racemis recurvatis, floribus adscendentibus, foliis trinerviis subserratis scabris. *Hort. nps.* 259. *Hort. cliff.* 409. *Gron. virg.* 97. *Rey. lugdb.* 161.
Virga aurea angustifolia, panicula speciosa canadensis. *Pluk. alm.* 389. t. 236. f. 1. *Morif. hist.* 3. p. 125.
Virga aurea americana, foliis serratis angustis subtus nervosis. *Morif. bles.* 322.
Habitat in Virginia, Canada. 2
- altissima.* 3. **SOLIDAGO** paniculato-corymbosa; racemis recurvis, floribus adscendentibus, foliis enerviis subintegerrimis. *Hort. nps.* 259.
Virga aurea altissima serotina, panicula speciosa patula. *Mart. cent.* 14. t. 14.
Habitat in America septentrionali. 2
Habitus precedenti simillimus, diversus magnitudine, tempore florendi, serraturis nervisque foliorum; ceterum eadem commiscet plantas vix genuinas, forte hybridas, ut vix limites reperias. Itaque conferantur.
Virga aurea novæ anglie altissima, paniculis nonnquam reflexis. *Boerb. lugdb.* 1. p. 97.
Virga aurea americana hirsuta, radice odorata. *Dill. elsb.* 410. t. 304. f. 391.
Virga aurea novæ anglie, rugosis foliis crenatis. *Dill. elsb.* 406. t. 308. f. 396. *Vir-*

SYNGENESIA: POLYGAM. SUPERFLUA. 879

Virga aurea americana aspera, foliis brevioribus serratis. *Dill. elsb.* 411. t. 305. f. 392.

Virga aurea marilandica spicis florum racemosis, foliis integris scabris: *Mart. cent.* 13. t. 13.

4. SOLIDAGO panicula corymbosa: racemis recurvis *laterifloris* adscendentibus, caule inferne ramoso floriferoque.

Habitas in America septentrionali. Kalm.

Planta *S. canadensi* dimidio minor. Folia non serrata, molliora, sed uno alterove dente interdum notata. Flores & omnia ut in precedentibus, et caulis infra medium Ramos emittit simplices, caule paulo breviores, apice corymbosos; nullos vero ramos inter corymbum terminalem caulis & medium seu ramos, quod in hac singulare.

5. SOLIDAGO panicula corymbosa, racemis supra caesia densioribus, caule glabro laevi.

Virga aurea marilandica caesia glabra. *Dill. elsb.* 414. t. 307. f. 395.

Virga aurea marilandica, foliis longis angustis acutis, ramis s. virgulis floriferis e foliorum alis exeuntibus longa serie. *Raj. supp.* 168.

Habitas in America septentrionali. &

6. SOLIDAGO caule obliquo, pedunculis erectis folio-*mexicana*, latis ramosis, foliis lanceolatis integerrimis. *Hort. cliff.* 409. *Roy. lugdb.* 161.

Virga aurea mexicana. *Bauh. pin.* 517,

Virga aurea, limonii folio, panicula uno versu disposita. *Tournef. inst.* 404.

Habitas in Mexico? &

7. SOLIDAGO caule flexuoso, foliis ovatis acuminatis *flexicaulis*, serratis, racemis lateralibus simplicibus. *Roy. lugdb.* 161.

Virga aurea montana, scrophulariae folio. *Pluk. alm.* 390. t. 235. f. 3.

Virga aurea canadensis, asteris folio. *Herm. par.* 244. t. 244.

Habitas in Canada.

8. SOLIDAGO caule erecto, foliis ovatis acuminatis *latifolia*, serratis, racemis lateralibus simplicibus.

Virga aurea, latissimo folio, canadensis glabra. *Pluk. alm.* 389. t. 235. f. 4.

Habitas in Canada. &

Ni-

880 SYNGENESIA: POLYGAM. SUPERFLUA.

Nimis affinis S. flexicauli. Caulis rectus, acute angulatus. Folia ovata, utrinque mucronata, serrata, supra laevia, subtus scabra. Racemi ex alis, foliopleurumque breviores.

- Virgaurea.* 9. **SOLIDAGO** caule subflexuoso angulato, racemis paniculatis erectis confertis.
Solidago caule erecto, racemis alternis erectis. *Hort. cliff.* 409. *Fl. suec.* 685. *Mat. med.* 393. *Roy. lugdb.* 161. *Hall. helv.* 729. *Dalib. parif.* 256. *Gmel. fib.* 2 p. 154.
Solidago floribus per caulem simplicem undique sparsis. *Fl. lapp.* 306.
Virga aurea vulgaris latifolia. Bauh. hist. 2. p. 162.
Virga aurea latifolia serrata. Bauh. pin. 268.
Habitat in Europæ paucis siccis. &
- rigida.* 10. **SOLIDAGO** foliis caulinis ovatis scabris, ramis alternis fastigiatis, corymbis terminalibus. *Hort. cliff.* 410. *Roy. lugdb.* 162.
Virga aurea novæ anglæ, lato rigidoque folio. Herm. par. 243. s. 243.
Habitat in Pennsylvania. &
- novboracensis.* 11. **SOLIDAGO** foliis radicalibus ovato-oblongis petiolatis, caule nudiusculo ramoso fastigiato.
Habitat in America septentrionali. &
Folia radicalia S. virgaurea, maxime scabra. Caulis pedalis, tribus quatuorve foliis minoribus, superne ramosus, fastigiatus. Flores solitarii ramos terminant, magni, lutei.
- alpina.* 12. **SOLIDAGO** foliis cordatis petiolatis dentatis. *Roy. lugdb.* 162.
Jacobæa alpina, foliis subrotundis ferratis. Bauh. pin. 131. *prodr.* 70. s. 71.
Jacobæa alpina laciniata, flore, bunththalmi. Bauh. pin. 131.
Jacobæa 3 latifolia 1. Clus. hist. 2. p. 23.
Senecio foliis petiolatis cordatis ferratis subtus incanis, floribus luteis ex summo caule racemosis. Hall. helv. 731.
Habitat in Alpibus Helvetiæ, Austriæ, Tyrol. &
- Doronicum.* 13. **SOLIDAGO** foliis lanceolatis subdentatis: caulinis amplexicaulibus, caule subunifloro.
Doronicum foliis lanceolatis alternis dentatis: caulinis am-

SYNGENESIA: POLYGAM. SUPERFLUA. 881

amplexicaulibus, caule ramoso. *Ray. lugdb. 160.*
Sauv. monsp. 84.

Doronium longifolium hirsutie asperum. Bauh. pin. 185.

Doronicum II, Austriacum I. Claf. hist. 2. p. 17.

β. *Tuffilago alpina, folio oblongo. Bauh. pin. 197. prodr. 101.*

γ. *Virga aurea montana serrata, magno flore. Bauh. pin. 268.*

Habitat in Alpibus Helveticæ, Austria, Italia, Monspelii.

14. **SOLIDAGO** foliis linearibus integerrimis, corym-
 bo simplici. *Hort. cliff. 410.*

Linaria aurea affinis. Bauh. pin. 213. prodr. 107.

Jacobæa, lini folio, hispanica & Italica. Bocc. mus. 2. p. 60. t. 49. Barr. rar. 97. t. 802.

Habitat in Hispania, Italia.

25. **SOLIDAGO** foliis lanceolatis serratis decurrentibus, *montana,*
 caule angulato. *Le Monier. Sauv. monsp. 84*

Conyza montana, foliis longioribus serratis, flore sulphureo albicante. Vaill. parij. 1725.

Habitat Monspelii.

INULA. 850

1. **INULA** foliis ovatis rugosis: subtus tomentosis ca-
lycum squamis ovatis. Amæn. acad. 1. p. 410. Mat. med. 392.

Aster foliis ovatis rugosis amplexicaulibus: subtus tomentosis, calycum squamis ovatis patulis. Hort. cliff. 407. Pl. suec. 695. Ray. lugdb. 166. Dalib. parij. 260. Gmel. fib. 2. p. 175.

Helenium vulgare. Bauh. pin. 266.

Helenium. Cam. epit. 35. Fuchs. hist. 242.

Habitat in Auglia, Belgio. &

2. **INULA** foliis dentatis hirsutis: radicalibus ovatis, cau-
 linis lanceolatis amplexicaulibus, caule paucifloro.

Aster luteus, radice odora. Bauh. pin. 266.

Asteris altera species apula. Col. ceph. 1. p. 251. t. 253.

Conyza altera apula. Moris. hist. 3. p. 113. f. 7. t. 21. f. 6.

Habitat in Italia, Galloprovincia, Narbona. &

3. **INULA** foliis oblongis integris hirsutis, caule piloso
 corymbofo, floribus confertis

K k k

Oculus Christi.
 Aster

882 SYNGENESIA: POLYGAM, SUPERFLUA.

Aster caule superne ramoso ampliato, foliis lanceolatis amplexicaulibus integris, calycibus laxis terminalibus.

Hort. cliff. 407.

Conyza pannonica lanuginosa. Bauh. pin. 265. *Morif. hist.* 3. p. 113. f. 7. s. 19. f. 1.

Conyza austriaca. Clus. hist. 2. p. 20.

Habitat in austria. 2

Britannica. 4. *INULA* foliis lanceolatis serratis subamplexicaulibus: subtus villosis, caule ramoso erecto.

α. *Conyza affinis. Bauh. pin.* 265.

Conyza palustris repens britannica dicta. Morif. hist. 3. p. 113. f. 7. s. 19. f. 8.

β. *Conyza aquatica, asteris flore aureo. Bauh. pin.* 266. *prodr.* 124. *Bursf. XV* 41.

Habitat in Eufatia, Bavaria, Scania.

Habitus 3. dysenterica, seu Caulis altior, erectior. Folia angustiora argutius serrata. 3. differt Foliis angustioribus & subtus magis villosis; caeterum radii petala angustissima & subfiliformia.

dysenterica. 5. *INULA* foliis oblongis, caule hirsuto paniculato, squamis calycinis setaceis. *Amæn. acad.* 1. p. 410. *Sauv. monsp.* 8;

Aster foliis semiamplexicaulibus oblongis, caule paniculato hirsuto, floribus latis, squamis calycinis setaceis. *Fl. suec.* 694. *Gvett. stamp.* 2. p. 355. *Dalib. paris.* 260. *Gmel. fib.* 2. p. 180.

Aster pratensis autumnalis, conyzae folio. *ist. goetl.* 273.

Conyza media, asteris flore luteo, f. tertia dioscoridis. Bauh. pin. 265.

Conyza 1. Tabern. hist. 1243.

Calaminthæ tertium genus. Fuchs. hist. 436.

Habitat in Europæ fossis subhumidis. 2

Pulicaria. 6. *INULA* foliis undulatis amplexicaulibus, caule prostrato.

Erigeron foliis undulatis amplexicaulibus. Fl. suec. 693.

Conyza ramis secundum flores gradatim enascentibus, foliis lanceolatis obtusis. Hort. cliff. 405,

Conyza minor, flore globofo. Bauh. pin. 266.

Conyza minima. Dod. pempt. 52.

Habitat ad vias & plateas Europæ temperatæ. ☉

Galiciua. 7. *INULA* foliis lanceolatis recurvis serrato-ciliatis, floribus solitariis, ramis angulatis. *Amæn. acad.* 1. p. 410. *Sauv. monsp.* 80. Aster

SYNGENESIA: POLYGAM. SUPERFLUA. 883

- Aster foliis lanceolatis amplexicaulibus serrato-ciliatis glabris acuminatis recurvis, floribus solitariis, caule striato. *Fl. Jacq.* 696. *Dalib. parif.* 259 *Gmel. fib.* 2. p. 177. t. 67.
- Aster montanus luteus, salicis glabro folio. *Bauh. pin.* 266
- Bubonium luteum 1. *Tabern. hist.* 716.
Habitat in Europæ borealis pratis uliginosis, asperis. & Differt ab Inulæ hirta, Caule glabro superne sulcato f. angulato, Foliis glabris, margine tantum scabris.
8. INULA foliis lanceolatis semiamplexicaulibus hispida hirta. dis, caulibus subunifloris teretibus.
 Aster foliis lanceolatis sessilibus integerrimis, caulibus simplicibus unifloris, squamis calycinis subulatis. *Dalib. parif.* 259.
 Aster luteus, hirsuto salicis folio. *Bauh. pin.* 266.
 Aster tertius pannonicus clusii luteus, folio hirsuto salicis. *Bauh. hist.* 2. p. 1049.
Habitat in Bavaria, Gallia, Geneva. & Habitus Inulæ salicinæ, sed folia latiora, obtusiora, vix manifesto serrata, venosa, scabra præsertim margine & carina. Caule foliisque pilis rigidiusculis adpersis, ubique teretibus, nec sulcato-angulatis.
9. INULA foliis lanceolatis recurvis subdentatis scabris, germanica, floribus subsæscuiatis.
 Aster foliis lanceolatis amplexicaulibus oris reflexis, ramis multifloris, calycibus oblongis laxis. *Gmel. fib.* 2. p. 181. t. 78. f. 1
 Conyzæ affinis germanica. *Bauh. pin.* 266. *Morif. hist.* 3. f. 7. t. 19. f. 26.
 Aster thuringiacus altissimus latifolius montanus, flore luteo parvo. *Hull. jen.* 181.
Habitat in Misnia, Pannonia, Sibiria. & Habitus J. Salicinæ. Folia eadem; at Flores cylindrici, conferti ad apicem caulis in corymbum.
10. INULA foliis linearibus acuminatis nervosis glabris ensifolia. sparsis, caule subunifloro.
 Aster luteus, linearæ rigido glabro folio. *Bauh. pin.* 267.
Habitat in Austria inferiore &
11. INULA foliis linearibus carnosissimis tricuspidatis. *crithmoides.*
 Aster flore terminali, foliis linearibus tricuspidatis. *Hort. cliff.* 409. *Roy. lugdb.* 168. *Guss. stamp.* 2. p. 464.
 Crithmum maritimum, colore asteris artici. *Bauh. pin.* 288 K k k 2 Crith-

884 SYNGENESIA: POLYGAM. SUPERFLUA.

Crithmum chrysanthemum. Dod. pempt. 706.

Habitat in Angliæ, Galliæ, Lusitanæ, Hispaniæ maritimis. &

provincialis. 12. *INULA* foliis subserratis subtus tomentosis: radicalibus petiolatis ovatis, caule erecto unifloro.

Jacobææ rotundifolia incana. Baub. pin. 131. prodr. 66. Burf. VI: 127.

Habitat in Collendaneis Galloprovinciæ. &

Radix fibris validis, Caulis pedalis, simplicissimus, erectus, villosus. Folia Radicalia ovata, petiolata, obtuse serrata, subtus villosa, obtusiuscula; Caulina paucica, lanceolata, sessilia, minus serrata, subtus villosa: summa linearia. Flos sat magnus, flavus.

montana. 13. *INULA* foliis lanceolatis pubescentibus integerrimis, caule unifloro.

Aster foliis lanceolatis sessilibus villosis, caule subunifloro simplicissimo. Roy. lugdb. 168.

Aster montanus, luteo magno flore. Baub. pin. 267.

Morif. bist. 3. p. 119. f. 7. t. 21. f. 12.

Habitat in Alpibus Helvetiæ, Valtellinæ. &

851

ARNICA.

montana. 1. *ARNICA* foliis ovatis integris: caulinis geminis oppositis.

Doronicum foliis caulinis oppositis. Fl. suec. 684. Mas. med. 395. Gmel. sib. 2. p. 153.

Doronicum foliis oblongo-ovalibus Fl. lapp. 304. Roy. lugdb. 160.

Doronicum, plantaginis folio, alterum. Baub. pin. 189.

Alisma. Matth. disc. 934.

Diuretica. Renss. spec. 118.

alpina. β. *Doronicum* foliis lanceolatis. *Fl. lapp. 305. Roy. lugdb. 160.*

Caltha alpina. Tabern. ic. 336.

Habitat in Alpibus & pratis Europæ frigidioris. &

scorpioides. 2. *ARNICA* foliis alternis serratis. *Hall. helv. 737. †*

Doronicum radice scorpii brachiata. Baub. pin. 184.

Doronicum brachiata radice. Dalech. bist. 1203. Lob. ic. 649.

β. *Doronicum* radice dulci. *Baub. pin. 184.*

Habitat in Helvetia. &

maritima. 3. *ARNICA* foliis lanceolatis: inferioribus serratis, caule folioso multifloro. †

Aste

SYNGENESIA: POLYGAM. SUPERFLUA. 885

After f. potius *Helenium maritimum fruticosum*, caule crasso, foliis fere *helenii officinalis*. *Gmel. fib. 2.*

p. 175.

Habitat in Kamtschatka & America septentrionali.

4. ARNICA foliis ovatis denticulatis.

crocea.

Gerbera foliis planis dentatis, flore purpureo. Burm.

af. 157. t. 56. f. 2.

Dens leonis, caulæ follo. Pet. mus. 393. Raj. suppl. 238.

Doronici forte species pumila, auriculæ urfi folio glabro, flore saturate croceo. Pluk. mant. 65.

Habitat in Æthiopia. &

5. ARNICA foliis pinnatifidis; lobis rotundatis.

Gerbera.

Gerbera foliis ex radice oblongis excavatis sinuosis subtus lanugine ruffa villosis. Burm. afr. 155. t. 56. f. 1.

After africanus monanthos, scolopendriæ folio, flore luteo. Raj. suppl. 161.

Doronicum africanum, folio asplenii. Buxb. cent. 5. t. 12. f. 24.

Scolopendriæ f. Asplenii facie planta æthiopica, summis pinnis acuminatis sericea lanugine ex ruffo candicante villosa. Pluk. alm. 336. t. 313. f. 5.

Habitat in Æthiopia.

6. ARNICA foliis pinnatis; laciniis linearibus.

coronopifolia.

Habitat in Æthiopia.

Præcedenti minis affinis.

DORONICUM. 852

1. DORONICUM foliis cordatis obtusis; radicalibus petiolatis; caulinis amplexicaulibus. *Mat. med. 394. pardaliancher.*

Doronicum foliis cordatis denticulatis, caule ramoso.

Hort. cliff. 411. Roy. lugdb. 159.

Doronicum maximum, foliis caulem amplexantibus.

Bauh. pin. 184.

Doronicum VII, austriacum 3. Clus. hist. 2. p. 19.

β. Doronicum radice scorpii. *Bauh. pin. 184.*

Doronicum latifolium. Clus. hist. 2. p. 16.

Aconitum Pardalianches. Dod. purg. 305.

Habitat in Alpibus Helvetiæ, Pannoniæ, Valleisæ. &

2. DORONICUM foliis ovatis acutis subdentatis, ramis alternis. *Hort. cliff. 411. Roy. lugdb. 160. Da- plintagine- win.*

lib. parif. 256.

K k k 3

Do-

886 SYNGENESIA: POLYGAM. SUPERFLUA.

Doronicum plantaginis folio *Bauh. pin.* 184.

Doronicum minus officinarum *Dalech. hist.* 1202.

β. *Doronicum*, plantaginis folio. lusitanicum. *Tournef. inst.* 488.

Habitat in Lusitania, Hispania, Gallia. 2

incanum.

3. *DORONICUM* foliis lanceolatis denticulatis subtus tomentosis, caule unifloro. *Roy. lugdb.* 160. *Sauv. meth.* 84. †

Doronicum helveticum incanum. *Bauh. pin.* 185. *prodr.* 97. *Scheuch. alp.* 33. & 333.

Habitat in Alpibus Helveticis, Pyrenæis. 2

An etjam semina radii in hoc & sequente papposa?

Bellidiastrum.

4. *DORONICUM* caule nudo simplicissimo unifloro. *Hort. cliff.* 500. *Roy. lugdb.* 160. †

Bellidiastrum alpinum, foliis brevioribus hirsutis, caule palmari, flore albo. *Mich. gen.* 32. t. 29.

Bellis sylvestris media, caule carens. *Bauh. pin.* 261.

β. *Bellis* caule pedali s. olpedali nudo, foliis magnis latis, floribus rubris & albis. *Menz. pug.* t. 8.

Habitat in Alpibus Helveticis, Italicis, Tyrolensibus. 2

853

HELENIUM.

autumnale.

1. *HELENIUM.*

Helenium foliis decurrentibus. *Hort. cliff.* 418. *Hort. upf.* 266. *Gron. virg.* 101. *Roy. lugdb.* 180.

Chrysanthemum americanum perenne, caule alato, folio angustato glabro. *Morif. hist.* 3. p. 24. f. 6. t. 6. f. 74.

Aster floridanus aureus, caule alato. *Pluk. amalsb.* 43. t. 372. f. 4.

Aster luteus alatus, *Corn. canad.* 52. t. 63.

Habitat in America septentrionali. 2

854

BELLIS.

perennis.

1. *BELLIS* scapo nudo.

Bellis scapo nudo unifloro. *Hort. cliff.* 418. *Hort. upf.* 265. *Fi. suec.* 707. *Mat. med.* 405. *Roy. lugdb.* 177.

Dalib. parif. 264.

Bellis sylvestris minor. *Eaub. pin.* 267.

Bellis sylvestris. *Dod. pempt.* 265.

hortensis.

β. *Bellis hortensis*, flore pleno. *Bauh. pin.* 261.

γ. *Bellis hortensis*, flore albo bullato *Tournef. inst.* 491.

δ. *Bellis hortensis rubra*, flore multiplici fistuloso. *Tournef. inst.* 491.

ε. *Bel-*

SYNGENESIA: POLYGAM. SUPERFLUA. 887

- ε. *Bellis hartenfis prolifera.*
Habitat in Europæ apricis pascuis. &
2. **BELLIS** caule subfoliofo. annua.
Bellis minima pratensis, caule folioso. Bocc. mus. 2. p.
96. t. 35.
Bellis minima annua. Triumph. obs. 80. t. 80.
Bellis minima, caule folioso pubescens. Cup. cath.
Bellis-Leucanthemum annuum italicum. Michx. gen. 34.
Habitat in Sicilia, Hispania, Monspelii. ☉

TAGETES. 855

1. **TAGETES** caule subdiviso diffuso. *Hort. cliff. 418. patula.*
Hort. nps. 267. Roy. lugdb. 183.
Tanacetum africanum f Flos africanus minor. Bauh.
pin. 132.
Flos africanus. Dod. pempt. 255.
- β. **Tagetes indicus minor, multiplicato flore. Tournes. inf.**
488.
Tagetes minor, flore fulvo maculato. Dill. elth. 273.
t. 279. f. 361.
Habitat in Mexico. ☉
2. **TAGETES** caule simplici erecto, pedunculis nudis u-erecta.
nifloris. Hort. cliff. 418. Hort. nps. 267. Roy. lugdb. 180.
Tanacetum africanum majus, simplici flore. Bauh.
pin. 133.
Cariophyllus indicus. Cam. epit. 406.
- β. **Tagetes maximus rectus, flore maximo multiplicato.**
Bauh. hist. 3 p. 100.
- γ. **Tagetes mexicanus. flore fistuloso, simplex. Col. ecphr.**
2. p. 47. t. 46. f. 1.
- δ. **Tagetes mexicanus alter polyanthos, fistuloso flore.**
Col. ecphr. 2 p. 48. t. 46.
Habitat in Mexico. ☉
3. **TAGETES** caule simplici recto, pedunculis squamo-minuta.
sis multifloris. Hort. cliff. 419.
Tagetes multiflora, minuto flore albicaute. Dill. elth. 374.
t. 280. f. 362.
Habitat in Chili. ☉

CHRYSANTHEMUM. 856

* *Leucanthema.*

1. **CHRYSANTHEMUM** fruticosum, foliis linearibus frutescens.
dentato-trifidis. Hort. cliff. 417. Roy. lugdb. 174.
K k k 4 Leu-

888 SYNGENESIA: POLYGAM. SUPERFLUA.

Leucanthemum canariense, sapore pyrethri. *Walzb. hort.* 31. t. 24.

Chamæmelum canariense ceratophyllum fruticosus. *Morif. hist.* 3. p. 35.

Bellis canariensis frutescens, foliis crassis pyrethri sapore. *Raj. suppl.* 221.

Buphthalmum canariense leucanthemum, cotulae foetidae crassioribus foliis. *Pluk. alm.* 73. t. 272. f. 6.

Habitat in Canariis insulis. 5

Strothnum.

2. **CHRYSANTHEMUM** foliis lanceolatis superne serratis utrinque acuminatis. *Hort. cliff.* 416. *Roy. lugdb.* 174. id. *Scan. append.*

Aster, foliis profunde dentatis & quasi laciniatis, ramosus. *Raj. suppl.* 162.

Bellis major, radice repente, foliis latioribus serratis. *Morif. hist.* 3. p. 29. f. 6. t. 9. f. 11. *bles.* 239. *Raj. hist.* 351.

Bellis americana frutescens ramosa. *Raj. hist.* 1865.

Bellis americana procerior serotina ramosa, flore amplissimo. *Pluk. alm.* 65. t. 17. f. 2.

Habitat forte in America septentrionali? cum sero flore. 2

Leucanthemum.

3. **CHRYSANTHEMUM** foliis amplexicaulibus oblongis: superne serratis, inferne dentatis. *Hort. cliff.* 416. *Fl. succ.* 700. *Mat. med.* 404. *Roy. lugdb.* 174. *Dalib. parif.* 261.

Chrysanthemum foliis oblongis serratis. *Fl. lapp.* 310.

Bellis sylvestris, caule folioso, major. *Bamb. pin.* 261.

Bellis major. *Fuchf. hist.* 148. *Cam. epit.* 635.

Habitat in pratis Europæ. 2

montanum.

4. **CHRYSANTHEMUM** foliis imis spatulato-lanceolatis serratis, summis linearibus. *Sauv. monsp.* 87. †

Leucanthemum montanum minus. *Tournef. inst.* 492.

Bellis montana minor. *Bamb. hist.* 3. p. 115. *Magn. monsp.* 36.

Habitat Monspelii.

graminifolium.

5. **CHRYSANTHEMUM** foliis linearibus subintegerrimis. *Sauv. monsp.* 87.

Leucanthemum gramineo folio. *Tournef. inst.* 493.

Bellis montana, gramineis foliis. *Magn. monsp.* 291. * *Hort.* 31. t. 31.

Habitat Monspelii. 2

6. **CHRY-**

SYNGENESIA: POLYGAM SUPERFLUA. 889

6. CHRYSANTHEMUM foliis imis palmatis: fo-
liolis linearibus pinnatifidis. *Sanv. monsp.* 304.
Bellis montana major, foliis chrysanthemi cretici angu-
stioribus. *Magn. monsp.* 306. *
Leucanthemum montanum, foliis chrysanthemi. *Tour-
nef. inst.* 492.
Habitat Monspeli. 2
7. CHRYSANTHEMUM foliis pinnatifidis: lacinis alpinum,
parallelis integris, caulibus unifloris.
Chrysanthemum foliis linearibus pinnatifidis. *Sanv.
monsp.* 304?
Chamæmelum alpinum. *Baub. pin.* 136.
Leucanthemum alpinum, foliis coronopi. *Tournef. inst.*
493?
Habitat in Alpibus Helvetiæ ad Thermas Piperinas.
* *Chrysanthema.*
8. CHRYSANTHEMUM foliis simplicibus ovatis si-
nuatis angulatis serratis acutis.
Matricaria sinensis, minore flore, petalis & umbone o-
chroleucis. *Pluk. alm.* 142. t. 430. f. 2.
β. Chrysanthemum maderaspatanum, oxyacanthæ foliis
cæsis ad marginem spinosis, calyce argenteo. *Pluk.
alm.* 101. t. 160. f. 6.
Matricaria indica, latiore folio, flore pleno. *Maris. hist.*
3. p. 33.
Matricaria sinensis, flore monstrosa. *Vaill. act.* 1720.
p. 368. *Fl. zeyl.* 421. *
Matricaria zeylanica hortensis, flore pleno. *Raj. suppl.*
224.
Habitat in India.
9. CHRYSANTHEMUM foliis simplicibus cuneifor-
mibus subpalmatis multifidis obtusis,
Pyrethrum foliis longe petiolatis palmatis supra dilata-
tis: ultimis lacinis trilobis. *Gmel. sib.* 2. p. 203. t. 84.
Habitat in Kamtschatka, America septentrionali.
Caulis debiles, diffusi, herbacei, inferne ramosi. Folia
petiolis longis, subcarnea, lævia, cuneiformia, triloba-
ta, sinuato-angulata, obtusa. Flores terminales, soli-
tarii, magnitudine C. coronarii, Pedunculo subvillo-
so.
10. CHRYSANTHEMUM foliis amplexicaulibus: su-
perne laciniatis; inferne dentato-serratis. *Hort. cliff.*
416. *Fl. juv.* 699. *Roy. lugdb.* 174. *Dalib. paris.* 261.
K k k 5 Chry-

890 SYNGENESIA: POLYGAM. SUPERFLUA.

Chrysanthemum segetum. *Clus. hist.* 1. p. 334.
Chrysanthemum segetum vulgare glaucum. *Morif. hist.*
 3. p. 15. f. 6. t. 4. f. 1.
Bellis lutea, foliis profunde incis, major. *Bauh. pin.* 262.
Habitat in Scaniae, Germaniae, Belgii, Angliae, Galliae
agris. ☉

corymbosum. 11. **CHRYSANTHEMUM** foliis pinnatifidis incis ex-
 trorsum lationibus. *Hort. cliff.* 416. *Hort. upf.* 263.
Roy. lugdb. 174.

Chrysanthemum foliis matricariae. *Bauh. pin.* 134.
Chrysanthemum creticum. *Clus. hist.* 1. p. 334. *Morif.*
hist. 3. p. 16. f. 6. t. 4. f. 2. 3.
Habitat in Creta, Sicilia. ☉

corymbosum. 12. **CHRYSANTHEMUM** foliis pinnatis inciso-fer-
 ratis, caule multifloro. *Roy. lugdb.* 174. *Savv.*
monsp. 267.

Tanacetum montanum inodorum, minore flore. *Bauh.*
pin. 132.

Tanacetum leucanthemum. *Tabern. hist.* 379.

β. *Tanacetum inodorum, flore majore.* *Bauh. pin.* 132.
Habitat in Thuringia, Bohemia, Helvetia, Sibiria. &

bipinnatum. 13. **CHRYSANTHEMUM** foliis bipinnatis ferratis vil-
 lotis.

Pyrethrum foliis duplicato-pinnatis: pinnulis incis, pe-
dunculis unifloris. *Emel. fib.* 2. p. 205. t. 85. f. 1.
Habitat in Sibiria. &

Habitat Anthemidis apricae. Folia bipinnata, pinna-
lis sublaevigatis, hirsutis, fere Tanacetii. Peduncu-
li axillares, nudi, uniflori, pubescentes, longitudine sa-
liorum. Flores lutei: radio disco longe brevior.

densiflorum. 14. **CHRYSANTHEMUM** flosculis omnibus unifor-
 nibus hermaphroditis. *Hort. cliff.* 417. *Roy. lugdb.*
 174.

Bellis major spinosa, petalis carens s. nuda. *Morif. hist.*
 3. p. 29. f. 6. t. 9. f. 16.

Bellis spinosa, foliis agerati. *Bauh. pin.* 262.

Bellis spinosa. *Alp. exos.* 327. t. 326.

Habitat in Africa.

MATRICARIA.

857
Parthenium. 1. **MATRICARIA** foliis compositis planis: foliolis o-
 vatis incis, pedunculis ramosis. *Hort. cliff.* 416.
Hort. upf. 263. *Mat. med.* 400. *Roy. lugdb.* 173.

Matri-

SYNGENESIA: POLYGAM. SUPERFLUA. 891

Matricaria vulgaris f. *fativa*. *Bauh. pin.* 133.

Matricaria. *Dod. pempt.* 35.

Habitat in Europæ cultis, ruderalis. 2. 8.

2. **MATRICARIA** receptaculis hemisphæricis, foliis bi-*maritima*.
pinnatis subcarnosis: supra convexis, subtus carinatis.

Chamæmelum maritimum perenne humilius, foliis brevioribus crassis obscure virentibus. *Raj. angl.* 3. p. 186.

t. 7. f. 1. *It. wgoth.* 148. *

Habitat in Europæ septentrionalis littoribus maris.

3. **MATRICARIA** receptaculis hemisphæricis, radiis *Chamomilla*,
patentibus, seminibus coronato marginatis, squamis
calycinis margine obsoletis. *Fl. suec.* 702. *Mat. med.*

402. *Dalib. parif.* 262. *Gmel. sib.* 2. p. 200.

Matricaria foliis supradecompositis setaceis, pedunculis
solitariis. *Hort. cliff.* 415. *Roy. lugdb.* 173.

Chamæmelum vulgare f. *Leucanthemum*, *dioscoridis*.

Bauh. pin. 135.

Anthemis vulgaris. *Lob. ic.* 770. *Fl. lapp.* 309.

Habitat in Europæ agris, cultis. ☉

4. **MATRICARIA** foliis bipinnatis, pedunculis solitariis, *argentea*,
Hort. cliff. 415. * *Roy. lugdb.* 173.

Matricaria monoleucanthemos, foliis argenteis plerumque
conjugatis. *Vaill. act.* 1720. p. 369.

Chamæmelum orientale incanum, millefolii folio. *Tournef. cor.* 37.

Habitat in Oriente.

5. **MATRICARIA** receptaculis conicis, radiis flexis, *verutina*,
seminibus nudis. squamis calycinis margine æqualibus. *Fl. suec.* 701. (non synonyma) *Gmel. sib.* 2.
p. 200.

Matricaria leucanthemos annua suaveolens, chamæmeli
folio, ovariis albicantibus. *Vaill. parif.* 1720.

Chamæmelum tenue incisum. foliis floris deorsum pro-
tensis: umbone extuberante, suavius olens. *Cup. cat-
bol. suppl.* 17.

Habitat in Europa. ☉

COTULA. 858

1. **COTULA** foliis pinnato-multifidis, corollis radio *anthemoides*
destitutis. *Hort. cliff.* 417. *Roy. lugdb.* 173.

Ananthocyclus chamæmeli folio V. *Dill. elib.* 26. t.

23. f. 25

Chry-

892 SYNGENESIA: POLYGAM. SUPERFLUA.

Chrysanthemum exoticum perpassillum nudum, foliis coronopi. Pluk. alm. 101. t. 274. f. ult.

Habitat in Insula Helenæ, Hispania.

Conf. Chamæmelum luteum, capite aphylo. Bæd. pin. 135. Annon huic propius, quam Anthemidi nobili?

- coronopifolia* 2. COTULA foliis lanceolato-linearibus amplexicaulibus pinnatifidis. *Hort. cliff. 417. Hort. upf. 266. Roy. lugdb. 173. Moehr. E. N. C. 1742. vol. 6. p. 298.*
Ananthocyclus coronopi folio V. Dill. elch. 27. t. 23. f. 26.
Chrysanthemum exoticum minus, capite aphylo, chamæmelli nudi facie. Breyn. cent. 156. t. 76.
Bellis annua, capite aphylo luteo, coronopi folio, caulibus procumbentibus. Herm. lugdb. 86. Moris. hist. 3. p. 30. f. 6. t. 6. f. ult.
Habitat in Æthiopia, nunc in Frisiæ inundatis prope Emdon. Mahringio teste. ☉
- viscosa.* 3. COTULA foliis lyrato-pinnatis, corollis radiatis. *Hort. cliff. 417. **
Jacobæa americana odorata viscosa, florum radiis brevissimis albis. Honst. nass.
Habitat in Vera Cruse.
- turbinata.* 4. COTULA receptaculis subtus inflatis turbinatis. *Hort. cliff. 417. Hort. upf. 266. Roy. lugdb. 173.*
Chamæmelum æthiopicum lanuginolum. Breyn. cent. 148. t. 73. Moris. hist. 3. p. 36. f. 6. t. 12. f. 14.
Habitat in Æthiopia. ☉

ANACYCLUS.

- 859
creticus. 1. ANACYCLUS foliis decompositis linearibus: laciniis divisis planis. *Hort. cliff. 417. Roy. lugdb. 171.*
Cotula cretica minima, chamæmeli folio, capite inflexo. Tournef. cor. 37.
Habitat in Creta.
- orientalis.* 2. ANACYCLUS foliis compositis setaceis acutis rectis. *Hort. cliff. 417. Roy. lugdb. 171.*
Chamæmelum orientale, foliis pinnatis. Tournef. cor. 37. Boerb. lugdb. 1. p. 110. t. 110.
Habitat in Oriente.
- valentinus.* 3. ANACYCLUS foliis decompositis linearibus: laciniis divisis teretiusculis acutis, floribus flosculosis. *Hort. cliff. 417. Roy. lugdb. 171.*
Chrysanthemum valentinum. Clus. hist. 1. p. 332.
Euph-

SYNGENESIA: POLYGAM. SUPERFLUA. 893

Buphthalmo tenuifolio simile. *Bauh. bist.* 3. p. 125.
Habitat ad Reg. Valentini agros & vias,
Confer Anthemidem valentinam.

ANTHEMIS. 860

* *Radio dicoloro f. albo.*

1. ANTHEMIS florum paleis rigidis pungentibus. *Roy. Cota. lugdb.* 172.

Anthemis italica arvensis annua major vulgarissima, flore maximo disco pungente. Mich. gen. 32.

Chamæmelum annuum ramosum, cotulæ foetidæ floribus amplioribus, capitulis spinosis. Moris. bist. 3. p. 36. f. 6. t. 8. f. 11. pro. 12.

Habitat in Italiæ arvis.

2. ANTHEMIS erecta, foliorum apicibus subspinosis. *altissima,*

Anthemis foliorum serraturis setaceis, radiis florum albis. Sauv. monsp. 265.

Chamæmelo affine Buphthalmum italicum segetum altissimum. Bauh. bist. 3. p. 120.

Chamæmelum Leucanthemum hispanicum, magno flore. Bauh. pin. 135. prodr. 70.

Habitat in Italiæ, Hispaniæ, G. Narbonensis agris. D. Sauvages. ☉

Caulis multiflorus, segete fere altior, striatus, erectus. Pedunculi superne parum incrassati. Calyces glabri. Paleæ latæ, subtruncatæ, terminatæ seta radicales longitudine calycis. Folia multifido-composita dentibus acuminatis: acuminibus pallidis, quasi spinosis.

3. ANTHEMIS foliis pinnatis denticulatis carnosis, caule ramoso. *maritima,*

Anthemis maritima perennis, foliis crassis & punctatis.

Mich. gen. 33.

Matricaria maritima. Bauh. pin. 134.

Chamæmelum maritimum. Bauh. bist. 3. p. 122.

Habitat Monspeliū, inque Italia.

4. ANTHEMIS foliis pinnatifidis obtusis planis, pedunculis hirsutis foliosis, calycibus tomentosis. *tomentosa,*

Hort. cliff. 415. * *Roy, lugdb.* 172.

Chamæmelum, coronopi folio, tomentosum. Vaill. acb.

1720. p. 412.

Chamæmelum maritimum incanum, folio absinthii crasso. Boerb. lugdb. 1. p. 110.

Habitat in Græciæ maritimis.

J. AN

mixta.

5. **ANTHEMIS** foliis simplicibus dentato-laciniatis.
Anthemis foliis singularibus pinnato-dentatis. *Gætt. stamp.* 146.
Anthemis maritima lanuginosa annua, foliis crassis pyrethri sapore, semistofculis albis juxta ditcum luteis. *Mieb. gen.* 32. t. 30. f. 1.
Chamæmelum annuum ramosum, coronopi folio, flore mixto. *Morif. hist.* 3. p. 36. f. 6. t. 18. f. 15.
Chamæmelum iustanicum latifolium, f. coronopi folio. *Breyn. cent.* 149. t. 74.
Habitat in Italia, Gallia. ☉
Folia non composita, sed laciniata, Radius florum albus, basi luteus.

chia.

6. **ANTHEMIS** foliis pinnatifidis laciniatis, pedunculis nudis subvillosis. *Hort. cliff.* 415. * *Roy. lugdb.* 172.
Habitat in Chio.

nobile.

7. **ANTHEMIS** foliis pinnato-compositis linearibus acutis subvillosis.
Anthemis foliis pinnato-decompositis: laciniis setaceis. *Hort. cliff.* 415. *Roy. lugdb.* 172.
Matricaria receptaculis conicis, radiis deflexis, seminibus nudis, squamis calycinis margine æqualibus. *Fl. succ.* 701. *Mat. med.* 401. *Dalib. parif.* 262.
Chamæmelum nobile f. *Leucanthemum* odoratus. *Baub. pin.* 135.
Chamæmelum odoratum. *Dod. pempt.* 270.
 8. *Chamæmelum* nobile, flore multiplici. *Baub. pin.* 135.
Chamæmelum romanum, flore multiplici. *Cam. epis.* 646.
Chamæmelum chium vernum. folio crassiore, flore magno. *Tournef. cor.* 37. *Roy. lugdb.* 172.
Habitat in Europæ pascuis apricis. 2

arvensis.

8. **ANTHEMIS** receptaculis conicis, paleis setaceis, seminibus coronato-marginatis. *Fl. succ.* 704. * *Dalib. parif.* 263.
Anthemis arvensis. *It. scan.* 177.
Chamæmelum inodorum. *Baub. pin.* 186? *Raj. angl.* 3. p. 185? *Rupp. jen.* 173.
Habitat in Europæ, præsertim Sueciæ agris. 8
Planta canescens. Caulis diffusus. Paleæ lanceolatae.

cotula.

9. **ANTHEMIS** receptaculis conicis: paleis setaceis, seminibus nudis. *Fl. succ.* 703. *Mat. med.* 403 *Dalib. parif.* 263.
Anthemis foliis pinnato-decompositis: laciniis linearibus, caulibus diffusis, radice annua. *Roy. lugdb.* 172.

Cha-

SYNGENESIA: POLYGAM. SUPERFLUA. 895

Chamæmelum foetidum. Baub. pin. 135.

Chamæmelum foetidum f. Cotula foetida Baub. hist. 3. p. 36.

*Habitat in Europæ uderatis, præcipue in Ucraina. ☉
Palea in hac minutissima sunt.*

10. **ANTHEMIS** foliis pinnato-multifidis planis: Jaci-
niis linearibus acutis trifidis, pedunculis, longissimis.
*Hort. cliff. 415. * Roy. lugdb. 172.*

*Chamæmelum orientale, foliis absinthii. Tournef. ser. 37.
Habitat in Creta:*

11. **ANTHEMIS** foliis simplicibus ovato-lanceolatis re-
pando-crenatis.

*Chrysanthemum lusitanicum, agerati folio. Tournef.
inst. 492.*

Habitat in Hispania.

Caules simpliciusculi, erecti. Folia ovato-oblonga, repando-crenata, petiolata, alterna. obtusa, vix villosa. Flores terminales, subsolitarii, globosi; Calyx minus imbricatus. Radius angustus; discus convexus flavus. Semina tetragona, coronata squamis quatuor, quarum due in fetas desinunt.

12. **ANTHEMIS** caulibus simplicibus unifloris decum-
bentibus, foliis pinnato-multifidis. *Hort. cliff. 414.
Hort. ups. 263. Mat. med 396. Roy. lugdb. 170.
Sauv. monsp. 265.*

*Bupthalmum creticum, cotulæ facie, flore luteo & al-
bo. Breyn. cent. 150. t. 75.*

*Chamæmelum specioso flore, radice longa fervida.
Schaw. afr. 138.*

Pyrethrum flore bellidis. Baub. pin. 148.

Habitat in Arabia, Syria, Creta, Apulia, Thuringia. 2

** Radio concolore f. luteo.*

13. **Anthemis** caule ramoso, foliis pinnato-multifidis seta
ceis, calycibus villosis pedunculatis. *Hort. cliff. 414.*

Bupthalmum cotulæ folio. Bauh pin. 134.

*Bupthalmum tenuifolium, folio fere millefolii. Baub.
hist. 3. p. 124.*

Bupthalmum alterum penæ. Dalech. hist. 863.

Habitat in G. Narbonensi

Affinis admodum Anacyclo valentino.

14. **ANTHEMIS** foliis trternatis. pedunculis terminali-
bus ramo longioribus. *Hort. cliff. 414.*

Chry-

500

896 SYNGENESIA: POLYGAM. SUPERFLUA.

Chrysanthemum palustre minimum repens, apii folio.
Sloan. jans. 126. *hist.* 1. p. 263. t. 155. f. 3. *Raj.*
suppl. 215.

Chrysanthemum humile, ranunculi folio. *Plum. spec.* 10.
Habitat in Jamaicae, Caribæarum pratis humidiusculis. ☉

- Millefolia.* 15. *ANTHEMIS* foliis bipinnatis: laciniis linearibus integris, pedunculis nudis longissimis. *Roy. lugdb.* 171.
Chamæmelum montanum, absinthii folio, parthenii odore. *Vaill. act.* 1720. p. 319.
Leucanthemum, folio absinthii, alpinum ciansi. *Boerb. lugdb.* 1. p. 107.
Pyrethrum foliis duplicato-pinnatis: pinnulis incisfis, pedunculis unifloris, caule procumbente. *Gmel. fib.* 2, p. 207. t. 86. f. 1. 2.
Habitat in Sibiria.
Habitus Millefolii. Folia *Millefolii*, sed paulo majora. Pedunculi in summis ramis, alterni, longissimi. Flores *Anthemidis aprieae*, sed parvi lutei.

- tinctoria.* 16. *ANTHEMIS* foliis bipinnatis serratis subtus tomentosis, caule corymbose.
Bupthalmum caule ramoso, foliis pinnatifidis: laciniis linearibus dentatis serratis, floribus pedunculatis. *Hort. cliff.* 414. *Fl. succ.* 698. *Roy. lugdb.* 171.
Bupthalmum tanacetii minoris foliis. *Bamb. piv.* 134.
Chrysanthemum foliis tanacetii. *Loef. pruss.* 47. t. 9.
aciumfeli. β. *Bupthalmum alpinum*, flore candido. *Trinms. obs.* 79. 1. 80.

Habitat in Sueciae, Germaniae apricis pratis siccis. 2
Planta β. differt radio perfecte explicato albo, basi tomentum inteo; reliqua conveniunt ut vix species diversa ernatur. Calycis squama interiores utrinque apice ciliata.

- arabica.* 17. *ANTHEMIS* caule decomposito, calycibus ramiferis. *Hort. cliff.* 413. t. 24. * *Hort. wps.* 264. *Roy. lugdb.* 170.
Asteriscus annuus trianthophorus Crassas arabibus dictus. *Shav. afr.* 58. t. 36. f. 58.
Habitat in Arabia. ☉

ACHILLEA.

* *Corollis flavis.*

- Sanctolina.* 1. *ACHILLEA* foliis setaceis dentatis: denticulis subintegris subulatis reflexis, *Hort. cliff.* 412.

Achil-

SYNGENESIA: POLYGAM. SUPERFLUA. 897

Achillea lutea tomentosa, santolinæ folio. Vaill. act.
1720. p. 417.

Ptarmica orientalis, santolinæ folio, flore majore. Tour-
nesf. cor. 37.

Habitat in Oriente. 2

2. **ACHILLEA** foliis lanceolatis obtusis acute serratis. *Ageratum,*
Hort. cliff. 413. Hort. ups. 265. Mas. med. 399.

Roy. lugdb. 176. Sauv. monsp. 88.

Ageratum foliis serratis. Bauh. pin. 221.

Balsamita minor. Dod. pems. 295. Moris. hist. 3. p.
38. f. 6. t. 1. f. 2.

Habitat in G. Narbonensi, Hetruria. 2

3. **ACHILLEA** foliis linearibus dentatis obtusis *plafalcata,*
nis: denticulis crenatis. Hort. cliff. 412. Roy. lugdb.
176.

Achillea incana, santolinæ foliis plerumque falcatis a-
speris, flore sulphureo. Vaill. act. 1720. p. 417.

Habitat in Oriente. 2

4. **ACHILLEA** foliis pinnatis hirsutis: pinnis linearibus *tomentosa,*
dentatis.

Achillea foliis linearibus pinnatitidis pubescentibus,
foliolis tripartitis transversalibus. media productiore.

Hort. cliff. 413. Hort. ups. 265. Roy. lugdb.
175.

Millefolium tomentosum luteum. Bauh. pin. 140.

Stratiotes millefolia, flavo flore. Clus. hist. 1. p. 330.

Habitat in G. Narbonensi, Valleria, Tararia. 2

5. **ACHILLEA** foliis pinnatis: foliolis lanceolatis incis *pubescens,*
serratis subtus lanigeris. Hort. cliff. 413. Roy. lugdb.
175.

Millefolium orientale, foliis tanacetii incanis, radiis palli-
de luteis. Boerb. lugdb. 1. p. 112.

Ptarmica orientalis, foliis tanacetii incanis, semifloscu-
lis florum pallide luteis. Tournef. cor. 37.

Matricaria tomentosa & incana, achilleæ folio, flore
aureo. Vaill. act. 1722. p. 286.

Habitat in Oriente. 2

6. **ACHILLEA** foliis pinnatis supradecompositis: laciniis *abrotanifolia*
linearibus distantibus. Roy. lugdb. 175.

Millefolium orientale altissimum luteum, abrotani fo-
lio. Tournef. cor. 37.

Habitat in Oriente.

* *Cerollis radio albis.*

macrophylla. 7. *ACHILLEA* foliis pinnatis planis inciso-serratis: extremis majoribus.

Achillea foliis bipinnatis: pinnaulis linearibus indivisis. *Sauv. meth.* 267.

Achillea foliis pinnatis, pinnis serratis maximis: extremis confluentibus. *Hall. helv.* 713.

Dracunculus alpinus, totius scabiosæ. *Bauh. pin.* 98. *prodr.* 39.

Ptarmica alpina, matricariæ foliis. *Trumf. obs.* 83. t. 23. *Bocc. mus.* 2. p. 150. t. 110. *Barr. rar.* 1119. t. 991.

Corymbifera millefolii umbella, folio alato & laciniato. *Roj. hist.* 345.

Habitat in Alpibus Helvetiæ, Italiæ.

Folia magnitudine Tanacetii.

impatiens. 8. *ACHILLEA* foliis pinnatis: pinnis distantibus linearilanceolatis: basi sursum acutis. *Roy. lugdb.* 176.

Achillea foliis pinnatis, pinnis longis acutis: inferioribus sursum dente auctis glaberrimis. *Gmel. sib.* 2. p. 197. t. 83. f. 1.

Habitat in Sibiria.

Planta albis gaudet floribus. Foliorum pinna remota et admodum distantes, distinguunt plantam a reliquis.

Clavenna. 9. *ACHILLEA* foliis pinnatifidis planis obtusis tomentosis.

Achillea foliis sinuato-laciniatis planis villosis nitidis obtusis. *Hort. cliff.* 412. *Roy. lugdb.* 176.

Abinthium alpinum umbelliferum latifolium. *Bauh. pin.* 139.

Abinthium alpinum umbelliferum. *Clus. hist.* 1. p. 340.

Dracunculus argenteus. *Morif. hist.* 3. p. 40. f. 6. t. 10. f. 5.

Habitat in Alpibus Austriæ, Pannoniæ, Corinthiæ. &

Ptarmica. 10. *ACHILLEA* foliis lanceolatis acuminatis argute serratis.

Achillea foliis integris minutissime serratis. *V. Hort. cliff.* 412. *Hort. ups.* 265. *Fl. succ.* 706. *Mat. med.* 398. *Roy. lugdb.* 176. *Hall. helv.* 713. *Dalib. paris.* 264. *Gmel. sib.* 2. p. 196.

Dracunculus pratensis, serrato folio. *Bauh. pin.* 98.

Ptarmica vulgaris. *Clus. hist.* 2. p. 12.

β. *Ptarmica* vulgaris, flore pleno. *Clus. hist.* 2. p. 12.

Habitat in Europa temperata. & 11. A-

SYNGENESIA: POLYGAM. SUPERFLUA. 899

11. **ACHILLEA** foliis lanceolatis dentato-ferratis: denti-^{alpinæ}culis tenuissime ferratis. *Hort. cliff.* 413. *Roy. lugdb.* 176.
Parmica alpina, foliis profunde iucifis. *Tournef. inst.* 497.
Parmica alpina, incanis ferratis foliis. *Herm. lugdb.* 694.
Habitat in Alpibus Sibiricæ. 2
an locus potuerat ex præcedenti formasse banc?
12. **ACHILLEA** foliis pinnulis pectinatis integriusculis, ^{arata}pedunculis villosis.
Achillea foliis pinnatis, pinnis longis acutis subhirsutis
raro dentatis. *Hall. helv.* 714.
Matricaria alpina, chamæmeli foliis. *Bauh. pin.* 134.
Parthertium alpinum. *Clus. hist.* 1. p. 336.
Habitat in Alpibus Helvetiæ, Valleisæ, Austriæ. 2
Margines calycis atræ & sphacelatæ sunt.
13. **ACHILLEA** foliis pinnatis dentatis hirsutis, floribus ^{nano}glomeratis.
Achillea foliis pinnatis lanugine obductis, floribus albis
umbellatis. *Hall. helv.* 713.
Millefolium alpinum incanum, flore specioso. *Bauh.*
hist. 3. p. 138. *Morif. hist.* 3. p. 39. f. 6. t. 11. f.
11. *Raj. hist.* 347.
Millefolium alpinum tomentosum odoratum nanum.
Bocc. mus. p. 166. t. 120.
Habitat in Alpibus Helvetiæ, Valleisæ.
Planta parva. Calyces spadicci.
14. **ACHILLEA** foliis bipinnatis nudis: laciniis linearibus ^{Millefolium}dentatis. *Hort. cliff.* 413. *Fl. suec.* 705. *Mat. med.*
397. *Roy. lugdb.* 175. *Gron. virg.* 101. *Dalib. paris.* 263.
Achillea foliis pinnato-pinnatis. *Fl. lapp.* 311.
Millefolium vulgare album. *Bauh. pin.* 140.
13. *Millefolium purpureum.* *Tabern. hist.* 130.
Habitat in Europæ pascuis pratisque. 2
15. **ACHILLEA** foliis bipinnatis: pinnis distantibus u-^{nobilis}trinque subseptenis obtusiusculis.
Millefolium nobile. *Trag. hist.* 476.
Tamacetum minus album, odore camphoræ. *Bauh. pin.*
132.
Habitat in Helvetia, Misnia, Bohemia, G. Narbo-
nenli, Tataria. 2
Folia ferme Anthemidis tinctoriæ.
16. **ACHILLEA** foliis linearibus: pinnis subrotundis re-^{cretica}trorsum imbricatis, caule tomentofo.

900 SYNGENESIA: POLYGAM. SUPERFULA.

Millefolium incanum creticum. Bauh. pin. 140. prodr. 72.

Millefolium creticum. Bauh. hist. 3. p. 139.

Habitat in Creta. 2.

Habitus A. Millefolii. Caulis tomentosus. Folia linearia, subtomentosa, retrorsum imbricata: pinnis reniformibus, supra quadrifariam imbricatis. Flores albi.

* *

- egyptiaca.* 17. *ACHILLEA* foliis pinnatis: foliolis obtuse lanceolatis serrato-dentatis. *Hort. cliff. 413. Roy. lugdb. 175.*
Parnica incana, pinnulis cristatis. Tournef. cor. 37. in. 1. p. 228. t. 87.
Absinthium santonicum ægyptiacum. Bauh. pin. 139.
Absinthium ægyptiacum. Dod. pempt. 25.
Habitat in Ægypto & Oriente.

- bipinnata.* 18. *ACHILLEA* foliis bipinnatis tomentosis: foliolis ovatis integris. *Hort. cliff. 413. * Roy. lugdb. 176.*
Parnica orientalis, santolinæ folio, flore minore. Tournef. cor. 37?
Habitat in Oriente. 2.

- inodora.* 19. *ACHILLEA* foliis dentato-laciniatis integerrimis: apice trifidis. *Roy. lugdb. 176.*
Ageratum, laciniatis foliis, inodorum africanum. Herm. flor. 55. Boerb. lugdb. 1. p. 125.
Habitat in Africa.

862

TRIDAX.

- procumbens.* 1. *TRIDAX. Hort. cliff. 418. **
Aster americanus procumbens, foliis laciniatis & hirsutis. Houst. mff.
Habitat in Vera Cruce.

863

SIGESBECKIA.

- orientalis.* 1. *SIGESBECKIA* petiolis sessilibus, calycibus involu-
 cratis.
*Sigesbeckia. Hort. cliff. 412. t. 23. * Hort. upf. 267. **
Bidenti similis foliis latissimis serratis. Buxb. cent. 3. p. 29. t. 52.
Cichorio affinis Lapsana sinica, menthastri foliis, calyce fimbriato hispido. Pluk. amaltb. 58. t. 380. f. 2.
Habitat in China, Media ad pagos. ☉
- occidentalis.* 2. *SIGESBECKIA* petiolis decurrentibus, calycibus nudis.

Ver-

SYNGENESIA: POLYGAM. SUPERFLUA. 901

Verbesina foliis ovatis petiolatis decurrentibus oppositis, flosculis radiorum solitariis. *Gron. virg.* 179. *
Habitat in Virginia.

VERBESINA.

864

1. VERBESINA foliis alternis decurrentibus undulatis *alwa* obtusis. *Hort. cliff.* 411. *Hort. ups.* 262.

Cannabina indica, foliis integris, alato caule. *Magn. hort.* 40. t. 40.

Chrysanthemum curassavicum, alato caule, floribus aurantiis. *Herm. par.* 125. t. 125.

Chrysanthemum americanum, caule alato, flore aphylo globoso aurantio, foliis baccharidis. *Comm. hort.* 1. p. 5. t. 13.

Chrysanthemum americanum bideus alatum, flore parvo aurantiaco. *Pluk. alm.* 100. t. 84. f. 3.

Chrysanthemum conyzoides curassavicum, abrotani fœminæ flore aurantio. *Volk. norib.* 106. t. 106.

Habitat in Curassao, Surinamo. &

A reliquis maxime differt habitu & structura.

2. VERBESINA foliis alternis petiolatis ovato-lanceola-*chinesis* tis serratis.

Habitat in China. Osbeck.

Caulis simplex, teres, subtomentosus. Rami simplices, ex alis superioribus foliorum. Folia ovato-lanceolata, subtomentosa, obtusiuscula, petiolata, serrato dentata, alterna. Flores terminales, solitarii, pedunculati. Calyx hemisphæricus, imbricatus, tomentosus. Corolla lutea. Flosculi Radii lanceolati, integri, multi; Disci vero copiosi, paleis interstincti. Semina coronata margine & setis 2 levibus.

3. VERBESINA floribus corymbosis, foliis lanceolatis *virginica* petiolatis. *Gron. virg.* 102. *

Habitat in Virginia.

4. VERBESINA foliis oppositis lanceolatis subserratis, *Pend. Ac. nulla,* pedunculis flore longioribus. *Fl. zeyl.* 308. *

Bidens zeylanica, flore luteo, melissæ folio, *Actinella* dicta. *Seb. thes.* 1. p. 19. t. 10. 11.

Chrysanthemum maderaspatanum latifolium, scabiosæ capitulis parvis. *Pluk. alm.* 99. t. 154. f. 4.

Habitat in Zeylona.

5. VERBESINA foliis oppositis lanceolato-ovatis petio- *Act. illa.* latis

L 113

902 SYNGENESIA: POLYGAM. SUPERFLUA.

- latis serratis, pedunculis unifloris dichotomizæ caulis.
Fl. zeyl. 309. * *Mat. med.* 406.
 Chrysanthemum bidens f. Bidens zeylanica, flore luteo,
 lamii folio, Acmella dicta. *Raj. suppl.* 228. *Breyn.*
alm. 48. t. 48.
 Senecio ind. orient. ocymi majoris folio profunde cre-
 nato. *Pluk. alm.* 343. t. 315. f. 2.
Habitat in Zeylona.
- Lavenia.* 6. VERBESINA foliis oppositis ovatis trinerviis glabris
 petiolatis, seminibus tricornibus. *Fl. zeyl.* 310. *
 Eupatorio-phalacron scrophulariæ aquaticæ foliis oppo-
 sitis. *Burm. zeyl.* 94. t. 42.
 Chrysanthemum aquaticum, scrophulariæ folio, Lavenia
 dictum. *Raj. suppl.* 217.
Habitat in Zeylona.
- alba.* 7. VERBESINA foliis oppositis lanceolatis serratis. *Hort.*
cliff. 500. *Gran. virg.* 103.
 Eupatorio-phalacron balsaminæ feminæ folio, flore al-
 bo discoide: V. *Dill. elsh.* 138. t. 113. f. 137.
 Scabiosa conyzoides americana latifolia, capitulis & flo-
 ribus albis parvis. *Pluk. alm.* 335. t. 109. f. 1. *Morif.*
bist. 3. p. 47. f. 6. t. 13. f. 16.
Habitat in Virginia, Surinamo. &
- prostrata.* 8. VERBESINA foliis oppositis lanceolatis serratis, flo-
 ribus alternis geminis subsessilibus.
 Eupatorio-phalacron menthæ arvensis folio. V. *Dill.*
elsh. 139. t. 113. f. 138.
 Chrysanthemum maderaspatanum, menthæ arvensis folio
 & facie, floribus bigemellis ad alas, pediculis-curtis.
Pluk. alm. 100. t. 118. f. 5.
Habitat in India.
- calendulacea* 9. VERBESINA foliis oppositis lanceolatis obtusifusculis,
 pedunculis longis unifloris. *Fl. zeyl.* 311. *
 Caltha flore solitario ex alis foliorum longissimo pedun-
 culo prodeunte. *Burm. zeyl.* 52. t. 22. f. 1.
Habitat in Zeylona.
- asteroides.* 10. VERBESINA foliis oppositis lineari-lanceolatis ob-
 tusis integerrimis tomentosis, pedunculo unifloro.
 Buphtalmum foliis oppositis lanceolato-linearibus obtu-
 sis integerrimis, calycibus subrotundis. *Hort. cliff.*
 415. *
Habitat - - -

Flos

SYNGENESIA: POLYGAM. SUPERFLUA. 903

Flos *Asteris* facie, luteus: radio violaceo, coroma seminatum ex 5 dentibus inspidis setaceis. Folia *lychnidis*.

TETRAGONOTHECA. 865

1. TETRAGONOTHECA

Tetragonotheca doronici maximi folio. *Dill. elib.* 378. *botanich. der.*

z. 283. *f.* 365. *Gron. virg.* 103.

Habitas in Virginia.

BUPHTHALMUM.

* *Asterisci* T. 862

1. BUPHTHALMUM foliis oppositis lanceolatis, peristemonioidis bidentatis. *Hort. cliff.* 415. *Gron. virg.* 101.

Roy. lugdb. 170.

Asteriscus frutescens, leucodi foliis sericeis & incanis.

Dill. elib. 44. *t.* 38. *f.* 44.

Chrysanthemum fruticosum maritimum, foliis glaucis oblongis, flore luteo. *Sloan. jam.* 125. *bist.* 1. p. 260.

Catesb. car. 1. p. 93. *t.* 93.

Habitat in Jamaica, Virginia. ☉

2. BUPHTHALMUM calycibus acute foliosis, ramis spinosis alternis, foliis lanceolatis amplexicaulibus integerrimis. *Hort. cliff.* 414. *Roy. lugdb.* 170. *Samv. monsp.* 87.

Aster luteus, foliis ad florem rigidis. *Bauh. pin.* 266.

Aster legitimus clusii alter s. spinosus luteus. *Barr. ic.* 351.

Habitat in G. Narbonensi, Hispanias Italia, ad margines agrorum. ☉

3. BUPHTHALMUM calycibus obtuse foliosis sessilibus axillaribus, foliis oblongis obtusis. *Hort. cliff.* 414. *Hort. upf.* 464. *Roy. lugdb.* 170. *Samv. monsp.* 87.

Asteriscus annuus lusitanicus odoratus. *B. Seb. thes.* 1. p. 47. *t.* 29. *f.* 7.

Chrysanthemum conyzoides lusitanicum. *Breyn. cent.* 157. *t.* 77.

Habitat in Creta, Lusitania, Massilia. ☉

4. BUPHTHALMUM calycibus obtuse foliosis pedunculatis, ramis alternis, foliis cuneiformibus. *Hort. cliff.* 414. *Hort. upf.* 264. *Roy. lugdb.* 170.

Aster supinus lignosus ficulus, conyzæ odore. *Bocc. mus.* 2. p. 161. *t.* 129.

904 SYNGENESIA: POLYGAM. SUPERFLUA.

Aster supinus luteus massilioticus. Barr. ic. 1151.

Aster 2 supinus. Clus. hist. 2. p. 13.

Aster luteus supinus. Baub. pin. 267.

Habitat Massiliæ, & in Sicilia ad littora maris. & ☉

* *Asteroidea T.*

salicifolium. 5. BUPHTHALMUM foliis lanceolatis subserratis villosis, calycibus nudis. Hort. cliff. 414. Roy. lugdb. 170.

Hall. belv. 711. Sauv. monsp. 87.

Aster luteus major, foliis succilæ. Baub. pin. 266.

Aster 3, austriacus 1. Clus. hist. 2. p. 13.

Habitat ad radices Alpium Austriæ, Stiria, Lusatia, Monspelii. &

*grandiflorus. 6. BUPHTHALMUM foliis lanceolatis subdenticulatis glabris, calycibus nudis. Hort. cliff. 415. * Hort. upf. 264. Roy. lugdb. 170.*

Aster luteus angustifolius. Baub. pin. 266.

Chrysanthemum perenne minus, foliis glabro folio, ramosum. Moris. hist. 3. p. 21. f. 6. t. 7. f. 52.

Habitat in Alpibus Austriæ, Italia, Monspelii. &

Helianthus. 7. BUPHTHALMUM calycibus foliosis, foliis cordatis serratis trinerviis basi hinc brevioribus. Hort. upf. 264.

Helianthus foliis ovatis acuminatis serratis, pedunculis longissimis. Gron. virg. 179.

Corona Solis caroliniana, parvis floribus, folio trinervi amplo aspero, pediculo alato. Mart. cent. 20. t. 20.

Chrysanthemum, scrophularia folio, americanum. Pluk. alm. 99. t. 22. f. 1.

Chrysanthemum virginianum, foliis glabris scrophularia vulgaris æmulis. Moris. hist. 3. p. 24. f. 6. t. 3. f. 69. Raj. synopl. 211.

Habitat in America septentrionali. &

Folia summa altero latere bascos longe breviora sunt.

Radius pistillis instructus & calycis folia obtusiora reddunt plantam distinctam ab Helianthemis, quibus simillima.

POLYGAMIA FRUSTRANEA.

HELIANTHUS.

867
HYMNEUS.

1. HELIANTHUS foliis omnibus cordatis; nervis po-
né basin unitis; extrorsum denudatis,

Heli

SYNGENESIA: POLYGAM. FRUSTRAN. 905

Helianthus radice annua. Vir. cliff. 88. Hort. cliff. 419.

Hort. upf. 268. Roy. lugdb. 180.

Helenium indicum maximum. Baub. pin. 276.

Herba maxima. Dod. pempt. 264.

Chrysis. Reneal. spec. 84. t. 83.

Habitat in Peru, Mexico. ☉

2. **HELIANTHUS** foliis inferioribus cordatis: nervis *multiflorus*,
pone basin unitis denudatis: superioribus ovatis.

Helianthus radice tereti-inflexa perenni. Hort. cliff. 419.

Hort. upf. 268. Roy. lugdb. 180.

Helenium indicum ramosum. Baub. pin. 277.

Corona Solis minor femina. Tabern. ic. 764.

Habitat in Virginia. ☿

Calyx constat squamis 40 a 50.

3. **HELIANTHUS** foliis, ovato cordatis: nervis *intuberosus*,
tra folium unitis.

Helianthus radice tuberosa. Hort. cliff. 419. Hort. upf.
268. Roy. lugdb. 180.

Helenium indicum tuberosum. Baub. pin. 277.

Chrysanthemum latifolium brasilianum. Baub. prodr. 70.

Flos Solis farnesianus. Col. ecphr. 2. p. 11. t. 13.

Habitat in Brasilia. ☿

4. **HELIANTHUS** caule inferne laevi, foliis lanceola- *decapetalus*,
to-cordatis, radlis decapetalis.

Habitat in Canada ex semine. D. Kalm.

Simillima H. multifloro, sed differt quod caulis, vix hu-
mana altitudinis, inferne glaber. Folia magis lan-
ceolata-ovata (nec cordata) nervis longius intra basin
coeuntibus. Corolla radiis non pluribus quam decem.
Calyx squamis viginti constat. Habitus distinctus.

5. **HELIANTHUS** radice fusiformi. *Hort. cliff. 420. strumosus.*

Roy. lugdb. 181.

Chrysanthemum canadense latifolium altissimum. Mo-
rif. blef. 250. bist. 3. p. 23.

Chrysanthemum canadense latifolium elatus. Bocc. sic.
52. t. 27. f. 4.

Chrysanthemum canadense strumosum vulgo. Herm.
*lugdb. 143. **

Habitat in Canada.

6. **HELIANTHUS** foliis lanceolatis scabris, caule stri- *giganteus*,
eto: inferne glabro.

Helianthus foliis lanceolatis sessilibus. Gron. virg. 104.

L 115

Chry.

906 SYNGENESIA: POLYGAM. FRUSTRAN.

Chrysanthemum virginianum altissimum angustifolium,
puniceis caulibus *Morif. hist.* 3. p. 24 f. 6. t. 7. f. 67.

Chrysanthemum virginianum elatius angustifolium,
caule hirsuto viridi. *Pluk. alm.* 99. t. 159. f. 5.

Habitat in Virginia, Canada. &

Radius flosculis 16. f. 18, *apice bifidis sterilibus. Caulis*
decem pedum.

levis. 7. *HELIANTHUS* foliis lanceolatis ferratis lævibus.
Gron. virg. 104.

Habitat in Virginia.

angustifolius 8. *HELIANTHUS* foliis linearibus. *Gron. virg.* 103.
Flos Solis marianus, foliis alternis angustissimis scabris.
Pet. mus. 103.

Habitat in Virginia.

atrorubens. 9. *HELIANTHUS* foliis oppositis sessilibus ovato-ob-
longis trinerviis, panicula dichotoma.

Chrysanthemum virginianum repens, foliis asperis bina-
tim sessilibus acuminatis. *Morif. hist.* 3. p. 22. f. 6.
t. 7. f. 66 *mala.*

Habitat in America septentrionali. &

Caulis vix humane altitudinis, superne lævissimus pi-
lis sparsis asperis, purpureus rore glauco. Flores in
panicula ramis trichotomis, flavi, parvi, peduncula-
ti: intermedio pedunculo simplicis.

atrorubens. 10. *HELIANTHUS* foliis ovatis crenatis trinerviis sca-
bris, squamis calycinis crectis longitudine disci. *Gron.*
virg. 103.

Corona Solis minor, disco atrorubente. *Dill. elsb.* 111.
t. 94. f. 110.

Habitat in Virginia. &

860
laciniata.

RUDBECKIA.

1. *RUDBECKIA* foliis compositis laciniatis. *Vir. cliff.*
88. *Hort. cliff.* 420. *Hort. ups.* 269. *Roy. lugdb.*
181. *Gron. virg.* 180.

Chrysanthemum americanum perenne, foliis divisis di-
latius virentibus, majus. *Morif. hist.* 3. p. 22. f. 6. t.
6. f. 53. & 54.

Doronicum americanum, laciniato folio. *Barb. pin.*
516.

Aconitum helianthemum canadense Gron. canad. 178.
t. 179.

Habitat in Virginia, Canada. &

2. *RUD-*

SYNGENESIA: POLYGAM. FRUSTRAN. 907

2. RUDBECKIA foliis inferioribus trilobis. superioribus *stellatis*, indivisis. *Hort. nps.* 269.
Rudbeckia foliis trilobis. *Gron. virg.* 180
Chrysanthemum cannabinum virginianum hirsutum, disco magno, petalis aureis radiato. *Pluk. alm.* 100. t. 22. f. 2.
Chrysanthemum annuum majus virginianum, foliis laciniatis & hirsutis, umbone nigricante. *Morif. hist.* 3. p. 19. n. 37.
Habitat in Virginia. ♂
3. RUDBECKIA foliis indivisis spatulato-ovatis; radii *hirta*, petalis emarginatis.
Rudbeckia, ramis indivisis unifloris, foliis ovato-lanceolatis, hirta. *Battm. canan.* 227. *
Rudbeckia foliis lanceolato-ovatis alternis indivisis, petalis radii integris. *Gron. virg.* 181.
Obeliscotheca integrifolia, radio aureo, umbone atro-rubente. *Dill. elth.* 295. t. 218. f. 285.
Chrysanthemum helenii folio, umbone floris gradusculo prominente. *Pluk. alm.* 99. t. 242. f. 2. *Morif. hist.* 3. p. 23. *Raj. suppl.* 210.
Habitat in Virginia, Canada. ♂. ♀
4. RUDBECKIA foliis lanceolato-ovatis alternis indivisis, petalis radii bifidis. *Gron. virg.* 104. 181. *
Chrysanthemum americanum, doronici folio, flore perfici coloris: umbone magno prominente ex atro purpureo viridi & aureo fulgente. *Pluk. alm.* 99. t. 21. f. 1. *Catesb. car.* 2. p. 59. t. 59.
Dracunculus virginianus latifolius, petalis florum longissimis purpurascens. *Morif. hist.* 3. p. 42. f. 6. t. 9. f. 1. *Raj. suppl.* 218.
Bobartia. *Pot. mus.* *Ann. herb.* 473.
Habitat in Virginia, Carolina.
5. RUDBECKIA foliis oppositis lanceolato-ovatis *ser-oppositifolia*, ratis, petalis radii bifidis. *Gron. virg.* 180.
Habitat in Virginia.
Calyx brevissimus foliolis ovatis. Radius flavus, longissimus. Discus convexus. flavus.

COREOPSIS. 869

1. COREOPSIS foliis decomposito-pinnatis linearibus. *verticillate*.
Coreopsis foliis verticillatis linearibus multifidis. *Gron. virg.* 105.

Cera-

908 SYNGENESIA: POLYGAM. FRUSTRAN.

Ceratocephalus delphinii foliis. *Vaill. act.* 1720.

Chrysanthemum marianum, scabiosæ tenuissime divisæ foliis, ad intervalla confertis. *Pluk. mant.* 48. s. 344. f. 4.

Habitat in Virginia. ♂

ripteris.

2. COREOPSIS foliis subternatis integerrimis. *Hort. nys.* 269. *

Rudbeckia foliis compositis integris. *Roy. lugdb.* 181.

Chrysanthemum virginianum, folio acutiore lævi trifoliato s. *anagyridis* folio. *Morif. hist.* 3. p. 21. f. 6. s. 3. f. 44. *Raj. suppl.* 215.

Habitat in Virginia. ♀

Folia pleraque ternata, sæpe etiam in eamle ferme pinata.

alba.

3. COREOPSIS foliis subternatis cuneatis ferratis.

Chrysanthemum americanum, ciceris folio glabro, bellidis majoris flore. *Herm. par.* 124. s. 124. flore albo. *Pluk. alm.* 101. s. 160. f. 3.

Habitat in Insula St. crucis. ♀

auriculata.

4. COREOPSIS foliis ovatis: inferioribus ternatis. *Gron. virg.* 105.

Chrysanthemum hirsutum virginianum, auriculato dolcamaræ folio, octopetalon. *Pluk. alm.* 101. s. 83. f. 5. & t. 242. f. 4. *Raj. suppl.* 212.

Chrysanthemum virginianum trifoliatum humilius, obscurioribus foliis hirtutis. *Morif. hist.* 3. p. 20. f. 6. s. 3. f. 45.

Habitat in Virginia.

lanceolata.

5. COREOPSIS foliis lanceolatis integerrimis.

Coreopsis foliis integerrimis. *Roy. lugdb.* 181.

Coreopsis. *Hort. cliff.* 420.

Bidens succisæ folio, radio amplo laciniato. *Dill. elsh.* 55. t. 48. f. 56.

Bidens caroliniana, florum radiis latissimis insigniter dentatis, semine alato per maturitatem convolutæ. *Mart. cent.* 26. t. 26.

Habitat in Carolina.

angustifolia.

6. COREOPSIS foliis linearibus integerrimis. *Gron. virg.* 181.

Habitat in Virginia.

Bidens.

7. COREOPSIS foliis lanceolatis ferratis oppositis amplexicaulis.

Core-

SYNGENESIA: POLYGAM. FRUSTRAN. 909

*Coreopsis bidentis. It. wgotb. 272. **

Hepatorium aquatile, folio non diviso, flore aureis petalis undique radiato. Triumf. absf. 65.

Chrysanthemum aquaticum bidens, folio non diviso, flore aureis petalis undique radiato. Moris. hist. 3. p. 17. f. 6. t. 5. f. 25.

Eupatorium cannabinum chrysanthemum. Barr. ic. 1209.

Habitat ad fossas Europæ. ☉

Si observationibus evincatur hanc BIDENTIS cernuæ solam esse varietatem, ad Bidentes reducatur, sin constans sit, permaneat.

8. COREOPSIS foliis lanceolatis serratis alternis petiolaris alternifolia, decurrentibus. *Hort. upf. 270.*

Coreopsis foliis serratis. Roy. lugdb. 181.

Chrysanthemum canadense bidens, alato caule. Moris. blef. 253. Raj. hist. 337.

Chrysanthemum virginianum, alato caule, bidens altissimum, folio aspero, flore minore serotino. Moris. hist. 3. p. 25. f. 6. t. 7. f. 75. & 76.

Chrysanthemum virginianum, caule alato, ramosius, flore minore. Pluk. alm. 100. t. 159. f. 3.

Habitat in Virginia, Canada. &

CENTAUREA. 870

* *Jaceæ calycibus lævibus inermibus.*

1. CENTAUREA calycibus inermibus: squamis linearibus subulatis, foliis pinnatis serratis. *Hort. cliff. 420. Hort. upf. 271. Roy. lugdb. 137. Sauv. monsp. 264.*

Chondrilla foliis laciniatis serratis, purpurascens flore. Baub. pin. 130.

Senecio-Carduus apulus. Col. ecphr. 1. p. 32. t. 34.

Habitat in Hetruria, G. Narbonensi. ☉

2. CENTAUREA calycibus inermibus subrotundis glabris: squamis ovatis, foliis sinuatis. *Hort. cliff. 421. Hort. upf. 271. Roy. lugdb. 138.*

Cyanus orientalis major moschatus, flore purpureo & albo. Moris. hist. 3. p. 135. f. 7. t. 25. f. 5.

3. *Cyanus orientalis major, foliis magis dissectis, flore luteo. Moris. hist. 3. p. 135. f. 7. t. 25. f. 9.*

Habitat in Tataria. ☉

3. CENTAUREA calycibus inermibus: squamis lanceolatis, foliis lanceolatis subdentatis. *Hort. cliff. 421. Roy. lugdb. 138.*

Ja-

Jacea foliis erucæ lanuginosis. *Tournef. inst.* 444.
 Amberboi, erucæ folio, majus. *Vaill. act.* 1718. p. 181.
 Stœbe major foliis erucæ mollibus lanuginosis. *Bauh.*
pin. 273.

Habitat - - - 2

- Lippi.* 4. CENTAUREA calycibus inermibus: squamis mucronatis, foliis pinnatifidis obtusis decurrentibus.
 Amberboi, erucæ folio, minus. *Isn. act.* 1719. p. 169.
t. 10.

Habitat in Ægypto inter Alexandriam & Rosette. ☉

- alpina.* 5. CENTAUREA calycibus inermibus: squamis ovatis obtusis, foliis pinnatis glabris integerrimis: impari serrato. *Hort. cliff.* 421. *Roy. lugdb.* 138.

Centaurium alpinum luteum. *Bauh. pin.* 117. *prodr.* 56.

Morif. hist. 3. p. 132. f. 7. t. 25. f. 5.

Centaurium majus luteum. *Corn. canad.* 69. t. 70.

Habitat in Baldo.

- Centaurium.* 6. CENTAUREA calycibus inermibus: squamis ovatis, foliis pinnatis: foliolis serratis decurrentibus.
Hort. cliff. 421. *Roy. lugdb.* 137.

Centaurium majus, folio in lacinias plures diviso. *Bauh.*
pin. 117.

Centaurium majus 1 vulgare. *Clus. hist.* 2. p. 10.

Habitat in Alpibus Gargano, Baldo, Tataria. 2

* Cyani calycinis squamis serrato-ciliatis.

- phrygia.* 7. CENTAUREA calycibus recurvato-plumosis, foliis indivisis.

Centaurea calycibus ciliatis: ciliis setaceis recurvatis.

Hort. cliff. 422. *Hort. ups.* 270. *Roy. lugdb.* 139.

Dalib. parif. 266.

- Jacea latifolia & angustifolia, capite hirsuto. *Bauh. pin.* 271.

Jacea 4 austriaca, capite villoso. *Clus. hist.* 2. p. 7.

β. Jacea alba, hirsuto capite. *Bauh. pin.* 271.

Habitat in Helvetia, Austria, Finlandia. 2

- capillata.* 8. CENTAUREA calycibus seta recurvato-plumosis, foliis imis pinnatis dentatis; summis lanceolatis.

Centaurea calycibus ciliatis oblongis, foliis scabris, caulium pinnatis: ramorum simplicissimis linearibus.

Gmel. sib. 2. p. 98. t. 43.

Cyanus foliis calycinis plumosis reflexis, foliolis pinnatis.

Act. goett. 1. p. 202. t. 6.

Habitat in Sibiria. Gmelinus.

9. CEN-

9. CENTAUREA calycibus pinnatis: squamula ovata: *nigra*.
ciliis capillaribus erectis.

Cyanus foliis imis ad basin pinnatis: superioribus integris
ovato-lanceolatis. *Hall. helv.* 691. t. 23. f. 1.

Jacea nigra laciniata. *Baub. pin.* 271.

Jacea austriaca V1. *Clus. hist.* 2. p. 7.

Habitat in Anglia, Helvetia, Austria.

Differt a *C. phrygia*, quod squamula ciliaris sit ovatae-
recto patens, nec recurva & jubulata.

10. CENTAUREA calycibus ciliatis verticillato-folio-pullata.
lis, foliis lyrato dentatis, caule foliis brevior.

Centaurea inermis, calycum ciliis pinnatis, foliis oblon-
gis villosis humifutis, floralibus verticillatis ligulatis.
Sauv. monsp. 268.

Jacea humilis alba, hieracii folio. *Baub. pin.* 271. *Mor-
ris. hist.* 3. p. 140 f. 6. t. 28. f. 18.

Jacea montepellana, cui in squamis fibræ nigrae, inter-
dum acaulis. *Baub. hist.* 3. p. 29.

Jacea pumila serpens acaulis ferme, flore albo. *Lob. ic.*
542.

Habitat in G. Narbonensi, Hispania. ☉

Calyx foliis cingitur sessilibus, intra haec folia Squamæ
calycine lineari-lanceolatae margine atro, apice sesis
pallidis ciliatæ. Folia in caule sæpe lyrata sunt.

11. CENTAUREA calycibus ferratis, foliis lanceolatis *montana*.
decurrentibus, caule simplicissimo. *Hort. cliff.* 422.

Hort. ups. 270. *Roy. lugd.* 138.

Cyanus montanus latifolius f. *Verbasculum cyanoides*.
Baub. pin. 273.

Cyanus montanus, caule folioso, capitulo oblongo.
Bocc. mus. 2. p. 20. t. 2.

Cyanus major. *Lob. ic.* 548.

12. Jacea integrifolia humilis. *Baub. pin.* 271. *prodr.* 127.
Burf. XV. 29.

Habitat in Alpibus Helveticis. ☿

Planta β. foliis minoribus, parum basi decurrentibus, nec
ad proximam fuitum, a vulgari differt.

12. CENTAUREA calycibus ferratis: foliis linearibus *Cyanus*,
integerrimis: infimis dentatis. *Hort. cliff.* 422. *Fl.*
jacq. 710. *Mat. med.* 408. *Roy. lugd.* 130. *Dalib.*
paris. 265.

Cyanus segetum. *Baub. pin.* 273.

Cyanus vulgaris. *Lob. ic.* 546.

Habitat inter Europæ segetes biennes. ☉

912 SYNGENESIA: POLYGAM. FRUSTRAN.

- paniculata.* 13. **CENTAUREA** calycibus ciliatis, foliis pinnatifidis linearibus, caule paniculato.
Centaurea calycibus ciliatis, foliis filiformibus bipinnatis, caule folioso ramoso. Sam. monsp. 264.
Jacea non spinosa, foliis magis divisis, elatior, capitulis minoribus non splendentibus. Moris. hist. 3. p. 140. f. 7. t. 28. f. 15.
Stœbe major, cauliculis non splendentibus. Bauh. pin. 273.
Centaurei majoris species tenuifolia. Bauh. hist. 3. p. 31.
Habitat in G. Narbonensi, Austria, Hispania, Verona, Sibiria. ☉
- sessiliflora.* 14. **CENTAUREA** calycibus ferratis ciliatisque, foliis lanceolatis dentatis, floribus sessilibus. *Hort. cliff. 422.**
Roy. lugdb. 139.
Habitat in G. Narbonensi.
- spinosa.* 15. **CENTAUREA** calyce subciliato, ramis spinosis *Hort. cliff. 422.**
Jacea cretica aculeata incana. Tournef. inst. 445.
Stœbe spinosa cretica. Moris. hist. 3. p. 126.
Cyanus spinosus. Alp. exot. 163. t. 162.
Habitat in Creta.
- ragusina.* 16. **CENTAUREA** calycibus ciliatis, foliis tomentosis pinnatifidis, foliolis obtusis ovatis integerrimis: exterioribus majoribus. *Hort. cliff. 422. Roy. lugdb. 139.*
Jacea cretica lutea, foliis cinaræ. Moris. hist. 3. p. 141. f. 6. t. 27. f. 22.
Jacea arborea argentea ragusina. Zan. bist. 107. t. 43.
Stœbe montana nivea, capite cardui, subrotundis foliorum lobis. Barr. ic. 309.
Habitat in Creta, Epidaurò. &
- cineraria.* 17. **CENTAUREA** calycibus ciliatis terminali sessilibus, foliis tomentosis bipinnatifidis: laciniis linearibus. *Hort. cliff. 422. Roy. lugdb. 139.*
Jacea montana candidissima, stœbes foliis. Bauh. pin. 272. prodr. 126. Moris. hist. 3. p. 141. f. 7. t. 26. f. 20.
 18. *Jacea cineraria laciniata, flore purpureo. Triumf. obs. 72. Moris. hist. 3. p. 141. Barr. ic. 348. 347.*
Habitat in Italia. &
- argentea* 18. **CENTAUREA** calycibus ciliatis villosis, foliis cuneiformibus superne serratis inferne dentatis. *Hort. cliff. 422.* Ja.

Jacea cretica laciniata argentea, flore parvo flavescente. *Tournef. cor.* 32. *Barr. ic.* 218.

Habitat in Creta.

19. **CENTAUREA** calycibus ciliatis, foliis tomentosis ^{*Sibirica.*} indivisis pinnatifidisque integerrimis, caule declinato.

Centaurea calycibus ciliatis subrotundis, foliis pinnatis & integris: foliolis simplicissimis: extremo maximo.

Gmel. fib. 2. p. 96. t. 42. f. 2.

Habitat in Sibiria. D. Gmelin. 2

Folia tomentosa, lanceolata: radicalia pleraque pinnatifida: Lobis sursum versis, lanceolatis, integerrimis, decurrentibus, extremo majore. Folia caulina saepius 6 f. 7, superioribus indivisis. Caulis simplicissimus, raro unico ramulo, pubescens, subsulcatus. Flos saepius unicus. Calyce ventricoso: foliolis auctis Squamula ovata, magnitudine folioli calycini, argute ciliatis, pallidis. Corolla purpurea; radio sterili magno.

20. **CENTAUREA** calycibus ciliatis, foliis lanceolatis: ^{*Semperotens.*} superne ferratis. *Hort. cliff.* 422. * *Roy. lugdb.* 139.

Jacea lusitanica sempervirens. *Morif. bist.* 3. t. 139. f. 7. t. 28. f. 9.

Habitat in Lusitania. 2

21. **CENTAUREA** calycibus margine ciliatis, foliis pinnatifidis: pinnis lanceolatis. *Hort. cliff.* 422. *Fl. succ.* 708. *Roy. lugdb.* 139. *Dalib. parif.* 266. *Gmel. fib.* 2. p. 93.

Scabiosa major, squamatis capitulis. *Baub. pin.* 269.

Scabiosa major. *Dalech. bist.* 1066.

Habitat in Europæ septentrionalis pratis.

22. **CENTAUREA** calycibus squama ciliatis, foliis pinnatifidis: pinnis lanceolatis. ^{*orientalis.*}

Centaurea calycibus ciliatis, foliis pinnatis glabris: foliolis lanceolatis integerrimis. *Hort. upf.* 271.

Cyanus foliis radicalibus: partim integris, partim pinnatis, bractea calycis ovali, flore sulphureo. *Hall. aet. angl.* 1745. vol. 43. n. 472. p. 94.

Habitat in Sibiria. 2

Differt a C. Scabiosa, cui simillima. non tantum Corollasulphurea, sed præcipue Calyce, cujus foliola terminata squama ovata, ferruginea, ciliata. ciliis longitudine ipsius squamæ, pectinatis, cum in C. Scabiosa foliorum calycis margo niger ciliato-lacerus sit, nec in squamam eductus.

M m m

23.

914 SYNGENESIA: POLYGAM. FRUSTRAN.

- Stœbe.* 23. *CENTAUREA* calycibus ciliatis oblongis, foliis pinnatifidis linearibus integerrimis. *Roy. lugdb.* 140.
Stœbe incana, cyano similis, tenuifolia. *Baub. pin.* 273.
Stœbe austriaca humilis. *Clus. hist.* 2. p. 10.
Habitat in Austria.
- acaulis* 24. *CENTAUREA* calycibus ciliatis, foliis lyratis, caule subnullo.
Crocodilium acaulon ferme, calcitrans folus crassis tomentosis. *Vaill. act.* 1718. p. 162.
Jacea acaulos lutea, erucæ folio, squamarum ciliis candido. *Shaw. spec.* 342. t. 342.
Habitat in Arabia.
 * *Rhapontica calycinis squamis membranaceis, aridis, integerrimis, muticis.*
- Behen,* 25. *CENTAUREA* calycibus squamosis, foliis radicalibus lyratis; lobis oppositis, caulinis amplexicaulibus.
Mat. med. 409.
Rhaponticoides lutea, foliis inferioribus dissectis, ceteris connatis. *Vaill. act.* 1718. p. 229.
Behen album. *Rauw. itin.* 288. t. 288. *Baub. hist.* 3. p. 36. t. 37. *Baub. pin.* 234.
Habitat in Asia minore, Libano. &
- Jacea.* 26. *CENTAUREA* calycibus squamosis, foliis lanceolatis; radicalibus sinuato-dentatis, ramis angulatis.
Hort. cliff. 421. *Fl. succ.* 709. *Roy. lugdb.* 142. *Dalib. parif.* 265.
Jacea nigra pratensis latifolia. *Baub. pin.* 271.
Jacea nigra. *Paul. dan.* t. 256.
Habitat in Europa septentrionali. &
- alba,* 27. *CENTAUREA* calycibus squamosis mucronatis, foliis pinnato-dentatis.
Jacea, calyculis argenteis, minor. *Tournef. inst.* 444.
Jacea alba. *Tabern. ic.* 153.
Stœbe, calyculis argenteis, minor. *Baub. pin.* 273.
Habitat in Hispania.
Caulis paniculatus, angulatus. Folia linearia, pinnato-dentata, acuminata: superiora tantum dentata; superiora linearia, simplicissima. Calyces terminales, ovati, parvi, Squamis membranaceis, laxis, mucronatis.
- splendens.* 28. *CENTAUREA* calycibus squamosis obtusis, foliis radicalibus bipinnatifidis; caulinis pinnatis dentibus laevicolatis. *Roy. lugdb.* 142. *Sauv. meth.* 289.

SYNGENESIA: POLYGAM. FRUSTRAN. 915

Jacea, calyculis argenteis, maior. *Tournef. inst.* 444.

Stœbe caliculis argenteis. *Bauh. pin.* 273.

Stœbe salmantica 3. *Clus. hist.* 2. p. 10.

Habitat in Helvetia, Hispania, Sibiria, &

29. CENTAUREA calycibus squamosis, foliis ovato-Rhapontica. oblongis deuticulatis integris petiolatis: subtus tomentosis. *Hort. cliff.* 421. *Roy. lugdb.* 142.

Rhaponticum folio helenii incano. *Bauh. pin.* 117.

Rha f. Rhei ut existimatur. *Dod. pempt.* 389.

Habitat in Alpibus Helveticis, Veronæ.

30. CENTAUREA calycibus squamosis, foliis indivisis glastifolia. integerrimis decurrentibus. *Hort. cliff.* 421. *Roy. lugdb.* 142. *Gmel. sib.* 2. p. 83

Centaurium majus orientale erectum, glasti folio, flore luteo. *Comm. rar.* 29 t. 39.

Habitat in Oriente, Sibiria. &

31. CENTAUREA calycibus squamosis, foliis tomentosis: radicalibus lanceolatis, caulinis pinatifidis, caule simplici. *Roy. lugdb.* 142. *Hort. ups.* 271. *Sauv. monsp.* 289.

Jacea montana incana, capite plini. *Bauh. pin.* 272.

Chamæleon non aculeatus. *Lob. ic.* 7.

Habitat Monspeli in saxosis glareosis. &

* Stœbe calycinis spinis palmatis.

32. CENTAUREA calycibus palmato-spinosis, foliis de-sentifolia. currentibus spinulosis repando-dentatis. *Hort. cliff.* 423. n. 22. *Roy. lugdb.* 140.

Jacea latifolia, caule alato, capite magno turbinato: multiplici aculeo deflexo e squamis cuspidatis orto. *Pluk. alm.* 191. t. 38. f. 1.

Jacea laciniata, fionchi folio, f. Jacea latifolia purpurea, capite spinoso. *Bauh. pin.* 272. *prodr.* 128. *Herm. lugdb.* 331. t. 675.

Habitat in maritimis maris Mediterranei. ☉

33. CENTAUREA calycibus palmato spinosis, foliis eridis. decurrentibus tomentosis oblongis: infimis sinuato-dentatis. *Sauv. meth.* 288.

Jacea, foliis seridis candicantibus, purpurea. *Bauh. pin.* 272.

Jacea purpurea hispanica muricata. *Bauh. hist.* 3. p. 33. *descr.*

M m m 2

Jacea

660

916 SYNGENESIA: POLYGAM. FRUSTRAN.

Jacea latifolia, spinosis capitulis pulchris, flore purpureo.

Bauh. bist. 3. p. 33. *Raj. bist.* 320.

Habitat in Hispania.

romana.

34. *CENTAUREA* calycibus palmato-spinosis, foliis decurrentibus inermibus: radicalibus pinnatifidis: impari maximo. *Hort. cliff.* 423. *Roy. lugdb.* 141.

Jacea spinosa cretica. *Zan. bist.* 141. t. 42.

Cyanus erucæ folio, flore rubro. *Barr. rar.* 87. t. 504.

Habitat in Campania Romana. ☉

sphærocephala.

35. *CENTAUREA* calycibus palmato-spinosis, foliis ovato-lanceolatis petiolatis dentatis. *Hort. cliff.* 423. *Roy. lugdb.* 140.

Jacea sphærocephala spinosa tingitana. *Herm. lugdb.* 332.

t. 333. *Morif. bist.* 3. p. 143. f. 7. t. 27. f. 9.

Habitat in Mauritania.

islandi.

36. *CENTAUREA* calycibus palmato-spinosis, foliis lyrato-dentatis hispidis subamplexicaulibus, floribus sessilibus terminalibus.

Calcitrapoides procumbens, cichorii folio, flore purpureascente. *Isn. parif.* 1719. p. 164. t. 9.

Habitat in Europa australi? &

napifolia.

37. *CENTAUREA* calycibus palmato-spinosis: setis æqualibus, foliis decurrentibus sinuatis spinulosis. *Roy. lugdb.* 141. *Hort. upf.* 272. *

Jacea cyanoides altera, alato caule. *Herm. par.* 189. t. 189. *bon.*

Jacea peregrina napifolia, echinatis capitulis, caule alato. *Pluk. alm.* 192. t. 94. f. 2. *Morif. bist.* 3. p. 154. f. 7. t. 26. f. 20.

Habitat in Creta.

aspera.

38. *CENTAUREA* calycibus squamis palmato-quinque-spinosis, foliis lanceolatis dentatis. *Sauv. monsp.* 289.

Centaurea calycibus spinosis, foliis lanceolatis dentatis. *Roy. lugdb.* 142.

Stœbe squamis asperis. *Bauh. pin.* 273. *Raj. bist.* 319.

Jacea, cum spinosis capitulis, purpurea tenuifolia. *Bauh. bist.* 3. p. 33.

Habitat Montipellii, inque Hetruria, Lusitania.

* *Calcitrapæ calycinis spinis compositis.*

triophora.

39. *CENTAUREA* calycibus duplicato-spinosis lanatis,

SYNGENESIA: POLYGAM. FRUSTRAN. 917.

tis, foliis semidecurrentibus integris sinuatisque. *Hort. upf.* 272. *Roy. lugdb.* 140.

Centaurea calycibus duplicato-spinosis, foliis decurrentibus integris. Hort. cliff. 423.

Calcitrapa lutea, alato caule, capite eriophoro. Vaill. act. 1718. p. 212.

Carduus lusitanicus canescens. alato caule, capite lanuginoso, Tournesf. inst. 441.

Habitat in Lusitania. ☉

40. *CENTAUREA calycibus subduplicato-spinosis. foliis sessilibus lateralibus pinnatifidis linearibus dentatis. Hort. upf.* 273. *

Centaurea calycibus subulato-spinosis sessilibus, foliis linearibus pinnatifidis. Hort. cliff. 423. *Roy. lugdb.* 140. *Gort. gelr.* 496. *Dalib. paris.* 266. *Sauv. monsp.* 288.

Carduus stellatus s. Calcitrapa. Bauh. hist. 3. p. 89.

Carduus stellatus, foliis papaveris erratici. Bauh. pin. 387. *Raj. hist.* 317.

Carduus muricatus vulgo Calcitrapa dictus. Clus. hist. 2. p. 7.

Hippophæstum. Col. phytob. 85. t. 24.

Habitat in Helvetia, Anglia & Europa australiori secus vias. ☉

Flos hinc purpureus, nec ut in Solstitiali, Melitenensi aut Siculo luteus.

41. *CENTAUREA calycibus duplicato-spinosis, foliis solstitialis, rameis decurrentibus inermibus lanceolatis; caulinis lyrato-pinnatifidis. Hort. cliff.* 423. n. 21. *Hort. upf.* 272. * *Roy. lugdb.* 140. *Dalib. paris.* 267. *Sauv. monsp.* 288.

Carduus stellatus luteus, foliis cyani. Bauh. pin. 387. *Raj. hist.* 317.

Carduus stellatus mitior apulus. Col. ecphr. 1. p. 30. t. 31.

Habitat in Gallia, Anglia, Italia. ☉

Flores terminales, sæpius conferti, foliis lanceolato-linearibus obvallati.

42. *CENTAUREA calycibus duplicato-spinosis subrotundis lateralibus, foliis decurrentibus lanceolatis sinuosis inermibus. Roy. lugdb.* 140. *Sauv. monsp.* 288.

Jacea melitenensis, capitulis conglobatis. Bocc. sic. 65. t. 35. *Morif. hist.* 3. p. 145. *Raj. hist.* 323.

Habitat in Melita. ☉

Folia viscida, mollia: ima sinuata, spærbulata. Sauv.

M m m 3

43.

918 SYNGENESIA: POLYGAM. FRUSTRAN.

- scabra.* 43. CEN TAUREA calycibus ciliato-spinosis. foliis ly-
ratis inermibus incanis, capitulis terminalibus. *Sauv.*
monsp. 290.
Jacea ficula, cichorii folio flore luteo, capite spinoso.
Magn. monsp. 138. *Morif. hist.* 3. p. 144 f. 7. t. 28 f. 26.
Habitat in Sicilia. 2
- Centauroides* 44. CEN TAUREA calycibus cillatis spinosis, foliis pin-
natifidis: laciniis superioribus majoribus. *Roy. lugdb.*
141. n. 27.
Jacea lutea spinosa centauroides. *Baub. pin.* 272.
Jacea laevis maxima centauroides apula. *Cul. cephr.* 1.
p. 33. t. 35.
Habitat in Italia.
- Collina.* 45. CEN TAUREA calycibus ciliatis spinosis, foliis bi-
pinnatifidis.
Jacea lutea, capite spinoso. *Baub. pin.* 272.
Jacea lutea VI. *Tabern. hist.* 436.
Habitat in Italia, Hispania.
- * *Crocodiloidea spinis simplicibus.*
- salmantica.* 46. CEN TAUREA calycibus setula reflexa spinosis gla-
bris, foliis dentato-pinnatifidis serratis. *Hort. cliff.*
421. *Roy. lugdb.* 138. *Sauv. monsp.* 287.
Jacea major, foliis cichoraceis mollibus, flore stamineo.
Morif. hist. 3. p. 140. f. 7. t. 26. f. 14.
Jacea major. foliis cichoraceis mollibus lanuginosis.
Baub. pin. 273.
Storae salmantica 1. *Clus. hist.* 2. p. 9. *Raj. hist.* 324.
Habitat in Europa australi. 2
Squamæ calycinae ita parvae sunt, ut vix manifestæ.
- panicata.* 47. CEN TAUREA calycibus simplicissime spinosis, fo-
liis inferioribus pinnatifidis, superioribus lanceolatis,
pedunculis fougissimis. *Roy. lugdb.* 141.
Jacea cyanoides, echinato capite. *Baub. pin.* 272. *Me-
rif. hist.* 3. p. 145.
Cyanoides flos. *Dod. pempt.* 251.
Habitat in Hispania. ☉
- peregrina.* 48. CEN TAUREA calycibus setaceo-spinosis, foliis
lanceolatis petiolatis; inferne dentatis. *Hort. cliff.* 423.
Roy. lugdb. 141.
Centaureum majus. folio molli acuto laciniato, flore
aureo magno, calyce spinoso. *Boerh. lugdb.* 1. p. 144.
Habitat in Europa australi. 3

SYNGENESIA: POLYGAM. NECESSARIA. 919

49. **CENTAUREA** calycibus simplicissime spinosis, foliis pinnatifidis integerrimis: lacinia extima majore dentata.

Cyanus purpureus, capite spinoso, syriacus. *Harr. rar.*
927. t. 503. bon.

Habitat in Syria.

50. **CENTAUREA** calycibus setaceo-spinosis, foliis decurrentibus sinuatis spinosis. *Hort. cliff.* 424. *Roy. lugdb.* 141. *Sauv. monsp.* 289.

Carduus tomentosus, capitulo minore. *Bauh. pin.* 382.

Carduus galactites. *Bauh. hist.* 3. p. 54.

- β. Carduus creticus non maculatus, caule alato. *Tournef. cor.* 31.

Habitat in Europa australi. &

POLYGAMIA NECESSARIA.

MILLERIA 871

1. **MILLERIA** foliis cordatis, pedunculis dichotomis. *quinqueflora*
Hort. cliff. 425. * *Roy. lugdb.* 182.

Milleria annua erecta, floribus spicatis luteis. *Mart. cent.*
41. t. 41.

- β. Milleria annua ramosior, foliis maculatis profundius serratis. *Mart. cent.* 47. t. 47. f. 1.

Habitat in Panama, Vera Cruce.

2. **MILLERIA** foliis ovatis, pedunculis simplicissimis. *biflora.*

Hort. cliff. 425. t. 25. * *Hort. upl.* 275. *Roy. lugdb.* 182.

Milleria annua erecta minor, foliis parietariæ, floribus ex foliorum alis. *Mart. cent.* 47. t. 47. f. 1.

Habitat in Campechia. ☉

SILPHIUM. 872

1. **SILPHIUM** foliis alternis pinnato-sinuatis.

laciniatum.

Bidens americana frutescens, sphondylii folio & facie.

Plum. spec. 10.

Habitat in America septentrionali, Mississipi. Collinson. &

Radix perennis. Caulis bis humane altitudinis, cassioe pollicis, simplicis-sinus, inferne laevis, superne ruberculis fuscis pilisque patulis albis scaber, teres. Folia alterna, petiolata, bipedalia longitudine, pedalia latitudine, basi amplexicaulis, pinnatifida: Laciniis utrinque 4 f. 5, distantibus, angustis, dentato-sinuatis

M m in 4

fco-

scabra, costis utrinque elevatis solidissimis; margo foliorum superiorum purpurascens. Cal. ex squamis 10, in magnas subulatas spinas exeuntibus. Petala radiis 30, longitudine calycis; Stylo bifido, tenui. Flosculi disci multi, flavi, paleis distincti, basi attenuati, Stylo simplici. Semina in radio membranacea, ovalia, emarginata, aristis 2 parvis. Semina disci columnaria, tetragona, abortientia.

- Asteriscus.* 2. **SILPHIUM** foliis indivisis sessilibus oppositis; inferioribus alternis.
Silphium foliis oppositis. Roy. Lugdb. 181.
Silphium. Hort. cliff. 494. Gron. virg. 106.
Asteriscus coronæ solis folio & facie. Dill. elsb. 42. t. 37. f. 42.
Habitat in Virginia, Carolina. 2
- Helianthoides.* 3. **SILPHIUM** foliis oppositis petiolatis serratis. *Gron. virg. 181.*
Chrysanthemum marianum, virgæ aureæ americanæ foliis, florum petalis tridentatis. Pluk. mant. 46.
Habitat in Virginia.
- Solidaginoides.* 4. **SILPHIUM** foliis oppositis lanceolatis petiolatis acutis serratis.
Habitat in Virginia. D. Gronovius.
Caulis glaber. Folia subtus glabra. Calyx serie duplici foliorum. Radix petala leviter trifida; Disci abortientia.
- trifoliatum.* 5. **SILPHIUM** foliis ternis. *Roy. Lugdb. 181.*
Chrysanthemum virginianum, foliis asperis tribus s. quaternis ad genicula litis. Moris. hist. 3. p. 24. f. 6. t. 3. p. 68. Raj. suppl. 211.
Habitat in Virginia. 2

CHRYSOGONUM.

- virginianum.* 1. **CHRYSOGONUM** petiolis folio longioribus.
Chrysogonum. Hort. cliff. 424. Gron. virg. 106.
Chrysanthemum virginianum. villosum, disco luteo: quinque petalis crenato. Pluk. alm. 100. t. 83. f. 4. & t. 242. f. 3. Raj. suppl. 213.
Habitat in Virginia
- peruvianum.* 2. **CHRYSOGONUM** foliis sessilibus amplexicaulibus, calycibus sessilibus squamosis.
Eupatoriophalacron folio trinervi subrotundo, flore minore luteo radiato. Faw. peruv. 2. p. 266. t. 50.

Ha-

Habitat in Peru. D. Jussieu. ☉

Caulis teres, erectus, compositus. Folia opposita, sessilia, amplexicaulia, cordato-oblonga, glabra, subtrimeria, integerrima, margine scabra. Calyx terminalis, sessilis, solitarius, glaber, cylindrico-oblongus, imbricatus squamis raris, obtusis. Corolla lutea; Radius pentapetalus, longitudine calycis, pistillis bifidis. Discus flosculus 5 paleis distinctus. Rami ex alis (non flori proximis) caulis similes.

MELAMPODIUM. 874

1. MELAMPODIUM. *Hort. cliff. 425.**

AMERICANUM

Habitat in Vera Cruce.

CALENDULA. 875

1. CALENDULA feminibus radii cymbiformibus echi-officinalis. natis: disci bicornibus. *Hort. cliff. 425. Hort. ups. 274. Fl. succ. 712. Mat. 407. Roy. lugdb. 177. Delib. paris. 268*

Caltha vulgaris. Bauh. pin. 275.

β. *Caltha arvensis. Bauh. pin. 275.*

γ. *Caltha polyanthos major. Bauh. pin. 275.*

δ. *Caltha floribus reflexis. Bauh. pin. 275.*

ε. *Caltha prolifera, majoribus s. minoribus floribus. Bauh. pin. 275.*

Habitat in Europæ arvis. ☉

2. CALENDULA foliis lanceolatis denticulatis, pedunculis filiformibus. *Hort. ups. 274.*

Calendula foliis dentatis. Roy. lugdb. 177.

Calendula feminibus radii obsolete serratis; disci cordatis. Hort. cliff. 425.

Calendula humilis africana, flore intus albo, foris violaceo simplici. Herm. lugdb. 104. t. 105. Breyn. ic. 26. t. 14. Stiff. bot. 59. t. 59.

Habitat in Æthiopia. ☉

3. CALENDULA foliis lanceolatis dentatis, pedunculis superne incrassatis. *Hort. cliff. 274.*

Calendula foliis radicalibus sinuatis, caulinis superne denticulatis. Roy. lugdb. 177.

Cardispermum africanum pubescens, foliis incis, parvo flore. Trant. act. paris. 1724. p. 39. t. 2.

Habitat in Æthiopia. ☉

M m m 5

4. CA-

922 SYNGENESIA: POLYGAM. NECESSARIA.

mediculis

CALENDULA foliis lanceolatis sinuato-dentatis, caule nudo.

Calendula africana, foliis scabiosæ argenteæ viridibus. *Herm. afr. 5.*

Bellis florum pediculis peneaphyllis, foliis incis. Comm. hort. 2. p. 66. t. 33.

Caltha africana, flore intus albo extus leviter violaceo, semine plano cordato. *Boerb. lugdb. 1. p. 113.*

Habitat in Æthiopia.

graminifolia 5. **CALENDULA** foliis linearibus subintegerrimis, caule subnudo.

Calendula foliis linearibus denticulatis & integerrimis Roy. lugdb. 177.

Calendula africana, foliis gramineis rarius dentatis aureis. *Herm. afr. 5.*

Caltha africana, foliis croci angustis, florum petalis externe purpurascens internis albis. *Boerb. lugdb. 1. p. 113.*

Dimorphotheca statice folio. Vaill. afr. 1720. p. 280.

Bellis africana, florum pediculis foliosis foliis angustis & integris. *Comm. hort. 2. p. 67. t. 34.*

Habitat in Æthiopia.

Anne h. e. posteriores 3. 4. 5. e C. pluviali olim oritur hinc? See snadet lucas, semina, color corolla, habitus.

576

ARCTOTIS.

trifidis.

1. **ARCTOTIS** flosculis radiantibus vicinis tripartitis.

Arctotis foliis pinnatifidis, caule herbaceo, petalis radii profunde trifidis. Roy. lugdb. 179.

Aneimonospermum arctotis, foliis & racie taraxaci incanis. Breyn. 11. 27. t. 15.

Habitat in Æthiopia. ☉

Calendula.

2. **ARCTOTIS** flosculis radiantibus duodenis subintegris.

Arctotis foliis ovatis dentatis: petiolis longissimis superne dentatis, caule ramoso. Hort. cliff. 412. Roy. lugdb. 178.

Aneimonospermum africana, jacobææ maritimæ foliis, flore sulphureo. Comm. rar. 36. t. 36.

Habitat in Æthiopia. ☉

aspera.

3. **ARCTOTIS** foliis pinnato-sinuatis laciniis oblongis dentatis. *Hort. cliff. 412. Hort. nup. 275. Roy. lugdb. 178.*

Ar-

SYNGENESIA: POLYGAM NECESSARIA. 923

Arctotheca jacobææ folio, radiis florum intus luteis, extus purpureis. *Vaill. alt.* 1720. p. 428.

Anemonospermus africana, foliis cardui beuedicti, florum radiis intus sulphureis. *Comm. hort.* 2. p. 43. t. 22.

- β. *Anemonospermus afra*, folio jacobææ tenuiter laciniato, flore aurantio pulcherrimo. *Boerh. lugdb.* 1. p. 100. t. 100.

Habitat in Æthiopia.

4. *ARCTOTIS* foliis lanceolato-linearibus integris den- *angustifolia* ticulatis. *Hort. cliff.* 412. *Roy. lugab.* 178.

Anemonospermus afra, folio serrato rigido, flore intus sulphureo extus puniceo. *Boerb. lugdb.* 1 p. 100.

Habitat in Æthiopia.

OSTEOSPERMUM. 877

1. *OSTEOSPERMUM* foliis oppositis palmatis. *Hort. Uvedalia* cliff. 424. *Hort. nps.* 274. *Roy. lugdb.* 179. *Gron. virg.* 106,

Chrysanthemum, angulosis platani foliis, virginianum *Pluk. alm.* 99. t. 83 f. 3.

Chrysanthemum pereune virginianum majus, platani orientalis folio. *Morif. hist.* 3. p. 22. f. 6. t. 7. f. 55.

Habitat in Virginia. 2. b

2. *OSTEOSPERMUM* spinis ramosis. *Hort. cliff.* 424. *spinosum* *Roy. lugdb.* 179.

Chrysanthemoides osteospermum africanum odoratum spinosum & viscosum. *Comm. hort.* 2. p. 85. t. 43.

Chrysanthemum africanum frutescens spinosum. *Volk. norib.* 105. t. 105.

Habitat in Æthiopia. b

3. *OSTEOSPERMUM* foliis ovalibus serratis. *Hort. moniliferum* cliff. 424. *Roy. lugab.* 179.

Chrysanthemoides atrum, populi albæ foliis. *T. Dill. elib.* 80. t. 68. f. 79.

Chrysanthemum africanum frutescens, telephii fere foliis crassis, osteocarpon. *Pluk. anatib.* 55. t. 382. f. 4.

Chrysanthemum arborescens æthiopicum, foliis populi albæ *Breyn. cent.* 145. t. 76.

Habitat in Æthiopia. b

4. O-

polygaloides. 4. *OSTEOSPERMUM* foliis lanceolatis imbricatis sessilibus. *Roy. lugdb.* 179.

Monilifera polygalæ foliis. *Vaill. act.* 1720. p. 374.

Chrysanthemum fruticosum, *polygoni* foliis, africanum, caulibus scabris, flore minore. *Pluk. mant.* 47. t. 382.

Habitat in Æthiopia. 5

878 OTHONNA.

* *Herbaceæ.*

gerifolia. 1. *OTHONNA* foliis reniformibus suborbiculatis denticulatis petiolatis.

Solidago foliis reniformibus suborbiculatis denticulatis. *Hort. cliff.* 410.

Jacobæa africana, *hederæ terrestris* folio, repens. *Comm. mel. hort.* 2. p. 145. t. 73.

Habitat in Æthiopia. 2

bulbosa. 2. *OTHONNA* foliis ovatis integerrimis, caule herbaceo.

Jacobææ affinis planta tuberosa capitis b. *spei. Breyn. cent.* 2. 66. *Morif. hist.* 3. p. 111. f. 7. t. 18. f. 33.

Habitat in Æthiopia. 2

semibifolia. 3. *OTHONNA* foliis amplexicaulibus sinuatis difformibus.

Jacobæa, *fonchi* folio, flore purpureo amplo, africana. *Breyn. ic.* 31. t. 21. f. 1.

Habitat in Æthiopia.

sibirica. 4. *OTHONNA* foliis subcordatis, caule subnudo simplicissimo. *Hort. ups.* 273.

Jacobæastrum *cacaliæ* folio. *Amm. ruth.* 221. t. 24.

Jacobæa orientalis, *cacaliæ* folio. *Tournef. cor.* 37.

Jacobæoides ari folio crenato. *Vaill. act.* 1720. p. 300.

Solidago foliis cordatis subrotundis dentatis petiolatis. *Gmel. sib.* 2. p. 169.

Habitat in Sibiria. 2

palustris. 5. *OTHONNA* foliis lanceolatis dentato-sinuatis.

Othonna palustris. *It. scan.* 241. 95. *

Solidago foliis inferioribus lanceolatis serrato-sinuatis; superioribus integris amplexicaulibus. *Hort. cliff.* 410.

Fl. sacc. 687. *Roy. lugdb.* 162. *Gmel. sib.* 2. p. 158. t. 72.

Jacobæa aquatica *elator*, foliis magis dissectis. *Morif. hist.* 3. p. 110. f. 7. t. 19. f. 24.

Cony.

SYNGENESIA: POLYGAM. NECESSARIA. 925

Conyza aquatica laciniata. Baub. pin. 266.

Habitat in Europæ paludibus maritimis. &

6. OTHONNA foliis lanceolatis subdenticulatis villosis, integrifolia. floribus umbellatis.

Solidago foliis lanceolatis sessilibus integris denticulatis, floribus umbellatis, involuero setaceo. Gmel. fib. 2. p. 154. t. 71.

Jacobæa montana lanuginosa angustifolia non laciniata. Baub pin. 131. Meris. hist. 3. p. 111. f. 7. t. 12. f. 28.

Jacobæa pannonica 2. Clus. hist. 2. p. 22.

Habitat in Alpibus Pyrenaicis, Helveticis, Austriacis, Sibiricis. &

7. OTHONNA foliis lanceolatis integris subdentatis vil-helenitis lois, floribus umbellatis.

Conyza incana. Baub. pin. 265.

Conyza helenitis mellita incana. Lob. ic. 347. sec. figuram.

Habitat in Sibiria. D. Gmelin. Gallia. &

Caulis sesquipedalis, simplicissimus. Folia lanceolata, sursum sensim minora, albis villis utrinque subhirsuta, margine minimis dentibus notata. Umbella terminalis: pedunculis simplicissimis, unifloris, intermedio brevissimo. Corolla lutea. Ex Sibiria.

8. OTHONNA foliis pinnatifidis subvillosis: laciniis si-Cineraria, nuatis, caule herbaceo.

Jacobæa maritima f. Cineraria latifolia. Baub. pin. 131. prodr. 69.

Habitat in Canada.

Similima O. maritimæ, at Folia non tomentosa, sed parum villosa præsertim subtus; Corollæ radius non recurvatus, sed patens. Caulis non fruticans. Calyx similimus maritimæ, sed apice parum sphucelatus, quod illa non habet.

** Fruticosæ.*

9. OTHONNA foliis pinnatifidis tomentosis, laciniis si-maritima, niis sinuatis, caule fruticoso. Hort. ups. 273.

Solidago foliis pinnatifidis: laciniis sinuatis, corymbis racemosis. Hort. cliff. 410.

Jacobæa maritima. Baub. pin. 131.

Cineraria. Dod. pempt. 642.

Habitat ad Maris inferi littora. &

926 SYNGENESIA: POLYGAM. NECESSARIA.

- postinata.* 10. OTHONNA foliis pinnatifidis: laciniis linearibus parallelis. *Hort. cliff.* 419. *Roy. lugdb.* 179.
Jacobæa africana frutescens, foliis absinthii umbelliferi incanis. *Comm. hort.* 2. p. 137. t. 69. *Raj. suppl.*
 β. *Jacobæa absinthites*, tomentosis cinerariæ foliis, æthiopica, calice integro summis oris dentato. *Pluk. alm.* 106.
Habitat in Æthiopia. ♀
- abrotanifolia* 11. OTHONNA foliis multifido-pinnatis linearibus. *Roy. lugdb.* 380.
Jacobæa africana fruticans, foliis abrotani f. crithmi, major & minor. *Volk narib.* 225. t. 225. *Raj. suppl.* 179.
Habitat in Æthiopia. ♀
- coronopifolia* 12. OTHONNA foliis infimis lanceolatis integerrimis, superioribus sinuato-dentatis. *Hort. cliff.* 419. *Roy. lugdb.* 179.
Jacobæa africana frutescens, coronopi folio. *Comm. hort.* 2. p. 139 t. 70. *Raj. suppl.* 175.
Habitat in Æthiopia. ♀
- cheirifolia.* 13. OTHONNA foliis lanceolatis integerrimis. *Hort. cliff.* 419. *Roy. lugdb.* 179.
Jacobæa africana frutescens, crassis & succulentis foliis. *Comm. hort.* 2. p. 147. t. 74. *Raj. suppl.* 174.
 β. *Aster fruticosus africanus* luteus, foliis thymelææ. *Raj. suppl.* 167
Habitat in Æthiopia. ♀
- dentata.* 14. OTHONNA foliis ovato-cuneiformibus dentatis. *Solidago* foliis oblongis dentatis glabris, floribus magnis umbellatis. *Burm. afr.* 164. t. 59.
Jacobæa africana lutea, folio obtuso dentato. *Herm afr.* 14.
Habitat in Æthiopia.

POLYMNIA.

- 579
canadensis. 1. POLYMNIA. *Gen. nov.* 1181. t. 1. f. 5. *
Habitat in Canada. *Kalm.* 2

ERIOCEPHALUS.

- 180
africanus. 1. ERIOCEPHALUS.
Eriocephalus foliis integris divisisque *Hort. cliff.* 424. *Roy. lugdb.* 178.
Eriocephalus sempervirens, foliis fasciculatis & digitatis. *Dill. elsb.* 132. t. 110. f. 134.

Abro-

SYNGENESIA: POLYGAM. NECESSARIA 927

Abrotannus africanum, folio tereti tridentato. *Walsh.*
hort. 1. t. 1.

Habitat in Æthiopia. b

FILAGO. 881

1. **FILAGO** floribus sessilibus terminalibus, foliis flora-pygmae-
libus majoribus.

Gnathaliu caule simplicissimo, foliis superioribus ma-
joribus. *Roy. lugdb. 147.*

Gnaphalium (species omnes). *Vaill. del. 1719. p. 314.*

Gnaphalium roseum hortensie. Baub. pin. 263. prodr.
122. Barr. ic. 127.

Habitat in Europa australi. c

2. **FILAGO** tomentosa, corymbo subramoso, foliis ob-maritima,
longis obtusis crenatis.

Santolina corymbo terminali subdiviso, foliis oblongis
integerrimis obtusis. *Hort. cliff. 398. Roy. lugdb. 147.*
Sauv. monsp. 80.

Gnaphalium maritimum. Baub. pin. 263.

Chrysanthemum perenne gnaphaloides maritimum. Mo-
rif. hist. 3. p. 81. f. 6. t. 4. f. 47.

Habitat in Europæ australis, marisque mediterranei
littoribus.

MICROPUS. 882

1. **MICROPUS. Hors. upf. 275. Roy. lugdb. 145. supinus.**

Gnaphalium supinum, echinato semine. Pluk. alm. 171.
t. 187. f. 6. Raj. suppl. 191.

Pseudo-Gnaphalium supinum, semine echinato. Morif.
hist. 3. p. 93.

Habitat in Lusitanie maritimis. c

SPHÆRANTHUS. 883

1. **SPHÆRANTHUS. Roy. lugdb. 145. Fl. zeyl. 312. indicus**

Sphæranthus purpurea alata ferrata. Burm. zeyl. 220. t.
94. f. 3.

Scabiola indica major, caule & pediculis foliatis. Pluk.
phys. 312. f. 6. t. 108. f. 7. Raj. suppl. 235.

Adaca-manjen. Rheed. mal. 10. p. 85. t. 43. Raj. suppl.
241.

Habitat in India

MO.

MONOGAMIA.

584

SERIPHIMUM.

cinereus

1. SERIPHIMUM floribus racemosis, foliis patentibus.
*Artemisia floribus simplicibus. Hort. cliff. 403. **
Breynia cineroides capensis. Pet. gaz. 2. t. 3. f. 9.
Tamariscus æthiopicus, coridis folio glabro, herbæ impiæ capitulis in spicam sessilibus. Pluk. mant. 178. t. 297. f. 1.
Frutex cinereus muscosus, herbæ impiæ capitulis in spicam sessilibus. Pluk. alm. 159. mant. 85.
Habitat in Æthiopia. ♀

plumosum.

2. SERIPHIMUM floribus spicatis, foliis linearibus.
Stœbe floribus spicatis, calycibus unitoris. Roy. lugdb. 145.
Breynia capensis, capitulis albis plumosis. Pet. gaz. 9. t. 5. f. 4.
Habitat in Æthiopia. ♀

fuscum.

3. SERIPHIMUM floribus capitatis, foliis imbricatis.
Eupatorium ericoides capit. bonæ spei. Breyn. cent. 2. 69. Moris. hist. 3. p. 97. f. 7. t. 18. f. 10.
Abrotanoides capensis, ericæ folio. Pet. gaz. 9. t. 5. f. 2.
Habitat in Æthiopia. ♀
Habitus Erica magnitudine, foliis, flosculis. Fruticulus Ramis lignosis, teretibus, inæqualibus. Folia acerosa, brevissima, filiformia, conferta, imbricata, tomentosa. Capitula florum terminalia, globosa ex pluribus, sessilia. Calyces glabri, nigricantes.

585

CORYMBIUM.

africanum.

1. CORYMBIUM. *Hort. cliff. 494.*
Corymbium foliis ad radicem longissimis liratis. Burm. afr. 189. t. 70. f. 1.
Bupleurifolia, semine papposo, valerianoides umbellata, cauliculo scabro. Pluk. alm. 73. t. 272. f. 5.
Bupleuri similis planta æthiopica, ad caulium nodos tomentosa. Pluk. alm. 73. t. 272. f. 4.
Habitat in Æthiopia. ♀

586

IASIONE.

montana.

1. IASIONE.

Jasione

Jasione foliis lineari-lanceolatis obsolete ferratis. *Hort. cliff.* 426. *Fl. suec.* 713. *Roy. lugdb.* 191. *Hall. belv.* 496. *Dalib. parif.* 268.

Rapunculus scabiosæ capitulo cæruleo. *Bauh. pin.* 92.

Rapuntium montanum capitatum leptophyllon. *Cal. ecphr.* 1. p. 226. t. 227.

Aphyllanthes. *Dalech. hist.* 804.

Habitat in Europæ collibus siccissimis. ☉

LOBELIA. 887

* *Foliis integerrimis.*

1. LOBELIA frutescens, foliis ovali-oblongis integer-plumierii. *Fl. zeyl.* 313.

Lobelia frutescens, portulacæ folio. *Plum. gen.* 21.

Catesb. car. 1. p. 79. t. 79.

Habitat in Indiis. h

Racemus dichotomus, bracteis subulatis, oppositis, interne ad basin villosis. Fructus bilocularis.

2. LOBELIA fruticosa, foliis linearibus confertis integer-pinifolia. *Roy. lugdb.* 241.

Rapunculus æthiopicus, violaceo galeato flore, foliis pinastri. *Breyn. cent.* 173. t. 87.

Rapunculus fruticosus, foliis nervosis acutis, floribus in ramulorum summo. *Burm. afr.* 111. t. 41. f. 2.

Rapunculus flore cæruleo galericulato, rosmarini folio angustiore. *Raj. suppl.* 384.

Rapuntium æthiopicum, violaceo galeato flore, foliis pinastri. *Raj. suppl.* 385.

Habitat in Æthiopia. h

3. LOBELIA foliis linearibus bilocularibus integerrimis, *Dortmanna.* caule subnudo. *Fl. suec.* 714.

Lobelia foliis bilocularibus subulatis. *Fl. lepp.* 279.

Hort. cliff. 426.

Dortmanna lacustris, floribus sparsis pendulis. *Rudb. act. nps.* 1720. p. 97. t. 2. *

Gladiolus stagnalis f. lacustris. *Rudb. elyf.* 2. p. 202. f. 15.

Gladiolus stagnalis dortmanni. *Clus. cur.* 40. *Raj. bist.* 1911. *

Habitat in Europæ frigidissimæ lacubus & ripis. &

4. LOBELIA foliis lanceolatis integerrimis, racemo spicato. *Tropa.*

Rapuntium spicatum, foliis auritis, vulgo *Trapa*. *Few. peruv.* 2. p. 739. t. 29.
Habitat in Peru.

kalmii.

5. *LOBELIA* caule erecto, foliis lanceolato-linearibus obtusiusculis alternis integerrimis, racemo terminali. *Rapuntium canadense pumilum*, linearæ folio. *Tournef. inst.* 164.

Habitat in Canada. Kalm. ☉

Planta pedalis, Caule tereti, vix ramoso. Folia linearilanceolata inferne, superne linearia, integerrima, obtusiuscula, nuda, sessilia. Racemus terminalis, laxis, Pedunculis alternis, unisloris, ex alis supremorum foliorum, foliolo instructis. Flores caules.

- paniculata.* 6. *LOBELIA* foliis linearibus integerrimis, panicula dichotoma.

Rapuntium, foliis linearibus acutis, floribus gemellis, polyanthos. *Burm. afr.* 100. t. 38. f. 3.

Habitat in Æthiopia.

cornuta.

7. *LOBELIA* foliis ovatis petiolatis, staminibus florum longissimis.

Habitat in Cayenna.

Plantam apud D. Royenum vidi, utram Folia integerrima e memoria excidit.

* *Caule erecto, foliis incis.*

Phytenma.

8. *LOBELIA* foliis ovatis crenatis, caule subnudo, antheris hirsutis distinctis.

Habitat in Æthiopia.

longiflora.

9. *LOBELIA* foliis lanceolatis dentatis, pedunculis brevissimis lateralibus, tubo corollæ filiformi longissimo. *Roy. lugdb.* 241.

Rapunculus aquaticus, foliis cichorei, flore albo: tubo longissimo. *Sloan. jam.* 58. *hist.* 1. p. 158. t. 101. f. 2. *Raj. suppl.* 383.

Habitat in Jamaica ad ripas.

Cardinalis.

10. *LOBELIA* caule erecto, foliis lanceolatis serratis, spica terminali. *Hort. cliff.* 426. *Hort. apj.* 276. *Roy. lugdb.* 241. *Gron. virg.* 107.

Rapuntium galeatum virginianum, coccineo flore majore. *Morif. hist.* 2. p. 466. f. 5. t. 5. f. 54.

Rapuntium maximum, coccineo spicato flore. *Herm. mex.* 879. t. 880.

Habitat in Virginia. 2

11. LO-

11. LOBELIA caule erecto, foliis ovato-lanceolatis *spkilanka*. crenatis, calycum sinibus reflexis. *Hort. cliff.* 426. *
Roy. lugdb. 241.
 Rapunculus galeatus virginianus, flore violaceo majore.
Morif. hist. 2. p. 466. f. 5. t. 5. f. 55.
 Trachelium americanum, flore cæruleo. *Lob. ic.*
Habitat in Virginia
12. LOBELIA caule erecto, foliis ovatis subserratis *pe-inflato*, dunculo longioribus, capsulis inflatis. *Hort. upf.* 276.
AE. upf. 1741. p. 23. t. 1.
 Lobelia caule erecto brachiato, foliis ovato-lanceolatis obsolete incisiss, capsulis inflatis. *Hort. cliff.* 500. *Roy. lugdb.* 528.
Habitat in Virginia, Canada. ☉
13. LOBELIA caule erecto, foliis cordatis obsolete den- *cliffortiano*, tatis petiolatis, corymbo terminali. *Hort. cliff.* 426. t. 26. * *Gron. virg.* 107.
 Rapuntium trachelii folio, flore purpurascens. *Plum. spec.* 5.
Habitat in Virginia. Canada. ☉
- * *Caule prostrato, foliis incisiss.*
14. LOBELIA caule erecto, foliis inferioribus subrotun- *arem*, dis crenatis, superioribus lanceolatis serratis, floribus racemosis.
 Lobelia foliis oblongo-ovatis, floribus laxè spicatis.
Goett. stamp. 1. p. 35. *Dalib. paris.* 268.
 Rapunculus galeatus blesensis f. soloniensis, flore violaceo minore. *Morif. hist.* 2. p. 407. f. 5. t. 5. f. 56.
 Rapuntium urens soloniense. *Morif. bles.* 300. *Bocc. sic.* 20. t. 11. *Raj. hist.* 746.
 Draba flore cæruleo galeato. *Bauh. prodr.* 53. pin. 110.
Habitat in Gallia. ☉
15. LOBELIA caule prostrato, foliis lanceolatis ovalibus *Laurentia*, crenatis, caule ramoso, pedunculis solitariis unifloris longissimis.
 Laurentia annua minima, flore cæruleo. *Mich. gen.* 18. t. 14.
 Rapunculus aquaticus repens, flore cæruleo inaperto.
Raj. suppl. 383. *Bocc. muf.* 35. t. 27.
Habitat in Italia. ☉

- Erans.* 16. LOBELIA caule patulo, foliis lanceolatis subdentatis, caule erecto ramoso, pedunculis longissimis. Campanula minor africana, erini facie, flore violaceo, caulibus erectis. *Herm. lugdb.* 110. t. 111.
Habitat in Æthiopia. ☉
Caulis *spithameus, filiformis, inferne ramosus.* Folia *radicalia ovalia, caulina lanceolata, omnia glabra, subdentata.* Pedunculi *axillares, nudi, foliis triplo longiores, uniflori.* Corollæ *cæruleæ.*
- erinoidea.* 17. LOBELIA caulibus procumbentibus, foliis lanceolatis serratis, pedunculis lateralibus. *Hort. cliff.* 426. Campanula minor africana, erini facie, caulibus procumbentibus. *Herm. lugdb.* 108. t. 109.
Habitat in Æthiopia. ☉
- ceylanica.* 18. LOBELIA caule repente, foliis ovatis serratis: inferioribus obtusis, superioribus acutis, pedunculis unisporis, capsulisque subvillosis. Campanula ceylanica, senecionis folio, flore purpureo. *Seb. thes.* 1. p. 37. t. 22. f. 12?
Habitat in China. *Osbeck.*
Planta *parva, facie Veronica chamaedryos.* Caulis *repens.* Folia *alterna, ovata, petiolata, fere glabra: inferiora obtusa crenata; superiora acuta, argute serrata.* Pedunculi *axillares, subsolitarii, folio longiores, uniflori, pubescentes.* Calyx *subvillosus.*
- hirta.* 19. LOBELIA caulibus procumbentibus, foliis ovatis subserratis, pedunculis solitariis glabris longioribus, germinibus hirsutis.
Habitat in Æthiopia.
Caules *procumbentes.* Folia *ovata, glabra, obtuse serrata, remota, petiolata.* Ex *alis singulis Ramulus* & Pedunculus *capillaris, longus, Germine piloso.*
- lutea.* 20. LOBELIA caulibus procumbentibus, foliis lanceolatis serratis, floribus sessilibus subspicatis. Rapuntium foliis oblongis dentatis, flore luteo spicato, flagellis procumbentibus. *Burm. afr.* 101. t. 39. f. 1. Rapunculus africanus ramosus, foliis dentatis, floribus gladioli laevibus. *Herm. afr.* 18. *Raj. suppl.* 248.
Habitat in Æthiopia.
- hirsuta.* 21. LOBELIA foliis ovalibus erenatis lanatis, floribus lateralibus solitariis. Rapuntium foliis subrotundis hirtis, flore ex alis foliario. *Burm. afr.* 105. t. 40. f. 2. Ra

Rapuntium africanum galeatum, foliis oblongo-rotundis dentatis villosis. *Raj. suppl.* 385.

Rapunculus galeatus æthiopicus cæruleus, hirsutis & subrotundis crenatis foliis. *Pluk. alm.* 316.

Habitat in Æthiopia.

22. LOBELIA foliis lanceolatis dentatis, pedunculis *coronopifolia* longissimis.

Rapuntium folio longo profunde dentato, florum petalis longissimis. *Burm. afr.* 98. t. 38. f. 1.

Rapuntium æthiopicum, flore cæruleo galeato, foliis coronopi. *Herm. afr.* 18. *Breyn. cent.* t. 88.

Habitat in Æthiopia. &

23. LOBELIA foliis linearibus dentatis, spica terminali *romosa* foliosa. *Hort. cliff.* 500.

Habitat in Æthiopia. ☉

24. LOBELIA caulibus erectis, foliis ovatis dentatis. *Cheiranthus*
Cheiranthus africana, flore luteo. *Comm. hort.* 2. p. 83. t. 42.

Habitat in Africa.

25. LOBELIA foliis inferioribus pedatis.

Rapuntium tuberosum, foliis multifidis, floribus *bulbosa* spicatis. *Burm. afr.* 99. t. 38. f. 2.

Habitat in Æthiopia.

VIOLA. 888

* *Acaules.*

1. VIOLA acaulis, foliis palmatis quinquelobis dentatis *palmate* indivisisque.

Viola foliis palmatis sinuatis, stolonum reniformibus.

Gron. virg. 182.

Viola virginiana, platani fere foliis parvis & incanis.

Pluk. mant. 187.

Viola alba, folio securis romanæ effigie, floridana. *Pluk.*

amalt. 209. t. 447. f. 1.

Habitat in Virginia. &

2. VIOLA acaulis, foliis pedatis septempartitis.

pedata

Viola foliis palmatis. *Gron. virg.* 107.

Viola virginiana tricolor, foliis multifidis, cauliculo aphylo. *Pluk. alm.* 388. t. 114. f. 7.

Viola mariana, folio digitato. *Pet. succ.* 20.

Habitat in Virginia. &

N n n 3

3. VIO.

- pinnata.* 3. *VIOLA* acaulis, foliis pinnatifidis.
Viola alpina, folio in plures partes dissecto. *Bauh. pin.* 199.
Viola montana, folio multifido. *Bank. hist.* 3. p. 544.
Habitat in Sibiria & Alpibus Europæ.
- lanceolata.* 4. *VIOLA* acaulis, foliis lanceolatis crenatis.
Viola acadiensis, folio longo sinuata. *Dierv.*
Habitat in Canada, Sibiria.
Planta Canadensis a D. Kalm. allata. Folia habuit angusta, longe petiolata, nervis alternis, parallelis, obsolete serrata; Sibirica a D. Gmelino; Folia habuit ovato-lanceolata, petiolis folio dimidio brevioribus minus striata & Flores majores.
- primulifolia.* 5. *VIOLA* acaulis, foliis oblongis subcordatis: petiolis membranaceis.
Viola acaulis, foliis ovatis crenatis glabris. *Gron. virg.* 182.
Habitat in Sibiria, Virginia.
Folia cordata, oblonga, obtusissima, crenata, basi decurrentia per petiolum, omnino ut in Primula officinali.
- hirta.* 6. *VIOLA* acaulis, foliis cordatis piloso-hispidis. *Fl. suec.* 718. *Dalib. parif.* 269.
Viola martia hirsuta inodora. *Morif. hist.* 2. p. 475. f. 5. t. 35. f. 4.
Viola trachelii folio. *Raj. hist.* 1051.
Viola. Brunff. herb. 1. p. 137. f. 3.
Habitat in Europæ frigidioris nemoribus. &
- palustris.* 7. *VIOLA* acaulis, foliis reniformibus. *Hall. belv.* 501. *Fl. suec.* 717. *Dalib. parif.* 270.
Viola foliis subrotundo-cordatis; pedunculis radicatis. *Fl. lapp.* 278.
Viola palustris rotundifolia glabra. *Morif. hist.* 2. p. 475. f. 5. t. 35. f. 5.
Habitat in Europæ frigidioris paludibus. &
- odorata.* 8 *VIOLA* acaulis, foliis cordatis, stolonibus reptantibus.
Viola acaulis, stolonibus teretibus repantibus, pedunculis radicalibus. *Hort. cliff.* 427. *Fl. suec.* 715. *Mat. med.* 410. *Dalib. parif.* 268.
Viola martia purpurea, flore simplici odoro. *Bauh. pin.* 119.
Viola odorata. *Revealm. spec.* 141. t. 140.
Viola nigra f. purpurea. *Dod. pempt.* 156. t. 1. 2.
 β. *Viola martia alba.* *Bauh. pin.* 199.
 γ. *Viola martia*, multiplici flore. *Bauh. pin.* 199.
Habitat in Europæ nemoribus. &

* Con-

* *Caulescentes.*

9. VIOLA caule demum adscendente, foliis oblongo-cordatis.
cordatis.
Viola caulibus adscendentibus floriferis, foliis cordatis.
Hort. cliff. 427. *Fl. succ.* 716. *Roy. lugdb.* 430. *Gron. virg.* 182. *Hall. belv.* 501. *Dalib. paris.* 269.
Viola foliis cordatis oblongis, pedunculis subradicatis.
Fl. lapp. 277.
Viola martia inodora sylvestris. *Bauh. pin.* 199.
Viola cærulea martia inodora sylvatica, in cacumine semen ferens. *Bauh. hist.* 3. p. 543.
Habitat in Europæ apricis. 2
10. VIOLA caulibus erectis, foliis cordatis oblongis. *montana.*
Viola foliis ovato-lanceolatis, caule erecto, stipulis dentatis. *Roy. lugdb.* 430. *Dalib. paris.* 269.
Viola martia arborescens purpurea. *Bauh. pin.* 199.
Viola erecta, flore cæruleo & albo. *Morif. hist.* 2. p. 475. f. 5. t. 7. f. 7.
Viola arborescens. *Cam. epit.* 911.
Habitat in Alpibus Lapponiæ, Aultriæ, Baldo. 2
11. VIOLA caule fruticoso foliis lanceolatis integerrimis. *arborescens.*
Viola hispanica fruticosa longifolia. *Tournef. inst.* 421.
Viola hispanica fruticans. *Barr. ic.* 568.
Habitat in Hispania. 3
12. VIOLA caule abbreviato, florum nectario subulato *calcarum.*
petalis longiore, foliis subovatis.
Viola alpina purpurea, exiguis foliis. *Bauh. pin.* 199.
Viola montana cærulea grandiflora. *Tournef. inst.* 420
Melanium montanum. *Dalech. hist.* 1204.
- β. VIOLA pyrenaica longius caudata, teucryi folio. *Tournef. inst.* 421.
Viola pyrenaica, folio teucryi, serotina. *Tournef. schol.* 131. *Raj. suppl.* 510.
Habitat in Pyrenæis.
Quæ demis natura ex herba V. tricolore, adjecit floribus in hac planta.
13. VIOLA caule triquetro diffuso, foliis oblongis incisis, stipulis dentatis. *trisoien*
Fl. succ. 721. *Dalib. paris.* 269.
Viola pedunculis caulinis angulatis, stipulis oblongis pinnato-dentatis, foliis ovato-oblongis crenatis. *Hort. cliff.* 527.

Viola bicolor arvensis. Bauh. pin. 200.

Trinitatis herba. Fuchsf. hist. 803.

Jacea altera. Cam epit. 913.

9. *Viola tricolor hortensis repens. Bauh. pin. 199.*

Viola tricolor. Renealm. spec. 144. t. 140.

Jacea f. Flos trinitatis. Cam. epit. 912.

Habitat in Europæ cultis. ☉

densf. 14. *VIOLA caule erecto, foliis cordatis acuminatis.*

*Viola caulescens, foliis cordatis oblongis acuminatis. Amœn. acad. 1. p. 157. **

Habitat in Canada. Kalm.

Caulis erectus, spithameus, teretiusculus. Folia alterna, reniformi-cordata, acuminata, serrata, lævia, subtus pallida, petiolis folio dimidio brevioribus, insummis adhuc minoribus. Stipulæ binæ, erectæ, lanceolatae, marcescentes. Pedunculi ex singulis alis solitarii, nudi, longitudine foliorum. Flores corollati. Differt a sequenti foliis magis mucronatis & floribus caulinis corollatis, Caulibus firmiteribus majoribus.

mirabilis. 15. *VIOLA caule triquetro, foliis reniformi-cordatis, floribus caulinis apetalis.*

*Viola floribus radicalibus corollatis abortientibus, caulinis apetalis semiuiferis. Hort. cliff. 427. It. æl. 81. * Fl. suec. 719.*

Viola montana latifolia, flores e radice, semina in cacumine ferens. Dill. elib. 408. t. 303. f. 390.

Habitat in Germaniæ, Sueciæ nemoribus. &

biflora. 16. *VIOLA caule bifloro, foliis reniformibus ferratis.*

Viola foliis reniformibus, pedunculis caulinis. Fl. lapp. 276. Fl. suec. 720. Hort. cliff. 427. Roy. lugdb. 431.

Viola alpina rotundifolia lutea. Bauh. pin. 199.

Viola rotundifolia montana major. Pluk. alm. 388. t. 234. f. 1.

Viola alpina rotundifolia minor lutea. Pluk. alm. 388. t. 233. f. 7.

Habitat in Alpibus Lapponiæ, Austriæ, Helvetiæ, Angliæ. &

uniflora. 17. *VIOLA caule unifloro, foliis cordatis dentatis.*

Habitat in Sibiria. &

Caulis vix spithameus, nudus, firmus, apice ferens Folia tria, subalterna, cordata, dentata, subvillosa, pennatis

tiolis brevissimis insidentia. Flos unicus, pedunculo brevissimo, luteus, foliis major.

18. VIOLA caule basi ramosissimo, foliis lanceolato-linearibus integerrimis distantibus, calycibus pone æqualibus. *Fl. zeyl.* 317. *

Herba indica, flore monopetalo labiato, capsula erecta trilaterali octosperma. *Raj. suppl.* 402.

Nelam-parenda *Rheed. mal.* 9. p. 117. t. 60.

Habitat in India. 2

19. VIOLA caule procumbente, foliis lanceolatis subsuffrutescentibus ferratis confertis, calycibus pone æqualibus. *Fl. zeyl.* 318. *

Habitat in India. 3

IMPATIENS.

* *Pedunculis unifloris.*

1. IMPATIENS pedunculis unifloris solitariis, foliis chinensis oppositis ovatis, nectariis arcuatis.

Habitat in China. ☉

Caulis alternatim ramosus, ruber. Folia opposita, sessilia, ovata, subserrata. Pedunculi axillares, solitarii, folio longiores, uniflori. Flos purpureus. Nectarium valde arcuatum, crassum.

2. IMPATIENS pedunculis unifloris solitariis, foliis obovatis: serraturis lanceolatis, nectariis flore longioribus.

Valli-onapu. *Rheed. mal.* 9. p. 91. t. 48.

Habitat in India. ☉

Folia lanceolata, alterna, crenata, e singula crena mucrone prominente. Pedunculi solitarii, uniflori, longitudine fere foliorum. Flos magnitudine f. Balsaminæ, at Calcar subulatum, longitudine fere pedunculi.

3. IMPATIENS pedunculis unifloris aggregatis, foliis oppositifolia oppositis linearibus. *Fl. zeyl.* 314. *

Kondam-pallu. *Rheed. mal.* 9. p. 57. t. 31?

Habitat in Zeylonæ arenosis.

4. IMPATIENS pedunculis unifloris aggregatis, foliis cornuta lanceolatis, nectariis flore longioribus. *Fl. zeyl.* 316. *

Balsamina latifolia, floris calcari longissimo. *Burm.* zeyl. 41. t. 16. f. 1.

Habitat in Zeylona. ☉

N u n s

5. IM-

Balsamina. 5. IMPATIENS pedunculis unifloris aggregatis, foliis lanceolatis, nectariis flore brevioribus. *Hort. ups.* 276. *Impatiens* pedunculis confertis unifloris. *Hort. cliff.* 428.

Balsamina femina. *Bauh. pin.* 306.

Balsamina. *Dod. pempt.* 671.

Habitat in India. ☉

* *Pedunculis multifloris.*

triflora. 6. IMPATIENS pedunculis trifloris solitariis, foliis angustis lanceolatis, *Fl. zeyl.* 315. *

Balsamina angustifolia, floribus ternis communi pedunculo ortis. *Burm. zeyl.* 41. s. 16. f. 2.

Balsamina erecta s. femina, persicæ angusto folio, zeylanica. *Herm. par.* 105. s. 105.

Habitat in Zeylonæ paludosis.

noli tangere. 7. IMPATIENS pedunculis multifloris solitariis, foliis ovariis, geniculis caulinis tumentibus. *Fl. suec.* 722. *Dalib. parif.* 270.

Impatiens pedunculis solitariis multifloris. *Hort. cliff.* 428. *Roy. lugdb.* 431. *Hall. herb.* 405. caule angulato. *Gort. petr.* 502.

Balsamina lutea s. Noli me tangere. *Bauh. pin.* 306.

Noli me tangere. *Col. ecphr.* 1. p. 149. s. 150.

Habitat in Europæ, Canadæ nemoribus. ☉

Clas-

Classis XX.

GYNANDRIA.

DIANDRIA.

ORCHIS.

860

Bulbis indivisis.

1. ORCHIS bulbis indivisis, nectarii alis amplioribus *clisulata*.
liatis.

Orchis amboinensis, floribus albis fimbriatis. *Herm.*
parad. 209. t. 209.

Habitat in Amboina.

2. ORCHIS bulbis indivisis, nectarii labio lanceolato *ciliato*.
liato: cornu longissimo. *Aff. nps.* 1741. p. 6.

Orchis nectarii labio lanceolato ciliato, seta germine in-
torto longiore. *Roy. lugab.* 15. *Gron. virg.* 183.

Orchis palmata elegans lutea americana, cum longis
calcaribus luteis. *Morif. hist.* 3. p. 499.

Orchis marilandica grandis & procera, floribus luteis,
calcare longissimo: lobulo fimbriato. *Raj. suppl.* 588.

Habitat in Virginia, Canada. 2

3. ORCHIS bulbis indivisis, nectarii labio lanceolato *bifolia*.
integerrimo: cornu longissimo petalis patentibus. *Aff.*

nps. 1740. p. 5. *Fl. suec.* 723. *Dalib. parif.* 273.

Mat. med. 411.

Orchis alba bifolia minor, calcari oblongo. *Bauh. pin.*
83. *Pail. parif.* 151. t. 20. f. 7. *Segu. ver.* t. 15,
f. 10.

Testiculi species V. *Cam. epis.* 625.

3. ORCHIS trifolia minor. *Bauh. pin.* 83.

4. ORCHIS bifolia altera. *Bauh. pin.* 82.

Orchis bifolia latissima. *Bauh. pin.* 82.

Habitat in Europæ pascuis asperis. 2

4. ORCHIS bulbis indivisis, nectarii labio trifido, petalis
confluentibus, caule nudo.

Orchis radice rotunda, cucullo tridentato: *Gmel. sib.*
1. p. 16. t. 3. f. 2.

Habitat in Sibiria. 2

Folia radicalia ovata, lino.

5. OR-

- pyramidalis*. 5. ORCHIS bulbis indivisis, nectarii labio trifido antice
bidentato: cornu longo, petalis acuminatis. *Aët. nps*
1740. p. 7. *Dalib. parif.* 273.
Orchis radicibus subrotundis, bracteis flore brevioribus, nectarii labio trifido, seta longissima. *Hors. cliff*
429.
Orchis spica densa brevi; petalis caudatis. *Hall. it. belv.*
46.
Orchis purpurea, spica congesta pyramidali. *Raj. angl.*
3. p. 377. t. 48. *Segu. ver. t. 15. f. 11.*
Cynorchis militaris montana, spica rubente conglomerata. *Bauh. pin.* 81. *prodr.* 28.
β. Orchis globofo flore. *Bauh. pin.* 81.
γ. Orchis latifolia, hiante cucullo, altera. *Bauh. pin.* 81.
δ. Cynorchis latifolia, spica compacta. *Bauh. pin.* 81.
Habitat in Helvetiæ, Belgii, Angliæ, Galliæ arenosis, cretaceis. &
- coriophora*. 6. ORCHIS bulbis indivisis, nectarii labio trifido reflexo crenato: cornu brevi, petalis conniventibus. *Aët. nps*
1740. p. 8. *Dalib. parif.* 272.
Orchis bulbis indivisis, labello trifido punctato: lacina media longiore, cornu germinibus brevioribus. *Sauv. monsp.* 23.
Orchis odore hirci minor. *Bauh. pin.* 82. *Vaill. parif.*
149. t. 31. f. 30. 31. 32. *Hall. it. berc.* 7.
Tragorchis minor, flore fuliginoso. *Raj. hist.* 1213.
Tragorchis minor & verior gemmæ. *Lob. ic.* 177.
β. Orchis odore hirci minor, spica purpurascens. *Bauh. pin.* 82.
Habitat in Europæ australioris pascuis. &
- cubitalis*. 7. ORCHIS bulbis indivisis, nectarii labio trifido filiformi: intermedia ovata: cornu germinibus brevioribus. *Fl. zéyl.* 320. *
Habitat in Zeylona. &
- morio*. 8. ORCHIS bulbis indivisis, nectarii labio quadrifido crenulato: cornu obtuso. *Aët. nps* 1740. p. 8. *Mat. med.* 412. *Fl. suec.* 724. *Dalib. parif.* 273.
Orchis radicibus subrotundis, nectarii labio quadrifido æquali crenulato: cornu obtuso. *Roy. lugd.* 15.
Orchis morio femina. *Bauh. pin.* 82. *Vaill. parif.* t. 31. f. 13. 14. *Segu. ver. t. 15. f. 7.*
Triorchis serapias mas. *Fuchs. hist.* 559.
β. Orchis angustifolia. *Bauh. pin.* 83. *Vaill. parif. t. 31. f. 7. Or-*
33. 34.

- γ. *Orchis papilionem referens angustifolia alba. Baub. pin.*
84.
- δ. *Orchis morio mas, foliis maculatis. Baub. pin. 81, masculus,*
Testiculis IV. Cam. epit. 624.
- ε. *Orchis foliis sessilibus non maculatis. Baub. pin. 82.*
Habiat in Europæ nemoribus. &
α. *Labium æquale, calcar emarginatum, galea con-*
nivens obtusa.
β. γ. *Labium medio brevius, calcar obtusum, galea*
alis reflexis obtusis.
δ. ε. *Labium æquale, calcar integrum, galea alis*
reflexis acutis.
9. ORCHIS bulbis indivisis, nectarii labio quadrifido pun-
ctis scabro: cornu obtuso, petalis distinctis. *Act. upf.*
1741. p. 11. *Fl. succ. 726. It. æl. 45. * Dalib. parif.*
272.
- Orchis militaris pratensis humilior. T. Vaill. parif. 149.*
t. 31. f. 35. 36. Segu. ver. t. 15. f. 4.
Orchis panonica IV. Clus. hist. 1. p. 268.
Cynosorchis militaris pratensis humilior. Baub. pin. 81.
Habiat in Europæ temperate pratis. &
10. ORCHIS bulbis indivisis, nectarii labio quinquefido *militaris.*
punctis scabro: cornu obtuso, petalis confluentibus.
*Act. upf. 1740. p. 11. Fl. succ. 725. It. æl. 45. * Da-*
lib. parif. 271.
Orchis galea & alis cinereis. Baub. hist. 2. p. 755.
Orchis mas latifolia. Fuchf. hist. 554.
Cynosorchis latifolia, hiante cucullo, major. Baub.
pin. 80.
- β. *Orchis magna, latis foliis, galea fusca vel nigricante.*
I. B. Raj. angl. 3. p. 378. t. 19. f. 2.
- γ. *Orchis latifolia, hiante cucullo, minor. T. Vaill. pa-*
ris. t. 31. f. 22. 23. 24.
Cynosorchis latifolia, hiante cucullo, minor. Baub. pin.
81.
- δ. *Orchis militaris major. Tournef. inst. 438. t. 247*
Vaill. parif. 148. t. 31. f. 21. 27. 28.
Cynosorchis militaris major. Baub. pin. 81.
Habiat in Europæ temperate pratis. &
** Bulbis palmatis.*
11. ORCHIS bulbis subpalmatis rectis, nectarii cornu co-
nico: labio trilobo lateribus reflexo, bracteis flore lon-
gioribus. *Act. upf. 1740. p. 15. Fl. succ. 728. It. æl*
*48. * Dalib. parif. 274.*
Or-

Orchis radicibus palmatis, bracteis flore longioribus, nectarii labio trifido, cornu geminibus brevior. *Hort. cliff.* 429.

Orchis palmata pratensis latifolia, longis calcaribus. *Bauh. pin.* 85. *Vaill. parif.* 1. 31. f. 1. 2. 3. 4. 5.

β. Orchis palmata palustris latifolia. *Bauh. pin.* 86.

γ. Orchis palmata montana altera. *Bauh. pin.* 86.

δ. Orchis palmata sambuci odore. *Bauh. pin.* 86.

ε. Orchis palmata palustris maculata. *Bauh. pin.* 86.

Habitat in Europæ pratis. &

Petala 2 lateralia sursum flexa; tria vero conniventia.

Nectarii labium lateribus reflexum.

maculata. 12. ORCHIS bulbis palmatis patentibus, nectarii cornu germinibus brevior: labio piano, petalis dorsalibus erectis. *Act. upf.* 1740. p. 14. *Fl. succ.* 729. *Is. al.* 48. * *Dalib. parif.* 274.

Orchis palmata pratensis maculata. *Bauh. pin.* 85.

Orchis palmata montana maculata. *Bauh. pin.* 86.

Vaill. parif. 1. 31. f. 9. 10.

Satyrium basilicum femina. *Dod. pempt.* 240.

Habitat in Europæ pratis succulentis. &

Petala 3 exteriora erecta; 2 interiora conniventia. Ne-

ctarii labium trifidum, planum: lobis lateralibus majoribus crenatis: intermedio angustissimo, integerrimo.

conopsea. 13. ORCHIS bulbis palmatis, nectarii cornu setaceo germinibus longior: labio crenato. *Act. upf.* 1740. p. 13. *Fl. succ.* 727. * *Dalib. parif.* 275.

Orchis radicibus palmatis, bracteis longitudine floris, nectarii labio trifido, seta germinibus longior. *Hort. cliff.* 429.

Orchis palmata minor, calcaribus oblongis. *Bauh. pin.* 85. *Vaill. parif.* 1. 30. f. 8. 8.

Satyrium basilicum mas. *Fuchs. hist.* 712.

β. Orchis palmata angustifolia minor. *Bauh. pin.* 85.

γ. Orchis palmata pratensis maxima. *Bauh. pin.* 85.

Habitat in Europæ pratis montosis. &

flava. 14. ORCHIS bulbis palmatis, nectarii cornu filiformi longitudine germinis, labio trifido integerrimo. *Gro. virg.* 183.

Habitat in Virginia &

Folia inferiora palmaria, lanceolata, magna. Spica angusta; Flores obsolete latei.

* *Bulbis fasciculatis.*

15. ORCHIS bulbis fasciculatis, nectarii cornu longitudine *fuscescente*,
dine germinis, labio ovato basi dentato.
Orchis radicibus multis, labello quasi alato, calcare germinis longitudine. *Gmel. sib. 1. p. 20. t. 4. f. 2.*
Habitat in Sibiria. 2.
16. ORCHIS bulbis fasciculatis, nectarii labio bilobo *intersternatis*,
gerrimo, cornu longitudine germinis. *Fl. zeyl. 319. ** *rs.*
Habitat in Zeylona. 2.
17. ORCHIS bulbis fasciculatis filiformibus, nectarii labio *abortiva*,
ovato integerrimo. *Alf. upf. 1740. p. 17. Dalib.*
parif. 275.
Orchis radicibus cylindricis, bracteis flore brevioribus,
labello trifido obtuso, seta germinibus brevioribus. *Sarrv.*
monsp. 23.
Orchis abortiva violacea. *Bauh. pin. 86.*
PseudoLimodorum. *Clus. hist. 1. p. 270.*
Habitat in Gallia, Helvetia, Anglia, Italia. 2.
18. ORCHIS nectarii cornu setaceo longitudine germinis: *psycoder*.
labio tripartito ciliari.
Orchis marilandica, spica brevi conferta, floribus parvis.
calcaribus longissimis. *Raj. supp. 582.*
Orchis floribus aureis, spica habitiore congestis, bracteis
longitudine floris, labio inferiore nectarii fimbriato ca-
pillaceo. *Gron. virg. 184*
Habitat in Canada. Kalm.
Habitus *Orchidis maculatae*. Caulis foliis linearibus, a-
cuminatis, sparsis. Spica longa, densa. Corollae peta-
la 5, conniventia: quorum tria exteriora ovata: duo
interiora oblonga, subciliata. Labium reflexum, tripar-
situm: laciniis basi angustissimis, cuneiformibus, a-
pice ciliatis.
19. ORCHIS nectarii cornu longitudine germinis, labio *spectabilis*,
ovali emarginato, caule aphylo, foliis ovalibus.
Orchis flore pulcherrimo magno specioso, nectarii ga-
lea saturate caerulea, calcari niveo, foliis amplis ob-
longo-ovatis saturate viridibus. *Clayt.*
Orchis foliis inferioribus ovatis: superioribus ovato-ob-
longis, floribus ex alis superioribus. *Gron. virg. 109.*
Habitat in Virginia. D. Gronovius.
Folia radicalia bina, ovalia, petiolata, obtusa, longitudi-
ne fere plantae. Caulis nudus. Spica ex. 5 f. 6. flo-
ribus. Bractea luxceolato-cuales, longitudine germi-
nis.

nis. Florum petala 2 exteriora patula, viridia; 3 conniventia teneriora. Labium ovale, emarginatum: apicibus divaricatis.

8591

SATYRIUM.

hercinum.

1. SATYRIUM bulbis indivisis, foliis lanceolatis, nectarii labio trifido: intermedia lineari obliqua præmorsa. *Aët. npsf. 1740. t. 18. Dalib. parif. 275.*
Orchis barbata foetida. Baub. bist. 2. p. 756. Vaill. parif. t. 30. f. 6. Segu. ver. t. 15. f. 1.
Orchis barbata, odore hirci, brevior latioreque folio. Baub. pin. 20. Morif. bist. 3. p. 491. f. 12. t. 12. f. 9.
Habitat in Galliz, Cantii campestribus. &

viride.

2. SATYRIUM bulbis palmatis, foliis oblongis obtusis, nectarii labio lineari trifido: intermedia obsoleta. *Aët. npsf. 1740. p. 18. Fl. suec. 730. Dalib. parif. 276.*
Satyrium foliis oblongis caulinis. Fl. lapp. 313.
Orchis palmata, flore viridi. Baub. pin. 86. prodr. 30.
Orchis palmata, flore galericulato dilute viridi. Læf. prussf. 182. t. 56.
β. Orchis palmata batrachites. Baub. pin. 86. Vaill. parif. 153. t. 31. f. 6. 7. 8.
Habitat in Europæ frigidioris asperis. &

nigrum.

3. SATYRIUM bulbis palmatis, foliis linearibus, nectarii labio resupinato trilobo: intermedia majore. *Aët. npsf. 1740. p. 19. Fl. suec. 731.*
Satyrium foliis linearibus. Roy. prodr. 14.
Orchis palmata angustifolia alpina, nigro flore. Baub. pin. 86.
Palma christi minor. Cam. epit. 627.
Palmata angustifolia, flore lupinato, calcare brevissimo. Hall it. helv. 47.
Habitat in Alpibus Helveticis, Lapponicis. &

albidum.

4. SATYRIUM bulbis fasciculatis, foliis lanceolatis, nectarii labio trifido acuto: intermedia majore. *Aët. npsf. 1740. p. 19. Fl. suec. 733.*
*Satyrium scanense. It. scan. 153. **
Orchis palmata alpina, spica densa albo viridi. Hall. It. berc. 68.
PseudoOrchis alpina, flore herbaceo. Mich. gen. 30. t. 26.
Limodorum montanum, flore albo dilute virescente. Chem. aët. parif. 1705. p. 517.
Habi-

Habitat in Scania, Germaniæ, Helvetiæ, Avernæ præsylvaticis. 2

5. SATYRIUM bulbo ramoso, caule vaginato, corollis Epipogium. resupinatis: nectarii labio indiviso.

Epipogium. *Gmel. fib. 1. p. 12. t. 2. f. 2.*

Habitat in Sibiria montibus arenosis. 4

6. SATYRIUM bulbis fibrosis, foliis ovatis radicalibus, *repens.*

floribus secundis. *Ad. nps. 1740. p. 20. fl. succ.*

732. *Dalib. paris. 278.*

Satyrion foliis ovatis radicalibus. *Fl. lapp. 314.*

Orchis minor, floribus albis, f. radice repente. *Cam. bore.*

111. t. 35.

PseudoOrchis. *Bauh. pin. 84.*

Pyrola angustifolia polyanthos, radice geniculata. *Laf.*

pruss. 210. t. 68.

Epipactis foliis ovatis radicalibus. *Gmel. fib. 1. p. 13.*

- β. Orchis radice repente, foliis maculis nigris & albis adspersis. *Mentz. pug. t. 3. f. 4. 5.*

Habitat in Suecia, Angliæ, Sibiria, Helvetiæ sylvis.

OPHRYS. 892

* *Bulbis ramosis.*

1. OPHRYS bulbis fibroso-fasciculatis, caule vaginato, *Nidus avis* nectarii labio bifido.

Neottia bulbis fasciculatis, nectarii labio bifido. *Ad.*

nps. 1740. p. 33. Fl. succ. 742. Dalib. paris. 277.

Orchis abortiva fusca. *Bauh. pin. 86.*

Nidus avis. *Lob. ic. 195.*

Habitat in Suecia, Germaniæ, Galliæ nemoribus.

2. OPHRYS bulbis ramosis flexuosis, caule vaginato, *Corallorhiza* nectarii labio trifido.

Neottia bulbis reticulatis, nectarii labio trifido. *Ad. nps.*

1740. p. 34. *Fl. succ. 743.*

Neottia radice reticulata. *Fl. lapp. 315.*

Orobanche radice coralloide. *Bauh. pin. 88.*

Orobanche sueconum radice coralloide, flore albo. *Rudb.*

elys. 2. p. 235. f. 16.

Orobanche spuria f. *Corallorhiza Rapp. jen. 284. t. 2.*

Habitat in Europa borealis desertis. 2

3. OPHRYS bulbis aggregatis oblongis, caule subfolio- *spivalla.* so, floribus secundis, nectarii labio indiviso. *Ad. nps.*

1740. p. 32. *Dalib. paris. 277.*

O o o

Tri-

Triorchis alba odorata minor. Baub. pin. 84.
Satyrion odoriferum. Brauss. herb. 1. p. 105.
Tesliculus odoratus. Lob. ic. 186.

β. *Triorchis* f. *Tetrorchis alba odorata major. Baub. pin. 84.*

γ. *Orchialstrum æstivum palustre album odoratum.*
Mich. gen. 30. t. 26.

Habitat in Italiæ, Galliæ, Angliæ graminosis.

cernua.

4. *OPHRYS* bulbis fasciculatis, caule folioso, floribus cernuis, nectarii labio oblongo integro acuto.

Habitat in Virginia, Canada.

Radix fibris plurimis, crassis. Folia radicalia, linearia, longa. Caulina vaginantis, brevissima. Spica densa, oblonga. Flores arcu recurvati, cernui. Corollæ parvæ petalis superioribus confluentibus; labio inferiore oblongo, acuto, integro.

ovata.

5. *OPHRYS* bulbo fibroso, caule bifolio, foliis ovatis, nectarii labio bifido.

Ophrys foliis ovatis. Hort. cliff. 429. Act. ups. 1740. p. 28. Fl. succ. 738. Roy. lugdb. 15. Hall. botv. 277. Gmel. sib. 1. p. 25. Dalib. parif. 278.

Ophris bifolia. Baub. pin. 87.

Ophris. Fuchf. hist. 565.

β. *Ophris trifolia. Baub. pin. 87.*

Habitat in Europæ subhumidis pratis. &

cordata.

6. *OPHRYS* bulbo fibroso, caule bifolio, foliis cordatis.

Ophrys foliis cordatis. Fl. lapp. 247. Fl. succ. 739. Act. ups. 1740. p. 29. Gmel. sib. 1. p. 25.

Ophrys minima. Baub. pin. 87. prodr. 31.

Bifolium minimum. Baub. hist. 3. p. 534.

β. *Ophris minima, floribus purpurco-croceis. Mentz. pug. t. 9. f. 2.*

Habitat in Europæ frigida sylvis humentibus.

* *Bulbis rotundis.*

lilifolia.

7. *OPHRYS* bulbo subrotundo, caule nudo, foliis lanceolatis, nectarii labio integro, petalis dorsaliibus linearibus.

Ophrys scapo nudo, foliis radicalibus ovato-oblongis dimidii scapi longitudine. Gron. virg. 185.

Ophrys radice ovatis tunicatis, scapo nudo. Hort. cliff. 429. (vix memini)?

Habitat in Virginie, Canada, Succie paludibus. &

Pala

Planta virginica sexies major nostrate, at structura eadem, notabilis flore: petalis exterioribus linearibus.

8. OPHRYS bulbo subrotundo, scapo nudo trigono, ne-Læselii.
ctarii labello ovato.

Ophris diphyllus bulbosa. Læf. pruss. 180. t. 58.

Habitat in Sueciæ, Borussia paludibus.

Radix bulbis adscendens ramo. Folia lanceolata, bina, longitudine fere scapi. Scapus nudus, superne triquet. Petala angusta reflexa. Nectarii labium ovatum magnum respectu reliquorum. Flores in scapo 5 ad 8, nec ultra.

9. OPHRYS bulbo subrotundo, scapo subnudo pentagono, foliorum apicibus scabris, nectarii labio integro.

Habitat in Sueciæ paludibus turfosis. Bergius.

Bulbus ovatus, incurvus, infra radicans. Scapus pentagonus, maxima parte nudus. Folia radicalia 3. f. 4, spathulata, alterna, apice præsertim in pagina interiore scabra, exteriora breviora. Flores in racemo plurimi, virides, flavescentes. Petala 2 lateralibus ovato-oblonga, reflexo-erecta, 2 interiora lateralibus linearia, recurva; 1 supremum rectum, fornicem staminum formans. Labium nectarii lunceolato-ovatum, reflexum, integrum. Bulbi coherant deorsum quasi catenatim ramulo interstincti. Differt ab O. Læselii floribus pluribus, labio floribus angustis; ab O. monophyllo foliis non ovatis apice scabris, pedicellis non capillatis.

10. OPHRYS bulbo rotundo, caule nudo, folio ovato, monophyllo, nectarii labio integro.

Ophris monophyllos bulbosa. Læf. pruss. 180. t. 57.

Monochris ophioglossoides. Menz. pug. t. 5. f. 1. 2.

PseudoOchris monophyllos. Clus. hist. 1. p. 269.

Habitat in Borussia, Medelpadiæ paludibus sylvaticis. 2

11. OPHRYS bulbo globoso, caule nudo, nectarii labio trifido. *Monochris.*

trifido. Aët. upf. 1740. p. 29. Fl. suec. 740.

Herminium bulbo supra radicato, nectarii labio trifido.

Fl. suec. 740. Aët. upf. 1740. p. 29.

Herminium radice globosa. Fl. lapp. 317.

Orchis radice subrotunda, labello hastato. Gmel. fib. 1.

p. 18. t. 4. f. 1.

Orchis odorata moschata f. Monorchis. Bauh. pin. 84.

O o o 2

Or-

Orchis coleo unico f. *Monorchis flosculis pallide viridibus*. *Læs. pruss.* 184. t. 61.

Monorchis. *Mich. gen* 30. t. 26. *Rupp. jen.* 421. t. 2.

β. *Orchis lutea*, hirsuto folio. *Baub. pin.* 84.

γ. *Triorchis lutea*, folio glabro. *Baub. pin.* 84.

δ. *Triorchis lutea altera*. *Baub. pin.* 84.

ε. *Monorchis bifolia*, floribus viridibus moschum olentibus, alpina. *Mentz. pug. t.* 5. f. 3. 4. 5.

Habitat in Europæ prasis uliginosis.

alpina. 12. *OPHRYS* bulbis ovatis, caule nudo, foliis subulatis, nectarii labio indiviso.

ChamæOrchis alpina, folio gramineo. *Baub. pin.* 81. *prodr.* 29.

Orchis radicibus subrotundis, calcare nullo, foliis longis & peranguttis. *Hall. belv.* 269.

Habitat in Alpibus Lappoune, Helvetiæ: &

Radix Bulbis duobus, subrotundis (nec uno infra radicato, ut præcedentes tres), supra radicatis. Foliaradicalia sæpius 4, filiformi, subulata, pollicem longa. Caulis nudus, longitudine foliorum; Spica floribus 4 f. 5 alternis. Petala 5, conniventia: barba ovato-oblonga, integra, vix petalis longiore. Vaginæ 2 obvertunt basin plantæ & includunt folia cum scapo inferne.

camtschatea. 13. *OPHRYS* bulbis - - - caule filiformi vaginato, racemo laxo, nectarii labio lineari bifido.

Neottia nectarii labio bifido lineari. *Amæn. acad.* 2. p. 361. t. 4. f. 24. *

Habitat in Sibiria.

anthropophora. 14. *OPHRYS* bulbis subrotundis, caule folioso, nectarii labio lineari tripartito: medio elongato bifido.

Neottia bulbis subrotundis, nectarii labio quadrifido.

Aff. nps. 1740. p. 32. *Dalib. paris.* 277.

Orchis, flore nudi hominis effigiem repræsentans, femina. *Baub. pin.* 82. *Vaill. paris.* 147. t. 31. f. 19. 20.

Orchis antropophora orchades. *Col. esphr.* 1. p. 320.

Habitat in Italia, Lusitania, Gallia.

infestifera. 15. *OPHRYS* bulbis subrotundis, caule folioso, nectarii labio subquinelobo.

Cypripedium bulbis subrotundis, foliis oblongis caulinis. *Aff. nps.* 1740. p. 26. *Fl. suec.* 737. *is. oel.* 44. 121.

myodes. α. *Orchis muscæ corpus referens*, minor, galca & alis herbidis. *Baub. pin.* 83. *Vaill. paris.* 147. t. 31. f. 17. 18.

Or-

- Orchis myodes prima*, floribus muscam exprimens.
Lob. ic. 181.
- β. *Orchis muscam referens major*. *Baub. pin.* 83.
- γ. *Orchis muscam referens lutea*. *Baub. pin.* 83.
- δ. *Orchis muscam cæruleam majorem representans*.
Breyn. Moris. hist. 3. p. 494. f. 12. t. 13. f. 11.
- ε. *Orchis myodes lutea lusitanica*. *Breyn. cent.* 5. *Moris. hist.* 3. p. 495. f. 12. t. 13. f. 15.
- ζ. *Orchis myodes fusca lusitanica*. *Breyn. cent.* 41.
- η. *Orchis araneum referens*. *Baub. pin.* 94. *Tournef. inst. adrachnites* 434. t. 247. f. C. *Vaill. parif. t.* 30. f. 10. 11. 12. 13.
- Orchis adrachnitis*. *Lob. ic.* 135.
- θ. *Orchis fucum referens: colore rubiginoso*. *Baub. pin.* 83. *Vaill. parif.* 146. t. 31. f. 15. 16.
- ι. *Orchis fucum referens major, foliolis superioribus candidis & purpureiscentibus*. *Baub. pin.* 83. *Vaill. parif.* 146. t. 30. f. 9.
- Orchis fucum referens, flore subvirente*. *Baub. pin.* 83.
- κ. *Orchis scarabæum referens*. *Breyn. Moris. hist.* 3. p. 494. f. 12. t. 13. f. 10.
- λ. *Orchis cercopithecum exprimens lusitanica*. *Breyn. Moris. hist.* 3. p. 494. f. 12. t. 13. f. 4.
- Habitat in Europa temperiori. &*
Varietates Myodes α. & Arachnites γ fugitivis oculis insipienti diversissimæ adparent, qui vero has confert cum congeneribus, omnesque varietates simul inspicit, primam originem ex una facile perspiciat, nec limites distinguendas obtinebit, quamvis constantes persistant.

SERAPIAS. 893

1. SERAPIAS bulbis fibrosis, nectarii labio obtuso crena- *Helleborine.*
 to petalis brevior. *Act. upf.* 1740. p. 21. *Fl. succ.* 734.
It. al. 144. *It. gest.* 225. 197. 230. *Dalib. parif.* 276.
- α. *Serapias caule multifolio multiflora*. *Hort. cliff.* 429. *latifolia.*
Gmel. lib. 1. p. 8.
- Helleborine latifolia montana*. *Baub. pin.* 186.
- Epipactis f. Helleborine*. *Cam. epit.* 839.
- β. *Helleborine flore carneis*. *Baub. pin.* 187.
- γ. *Helleborine montana angustifolia spicata*. *Baub. pin.* 187.
- δ. *Helleborine montana angustifolia purpurascens*. *Baub. pin.* 187.
- ε. *Helleborine altera, strorubente flore*. *Baub. pin.* 186.

O o o 3

ζ Se-

- longifolia.* γ. *Serapias radicebus fibrosis, nectarii labio obtuso petalis brevioribus, foliis ensiformibus. It. æl. 144.*
Helleborine foliis prælongis angustis, floribus candidis. Vaill. parif. 97.
- palustris.* η. *Serapias angustifolia tomentosa, floribus herbaceis intus albis. Sauv. monsp. 42.*
Helleborine palustris f. pratensis nostras. Morif. hist. 3. p. 487. Raj. hist. 1231.
Helleborine angustifolia palustris f. pratensis. Bamb. pin. 87.
Habitat in Europæ asperis. &
Limites inter latifoliam & longifoliam & palustrem non reperi, diversitatem vidi.
- Lingua.* α. **SERAPIAS** bulbis subrotundis, nectarii labio trifido acuminato petalis longiore. *Act. nps. 1740. p. 23.*
*Satyrion radicebus subrotundis, cucullo clauso monopetalo; labello ovato-lanceolato. Sauv. monsp. 24.**
Orchis montana italica, flore ferrugineo, lingua oblonga. Bamb. pin. 84. prodr. 29.
 β. *Orchis montana italica, lingua oblonga altera. Bamb. pin. 84.*
Orchis macrophylla. Col. ecphr. 2. p. 321. t. 322.
Habitat in Italia, Lusitania. &

544
LIMODORUM.

- tuberosum.* α. **LIMODORUM.** *Rey. lugdb. 16. Gron. virg. 110.*
Act. nps. 1740. p. 21.
Helleborine americana, radice tuberosa, foliis longis angustis, caule nudo, floribus ex rubro pallide purpureiscentibus. Mart. cent. 40. t. 50.
Habitat in America septentrionali. &
Plumierii species americana australis plurimas non vidi.

895
ARETHUSA.

- bulbosa.* γ. **ARETHUSA** radice globosa, scapo vaginato, spatha diphylla. *Gen. nov. 1082.*
Arethusa. Gron. virg. 184.
Serapias bulbis subrotundis, caule unifloro. Act. nps. 1740. p. 24.
Orchidi affinis aquatica verna exigua. Gron. virg. 110.
Helleborine mariana monanthos, flore longo purpureiscente lilaceo. Pluk. mant. 100. t. 348. f. 7.
Habitat in Virginia, Canada. &

α. ARE-

2. ARETHUSA radice fibrosa, scapi folio ovali, folioso *ophioglossi-*
 spathaceo lanceolato. *Gen. nov.* 1083.
 Cypripedium folio caulino ovato-oblongo terminali lan-
 ceolato piano. *Hort. cliff.* 430. *Gron. virg.* 110.
 Helleborine virginiana, ophioglossi folio. *Pluk. alm.*
 182. t. 93. f. 2.
 Helleborine virginiana diphylla. *Morif. hist.* 3. p. 488.
 f. 12. t. 11. f. 15. mala.
Habitat in Virginia, Canada. &

3. ARETHUSA radice subpalmata, scapi folio folioloso *divaricata.*
 que spathaceo lanceolatis petalis, exterioribus adscen-
 dentibus.
 Serapias radicibus palmato-fibrosis, caule unifloro.
Gron. virg. 184.
 Helleborine lillii folio caulem ambiente, flore unico he-
 xapetalo: tribus petalis longis angustis obscure purpu-
 reis, cæteris brevioribus rosaceis. *Catesb. car.* 1. p.
 58. t. 58.
Habitat in Americæ borealis paludosis. &

CYPRIPEDIUM. 896

1. CYPRIPEDIUM radicibus fibrosis, foliis ovato-lan- *Calceolus.*
 ceolatis caulinis. *Aët. nps.* 1740. p. 24. *Fl. suec.* 735.
 Cypripedium foliis ovato-lanceolatis. *Fl. lapp.* 318.
Gron. virg. 111.
 Calceolus foliis ovato-lanceolatis. *Gmel. fib.* 1. p. 2. t. 1.
 Calceolus marianus. *Dod. pempt.* 180. f. 1. 2.
 Helleborine flore rotundo f. Calceolus. *Bauh. pin.* 187.
 β. Helleborine Calceolus dicta mariana, caule folioso, flo-
 re luteo minore. *Pluk. mant.* 101. t. 418. f. 2.
 Helleborine virginiana f. Calceolus flore luteo majore.
Morif. hist. 3. p. 488. f. 12. t. 11. f. 15.
 γ. Calceolus marianus canadensis. *Corn. canad.* 204.
 Helleborine flore majore purpureo. *Morif. hist.* 3. p. 488.
 f. 12. t. 11. f. 17.
 δ. Calceolus minor, flore vario. *Amm. ruth.* 133. t. 22.
 Calceolus foliis ovatis binis. *Gmel. fib.* 1. p. 5.
Habitat in Europæ, Asiæ, Americæ septentrionalibus.
 2. CYPRIPEDIUM bulbo subrotundo, folio subrotun- *bulbosum.*
 do radicali. *Aët. nps.* 1740. p. 25. *Fl. suec.* 736. *
 Cypripedium folio subrotundo. *Fl. lapp.* 319. t. 12. f. 5.
 Serapias scapo unifloro. *Gmel. fib.* 1. p. 7. t. 2. f. 3.
 Orchis lapponensis monofolia. *Rudb. elys.* 2. p. 209. f. 10.
Habitat in Lapponia, Russia, Sibiria. &

EPIDENDRUM.

* *Scandentes.**Vanilla.*

899

1. EPIDENDRUM foliis ovato-oblongis nervosis sessilibus caulinis, cirrhis spiralibus. *Roy. lugdb.* 13. *Act. upf.* 1740. p. 13. *Mat. med.* 418.
Vanillas piperis arbori jamaicensis innascens. Pluk. alm. 381. t. 320. f. 4.
Vanille. Merian. surin. 25. t. 25.
Volubilis siliquosa, plantaginis folio. Catesb. car. 3. p. 7. t. 7.
Lobus aromaticus subfuscus terebinthi corniculalis simili. Baub. pin. 404.
Habitat in America australi. ?

- domesticum.* 2. EPIDENDRUM caule scandente tereti subramoso, foliis lanceolatis, petalis lanceolatis. *Act. upf.* 1740. p. 37.
Angurèk warna. Kämpf. amœn. 867. t. 867.
Habitat in Indiæ orientalis Bataviæ arboribus, parasitica.

- Flos æris.* 3. EPIDENDRUM caule scandente tereti subramoso, foliis lanceolatis, petalis linearibus obtusis. *Act. upf.* 1740. p. 37.
Kolong ging. Kämpf. amœn. 868. t. 868.
Habitat in Java, parasitica.

* *Caule erecto folioso.*

- tennifolium.* 4. EPIDENDRUM foliis caulinis subulatis canaliculatis. Tserou-mau-maravara. *Rheed. mal.* 12. p. 11. t. 5.
Habitat in India.

- spiculatum* 5. EPIDENDRUM foliis caulinis oblongis alternis obtusis.
Ponnampou-maravara. Rheed. mal. 12. p. 7. t. 3. *fibrosa.*
Habitat in India.
Confer Viscum caryophylloides, foliis longis in apice incis, floris labello albo trifido, petalis luteis longis angustissimis. Catesb. car. 2. p. 68. t. 68. *bulbosum.*

- ovatum.* 6. EPIDENDRUM foliis caulinis ovatis acutis amplexicaulibus.
Anantali-maravara. Rheed. mal. 12. p. 15. t. 7.
Habitat in India.

* *See-*

* *Scapo nudo, foliis radicalibus.*

7. EPIDENDRUM foliis radicalibus subulatis acutis: *nodosum*.
 nodo radicatis. *Aët. upf.* 1740. p. 36.
 Epidendron foliis subulatis. *Hort. cliff.* 430.
 Epidendron curassavicum orchidi affinis, folio crasso
 sulcato. *Herm. parad.* 187. t. 187.
 Viscum arboreum f. Epidendron, flore albo specioso, a-
 mericanum, folio forma siliquarum nerii. *Pluk. alm.*
 390. t. 117. f. 6.
 Viscum delphinii flore minus, petalis angustioribus, radice
 fibrosa. *Sloan. jam.* 120. *hist.* 1. p. 251. t. 125. f. 3.
 β. Viscum radice bulbosa minus, delphinii flore rubro spe-
 cioso. *Sloan. jam.* 119. *hist.* 1. p. 250. t. 121 f. 2.
Habitat in America meridionali, parasitica.
8. EPIDENDRUM foliis oblongis obtusis compressis ar- *carinatum*.
 ticulatis. *Aët. upf.* 1740. p. 36.
 Bontia luzonica, geniculis interioribus carinulatis. *Pet.*
gaz. 44. f. 10.
Habitat in Insulis Luzonum, parasitica.
9. EPIDENDRUM foliis radicalibus oblongis obtusis su- *aloifolium*.
 perne latoribus. *Aët. upf.* 1740. p. 36.
 Orchis abortiva, flore majore rubro, folio aloes. *Rudb.*
elys. 2. p. 224.
 Kantijram-maravara. *Rheed. mal.* 12. p. 17. t. 8. *Raj.*
suppl. 572.
Habitat in Malabariz arboribus, parasitica.
10. EPIDENDRUM foliis radicalibus lanceolatis cana- *guttatum*.
 lieularis, petalis cuneiformibus retusis.
 Epidendrum caule tereti erecto nudo, foliis lanceolatis,
 petalis cuneiformibus obtusis. *Aët. upf.* 1740. p. 37.
 Viscum, delphinii flore albo guttato, minus, radice fibro-
 sa. *Sloan. jam.* 120. *hist.* 1. p. 251. t. 148. f. 2.
Habitat in jamaica, parasitica.
11. EPIDENDRUM foliis radicalibus linearibus apice *retusum*.
 bifariam retusis.
 Angeli-maravara. *Rheed. mal.* 12. p. 1. t. 1.
 β. Biti-maram-maravara. *Rheed. mal.* 12. p. 5. t. 2.
Habitat in India.
12. EPIDENDRUM foliis radicalibus lato-lanceolatis, *amabile*.
 petalis lateralibus orbiculatis.
Habitat in India. Osbeck.
 Radices crassæ, juniformes, supra arborea scandentes. Fo-
 lla

○ ○ ○ 5

lia Crini f. Scille officinalis, lata, lanceolata, carnosae, semipedalia. Culmus bipedalis, nudus, cinctus aliquot squamis acutis brevissimis. Flores nivei Orchidis Susannae emuli, f. magnitudine narcissi: constanter petalis 5, quorum 2 lateralia orbiculata, reliqua ovata. Cucullus alter triphyllus: lateralibus oblongis, intermedio bastato, bifido setis duabus subulatis.

enfilolum. 13. EPIDENDRUM caule tereti lævi, foliis ensiformibus petalis lanceolatis glabris.

Habitat in China. Osbeck.

Planta terrestris nec parasitica. Folia angusta, striata, acuta, numerosa, non carnosae. Scapus anceps, bipedalis, teres, nudus. Bracteis aliquot, alternis, membranaceis, amplexicanlibus, acuminatis. Flores alterni, ex alis bractearum solitarii, pedunculati, Pedunculis longitudine floris. Corolla ex petalis 5, lanceolatis, patentibus, lævibus, sexto latiore recurvato. Flos fragrantissimus.

moniliforme 14. EPIDENDRUM caule tereti articulado striato moniliformi nudo simplicissimo. *Art. nps. 1740. p. 37.*

Fu-Ran. Kempf. amon. 864. t. 864.

Habitat in japoniæ rupibus & arboribus parasitica.

TRIANDRIA

SISYRINCHIUM.

Bermudiana 1. SISYRINCHIUM caule foliisque ancipitibus. *Hort. cliff. 431. Hort. nps. 278. Gron. virg. 111. Roy. lngdb. 19.*

Sisyrrinchium cæruleum parvum, gladiato caule, virginianum. *Pluk. alm. 348. t. 61. f. 1*

Bermudiana graminea, flore minore cæruleo. *Dill. elth. 49. t. 41. f. 49.*

β. Sisyrrinchium bermudienſe, floribus parvis ex cæruleo & aureo mixtis. *Pluk. alm. 348. t. 61. f. 2.*

Bermudiana iridis folio, radice fibrosa. T. *Dill. elth. 48. t. 41. f. 48.*

Habitat α in Virginia, β in Bermudis. 2

Planta α, in loco natali, Folia sepe & scapum filiformia gerit, at si probe examinetur, tamen ancipitia; hæc per gradus ad cultæ staturam adscendere observatur.

TE.

TETRANDRIA.

NEPENTHES.

898

1. NEPENTHES. *Hort. cliff.* 431. * *Fl. zeyl.* 321. *distillatoria*.
Utricularia vegetabilis zeylanensium. *Pluk. alm.* 394. t.
237. f. 3.
Bandura zeylanica, in extremo foliorum folliculum
peniformem expansum habens. *Burm. zeyl.* 42. t. 17.
Planta mirabilis destillatoria. *Grim. E. N. C. an.* 1.
Dec. 2. p. 363. f. 363.
Habitat in Zeylona.

PENTANDRIA.

PASSIFLORA

900

* *Folii indivisi.*

1. PASSIFLORA foliis indivisis serratis. *Amæn. acad. serratifolia*.
1. p. 217. f. 1. *
Passiflora foliis ovato-lanceolatis integris serratis. *Hort.*
cliff. 431. *Ray. lugdb.* 260.
Granadilla americana, folio oblongo leviter serrato, pe-
talis ex viridi rubescentibus. *Mart. cent.* 36. t. 36.
Habitat Surinami.
2. PASSIFLORA foliis indivisis ovatis integerrimis: pe-
talis biglandulosis. *Amæn. acad.* 1. p. 218. f. 2. *
Clematis indica alia, flore minore pallido. *Plum. a-*
mer. 74. t. 89.
Clematis Murucuja pyriformis minor. *Morif. bist.* 3.
p. 7. f. 1. t. 2. f. 4.
Habitat in Dominica, Brasilia.
3. PASSIFLORA foliis indivisis ovatis integerrimis: pe-
talis æqualibus. *Amæn. acad.* 1. p. 219. f. 3. *
Granadilla flore cupreo, flore oliviformi. *Dill. elth.*
165. t. 138. f. 165.
Granadilla americana, fructu subrotundo, corolla flo-
ris erecta, petalis amœne fulvis, foliis integris. *Mart.*
cent. 37. t. 37.
Granadilla foliis sarsaparillæ trinerviis, flore purpureo,
fructu olivæformi cæruleo. *Catesb. car.* 2. p. 93. t.
93.
Habitat in Providentia, Bahama.

4. PAS-

- zittifolia.* 4. PASSIFLORA foliis indivisis cordatis integerrimis: petiolis æqualibus. *Amæn. acad.* 1. p. 219. f. 4.
Granadilla pomifera, tilix folio. *Few. peruv.* 1. p. 720. t. 12.
Habitat in Peru, Lima.
- maliformis.* 5. PASSIFLORA foliis indivisis cordato-oblongis integerrimis: petiolis biglandulosis, involucris integerrimis. *Amæn. acad.* 1. p. 220. f. 5. *
Passiflora foliis cordato-oblongis integerrimis, floribus solitariis, involucro tripartito integerrimo. *Roy. lugdb.* 261.
Clematis indica latifolia, flore clavato, fructu maliformi. *Plum. amer.* 67. t. 82. *Raj. suppl.* 340.
Habitat in Dominica & de la Tortue.
- laurifolia.* 6. PASSIFLORA foliis indivisis integerrimis, involucris dentatis. *Amæn. acad.* 1. p. 220. f. 6. *
Passiflora foliis solitariis oblongis integerrimis, floribus solitariis, involucro tripartito dentato. *Roy. lugdb.* 532.
Passiflora arborea, laurinis foliis, americana. *Pluk. alm.* 282. t. 211. f. 3.
Clematis indica, fructu citrifolmi, foliis oblongis. *Plum. amer.* 64. t. 80. *Raj. suppl.* 341.
Marquias. *Mer. surin.* 21. t. 21.
Habitat in Surinamo.
- multiflora.* 7. PASSIFLORA foliis indivisis oblongis integerrimis, floribus confertis. *Amæn. acad.* 1. p. 221. f. 7. *
Passiflora, laurinis foliis, polyanthos americana. *Pluk. alm.* 282.
Granadilla, flore minore corymbofo. *Plum spec.* 7.
Clematis indica polyanthos odoratissima. *Plum. amer.* 75. t. 90. *Raj. suppl.* 343.
Habitat in Dominica.
- * *Foliis bilobis.*
- perfoliata.* 8. PASSIFLORA foliis bilobis oblongis transversis amplexicaulibus. *Amæn. acad.* 1. p. 222. f. 8. *
Flos passionis perfoliatus f. periclymeni perfoliati folio. *Sloan. jam.* 104. *hist.* 1. p. 230. t. 142. f. 3. 4. *Raj. suppl.* 342.
Habitat in Jamaica sylvis saxosis.
- vulva.* 9. PASSIFLORA foliis bilobis basi emarginatis petiolatis. *Amæn. acad.* 1. p. 222. f. 9. *

Cle-

- Clematis indica, flore clavato suaverubente, fructu hexagono coccineo, folio bicorni. *Plum. amer.* 68. t. 83. *Raj. suppl.* 341.
 Flos passionis folii media lacinia quasi abscissa, flore minore carneo. *Sloan jam.* 104. *bist.* 1. p. 229.
Habitat in Jamaica, Dominica, Martinica, Cayenna.
10. PASSIFLORA foliis bilobis obtusis basi indivisis, ne- *Murucnja.*
 ctariis monophyllis. *Amæn. acad.* 1. p. 223. f. 10. *
 Passiflora foliis bilobis obtusis, nectariis indivisis. *Roy.*
lugdb. 261.
 Clematis indica, flore puniceo, folio lunato. *Plum. amer.* 72. t. 87. *Raj. suppl.* 342.
Habitat in Dominica.
11. PASSIFLORA foliis bilobis cuneiformibus basi bi- *vespertilio.*
 glandulosis: lobis acutis divaricatis. *Amæn. acad.* 1. p. 223. f. 11. *
 Passiflora foliis obverse lunulatis: punctis duobus mel-
 liferis sub basi. *Virid. cliff.* 91. *Hort. cliff.* 431. *Roy.*
lugdb. 260.
 Granadilla bicornis, flore candido, filamentis intortis.
Dill. elth. 161. t. 137. f. 164.
Habitat in America.
12. PASSIFLORA foliis bilobis cordatis oblongis pe- *capsularis.*
 tiolatis.
 Granadilla fructu rubente, folio bicorni. *Plum. spec.* 6.
Barr. obs. præf. 1. *titul.* f. 1.
Habitat in Gallia æquinoctiali.
 * *Foliis trilobis.*
13. PASSIFLORA foliis subtrilobis obtusis subrotun- *rotundifolia.*
 dis.
 Granadilla folio hederaceo, flore albo, fructu globoso
 villoso. *Plum. spec.* 6. *Barr. obs. præf.* 1. *titul.* f. 2.
Habitat in America australiori.
14. PASSIFLORA foliis trilobis oblongis subtus pun- *punctata.*
 ctatis: intermedio minore. *Amæn. acad.* 1. p. 224. f. 12. *
 Passiflora foliis bilobis obtusis, nectario multifido. *Roy.*
lugdb. 260.
 Granadilla folio tricuspidi obtuso & oculato. *Few. peruv.* 1. p. 718. t. 11. *Hort. angl.* 25.
Habitat in Peru. 2

lucida.

15. PASSIFLORA foliis trilobis cordatis æqualibus obtusis glabris integerrimis. *Amæn. acad.* 1. p. 224. f. 13.

Passiflora foliis cordatis trilobis integerrimis glabris: lateribus angulatis. *Hort. cliff.* 431. *Roy. lugdb.* 261.

Passiflora foliis trilobis integerrimis: laciniis semiovatis acutis integerrimis glaucis. *Græn. virg.* 112.

Clematis passionalis triphyllus, flore luteo. *Morif. bist.* 1. p. 7. f. 1. t. 2. f.

Flos passionis minor, folio in tres lacinias non serratas, minus profundas diviso. *Sloan. jam.* 104. *bist.* 1. p. 221.

Habitat in Virginie, Jamaicae glareosis, saxosis. &

suberosa.

16. PASSIFLORA foliis trilobis subpeltatis, cortice suberoso. *Amæn. acad.* 1. p. 226. f. 14.

Passiflora foliis trilobis peltatis integris: glandulis duabus sub dorso basi convexis. *Roy. lugdb.* 261.

Clematis indica, folio hederaceo, major, fructu olivæformi. *Plum. amer.* 70. t. 84. *Raj. suppl.* 339.

Passifloræ affinis, hederæ folio, americana. *Pluk. alm.* 202. t. 210. f. 4.

- β. Clematis indica, folio angusto trifido, fructu olivæformi. *Plum. amer.* 70. t. 85.

Flos passionis minor, foliis in tres lacinias non serratas profundius divisis, flore luteo. *Sloan. jam.* 104. *bist.* 1. p. 320.

Habitat in Dominica, Antillis. β

Caulis perennis: Cortice suberoso, fisso, albo. Flores axillares, gemini, albi, apetalæ. Bacca violacea.

holosericea.

17. PASSIFLORA foliis trilobis: basi utrinque denticulo reflexo. *Amæn. acad.* 1. p. 226. f. 15. *

Passiflora foliis cordato-trilobis integerrimis: basi utrinque denticulo reflexo. *Hort. cliff.* 432. *Roy. lugdb.* 261.

Granadilla folio hastato holosericeo, petalis candicantibus: fimbriis ex purpureo & luteo variis. *Mart. cent.* 51. t. 51.

Habitat in Vera Cruce.

birsuta.

18. PASSIFLORA foliis trilobis villosis, floribus oppositis. *Amæn. acad.* 1. p. 227. f. 16. *

Passiflora non crenata americana triphylla. *Pluk. alm.* 282. t. 212. f. 1.

Clematis indica, flore minimo pallido. *Plum. amer.* 73. t. 88.

Flos

Flos passionis albus, folio ibisci sericeo trilobato. *Herm. par.* 176. t. 176.

Habitat in Dominica, Curassao.

19. PASSIFLORA foliis trilobis cordatis pilosis, involucribus multifido-capillaribus. *Amœn. acad.* 1. p. 228. f. 17. * *Phil. bot.* 260. *

Passiflora florum involucribus triphyllis multifido-capillaribus. *Hort. cliff.* 431. *Roy. lugdb.* 261.

Flos passionis albus reticulatus. *Herm. par.* 173. t. 173.

Clematis indica hirsuta foetida. *Plum. amer.* 71. t. 86.

- β. Passiflora vesicaria hederacea, foliis lanuginosis, odore retro, filamentis florum ex albo & purpureo variegatis. *Pluk. alm.* 382. t. 104. f. 1.

Habitat in Dominica, Martinicana, Curassao. ☉

20. PASSIFLORA foliis trilobis integerrimis: lobis sub-minima lanceolatis: intermedio productiore. *Amœn. acad.* 1. p. 229. f. 18.

Passiflora f. Flos passionis curassavicus, folio glabro trilobato & angusto, flore flavescente omnium minimo.

Pluk. alm. 282. t. 210. f. 3.

Habitat in Curassao.

21. PASSIFLORA foliis trilobis serratis. *Amœn. acad. incarnata.* 1. p. 230. * *Hort. nps.* 278.

Passiflora foliis semitrifidis serratis: basi duabus glandulis convexis: lobis ovatis. *Hort. cliff.* 432. *Gron. virg.* 112. *Roy. lugdb.* 260.

Granadilla hispanis, Flos passionis italis. *Herm. mex.* 888. t. 888. *Raj. hist.* 649.

Clematis trifolia, flore roseo clavato. *Baub. pin.* 301.

Clematis trifolia, f. Flos Passionalis, flore viridi. *Morif. hist.* 2. p. 6. f. 1. t. 1. f. 9.

Balsamina indica repens triphylla. f. folio hastato. *Ambr. phyt.* 89. t. 90.

Habitat in Virginia, Brasilia, Peru. &

* *Foliis multifidis.*

22. PASSIFLORA foliis palmatis integerrimis. *Amœn. carulea.* acad. 1. p. 231. f. 20. *Hort. nps.* 278.

Passiflora foliis palmatis quinquepartitis integerrimis, involucribus cordatis triphyllis. *Hort. cliff.* 432. *Roy. lugdb.* 260.

Clematis quinquefolia americana f. Flos passionis. *Rob. ic.*

Flos

Flos passionis major pentaphyllus. *Sloan. jam.* 104.
hist. 1. p. 229. *Raj. suppl.* 339.

Habitat in Brasilia. ♀

serratedigita. 23. PASSIFLORA foliis palmatis ferratis. *Amæn. acad.*
 1. p. 232. f. 27.

Clematis indica polyphylla major, flore clavato, fructu
 colocynthis. *Plum. amer.* 62. t. 79. *Raj. suppl.*
 340.

Habitat in Martinica.

pedata. 24. PASSIFLORA foliis pedatis ferratis. *Amæn. acad.*
 1. p. 233. f. 22. *

Clematis indica polyphylla, flore crispato. *Plum. amer.*
 66. t. 81. *Raj. suppl.* 341.

Habitat in Dominica.

901

HEXANDRIA.

ARISTOLOCHIA.

bilobata. 1. ARISTOLOCHIA foliis bilobis.

Aristolochia longa scandens, foliis ferri equini effigie.

Plum spec. 5. *amer.* 91. t. 106. *Raj. suppl.* 395.

Habitat in Dominica.

trilobata. 2. ARISTOLOCHIA foliis trilobis.

Aristolochia folio hederaceo trifido, maximo flore, radice
 repente. *Plum. spec.* 5.

Ambuyaembo. *Marcgr. bras.* 15.

Habitat in America meridionali.

peltata. 3. ARISTOLOCHIA foliis reniformibus peltatis.

Aristolochia asari folio umbilicato, flore longissimo, radice
 repente. *Plum. spec.* 5.

Habitat in America meridionali.

arborescens. 4. ARISTOLOCHIA foliis cordato-lanceolatis, caule
 erecto fruticoso.

Aristolochia caule erecto fruticoso, foliis cordato-lanceolatis. *Hort. cliff.* 433.

Aristolochia polyrhizos, auriculatis foliis, virginiana.

Pluk. alm. 50. t. 78. f. 1.?

Habitat in America. ♀

indica. 5. ARISTOLOCHIA foliis cordato-oblongis, caule volubili,
 pedunculis multifloris. *Fl. zeyl.* 323. *

Cate.

Cateæ-vegon. *Rheed. mal.* 8. p. 48. t. 25. *Raj. suppl.* 394.

5. *Aristolochia* caule volubili, foliis cordato-oblongis, fructu pendulo, pedunculis ramosis. *Hort. cliff.* 433.
Aristolochia scandens odoratissima, floris labello purpureo, semine cordato. *Sloan. jarn.* 60. *bist.* 1. p. 162. t. 104. f. 1. *Raj. suppl.* 394.
Aristolochia mexicana, folio acutiore. *Morif. bist.* 3. p. 508. f. 12. t. 17. f. 7.
Habitat in India. 2

6. *ARISTOLOCHIA* foliis cordatis acutiusculis, caule *batica* volubili, pedunculis solitariis petiolo longioribus.
Aristolochia Clematidis serpens. *Baub. pin.* 307. *Morif. bist.* 3. p. 509. f. 12. t. 17. f. 6.
Aristolochia Clematidis *batica*. *Clus. hist.* 2. p. 71.
Habitat in Hispania, Creta, arbores scandens.

7. *ARISTOLOCHIA* hirta, floribus solitariis pendulis *hirta*, recurvatis sublabiatis.
Aristolochia longa subhirsuta, folio oblongo, flore (minimo *male Tournef. cor.* 8.) maximo. *Tournef. itin.* 1. p. 147. t. 147.
Habitat in Chio. 2
Limbus corollæ non elongatus in labium.

8. *ARISTOLOCHIA* foliis cordato oblongis undatis, *sempervirens*, caule infirmo, floribus solitariis.
Aristolochia caulibus infirmis, foliis cordato-oblongis undularis, floribus recurvis solitariis pendulis. *Hort. cliff.* 432. *Sauv. monsp.* 111.
Pistolochia cretica. *Baub. pin.* 307. *Morif. bist.* 3. p. 510. f. 12. t. 17. f. 16.
Pistolochia altera. *Clus. bist.* 2. p. 260.
Habitat in Creta. 5

9. *ARISTOLOCHIA* foliis cordato-oblongis planis, *Serpentaria*, caulibus infirmis flexuosis teretibus, floribus solitariis.
Aristolochia caulibus infirmis angulosis flexuosis, foliis cordato-oblongis planis, floribus recurvis solitariis. *Gron. virg.* 112. *Mat. med.* 416.
Aristolochia *Pistolochia* f. *Serpentaria* virginiana, caule nodoso. *Pluk. alm.* 50. t. 148. f. 5. *Catesb. car.* 1. p. 29. t. 29.
Aristolochia *Pistolochia* caule nodoso; f. *Serpentaria* virginiana. *Raj. suppl.* 394.

P P P

Ari-

Aristolochia polyrrhizos virginiana, fructu parvo pentangulari. *Morif. hist.* 3. p. 510. f. 12. t. 17. f. 14.
Habitat in Virginia 2

Pistolochia. 10. **ARISTOLOCHIA** foliis cordatis crenulatis petiolatis, floribus solitariis.

Aristolochia caule infirmo ramoso, foliis cordatis obsolete dentatis, floribus solitariis erectis. *Hort. cliff.* 433. *Sauv. monsp.* 111.

Aristolochia Pistolochia dicta. *Bauh. pin.* 307.

Pistolochia. *Clus. hist.* 2. p. 72. *Dod. pempt.* 525.

Habitat in Hispania, G. Narbonensi. 2

rotunda. 11. **ARISTOLOCHIA** foliis cordatis subsessilibus obtusis, caule infirmo, floribus solitariis.

Aristolochia, caule infirmo ramoso, foliis cordatis integerrimis, floribus solitariis erectis. *Hort. cliff.* 432. *Mat. med.* 414. *Sauv. monsp.* 111.

Aristolochia rotunda, flore ex purpura nigro. *Bauh. pin.* 307.

Aristolochia rotunda 1. *Clus. hist.* 2. p. 70.

Aristolochia. *Cam. epis.* 419.

β. *Aristolochia rotunda*, flore ex albo purpurascente. *Bauh. pin.* 307.

Habitat in Italia, Hispania, G. Narbonensi. 2

longa. 12. **ARISTOLOCHIA** foliis cordatis petiolatis integerrimis obtusiusculis, caule infirmo floribus solitariis.

Aristolochia caule infirmo ramoso, foliis cordatis integerrimis longe petiolatis, floribus solitariis. *Sauv. monsp.* 111.

Aristolochia longa vera. *Bauh. pin.* 307.

Aristolochia longa. *Clus. hist.* 2. p. 70.

β. *Aristolochia longa hispanica.* *Bauh. pin.* 307.

Habitat in Hispania, Italia.

clematidis. 13. **ARISTOLOCHIA** foliis cordatis, caule erecto, floribus axillaribus confertis. *Hort. nps.* 279. *Mat. med.* 415. *Dalib. paris.* 279.

Aristolochia caule erecto simplicissimo, foliis cordatis petiolatis, floribus lateralibus confertis. *Hort. cliff.* 432. *Sauv. monsp.* 111.

Aristolochia Clematidis recta. *Bauh. pin.* 307.

Aristolochia Clematidis vulgaris. *Clus. hist.* 2. p. 71.

Aristolochia rotunda. *Fuchj. hist.* 90.

Habitat in Gallia, Tataria. 2

PISTIA. 902

1. PISTIA foliis obcordatis. *Fl. zeyl.* 322. *Stratiotes.*
 Koddapail palustris, folio oblongo specioso. *Plum.gen.*
 30.
 Kodda-pail. *Rheed. mal.* 11. p. 63. t. 32.
 6. Stratiotes ægyptia. *Bauh. hist.* 3. p. 787. *Morif. hist.* 3.
 p. 619. f. 15. t. 4. f. infer. *Vestl. ægypt.* 44. t. 45.
Raj. hist. 1324.
 Stratiotes Hai-alein maovi. *Alp. ægypt.* 106. t. 108.
 Lenticula palustris VI, f. ægyptia, 1. Stratiotes aquatica,
 foliis sedo majore latioribus. *Bauh. pin.* 362. *Sloan.*
jam. 11. *hist.* 1. p. 15. t. 2. f. 2.
Habitat in Asiæ, Africæ, Americæ meridionalibus,
natant.
 Flores sicci mihi satis non fecere, examinent autoptæ
 accuratius.

DECANDRIA. 903

HELICTERES.

1. HELICTERES foliis cordatis serratis, fructu com-*Isora.*
 posito contorto.
 Helicteres. *Hort. cliff.* 433.
 Helicteres arbor indiæ orientalis, filiqua varicosa & ru-
 niculi in modum contortuplicata. *Pluk. alm.* 181.
 t. 245. f. 2.
 Frutex indicus, fructu e styli apice egresso sextuplici ru-
 niculo in spiram convoluto constante. *Raj. hist.* 1765.
Isora Murri. Rheed. mal. 6. p. 55. t. 30.
 2. Helicteres arbor Indiæ occidentalis, fructu majore.
Pluk. alm. 181. t. 245. f. 3.
 Abulito affinis arbor, althææ folio, cujus fructus est
 styli apex acutus quatuor s. quinque sillquis hirsutis,
 funis ad instar in spiram convolutis. *Sloan. jam.* 97.
hist. 1. p. 22.
Habitat in Malabarâ, Jamaica. h
 2. HELICTERES foliis lanceolatis integerrimis, fructu angustifolia
 ovato recto.
Habitat in China. Osbeck. h
Arbor. Folia lanceolata, supra nuda, subtus tomentosa,
petiolata, integerrima, acuminata. Flores petiolati,
sparsi. Calyx tomentosus. Petala quinque, ad basin la-
minarum utrinque dente anctâ. Receptaculum longi-
P p p 2 tudi-

tudine floris, filiforme, cujus apici stamina 10 parva, & Stylus unicus. Capsula ovata, tomentosa, minime contorta, sed ex Capsulis 5, parallelis, rectis, approximatis composita.

POLYANDRIA.

GREWIA.

964

occidentalis

1. GREWIA foliis subovatis.

Grewia corollis acutis. Hort. cliff. 433.

Ulmifolia arbor africana baccifera, floribus purpureis.

Pluk. alm. 393. t. 237. f. 1.

Ulmi facie arbutula æthiopica, ramulis alatis, floribus purpurascens. Comm. hort. 1. p. 165. t. 85. Seb.

thes. 1. p. 46. t. 29. f. 3. Raj. dendr. 13.

Habitat in Æthiopia, Curacao. 5

orientalis.

2. GREWIA foliis sublanccolatis.
- Fl. zeyl. 324. **

Grewia corollis obtusis. Roy. lugdb. 476. Hort. cliff. 433.

Pai-paroa f. Couradi. Rheed. mal. 5. p. 91. t. 46. Raj. hist. 1624.

Habitat in India. 5

965

ARUM.

* *Acaulia: foliis compositis.*

Dracunculus.

1. ARUM foliis pedatis: foliolis lanceolatis integerrimis

æquantibus spatham spadice longiorem. Roy. lugdb. 7.

Arum foliis palmatis: foliolis undecim lanceolatis integerrimis: intermediis majoribus, spadice clavato.

Hort. cliff. 344.

Dracunculus polyphyllus. Baub. pin. 195. Moris. hist. 3. p. 548. f. 13. t. 5. f. 46.

Dracontium. Dod. pempt. 329.

Habitat in Europa australi.

Dracunculum

2. ARUM foliis pedatis: foliolis lanceolatis integerrimis

superantibus spatham spadice brevioribus. Roy. lugdb. 7.

Arum polyphyllum minus & humilior. Herm. lugdb. 60.

β. Arum f. Arisarum virginianum, dracontii folio, pene viridi longo acuminato. Pluk. alm. 52. t. 271. f. 1.

Habitat in America.

pentaphyllum.

3. ARUM acaule, foliis quinatis.

Arum

Arum pentaphyllum indicum. Moris. bist. 3. p. 540. f. 13. t. 5. f. 27. mal.

Romphal planta indiæ orientalis. Zan. bist. 205. fig. mal.

Habitat in India.

4. ARUM acaule, foliis ternatis. *Gron. virg. 113. triphyllum.*
Dracunculus f. Serpentaria triphylla brasiliiana. Baub. pin. 195. prodr. 101. Dod. mem. 81.

β. *Arum minus triphyllum f. Arisarum, pene viridi, virginianum. Moris. bist. 3. p. 547. f. 13. t. 5. f. 43.*

γ. *Arum f. Arisarum triphyllum minus, pene atrorubente, virginianum. Pluk. alm. 52. t. 77. f. 5.*

Habitat in Virginia, Brasilia. 2

Plantæ brasiliianæ foliola lateralìa extrorsum lobata, virginianæ autem tantum gibba, eandem tamen suadet Floris structura.

* *Acaulia, foliis simplicibus.*

5. ARUM acaule, foliis trilobis, flore sessili. *Fl. zeyl. trilobatum. 326. **

Arum, trilobato folio, humilius & minus zeylanicum.

Herm. par. 78. t. 78. Raj. suppl. 575.

Arum humile zeylanicum latifolium, pistillo coccineo.

Comm. hort. 1. p. 97. f. 51.

Habitat in Zeylona. 2

6. ARUM acaule, foliis peltatis ovatis repandis: basi semibifidis. *Hort. cliff. 434. n. 4. Hort. upf. 280. Roy. lugdb. 7. Colocasia.*

Arum maximum ægyptium quod vulgo Colocasia.

Baub. pin. 195.

Arum ægypticum. Col. ecphr. 2 p. 1. t. 1.

Colocasia. Clus. bist. 2. p. 75.

Habitat in Cretæ, Cypri, Syriæ, Ægypti aquosis. 2

7. ARUM acaule, foliis peltatis ovatis integerrimis: basi esculentum. semibifidis. *Hort. cliff. 435. Hort. upf. 880. Roy. lugdb. 7.*

Arum minus, nymphææ folio, esculentum. Sloan. jam.

62. bist. 1. p. 167. t. 106. f. 1.

Brassica brasiliiana, foliis nymphææ. Baub. pin. 111.

Habitat in America. 2

8. ARUM acaule, foliis peltatis cordatis repandis: basi bipartitis. *Fl. zeyl. 327. Roy. lugdb. 7. macrorrhizon.*

P P P 3

Arum

- Arum maximum macrorrhizon zeylanicum. Herm. par.*
73. t. 73. *Raj. suppl.* 574.
Habitat in Zeylona. 2
- peregrinum.* 9. *ARUM* acaule, foliis cordatis obtusis mucronatis, angulis rotundatis. *Hort. cliff.* 435. *
Habitat in America. 2
- divaricatum* 10. *ARUM* acaule, foliis cordatis angulatis divaricatis. *Arum* acaule, foliis subhastatis. *Fl. zeyl.* 325. *
Neienschena major. Rheed, mal. 11. p. 39. t. 20.
Habitat in India. 2
- sagittifolium.* 11. *ARUM* acaule, foliis sagittatis triangularibus: angulis divaricatis acutis, *Hort. cliff.* 345. *Raj. lugdb.* 8.
Arum minus esculentum, sagittariæ foliis viridi nigricantibus. Sloan. jam. 63. *hist.* 1 p. 167, t. 106. f. 2.
Arum minus, sagittariæ foliis. Pluk. phyt. 149. f. 2.
Habitat in Brasilia, Jamaica, Barbados. 2
- maculatum,* 12. *ARUM* acaule, foliis hastatis integerrimis, spadice clavato. *Hort. cliff.* 8. *Hort. ups.* 279. *Mat. med.* 417. *Raj. lugdb.* 8. *Dalib. parif.* 279.
Arum vulgare non maculatum. Bauh. pin. 195.
Arum. Fuchs. hist. 69. *Lob. ic.* 597.
Arum venis albis. Bauh. pin. 195.
β. *Arum maculatum, maculis candidis & nigris. Bauh. pin.* 195.
Habitat in Europa australiore. 2
- virginicum.* 13. *ARUM* acaule, foliis hastato-cordatis acutis: angulis obtusis. *Hort. cliff.* 434. * *Gron. virg.* 112.
Habitat in Virginia. 2
- probofci-*
denm. 14. *ARUM* acaule, foliis hastatis, spatha declinata filiformi-tubulata. †
Arisarum flore in tenuem caudam abeunte. Tournes. inst. 161.
Arisarum minus probofci-deum. Bocc. mus. 2. p. 61. t. 50.
Arisarum latifolium minus repens cespitosum. Barr. rar. 182. t. 1150.
Habitat in Apenninis,
- Arisarum.* 15. *ARUM* acaule, foliis cordato-oblongis, spatha bifida, spadice incurvo. *Hort. cliff.* 435. *Raj. lugdb.* 7.
Arisarum latifolium majus. Bauh. pin. 196.
Arisarum latifolium alterum. Bauh. pin. 196.

Arisarum latifolium. *Clus. hist.* 2. p. 73. *Lob. ic.* 598.
Habitat in Mauritaniæ, Italiæ, Lusitanæ, Hispaniæ, Gal-
loprovinciæ nemoribus.

16. ARUM acaule, foliis ovato-oblongis. ovatum.

Karinpola. Rheed. mal. 11. p. 45. t. 23.

Habitat in India.

17. ARUM acaule, foliis lanceolatis, spadice setaceo de- renatifolium.
 clinato. *Hort. cliff.* 345. *Roy. lugdb.* 8. *Sauv. monsp.*
 16.

Arum angustifolium. Bauh. pin. 196. *Clus. hist.* 2. p.

74. *Morij. hist.* 3. p. 545. f. 13. t. 6. f. 21.

Habitat in Dalmatia, Romæ, Monspelii. &

* *Caulescens.*

18. ARUM caulescens, foliis sagittatis. arborescens.

Arum arborescens, sagittariæ foliis. Plum. amer. 44. t.

51. g. 5. 60.

Habitat in America meridionali. &

Numerosa cohors Ari apud Plumierum, Rheede, Her-
mannum, nondum sufficienter genere distincta est.

DRACONTIUM. 966

1. DRACONTIUM scapo brevissimo, petiolo radicato *polyphyllum*
 lacero, foliolis tripartitis: laciniiis pinnatifidis. *Hort.*
cliff. 434. * *Roy. lugdb.* 6

Arum polyphyllum, caule scabro punicante. Herm.
par. 93. t. 93.

Arum polyphyllum sirinamense, caule atrorubente gla-
bro & eleganter variegato. Pluk. alm. 52. t. 149.
 f. 1.

Habitat Surinami. &

2. DRACONTIUM foliis sagittatis, pedunculis petio- *spinosum.*
 lisque aculeatis. *Fl. zeyl.* 328.

Arum zeylanicum spinosum, sagittæ foliis. Herm. par.

75.

Arum minus zeylanicum, sagittariæ foliis. Raj. suppl.

575.

Habitat in Zeylona.

3. DRACONTIUM foliis subrotundis. *Cold. noveb.* 214. * *fastidum.*

Calla aquatilis odore allii vehemente prædita. Gron. virg.

186. *

Habitat in Virginia.

sanasthet-
censq.

4. DRACONTIUM foliis lanceolatis. *Ameri. acad.* 2.
p. 360. *
Habitat in Sibiria.

pertusum.

5. DRACONTIUM foliis pertusis, caule scandente.
Arum hederaceum, amplis foliis perforatis. *Plum. amer.*
40. t. 56. 57.
Habitat in America meridionali. h

907

CALLA.

æthiopica.

1. CALLA foliis sagittato-cordatis, spatha cucullata, spadice superne masculo. *Hort. cliff.* 435.
Arum æthiopium, flore albo odorato moschum olente.
Comm. hort. 1. p. 95. t. 50.
Habitat in Æthiopia.

palustris.

2. CALLA foliis cordatis, spatha plana, spadice undique hermaphrodito. *Hort. cliff.* 436. *Fl. succ.* 744.
Calla foliis cordatis. *Fl. lapp.* 320.
Dracunculus palustris s. radice arundinacea. *Baub. pin.* 195.
Dracunculus aquatilis. *Dod. pempt.* 330.
Provenzialia palustris. *Petit. gen.* 45.
Habitat in Europa borealis paludibus.

708

POTHOS.

scandens.

1. POTHOS. *Fl. zeyl.* 329. *
Potha. *Burm. zeyl.* 197.
Habitat in Zeylona. h

709

ZOSTERA.

marina.

1. ZOSTERA. *It. wgoth.* 166. t. 4. f. 1.
Ruppia foliis linearibus obtusis. *Mæbr. act. angl.* 1741.
p. 217.
Alga angustifolia vitriariorum. *Baub. pin.* 464? *Fl. succ.* 1137.
Fructus s. Alga marina graminea angustifolia seminifera ramosior. *Raj. angl.* 3. p. 52.
Habitat in mari Balthico, Oceano.

Clas

Classis XXI.

MONOECIA.

MONANDRIA.

ZANNICHELLIA. 910

1. ZANNICHELLIA. *Fl. lapp.* 321. *Fl. suec.* 745. *palustris.*
Hort. cliff. 437. *Koy. lugdb.* 9.
Zannichellia palustris major, foliis gramineis. *Micb.*
gen. 71. 2. 34. *f.* 1. 2.
Algoides vulgaris. *Vaill. act.* 1719. *p.* 15. 2. 1. *f.* 1.
Potamogeton capillaceum, capitulis ad alas trifidis.
Bomb. pin. 193. *prodr.* 101.
Potamogetoni similis graminifolia ramosa & ad genicula
polyceraia. *Pluk. alm.* 305. 2. 102. *f.* 7.
Habitat in Europæ, Virginie fœssis, fluviss. ☉

CALLITRICHE. 911

1. CALLITRICHE. *Gen. nov.* 1104. *palustris.*
Callitriche plinif. *Col. ecphr.* 315. 2. 316.
Corspermum foliis oppositis. *Fl. lapp.* 2. *Fl. suec.* 3.
Hort. cliff. 3. *Græn. virg.* 3. *Koy. lugdb.* 205.
Stellaria foliis omnibus subrotundis. *Hall. belv.* 199. minima.
Lenticula palustris bifolia, fructu tetragono. *Bomb. pin.*
 362.
 β. *Stellaria foliis omnibus angustis: apice resecto.* *Hall. bifide*
belv. 198.
Lenticula palustris angustifolia, folio in apice dissecto.
Lœf. pruss. 140. 2. 38.
 γ. *Stellaria foliis ad caulem angustis; in summitate subro-*
tundis. *Hall. belv.* 108.
Stellaria aquatica. *Bomb. pin.* 141.
Habitat in Europæ fœssis paludibus. ☉

CERATOCARPUS. 912

1. CERATOCARPUS. *Amæn. acad.* 1. *p.* 412. *Hort. arenarii*
usp. 281. * *Bomb. act.* 1. *p.* 241.
Habitat in Tatarie arenosis. ☉

CYNOMORIUM.

913
cynomoriu.1. CYNOMORIUM. *Mat. med.* 534.*Cynomorium purpureum officinarum. Mich. gen.* 17. t. 12.*Fungus mauritanicus verrucosus ruber. Petio. gaz.* t. 39. f. 8*Fungus typhoides coccineus tuberosus melitenus. Bocc. mus.* 2. p. 69. sic. 80. t. 81.*Fungus typhoides liburnensis. Till. pis.* 64. t. 25.*Habitat in Sicilia, Melita, Mauritania parasiticum terrestre.*

DIANDRIA.

LEMNA

914
trifolca.1. LEMNA foliis petiolatis lanceolatis. *Fl. lapp.* 470. *Fl. suec.* 1000. *Hort. cliff.* 417. *Roy. lugdb.* 516.*Lemna foliis lanceolatis: extremis decussatis. Sauv. monsp.* 91.*Lenticularia ramosa monorhiza, foliis oblongis pediculis longioribus donatis. Mich. gen.* 16. t. 11. f. 5.*Lenticula aquatica trifolca. Baub. pin.* 362.*Hederula aquatica. Lob. ic.* 2. p. 36.*Habitat in Europa sub aquis pigris parvis.*

minor.

2. LEMNA foliis sessilibus utrinque planiusculis, radicibus solitariis. *Hort. cliff.* 417. *Fl. suec.* 999. *Roy. lugdb.* 516.*Lenticula major monorhiza, foliis subrotundis utrinque viridibus. Mich. gen.* 16. t. 11. f. 3.*Lens palustris. Cam. epit.* 852. *Raj. angl.* 3. p. 129. t. 4. f. 1*Habitat in Europæ aquis quietis.*

gibba.

3. LEMNA foliis sessilibus subtus hemisphæricis, radicibus solitariis. *Hort. cliff.* 417. *Fl. suec.* 998.*Lenticula palustris major: inferne magis convexa, fructu polyspermo. Mich. gen.* 15. t. 11. f. 1.*Habitat in Europæ aquis stagnis.*

polyrhiza.

4. LEMNA foliis sessilibus, radicibus confertis. *Hort. cliff.* 417. *Fl. suec.* 997. *Roy. lugdb.* 516.*Lenticularia major polyrhiza inferne atropurpurea. Mich. gen.* 16. t. 11. f. 1.

Len-

Lenticula palustris major. Raj. angl. 3. p. 129. t. 4 f. 2.
Habitat in Europæ paludibus fossis.

TRIANDRIA.

TYPHA.

915

1. TYPHA foliis subensiformibus, spica mascula femi- latifolia.
 neaque approximatis. *It. scan. 168.*
Typha palustris major. Baub. pin. 20. Moris. hist. 3.
p. 246. f. 8. t. 13. f. 1.
Habitat in paludibus Europæ.
2. TYPHA foliis semicylindricis, spica mascula feminea- angustifolia.
 que remotis. *It. scan. 198.*
Typha. Hort. cliff. 439. Fl. succ. 772. Roy. lugdb. 73.
Gmel. fib. 1. p. 133. Fuchs. hist. 822.
Typha palustris, clava gracili. Baub. pin. 20.
Typha palustris media, Moris. hist. 3. p. 246. f. 8.
t. 13. f. 2.
Habitat in Europæ paludibus, &

SPARGANIUM.

916

1. SPARGANIUM foliis erectis triquetris. *Fl. lapp. erectum.*
345. Fl. succ. 770. Hort. cliff. 439. Roy. lugdb. 73.
Gron. virg. 114. Gmel. fib. 1. p. 133.
Sparganium ramosum. Baub. pin. 15. theatr. 228.
Platanaria f. Butomon. Dod. pempt. 601.
- β. Sparganium non ramosum. *Baub. pin. 15. theatr. 231.*
Habitat in Zonæ frigida septentrionalis aquosis. &
2. SPARGANIUM foliis decumbentibus planis. natans.
Sparganium foliis natantibus plano-convexis. Fl. lapp.
345. Fl. succ. 771.
Sparganium non ramosum minus. Dill. giff. 130.
spec. 58.
Sparganium minimum. Raj. hist. 1910. angl. 3. p. 437.
Habitat in Europæ borealis lacubus, paludibus. &

ZEA.

917

1. ZEA. *Hort. cliff. 437. Hort. mps. 281. Roy. lugdb. 73. Mays.*
Frumentum indicum Mays dictum, Baub. pin. 25.
theatr. 490.

Fru-

Frumentum indicum. *Cam. epit.* 196. *Dod. pempt.* 509.
Habitat in America. ☉
Varietates hujus plurimæ existunt.

918
Lacryma
Jobi.

COIX.

1. COIX seminibus ovatis. *Hort. cliff.* 437. *Hort. ap.* 281. *Fl. zeyl.* 330. *Gron. virg.* 114. *Roy. lugdb.* 7. *Lithospermum arundinaceum.* *Bauh pin.* 258. *Lacryma Jobi.* *Clas. hist.* 2. p. 216. *Habitat in Indiis.* 2
- dactyloides.* 2. COIX seminibus angulatis. *Hort. cliff.* 438. *Roy. lugdb.* 72. *Gramen Dactylon maximum americanum.* *Pluk. alt.* 178 t. 190. f. 2. *Gramen Dactylon indicum esculentum, spica articulata.* *Ambr. phyt.* 1. p. 545. t. 546. 547. *Morif. hist.* p. 185. f. 8. t. 3. f. 11. *Scheuch. gram.* 108. *Sesamum perenne indicum, spica trumentacea.* *Za.* *hist.* 181. t. 68. *Habitat in America.* 2

949

CAREX.

* *Spica unica simplicis.*

- dioica.* 1. CAREX spica simplici dioica. *Hort. cliff.* 438. *M. succ.* 746. *Cyperoides parvum, caulibus & foliis tenuissimis triangularibus, spica longiore (& subrotunda), capsulis oblongis.* *Mich. gen.* 56. t. 32. f. 1. 2. *Gramen cyperoides minimum, spica simplici capsulis.* *Morif. hist.* 3. p. 244. f. 8. t. 12. f. ult. (*mas*). *Scheuch. gram.* 497. t. 11. *Gramen cyperoides minimum, ranunculi capitulo simplici asperiore rotundo.* *Morif. hist.* 3. p. 245. f. t. 12. f. 36. *Habitat in Europæ pratis humidis.* 2
- bulicaris.* 2. CAREX spica simplici androgyna: superne masculis capsulis divaricatis retroflexis. *Carex spica simplici androgyna.* *Hort. cliff.* 438. *M. succ.* 747. *It. gutt.* 240. *Dalib. paris.* 287. *Gmel. f.* 1. p. 144. *Carex spica unica.* *Fl. lapp.* 747. *Carex minima, caulibus & foliis capillaceis, capitulis singulis.*

singulari tenuiori, capsulis oblongis utrinque acuminatis & deorsum flexis. *Mich. gen.* 66. t. 33. f. 1.
Gramen cyperoides minimum, seminibus deorsum reflexis puliciformibus. *Raj. hist.* 1298. *Pluk. phyt.* 34. f. 10. *Morif. hist.* 3. p. 244. f. 8. t. 12. f. 21.

Habitat in Europæ paludibus limosis. &

3. CAREX spica simplici androgyna: inferne mascula, *squarrosa*. capsulis imbricatis horizontalibus.

Habitat in Canada. Kalm.

Spica oblonga, cylindrica, crassa, constans Capsulis horizontalibus, imbricatis, apice lineari longitudine ipsius capsulae prominentibus; inferne Spica paleis exaridis masculis segitur. Inter Carices majores numeranda.

* *Spicis androgynis.*

4. CAREX spica composita, spiculis androgynis: inferioribus remotioribus, foliolo longiori instructis, culmo triquetro. *Is. ar.* 139. *Fl. suec.* 749. *

Carex maritima humilis, radice repente, caule triquetro, spica spadicea. *Mich. gen.* 67. t. 33. f. 4.

Gramen cyperoides spica simplici compressa disticha. *Scheuch. gram.* 498.

Gramen cyperoides minus repens, spica divisa. *Læf. pruss.* 116. t. 31.

Habitat in Europæ arena, præsertim mobili. &

5. CAREX spica composita, spiculis androgynis: inferioribus remotioribus foliolo longiori instructis, culmo tereti. *Fl. suec.* 748.

Habitat in Sueciæ paludibus sylvaticis.

6. CAREX spica composita, spiculis ovatis sessilibus *leporina*, approximatis alternis androgynis nudis. *Fl. suec.* 751.

Gmel. fib. 1. p. 147.

Carex spicis ternis sessilibus confertis androgynis. *Fl. lapp.* 322.

Gramen Cyperoides, spica e pluribus spicis mollibus composita. *Scheuch. gram.* 456.

Gramen Cyperoides palustre majus, spica divisa. *Bauh. pin.* 6. *Morif. hist.* 3. p. 244. f. 8. t. 12. f. 29.

Habitat in Europæ pratis nudis. &

7. CAREX spica supradecomposita inferne laxiore: spiculis androgynis ovatis sessilibus glomeratis: superne masculis

masculis. *Hort. cliff.* 438. *Fl. suec.* 750. *Dalib. parif.* 286. *Roy. lugdb.* 74. *Gmel. sib.* 1. p. 146. t. 32.

Carex palustris major, radice fibrosa, caule exquisite triangulari, spica brevi habitiore compacta. *Mich. gen.* 69. t. 33. f. 13. 14.

Gramen Cyperoides palustre majus, spica compacta.

Bauh. pin. 6. *Morif. hist.* 3. p. 244. f. 8. t. 12. f. 14.

Habitat in Europæ paludibus nemorosis.

muricata.

8. *CAREX* spiculis subovatis sessilibus remotis androgynis, capsulis acutis divergentibus spinosis. *Fl. suec.* 751. *Gmel. sib.* 1. p. 148.

Carex spica androgyna, spiculis compactis obscure alternis, culmo nudo compresso. *Gwett. stamp.* 1. p. 149.

Carex nemorosa, fibrosa radice, angustifolia minima, caule exquisite triangulari, spica brevi interrupta. *Mich. gen.* 69. t. 33. f. 11.

Gramen Cyperoides spicatum minimum, spica divulsâ aculeata. *Scheuch. gram.* 485.

Gramen nemorosum, spicis parvis asperis. *Bauh. pin.* 7. *Morif. hist.* 3. p. 244. f. 8. t. 12. f. 27.

Habitat in Europæ nemoribus humantibus. &

loliacea.

9. *CAREX* spiculis subovatis sessilibus remotis androgynis, capsulis ovatis teretiusculis mucicis divaricatis.

Habitat in Svecia

Radix repens. Folia graminea, tenera, levia. Culmus lavis, superne nudus. Spiculae 4 f. 8, ovatae, parvae, ad apicem culmi sparsae, albidae. Semina ovata, obtusa, mucica demto pistillo, subius rotundata. Differt a *C. muricata*, cui simillima, quod dimidio minor, quod capsula minus divaricata, obtusa, nec margine acuta; ad hanc nostram propius accedunt figura. *Morif. hist.* f. 8. t. 12. f. 26. *Pluk. phyt.* 34. t. 4. quam ad alias.

elongata.

10. *CAREX* spiculis oblongis sessilibus remotis androgynis, capsulis ovatis acutis. *Fl. suec.* 753.

Cyperoides polystachyon, spicis laxis paniculam veluti componentibus. *Scheuch. gram.* 487. t. 11. f. 4.

Habitat in Europa. &

canescens.

11. *CAREX* spiculis subrotundis remotis sessilibus obtusis androgynis, capsulis ovatis obtusiusculis. *Fl. suec.* 745. *Dalib. parif.* 287.

Carex spica androgyna, spiculis compactis obscure alternis, culmo nudo compresso. *Gwett. stamp.* 149.

Ca.

Carex spicis plurimis remotis sessilibus subrotundis tur-
gidis. *Fl. lapp.* 332.

Gramen Cyperoides spicis curtis divulsis. *Læs. pruss.*
117. t. 32.

Habitat in Europa septentrionali. &

* *Spicis sexu distinctis: femineis sessilibus.*

12. CAREX spicis confertis subseffilibus subrotundis: *stava*.
mascula lineari, capsulis acutis recurvis. *Fl. lapp.* 755.
Dalib. parif. 284.

Carex spicis ad apicem tribus seffilibus: femineis sub-
rotundis, capsulis acutis recurvis. *Fl. lapp.* 334.

Gramen Cyperoides palustre aculeatum, capitulo bre-
viore. *T. Scheuch. gram.* 426.

Gramen Cyperoides aculeatum germanicum s. minus.
Bauh. pin. 7. *Morif. hist.* 3. p. 243. f. 8. t. 12. f. 19.

Habitat in Europæ paludibus.

13. CAREX spicis remotis, femineis subseffilibus, ca-
psulis hirtis. *Fl. lapp.* 335. *Fl. suec.* 757. *Dalib. parif.*
282. *Roy. lugdb.* 74.

Carex pilosa, spicis erectis: masculis tribus seffilibus:
secunda minore; femineis duabus remotis. *Hort.*
cliff. 438.

Cyperoides polystachyon lanuginosum. *Morif. hist.* 3.
p. 243. f. 8. t. 12. f. 10. *Pluk. alm.* 178. t. 34 f. 6.

Habitat in Europæ sabulosis.

14. CAREX spicis linearibus erectis: mascula breviora *digitata*.
inferioreque, bracteis aphyllis, capsulis distantibus.
Fl. suec. 758. *Hort. cliff.* 438. *Dalib. parif.* 285.
Roy. lugdb. 75.

Carex spica mascula terminali, quam feminea suprema
breviora, capsulis distantibus. *Fl. lapp.* 338.

Cyperoides montanum nemorosum, caule triquetro
compresso, spicis ferrugineis tenuioribus inter se di-
stantibus. *Mich. gen.* 65. t. 32. f. 9.

Gramen caryophyllum montanum, spica varia. *Bauh.*
pin. 4. *prodr.* 9. *Scheuch. gram.* 448.

Gramen caryophyllum polycarpon, fructu triangulari.
Læs. pruss. 112. t. 27.

Habitat in Europæ nemoribus. &

15. CAREX spicis femineis seffilibus subsolitariis ova-
tis masculæ approximatis, culmo nudo, seminibus
glabris.

Cype-

Cyperoides montanum, foliis angustis parvis, spica spadicea divisa. *Scheuch. gram.* 419?

Cyperoides alpinum saxatile, capillaceo folio, caule rotundo triquetro, spica feminali nigricante plerumque unica. *Mich. gen.* 64. t. 32. f. 3.

Habitat in Europæ montanis apricis. &

Folia mollia, congenerum magis flavescentia.

- globularis.* 16. CAREX spica mascula oblonga, feminea sessili ovata, foliolo florali breviori approximata. *Fl. suec.* 759. Carex spica mascula a feminea in ala folioli remota. *Fl. lapp.* 336.

Habitat in Europa frigidiore.

- filiformis.* 17. CAREX spica mascula oblonga, femineis sessilibus oblongis, inferiore foliolo proprio brevior. *Fl. suec.* 760.

Cyperoides sylvaticum tenuifolium, spicis parvis tenuibus spadiceo-viridibus. *Scheuch. gram.* 425. t. 10. f. 11

Habitat in Europæ nemoribus.

- gyni-fera.* 18. CAREX spicis terminalibus confertis subrotundis: masculina oblonga. *Roy. lugdb.* 75.

Gramen cyperoides tenuifolium, spicis ad summmitatem caulis sessilibus globulorum æmulis. *Pluk. alm.* 178 t. 91. f. 8.

Habitat in Europa.

- saxatilis.* 19. CAREX spicis tribus ovatis sessilibus alternis; mascula oblonga. *Fl. suec.* 756.

Carex spicis tribus ad apicem sessilibus, femineis ovatis atris. *Fl. lapp.* 337.

Cyperoides alpinum saxatile, caule cubitali rotundo-triquetro nudo, spicis brevibus binis s. ternis confertim nascentibus, squamis obtuse mucronatis. *Mich. gen.* 63. t. 32. f. 4?

Habitat in Europæ alpihus.

* *Spicis sexu distinctis: femineis pedunculatis.*

- acutata.* 20. CAREX spicis androgynis terminalibus pedunculatis: florentibus erectis; fructiferis pendulis. *Hort. cliff.* 438. *Gmel. sib.* 1. p. 149.

Carex spicis ad apicem culmi pendulis androgynis. *Fl. lapp.* 324. *Roy. lugdb.* 74.

Gramen alpinum pulchrum, foliis caryophyllæis, spicis atris & tumentibus. *Scheuch. gram.* 481.

Gramen

Gramen Cyperoides alpinum majus, spicis atris plurimis simul junctis. *Rudb. elyf. 1. p. 55. t. 23.*

Habitat in Alpibus Europæ.

21. CAREX spicis ovatis pendulis: mascula longiore *et limosa* rectiore, radice repente. *Hort. cliff. 439. n. 8. Fl. suec. 762. Gmel. sib. 1. p. 140.*

Carex spicis versus apicem pendulis: masculis & feminis. *Fl. lapp. 325.*

Carex spicis seminiferis latis ovalibus pendulis. *Hall. helv. 239.*

Cyperoides spica pendula brevior: squamis e spadiceo f. fusco rutilante viridibus. *Scheuch. gram. 442. t. 10. f. 15.*

Habitat in Europæ frigida paludibus sylvaticis.

22. CAREX spicis pendulis: mascula erecta, femineis ob- *capillari* longis distichis, capsulis nudis acuminatis. *Fl. suec. 763. * It. gozl. 240.*

Carex spicis ex apice pendulis: masculina & femineis albis. *Fl. lapp. 326.*

Gramen Cyperoides pusillum, spica flavescente. *Rudb. elyf. 1. p. 58. t. 40.*

Habitat in Sueciæ pratis humidis.

23. CAREX spicis pendulis: mascula erecta, femineis o- *pallidiusculis* vatis imbricatis, capsulis confertis obtulis. *Fl. suec. 764. * Gmel. sib. 1. p. 139.*

Carex spicis tribus pedunculatis erectis remotis: mascula & femineis. *Fl. lapp. 327. Roy. lugdb. 75.*

Cyperoides polystachion flavicans, spicis brevibus prope summitatem caulis. *Pluk. alm. 178. t. 34. f. 5. Mich. gen. 61. t. 39. f. 13.*

Habitat in Europæ paludibus. &

24. CAREX spicis pedunculatis erectis remotis: femineis *panicæ* linearibus, capsulis obtusiusculis inflatis. *Fl. suec. 765. Dalib. parif. 283. Gmel. sib. 1. p. 136.*

Carex spicis remotis sessilibus, capsulis globosis. *Fl. lapp. 333.*

Carex spicis maribus sessilibus, feminis turgidis & erectis. *Hall. helv. 240.*

Cyperoides foliis caryophyllæis, spicis e rarioribus & tumidioribus vesicis compositis. *Mich. gen. 61. t. 32. f. 11. Pluk. alm. 178. t. 91. f. 7.*

Habitat in Europæ uliginosis. &

filiculata. 25. CAREX spicis terminalibus pedunculatis: mascula femineaque, capsulis subulatis longitudine spicæ.

Habitat in Canada. Kalm.

Culmus triquetus, bipedalis, vestitus Foliis angustioribus. Spica terminalis mascula, oblonga, pedunculata. In eadem ala Spica feminea ex Capsulis 7-8, conico-subulatis, longitudine ipsius spicæ, in fundo semine factis, parulis.

PseudoCyperus. 26. CAREX spicis pendulis, pedunculis geminatis.

rus.

Carex spicis pendulis: omnibus femineis: unica androgyna inferne mascula. *Hort. cliff.* 439. *Fl. suec.* 766. *Roy. lugdb.* 75. *Gron. virg.* 114. *Dalib. parif.* 281.

Cyperoides spica pendula brevior. *Schench. gram.* 440. Gramen Cyperoides spica pendula brevior. *Baub. pin.*

6. *Morif. hist.* 3. p. 242. f. 8. t. 12. f. 5.

PseudoCyperus. Dod. pempt. 339.

Habitat in Europæ fossis.

caespitosa. 27. CAREX spicis erectis cylindricis ternis subseffilibus: mascula terminali, culmo triquetro *Fl. suec.* 767.

Gmel. fib. 1. p. 137.

Gramen Cyperoides cariophyllatum, folio longiore & angustiore, spicis seffilibus compactis erectis. *Morif. hist.* 3. p. 243. f. 8. t. 12. f. 14.

Cyperoides sylvaticum angustifolium, spicis arvis tenuibus spadiceo viridibus. *Schench. gram.* 425?

Habitat in Europæ paludibus turfosiss. &

* *Spicis sexu distinctis: masculis pluribus.*

acuta. 28. CAREX spicis masculis pluribus, femineis subseffilibus, capsulis obtusiusculis. *Fl. suec.* 769. *Dalib. parif.* 281.

Carex spicis erectis oblongis seffilibus alternis foliolo florali brevioribus: superioribus masculis. *Hort. cliff.* 439. *Roy. lugdb.* 74.

nigra. α. Carex nigra verna vulgaris. *Fl. lapp.* 330.

Gramen Cyperoides, foliis cariophyllæis, vulgarissimum. *Raj. hist.* 1292.

rufa. β. Carex spicis masculis superioribus numerosis, femineis inferioribus: intermedia unica androgyna *Roy. lugdb.* 75.

Carex maxima, spicis plurimis remotis longis. *Fl. lapp.* 329.

Cyperoides foliis caryophyllæis, spicis habitioribus: squamis curtis. *Mich. gen.* 62. t. 32. f. 12.

Cype-

Cyperoides latifolium, spica ruffa f. caule triangulo.
Scheuch. gram. 458.

Gramen Cyperoides spica ruffa, f. caule triangulo. *Bauh. pin.* 6.

Habitat in Europa ubique: α. in siccioribus, β. in aquosis. &

29. CAREX spicis masculis pluribus, femineis peduncu-
latis, capsulis inflatis acuminatis. *Fl. suec.* 768. *Dalib. parif.* 282. *Gmel. fib.* 1. p. 143.

Carex spicis plurimis: florescentibus tenuibus, fructiferis
crassis. *Fl. lapp.* 330.

Cyperoides vesicarium, spicis viridantibus & subfuscis.
Scheuch. gram. 470.

Gramen Cyperoides angustifolium, spicis longis ere-
ctis. *Bauh. pin.* 6.

β. Carex culmo longissimo, spicis tenuibus remotis. *Fl. lapp.* 328. *Roy. lugdb.* 75.

Gramen Cyperoides sylvarum tenuius spicatum. *Raj. hist.* 1295. *Morif. hist.* 3. p. 243. f. 8. t. 12. f. 9.
Scheuch. gram. 418.

γ. Carex spicis femininis oblongis pedunculatis breviori-
bus: masculis sessilibus, ultima longissima, capsulis
inflatis acuminatis. *Roy. lugdb.* 76.

Gramen Cyperoides majus præcox, spicis rurgidis tere-
tibus flavescentibus. *Morif. hist.* 3. p. 242.

Habitat in Europæ nudis sylvaticis.

AXYRIS.

920

1. AXYRIS foliis lanceolatis tomentosis, floribus femi-
neis lanatis. *Gen. nov.* 1105. *Ceratoides.*

Urtica foliis lanceolatis. femininis hirsutis. *Roy. lugdb.* 210.

Ceratoides orientalis fruticosa, elæagni folio. *Tournef. cor.* 52.

Habitat in Tataria, Moravia. β

2. AXYRIS foliis ovatis, caule erecto, spicis simplici-
bus. *Gen. nov.* 1105. *Amaranthoides.*

Atriplex montana incanescens, latiusculo folio non si-
nuato, fructu in thyrsulis squarrosis nigricante patu-
lo. *Amm. rub.* 230?

Habitat in Dauria. ☉

Folia scabra pilis stellatis. Rami fructiferi ad basin lon-
ge nudi. Spica minima, subsessilis, simplicissima, ra-
mos terminans.

Q q q 2

3. A-

- hybrida.* 3. *AXYRIS* foliis ovatis, caule erecto, spicis conglomeratis. *Gen. nov.* 1105.
Habitat in Sibiria. ☉
Simillima precedenti, a qua differt Spica longe pedunculata, conglomerata s. secunda, torta, ramis fructiferis juxta caulem congestis Foliis, magisque asperis.
- prostrata.* 4. *AXYRIS* foliis obovatis, caule subdiviso, floribus capitatis. *Gen. nov.* 1105.
Habitat in Sibiria. ☉
Caulis ramosissimus, spithameus, diffusus. Folia obtuse ovata, petiolata. Flores ad apices ramorum conglomerati, immixtis foliolis numerosis.

921
 TRAGIA.

- volubilis.* 1. *TRAGIA* foliis cordato-oblongis, caule volubili.
Tragia scandens, longo betonicæ folio. Plum. gen. 14.
Urtica racemosa scandens angustifolia, fructu tricocco. Sloan. jam. 38. *bist.* 1. p. 123. t. 82. f. 1. *Raj. suppl.* 105.
Habitat in Jamaica. 2. ♀
Caulis volubilis, contra solem scandens. Folia cordata, oblongo-acuminata, argute serrata, petiolata, pilis rigidis adpersa. Racemi axillares, solitarii, masculi. Flosculis viridibus, trifidis. Antheis 3, luteis. Femineus flos axillaris, solitarius, insidens Pedunculo longo, articulato, incurvo. Capsula tricocca, setis rigidis urentibus tecta, singulo loculo extus ad basin notato punctis duobus prominentibus.
- involuta.* 2. *TRAGIA* involucris femineis pentaphyllis pinnatifidis.
Acalypha involucris femineis pentaphyllis pinnatifidis. Fl. zeyl. 34.
Croton foliis ovato-lanceolatis serratis hispidis, caule fruticoso. Roy. lugdb. 201.
Ricinocarpus zeylanicus hirsutus, foliis lanceolatis serratis. Burm. zeyl. 202. t. 92.
Schorigeram. Rheed. mal. 2. p. 73. t. 39. *Raj. bist.* 160.
Habitat in India. ♀
- Mercurialis.* 3. *TRAGIA* foliis ovatis. *Fl. zeyl.* 334.
Mercurialis maderaspatensis tricoccus acetabulis destituta. Pluk. alm. 248. t. 205. f. 4.

Pee-

Pecupameni. *Rheed. mal.* 10. t. 82. *Raj. suppl.* 205.
Habitat in India.

4. *TRAGIA* foliis lanceolatis obtusis integerrimis. *Fl. Chamalea.*
zeyl. 335. *
Chamalea foliis linearibus, flosculis spicatis, echinato
 fructu. *Burm. zeyl.* 59. t. 25.
Codi-avanacu. *Rheed. mal.* 2. p. 63. t. 34. *Raj. bist.*
 1710.
Habitat in India.

HERNANDIA. 922

1. *HERNANDIA.* *Hort. cliff.* 485. t. 23. *Fl. zeyl. senora.*
 423. *
Hernandia amplo hederæ folio umbilicato. *Plum. gen.* 6.
Nux vesicaria oleosa, foliis umbilicatis. *Pluk. alm.* 266.
 t. 208. f. 1.
Habitat in Indiis. b

PHYLLANTHUS. 923

1. *PHYLLANTHUS* caule arboreo, foliis ovatis ob-*grandifolia*,
 tusis integerrimis. *Hort. cliff.* 440.
Habitat in America. b
2. *PHYLLANTHUS* foliis lanceolatis serratis: crenis *Epiphyllanthos*,
 floriferis. *Hort. cliff.* 439. *Roy. lugdb.* 200.
Phyllanthus americana planta, flores e singulis folio-
 rum crenis proferens. *Comm. hort.* 1. p. 199. t. 102.
Seb. thes. 1. p. 21. t. 13. f. 2. *Catesb. car.* 2. p. 26.
 t. 26.
Filicifolia hemionitidi affinis americana *epiphyllanthos*,
 angustiori & longiori folio, ramosa caulescens. *Pluk.*
alm. 134. t. 247. f. 4.
Habitat in America calidiore, Surinamo, Jamaica, Ca-
rolina, Porto Rico. b
3. *PHYLLANTHUS* foliis pinnatis floriferis, floribus *Niru-*
 pedunculatis, caule herbaceo erecto. *Fl. zeyl.* 331.
Hort. nsp. 282.
Phyllanthus foliis alternis alternatim pinnatis, floribus
 dependentibus ex alis foliorum. *Hort. cliff.* 440.
Urinaria indica erecta vulgaris. *Burm. zeyl.* 230. t. 93.
 f. 2.
Fruticulus capsularis hexapetalos, casia poetarum foliis
 brevioribus. *Pluk. alm.* 159. t. 183. f. 5.

Q q q 3

Niru-

- Niruri* barbadense, folio ovali subtus glauco, petiolis
florum brevissimis. *Mart. cent.* 9. t. 9.
Kirganeli. Rheed. mal. 10. p. 29. t. 15. *Raj. dendr.* 29.
Habitat in Indiis. ☉
- spinaria.* 4. **PHYLLANTHUS** foliis pinnatis floriferis, floribus
sessilibus, caule herbaceo procumbente. *Fl. zeyl.* 332.
Fruticulus capularis hexapetalos, brevioribus foliis &
angustis. *Pluk. alm.* 159. t. 183. f. 6.
Tieru-kirganeli. Rheed. mal. 10. p. 8. t. 16. *Raj.*
dendr. 29.
Habitat in India. ☉
Folia in hac erecta, nec ut in antecedenti horizontalia
- maderaspa-* 5. **PHYLLANTHUS** foliolis alternis cuneiformibus
tenis. mucronatis. *Roy. lugdb.* 200.
Nirouri maderaspat. sennæ folio longiore. Pet. gaz.
Habitat in India.
- Emblica.* 6. **PHYLLANTHUS** foliis pinnatis floriferis, caule
arboreo, fructu baccato. *Fl. Zeyl.* 333.
Myrobalanus Emblica. Bauh. pin. 445.
Nellika. Zan. hist. 159. t. 61.
Nelli-camarum. Rheed. mal. 1. p. 69. t. 38. *Raj. hist.*
1556.
Habitat in India. ☿

924

TETRANDRIA.

BETULA.

- alba.* 1. **BETULA** foliis ovatis acuminatis serratis. *Hort. cliff.*
442. *Fl. succ.* 776. *Amœn. acad.* 1. p. 2. *Roy. lugdb.*
85. *Hall. belv.* 158. *Gmel. sib.* 1. p. 166.
Betula foliis cordatis serratis. Fl. lapp. 341.
Betula. Bauh. pin. 427. *Cam. epis.* 69.
Habitat in Europa frigidior. ☿
- nigra.* 2. **BETULA** foliis rhombeo-ovatis acuminatis duplica-
to-serratis.
Betula foliis ovatis oblongis acuminatis serratis. Gron.
virg. 188.
Betula nigra virginiana. Pluk. alm. 67. *Raj. dendr.* 12.
Habitat in Virginia, Canada. ☿
Folia ovata, serrata: serraturis majoribus profundioribus
remotioribus & insuper minimis tenuissimis confertis:
basis foliorum ex angulo obtuso integerrima. Ramuli
pubescentes. Petioli villosi.
3. BE-

3. BETULA foliis cordatis oblongis acuminatis serratis. *lenta*.
Betula julifera, fructu conoide, viminibus lentis. *Gron.*

virg. 115.

Habitat in Virginia, Canada. b

Folia cordato-ovata, acuminata, tenuissime & argute
serrata, glabra. Amentum fructus ovatum, sessile:
squamis acuminatis, integris.

4. BETULA foliis orbiculatis crenatis. *Fl. lapp.* 266. *t. nana.*

6. f. 4. *Fl. suec.* 777. *Amœn. acad.* 1. p. 4. t. 1.

Hort. cliff. 442. *Roy. lugdb.* 85. *Hall. belv.* 158.

Gmel. fib. 1. p. 170.

Betula pumila, foliis subrotundis. *Amm. alt.* 9. p. 314.
t. 14. *ruth.* 209.

Habitat in Alpibus Lapponicis, paludibus Sueciæ, Rus-
siæ. b

5. BETULA pedunculis ramosis.

Alnus.

Alnus. Fl. lapp. 340. *Fl. suec.* 775. *Hort. cliff.* 441.

Roy. lugdb. 85. *Gmel. fib.* 171.

α. *Alnus rotundifolia glutinosa viridis. Bauh. pin.* 428. *glutinosa.*

Alnus. Cam. epit. 68. *Læf. pruss.* 10. t. 1.

β. *Alnus folio incano. Bauh. pin.* 428. *incana.*

Habitat in Europa. b

Limites inter Alnum & Betulam nullos a natura po-
sitos esse docuere Species 2 & 3.

BUXUS. 925

1. BUXUS. *Hort. cliff.* 441. *Hort. upf.* 283. *Mat. med. semperi-*
423. *Roy. lugdb.* 400. *vens.*

α. *Buxus arborescens. Bauh. pin.* 471. *arborescens*

Buxus. Dod. pempt. 782.

β. *Buxus humilis. Dod. pempt.* 782. *suffruticosa.*

Buxus foliis rotundioribus. Bauh. pin. 471.

Habitat in Europa australi.

URTICA. 926

* *Oppositifoliæ.*

1. URTICA foliis oppositis, amentis fructiferis globosis. *pilulifera.*

Hort. cliff. 440. *Hort. upf.* 282. *Roy. lugdb.* 209.

Mat. med. 420. *Samv. monsp.* 307.

Urtica urens pilulas ferens. Bauh. pin. 232.

Urtica 1. d'oscoridis semine lini. Dod. mem. 4. p. 323.

Urtica romana. Lob. ic. 522.

- β. *Urtica altera pilulifera, parietariæ foliis. Dod. mem.* 4. p.
323. *Q q q 4* *Urtica*

Urtica pilulifera, parietariæ facie, semine lini. *Merisbist.* 3. p. 435.

Habitat in Europa australi. ☉

pumila.

2. *URTICA* foliis oppositis ovatis, racemis bipartitis. *Urtica* foliis lanceolato-ovatis petiolorum longitudine, racemis dichotomis petiolo brevioribus. *Gron. virg.* 114.

Habitat in Canada.

Radix fibrosa. Caulis *digiti altitudine, simplex.* Folia *opposita, ovata, trinervia, unda, serrata, petiolis ferè longitudinè digiti.* Racemi axillares, bipartiti vel etiam dichotomi, longitudine petiolorum.

urens.

3. *URTICA* foliis oppositis ovalibus. *Urtica androgyna*, foliis ovalibus. *Hort. cliff.* 440. *Fl. succ.* 773. *Roy. lugdb.* 210. *Dalib. parif.* 288. *Urtica* foliis ovatis, amentis cylindraceis androgynis. *Fl. lapp.* 375. *Hall. helv.* 178. *Urtica urens minor.* *Bauh. pin.* 232. *Urtica urens minima.* *Dod. pempt.* 152. *Habitat in Europæ cultis.* ☉

dioica.

4. *URTICA* foliis oppositis cordatis, racemis geminis. *Urtica dioica*, foliis oblongo-cordatis. *Hort. cliff.* 440. *Fl. succ.* 774. *Mat. med.* 419. *Roy. lugdb.* 210. *Hall. helv.* 177. *Dalib. parif.* 289. *Urtica* foliis cordatis, amentis cylindraceis, sexu distincta. *Fl. lapp.* 374. *Urtica urens maxima.* *Bauh. pin.* 232. *Urtica urens altera.* *Dod. pempt.* 151. *Habitat in Europæ ruderatis.* ♀

cannabina.

5. *URTICA* foliis oppositis tripartitis incis. *Hort. xps.* 282. *Urtica* foliis profunde laciniatis, semine lini. *Amm. rath.* 249. t. 25. *Habitat in Sibiria.* ♀

cylindrica.

6. *URTICA* foliis oppositis oblongis, amentis cylindricis solitariis indivisiis. *Urtica* foliis oblongis serratis nervosis petiolatis. *Gron. virg.* 187. * *Urtica racemosa humilior iners.* *Sloan. jans.* 38. *bist.* 1. p. 124. t. 82. f. 2. *Habitat in Jamaica, Virginia, Canada.* ♀

7. UR.

7. *URTICA* foliis oppositis lanceolatis integerrimis: hinc *parietaria*, angustioribus. *Roy. lugdb.* 210.
Parietaria foliis ex adverso nascentibus, urticæ racemiferæ flore. *Sloan. jam.* 50. *bist.* 1. p. 144. t. 93. f. 1.
Habitat in Jamaica.

* *Alternifoliæ.*

8. *URTICA* foliis alternis cordatis, glomerulis spicatis. *capitata.*
Habitat in Canada. *Kalm.*
Caulis erectus, angulatus, nudus. Folia alterna, cordata, acuminata, equaliter serrata, nuda, petiolis duplo longiora, subtrinervia. Racemi glomerati plures in spicam simplicem dispositi, sessilem, solitariam. Differt a sexta Foliis latioribus alternis.
9. *URTICA* foliis alternis ovatis, racemis compositis *divaricata*, divaricatis.
Urtica racemosa major virginiana mitior f. minus urens. Pluk. alm. 393. t. 237. f. 2.
Habitat in Virginia Canada. 2
10. *URTICA* foliis alternis cordato-ovatis, amentis *ra-canadensis*, mosi distichis erectis. *Hort. cliff.* 441. *Roy. lugdb.* 210.
Urtica canadensis racemosa mitior f. minus urens. Morris. bist. 3. p. 434. f. 11. t. 25. f. 2.
Habitat in Canada, Sibiria. 2
11. *URTICA* foliis alternis ovato-cordatis serratis *pe-interrupta*, tiolo subbrevioribus, spicis solitariis interruptis. *Fl. zeyl.* 336.
Urtica fativa spicata, foliis floribusque petiolis longissimis donatis. Burm. zeyl. 232. t. 110. f. 1.
Urtica pilulifera, foliis majoribus longissimis pediculis: minoribus pediculis donatis. Burm. zeyl. 232. t. 110. f. 1.
Lupulo vulgari similis ind. oriental. floribus in spicam ex origine foliorum prodeuntem. Pluk. alm. 229. t. 201. f. 5.
Batti-Schorigeram. Rheed. mal. 1. p. 75. t. 40.
Habitat in India.
12. *URTICA* foliis alternis orbiculato utrinque acutis *niva*, subtus tomentosis. *Hort. cliff.* 441. *Roy. lugdb.* 210.
Urtica racemifera maxima sinarum, foliis subtus argentea lanugine villosis. Pluk. analzb. 212.
Habitat in Chinæ muris. 2

927

MORUS.

alba.

1. MORUS foliis oblique cordatis lævibus. *Hort. cliff.* 441. *Hort. upf.* 283. *Roy. lugdb.* 211. *Dalib. paris.* 290.
 Morus fructu albo. *Baub. pin.* 459.
 Morus candida. *Dod. pempt.* 810.
Habitat in China. ♀

nigra.

2. MORUS foliis cordatis scabris. *Hort. cliff.* 441. *Hort.*
upf. 283. *Mat. med.* 422. *Roy. lugdb.* 211. *Dalib.*
paris. 290.
 Morus fructu nigro. *Baub. pin.* 459.
 Morus. *Dod. pempt.* 810.
Habitat in Italiæ maritimis. ♀

papyrifera

3. MORUS foliis palmatis, fructibus hispidis.
 Morus sativa, foliis urticæ mortuæ, cortice papyrifera.
Kæmpf. amæn. 471. t. 472.
 Morus papyrifera sativa japonica. *Seb. thes.* 1. p. 44. t.
 28. f. 3.
Habitat in Japonia. ♀

rubra.

4. MORUS foliis cordatis subtus villosis, amentis cylindricis.
 Morus virginienſis arbor, loti arboris inſtar ramoſa, foliis ampliffimis. *Pluk. alm.* 253. t. 246. f. 4.
Habitat in Virginia. ♀
Amenta longitudine Betule albæ. Folia recentia subtus maxime tomentosa; ſæpe etjam palmata ſunt folia.

indica.

5. MORUS foliis ovato-oblongis utrinque æqualibus inæqualiter ſerratis. *Fl. zeyl.* 337. *
 Tinda-parua. *Rheed. mal.* 1. p. 87. t. 49.
Habitat in India. ♀

tatarica.

6. MORUS foliis ovato-oblongis utrinque æqualibus æqualiter ſerratis.
Habitat ad Aſſoff. ♀
Præcedenti maxime affinis, ſed. pedunculis petiolisque longioribus, ſerraturis foliorum magis diſtinctis; fructus M. nigre.

sinctoria.

7. MORUS foliis ovatis hirsutis.
 Morus fructu viridi, ligno ſulphureo tinctorio. *Sloan. jam. hiſt.* 2. p. 3. t. 158. f. 1. *Raj. dendr.* 14.
 Zanthoxylum aculeatum, carpini foliis, americanum, cortice cinereo. *Pluk. alm.* 596. t. 239. f. 3.
Habitat in Braſilia, Jamaica. ♀

PEN-

PENTANDRIA.

XANTHIUM.

928

1. XANTHIUM caule inerme. *Hort. cliff.* 443. *Hort. strumarium.* *apsf.* 284. *Fl. succ.* 778. *Fl. zeyl.* 569. *Mat. med.* 424. *Roy. lugdb.* 85. *Gron. virg.* 188. *Hall. belv.* 161. *Dalib. parif.* 290.
Xanthium. Fuchs. hist. 579.
Lappa minor f. *Xanthium dioscoridis.* *Baub. pin.* 198.
- β. Xanthium elatius & majus americanum, fructu spinulis aduncis armato. *Morif. hist.* 3. p. 604. f. 15. t. 2. f. 2.
Xanthium majus canadense. *Herm. lugdb.* 635.
Habitat in Europa, Canada, Virginia, Jamaica, Zeylona, Japonia. ☉
2. XANTHIUM spinis ternatis. *Hort. apf.* 283. *spinosum.*
Xanthium spinis triplcibus. *Hort. cliff.* 443. *Roy. lugdb.* 85.
Xanthium spinosum, atriplcicis folio. *Morif. hist.* 3. p. 604. f. 15. t. 2. f. 3.
Xanthium lusitanicum spinosum. *Pluk. alm.* 206. t. 239. f. 1. *Herm. parad.* 246. t. 246. *Magn. hort.* 208. t. 208.
Habitat in Lusitania. ☉
Spina trifurcata sunt stipulae, quarum altera fit fructus.

AMBROSIA.

929

1. AMBROSIA foliis trilobis serratis. *Hort. apf.* 284. *trifida.*
Ambrosia foliis palmatis: laciniis lanceolatis serratis. *Hort. cliff.* 443. *Gron. virg.* 188. *Roy. lugdb.* 86.
Ambrosia virginiana maxima, platini orientalis folio. *Morif. hist.* 3. p. 4. f. 6. t. 1. f. 4.
- β. Ambrosia gigantea inodora, foliis asperis trifidis. *Raj. suppl.* 109.
Habitat in Virginia, Canada. ☉
Communis in Hortis planta foliis trilobis est, at varietas major, foliis inferioribus quinquelobis gaudet, cujus fructificationem non vidi.
2. AMBROSIA foliis bipinnatifidis: racemis panicula-elatior. *Hort. apf.* 284.
Ambrosia foliis composito-multifidis: internodiis remotissimis. *Gron. virg.* 188. *Cold. noveb.* 225.
Ambrosia maritima, foliis artemisiæ inodoris, elatior. *Herm. lugdb.* 32. *Raj. suppl.* 109.

Am

Ambrosia virginiana elatior & viridior hortensis facit

Morif. hist. 3. p. 4.

Habitat in Virginia, Canada. ☉

artemisiifolia.

3. *AMBROSIA* foliis bipinnatifidis: primoribus ramulorum indivisis integerrimis.

Ambrosia maxima inodora, marrubii aquatici foliis tunc nunitur laciniatis, virginiana. *Pluk. alm.* 27. t. 10.

5. mala. *Morif. hist.* 3. p. 4. *Raj. suppl.* 108.

Habitat in Virginia, Pennsylvania. ☉

Diversa a precedenti: Folia prima ad exortum ramulorum indivisa, oblonga, integerrima. Caulis primarius vix pede altior, magis diffusus. Rami quadripedales, ascendentes supra racemos, ut racemi laterales evadant, nec terminent plantam, contra in precedenti.

maritima.

4. *AMBROSIA* foliis multifidis, racemis solitariis pilosis.

Ambrosia foliis multifidis tripartitis. *Hort. cliff.* 443. *Roy. lugdb.* 85.

Ambrosia maritima. *Bauh. pin.* 138.

Ambrosia. *Dod. pempt.* 35.

Habitat in Hetruriz, Cappadociæ maritimis arenosis. ☉

PARTHENIUM.

Hypocrepis.

1. *PARTHENIUM* foliis composito-multifidis. *Hort. cliff.* 442. *Hort. upf.* 285. *Roy. lugdb.* 86.

Partheniistrum americanum, ambrosiæ folio. *Niff. aeth.* 1711. p. 423. t. 13. f. 2.

Abinthium crysini folio, achoyari alpini quoddammodo accedens. *Pluk. alm.* 8. t. 45. f. 3. mal.

Habitat in Jamaica glareosis ☉

integrifolium.

2. *PARTHENIUM* foliis ovatis crenatis. *Hort. cliff.* 442. *Gron. virg.* 115.

Partheniistrum helenii folio. *Dill. elsh.* 302. t. 225. f. 292.

Parmica virginiana, scabiosæ austriacæ foliis dissectis. *Pluk. alm.* 308. t. 53. f. 5. & t. 219. f. 1.

Habitat in Virginia. ☉. &

Ingruente nocte comam reflectit.

IVA.

anapa.

1. *IVA* foliis lanceolato-ovatis, caule herbaceo. *Hort. upf.* 285. *Gen. nov.* 1106.

Tar-

Tarchonanthos foliis cordatis serratis trinerviis, *Roy.*
lugdb. 538.

Habitat in America meridionali. D. B. Jussien. ☉

2. *IVA* foliis lanceolatis, caule fruticoso. *Gen. nov.* 1106. *frutescens*,

Parthenium foliis lanceolatis serratis. *Hort. cliff.* 443.

Gron. virg. 115. *Roy. lugdb.* 86.

Agerato affinis peruviana frutescens. *Pluk alm.* 12. t.
27 f. 1.

Elichryso affinis peruviana frutescens. *Herm. lugdb.* 666.

Pseudo-Helichrysum frutescens peruvianum, foliis longis serratis. *Morif. bist.* 3. p. 90.

Habitat in Virginia, Peru. ☿

AMARANTHUS. 932

* *Triandri.*

1. *AMARANTHUS* capitellis triandris subrotundis *tricolor*,
amplexicaulibus, foliis lanceolatis acuminatis.

Amaranthus capitulis subrotundis lateralibus, foliis lanceolatis acuminatis. *Hort. cliff.* 444. *Hort. upf.* 286.

Roy. lugdb. 419.

Amaranthus folio variegato. *Bauh. pin.* 121.

Amaranthus tricolor. *Lob. ic.* 252.

Habitat in India. ☉

2. *AMARANTHUS* capitellis triandris subrotundis sessilibus, foliis lanceolatis spartis. *melancholicus*.

Amaranthus folio variegato C. B. colore obscuriori f.
mas. *Tournef. inst.* 236.

Amaranthus verticillatus melancholicus, macula nigra
in medio foliorum infectus. *Pluk. alm.* 27.

Habitat in India. ☉

*Differt a precedente: Foliis supremis minime in rosam
confertim explicatis. Foliis purpurascentibus basisan-
guineis absque luteo colore. Caule altiore, duriore,
frigoris magis patiente. Florum capitellis neque am-
plexicaulibus, neque confluentibus.*

3. *AMARANTHUS* capitellis triandris subrotundis axillaribus, spicaque glomerata terminali, foliis ovato-cordatis petiolo brevioribus. *nifis*.

Amaranthus inermis, floribus lateralibus sessilibus confertis, foliis ovatis. *Roy. lugdb.* 419?

Habitat in China. ☉

Caulis inferne ramosus. Florum capitula sessilia, ad axillas glomerata, nec non plurima glomerata in spica

cam terminalem laxam. Flosculi aristati. Folia parva, viridia, ovato-cordata, obtusa, longis petiolis infidentia.

Rivinus.

4. **AMARANTHUS** spicis triandris erectis conglomeratis, foliis ovatis retusis.

Amaranthus floribus virentibus densissima spica congestis, foliis amplis rugosis, caule rubro stricto. *Gron. virg.* 116.

Amaranthus maximus, paniculis brevibus gracilissimis surrectis obsolete rubris. *Boerb. lugdb.* 1. p. 97.

Blitum pulchrum rectum magnum rubrum. *Bamb. hist.* 2. p. 966. an?

Habitat in Virginia. ☉

Planta alia colore hepatico-ruffo. Spicæ axillares nullæ. Folia ovata: inferiora retusa j. emarginata; floralia acuta. Flores pallidi, nec rubri. Fructus major, compressus, obtusus.

gracianus.

5. **AMARANTHUS** floribus triandris conglomeratis axillaribus, foliis lanceolatis obtusis.

Amaranthus floribus lateralibus congestis, foliis lanceolatis obtusis. *Gron. virg.* 116.

Habitat in Virginia. ☉

Blitum.

6. **AMARANTHUS** floribus glomeratis lateralibus, foliis ovatis retusis, caule diffuso.

Amaranthus foliis ovatis emarginatis. *Hort. cliff.* 444. *Fl. succ.* 779. *Roy. lugdb.* 420. *Hall. belv.* 176. *Dalib. parif.* 290.

Blitum album minus. *Bauh. pin.* 118. *Cam. epit.* 236.

Habitat in Europa temperatiore. ☉

* *Pentandri.*

caudatus.

7. **AMARANTHUS** racemis pentandris cylindricis pendulis longissimis. *Hort. cliff.* 443. *Hort. nps.* 285.

Fl. zeyl. 563. *Roy. lugdb.* 419.

Amaranthus maximus. *Bauh. pin.* 120.

Blitum majus peruvianum. *Clas. hist.* 2. p. 81.

Habitat in Peru, Persia, Zeylona. ☉

hybridus.

8. **AMARANTHUS** racemis pentandris cylindricis horizontalibus, caule inermi.

Amaranthus sylvestris maximus novæ anglæ, spicis viridibus. *Raj. hist.* 201.

Amaranthus major, virentibus spicarum paniculis, semine nigro. *Barr. ic.* 645.

Habitat in Virginia. ☉

9. A-

MONOECIA HEXANDRIA & HEPTAND. 991

9. **AMARANTHUS** racemis pentandris cylindricis e *hypochondri-*
rectis, caule inermi. *Hort. cliff.* 444 *Hort. ups.* 286. *aqu.*
Amaranthus sylvestris maximus novæ anglizæ, spicis
purpureis. *Tournef. inst.* 235.
Habitat in Virginia. ☉
Caulis viridis. Folia valde acuta, ruffo-lurida, præser-
tim subtus. Spicæ erectæ flosculis purpuratissimis, Sta-
mina 5, flava. Spicæ etjam ex alis longæ.
10. **AMARANTHUS** racemis pentandris lateralibus *retroflexus.*
terminalibusque, caule flexuoso: ramis retrocurvatis.
Habitat in Pennsylvania. *Kalm.* ☉
Planta pedalis Caule firmo, subvillosa, flexuoso a folio
ad folium. Rami inferiores primum recurvi, dein eri-
guntur (sed basi divaricata), hinc retroflexi. Folia o-
vata, acuta *A. Bht.* Spicæ terminales & axillares,
virides. Habitus *A. tristis.* Spicæ crassæ, erectæ. Ra-
mi sub dio minus basi divaricati & retroflexi, quam
in ollis Hortulanorum.
11. **AMARANTHUS** racemis pentandris cylindricis e-spinosus.
rectis, axillis spinosis. *Hort. cliff.* 444. *Fl. zeyl.* 338.
Gron. virg. 188. *Roy. lugdb.* 419.
Amaranthus indicus spinosus, spica herbacea. *Herm.*
lugdb. 31. t. 33.
Habitat in Indiis. ☉

HEXANDRIA.

ZIZANIA.

1. **ZIZANIA** panicula effusa. *933 aquatica.*
Zizania. *Gron. virg.* 189.
Arundo alta gracilis, foliis e viridi cæruleis, locustis mi-
noribus. *Sloan. jam.* 33. *bist.* 1. p. 110. t. 67.
Habitat in Jamaica, Virginiz inandasis.
2. **ZIZANIA** panicula subracemosa. † *terrestis.*
Katou-Tsfolam. Rheed. mal. 12. p. 113. t. 60.
Habitat in Malabariz siccis.

HEPTANDRIA.

GUETTARDA.

1. **GUETTARDA.** *934 speciosa.*
Habitat in Java. ‡

As-

Arbor habitu *Hernandia*. Folia amplissima, ovata s. obovata, apice rotundata cum acumine, nuda, integerrima, venis alternis, petiolata: Petiolis folio longe brevioribus, compressis. Pedunculus e regione petioli, aut etiam Pedunculi (in summis ramis oppositi), terminatus Cyma brevissima, dichotoma. Flores masculi sessiles, alterni, a latere tantum superiore cymæ. Calyces subtoментosi, cylindrici, integerrimi, vix manifeste bilobi. Corolla infundibuliformis: Tubo longo, tomentoso, cylindrico. Limbo septempartito: lobis ovali-oblongis, tubo triplo brevioribus. Stamina 7: Filamentis brevissimis, in fauce corollæ. Antheris oblongis, linearibus. Pistilli germen nullum. Stylus filiformis, tubo dimidio brevior. Stigma cylindrico-capitatum, obtusum. Feminei flores in eadem planta, masculis fere similes. Drupa sicca, subrotunda, torosa, ex Seminibus sex, magnis, lignosis, connexis.

POLYANDRIA

CERATOPHYLLUM.

- 935
demersum. 1. CERATOPHYLLUM. Hort. cliff. 446. Fl. suec. 783. Roy. lugdb. 214. Dalib. parif. 292.
 Hydroceratophyllum folio aspero quatuor cornibus armato. Vaill. act. 1719, p. 21. t. 2. f. 2.
 Equisetum sub aqua repens, foliis bifurcis. Læs. pruss. 67. t. 12.

β. Hydroceratophyllum folio lævi octo cornibus armato. Vaill. ibid.

Habitat in Europæ fossis majoribus sub aqua.

MYRIOPHYLLUM.

- 936
spicatum. 1. MYRIOPHYLLUM floribus masculis interrupte spicatis. Hort. cliff. 446. Fl. suec. 781. Roy. lugdb. 312. Dalib. parif. 292. Sauv. monsp. 160.
 Myriophyllum. Fl. lapp. 343.
 Millefolium aquaticum pennatum spicatum. Bauh. pin 141. prodr. 73. t. 73.
 Habitat in Europæ aquis quietis. &

- verticillatum.* 2. MYRIOPHYLLUM floribus omnibus verticillatis. Hort. cliff. 446. Fl. suec. 782. Roy. lugdb. 213. Dalib. parif. 292. Sauv. monsp. 160.
 Myriophyllum aquaticum minus. Clæs. hist. 2. p. 252.

Mil-

Millefolium aquaticum, flosculis ad foliorum nodos.

Baub. pin. 141.

Habitat in Europæ inundatis. &

Vidi plantas floribus masculis & femineis distinctas, vidi & rarius floribus hermaphroditis.

SAGITTARIA. 9 7 7

1. SAGITTARIA foliis sagittatis acutis. *Fl. lapp.* 344. *sagittifolia.*

Fl. suec. 780. *Roy. lugdb.* 493. *Gron. virg.* 116.

Hall. herb. 300. *Dalib. parif.* 297.

Sagitta minor. Dcd. pempt. 588.

Sagitta aquatica minor latifolia. Baub. pin. 194.

2. *Sagitta aquatica minor angustifolia. Baub. pin.* 194.

3. *Sagitta aquatica major. Baub. pin.* 194.

4. *Sagitta aquatica foliis variis. Læf. pruss.* 234. t. 74.

Gramen bulbosum aquaticum. Baub. prodr. 4.

Habitat in Europæ, Americæ fluviis lacubus argillofis.

An sufficienter distincta SAGITTARIA foliis cordatis, quæ a Moris. hist. 3. f. 15. t. 4. f. 6. delineata.

2. SAGITTARIA foliis sagittatis obtusis. *obtusifolia.*

Sagittariæ foliis planta, glomerato fructu monopyrene, coriandri figura. Pluk. alm. 326. t. 200. f. 7.

Culitamara. Rheed. mal. 11. p. 93. t. 45.

Habitat in Asia.

Differt a præcedente, Foliorum antica parte duplo latiore.

3. SAGITTARIA foliis ternatis. *trifolia.*

Sagittaria chinensis, foliis ternis longissimis. Pet. gaz.

29. t. 19. f. 3.

Habitat in China.

THELIGONUM. 9 3 8

1. THELIGONUM. *Sauv. monsp.* 129. *Cynocrambe.*

Parietaræ affinis Cynocrambe dicta, floribus muscosis.

Pluk. alm. 280.

Cynocrambe dioscoridis. Baub. pin. 122. *prodr.* 59. t.

59. *Magn. char.* 150.

Cynocrambe alifines foliis. Barr. ic. 335.

Alifines facie planta nova. Col. phys. 2. p. 28.

Alifina foetida fabio columnæ dioscoriden. Baub. hist.

3. p. 365. *Raj. hist.* 206.

Habitat in Italia, Monspelii. &

POTERIUM.

- 929
Sanguiforba. 1. POTERIUM inerme, caulibus subangulosis. *Hort. cliff.* 446. *Hort. nps.* 288.
Poterium inerme, filamentis longissimis. *Roy. lugdb.* 129.
Dalis. paris. 291.
Pimpinella Sanguiforba minor hirsuta. *Bauh. pin.* 160.
Pimpinella Sanguiforba. *Cam. epit.* 777.
β. Pimpinella Sanguiforba minor lævis. *Bauh. pin.* 160.
γ. Pimpinella sanguiforba inodora. *Bauh. pin.* 160, *prodr.* 84.
Habitat in Europæ australioris asperis. &
- hybridum. 2. POTERIUM inerme, caulibus teretibus strictis.
Poterium inerme, filamentis florem vix superantibus.
Roy. lugdb. 280
Poterium agrimonoides *Hort. nps.* 288.
Pimpinella agrimonoides. *Morif. bles.* 294. *nmb.* 57. *bist.* 3. p. 264. f. 8. t. 18. f. 9. *Bocc. sic.* 57. t. 29.
Pimpinella agrimonoides odorata. *T. Barr. rar.* 18. t. 632.
Habitat Monspelii. &
- spinosum. 3. POTERIUM spinis ramosis. *Hort. cliff.* 445. *Roy. lugdb.* 240.
Pimpinella spinosa. *Morif. bist.* 3. p. 262. f. 8. t. 18. f. 5.
Poterio affinis, foliis pimpinellæ, spinosa. *Bauh. pin.* 188.
Stœbe legitima dioscoridis. *Clus. bist.* 2. p. 303.
Habitat in Creta, Libano. &

QUERCUS.

- 940
Phecos. 1. QUERCUS foliis lanceolatis integerrimis glabris.
Gron. virg. 117.
Quercus f. Ilex maxilandica, folio longo angusto salicis. *Raj. dendr.* 8. *Catesb. car.* 1. p. 16. t. 16.
β. Quercus foliis oblongis non sinuatis. *Catesb. car.* 1. p. 17. t. 17.
γ. Quercus humilis, salicis folio brevior. *Catesb. car.* 1. p. 22. t. 22.
Habitat in America septentrionali. &
- Smilax. 2. QUERCUS foliis oblongo ovatis subtus tomentosis integerrimis. *Roy. lugdb.* 81. *Sauv. monsp.* 96.
Ilex folio angusto non serrato. *Bauh. pin.* 424.
Smilax dalech. *Bauh. bist.* 2. p. 101. fig.

Habitat in Hispania & adjacentibus.
Admodum affinis sequenti.

3. QUERCUS foliis ovato-oblongis indivisis serratisque, *Ilex*.
 cortice integro. *Hort. cliff.* 448. *Roy. lugdb.* 81.

Ilex oblongo serrato folio. *Bauh. pin.* 424.

Ilex arborea. *Bauh. hist.* 1. p. 95. *

Habitat in Europa australi. †

Variat foliis integerrimis & serratis, sepius etiam in eadem arbore; difficile sine cortice distinguitur a Subere.

4. QUERCUS foliis ovato-oblongis indivisis serratis *Suber*.
 subtus tomentosus, cortice rimoso fungoso. *Hort. cliff.*
 448. *Roy. lugdb.* 81. *Mat. med.* 427. *Sauv. monsp.*
 96.

Suber latifolium sempervirens. *Bauh. pin.* 424.

Habitat in Europa australi. †

5. QUERCUS foliis oblongo-ovatis fimbriato-spinosis *gramuntia*.
 subtus tomentosus, glandibus pedunculatis. *Sauv.*
monsp. 96. †

Ilex foliis rotundioribus & spinosis, e loco *gramuntio*.

Magn. monsp. 140.

Habitat Monspelii. †

6. QUERCUS foliis ovatis indivisis spinoso-dentatis glabris. *Coccifera*
bris. *Roy. lugdb.* 8.

Quercus foliis ovatis dentato-spinosis, glandibus sessilibus. *Sauv. monsp.* 96.

Ilex aculeata cocciglandifera. *Bauh. pin.* 425.

Ilex coccifera. *Cam. epit.* 774.

Habitat in G. Narbonensi, Hispania. †

7. QUERCUS foliis obovatis utrinque acuminatis sinuato-serratis: denticulis rotundatis uniformibus. *Prinus*.
cliff. 448. *Gron. virg.* 117. *Roy. lugdb.* 80.

Quercus, castaneæ foliis, procera arbor, virginiana *Pluk.*

alm. 309. t. 54. f. 3. *Raj. hist.* 1916. *Catesb. car.*

1. p. 18. t. 18.

Habitat in America boreali. †

8. QUERCUS foliis cuneiformibus obsolete trilobis. *nigra*.
Gron. virg. 117.

Quercus folio non serrato in summitate quasi triangulo.

Catesb. car. 1. p. 20. t. 20.

R r r 2

B. Quer-

β. *Quercus marilandica* folio trifido ad *lassafras* accedente. *Raj. Catesb. car.* 19. t. 19.
Habitat in America septentrionali. h

rubra.

9. **QUERCUS** foliis obtuse-sinuatis setaceo-mucronatis.

Quercus foliorum sinubus obtusis: angulis lanceolatis seta terminatis integerrimis vix divisis. *Gron. virg.* 117.

Quercus *esculi* divisura, foliis amplioribus aculeatis. *Pluk. alm.* 309. t. 54. f. 4. *Catesb. car.* 1. p. 23. t. 23.

β. *Quercus* foliorum sinubus obtusis: angulis acutis seta terminatis: intermediis vix tridentatis margine integerrimo. *Hort. cliff.* 448. *Roy. lugdb.* 80.

Quercus *carolinensis*, virentibus venis muricata. *Catesb. car.* 1. p. 21. t. 21. f. 1.

Quercus *virginiana* venis rubris muricata. *Pluk. alm.* 309. t. 54. f. 1.

Habitat in Virginia, Carolina. h

alba.

10. **QUERCUS** foliis oblique pinnatifidis: sinubus angulisque obtusis.

Quercus foliis superne latioribus opposite sinuatis, sinubus angulisque obtusis. *Gron. virg.* 117..

Quercus *alba* *virginiana*. *Catesb. car.* 1. p. 21. t. 21. f. 2
Habitat in Virginia. h

Esculus.

11. **QUERCUS** foliis pinnato-sinuatis lævibus, fructibus sessilibus. *Roy. lugdb.* 80.

Quercus *parva* s. *Fagus* *græcorum* & *Esculus*. *Bauh. pin.* 420.

Habitat in Europa australi. h

Robur.

12. **QUERCUS** foliis deciduis oblongis superne latioribus: sinubus acutioribus; angulis obtusis. *Hort. cliff.* 448. *Fl. succ.* 784. *Mat. med.* 426. *Hall. belv.* 152.

Roy. lugdb. 80. *Dalib. paris.* 293. *Gmel. sib.* 150.

Quercus cum longo pediculo. *Bauh. pin.* 420.

Quercus. *Fuchf. hist.* 229. *Læf. pruss.* 211. t. 69.

Habitat in Europa. h

Ægilops.

13. **QUERCUS** foliis ovato-oblongis glabris serrato-repandis.

Quercus calyce echinato, glande majore. *Bauh. pin.* 420.

Cerri glans *Ægilops* aspris. *Bauh. hist.* 1. p. 77. *fructus.*

Ha-

Habitat in Hispania. ♀
Noscitur calyce maximo.

14. QUERCUS foliis oblongis lyrato-pinnatifidis: laci-*Corvis.*
 nis transversis acutis; subtus subtomentosis.
 Quercus calice hispido, glande minore. *Bauh. pin.* 420.
 Phagus s. Esculus. *Bauh. hist.* 1. p. 74. fig.
Habitat in Hispania. ♀

JUGLANS. 941

1. JUGLANS foliolis ovalibus glabris subserratis sub-*regia.*
 æqualibus. *Hort. cliff.* 449. *Hort. nps.* 286. *Mat. med.*
 430. *Roy. lugdb.* 81. *Dalib. parif.* 293.
 Nux juglans s. regia vulgaris. *Bauh. pin.* 417.
 Nux juglans. *Dod. pempt.* 816.
 2. Nux juglans fructu maximo. *Bauh. pin.* 417.
 3. Nux juglans fructu teneri & fragili putamine. *Bauh.*
pin. 417.
 4. Nux juglans bifera. *Bauh. pin.* 417.
 5. Nux juglans fructu serotino. *Bauh. pin.* 417.
Habitat - - - ♀

2. JUGLANS foliolis lanceolatis serratis: exterioribus alba.
 latioribus.
 Juglans alba, fructu ovato compresso profunde inscul-
 pto durissimo: cavitate intus minima. *Gron. virg.* 190.
 Nux juglans alba virginienfis. *Park. theatr.* 1414. *Ca-*
tesb. car. 1. p. 38. t. 38.
 Nux juglans virginiana alba minor, fructu nucis mo-
 schatæ simili: cortice glabro summo fastigio velut
 in aculeum producto. *Pluk. alm.* 254. t. 309. f. 2.
Habitat in Virginia. ♀
 Folia ex foliolis *sæpius* quinis.

3. JUGLANS foliolis lanceolatis argute serratis; exteri-*nigra.*
 oribus minoribus. *Hort. cliff.* 449. *Hort. nps.* 287.
Gron. virg. 189. *Roy. lugdb.* 82.
 Nux juglans virginiana nigra. *Herm. lugdb.* 452. t. 453.
Catesb. car. 1. p. 67. t. 67.
Habitat in Virginia, Marilandia. ♀
 Folia ex foliolis *sæpius* tredecim.

FAGUS. 942

1. FAGUS foliis lanceolatis acuminato-serratis subtus *Castanea.*
 nudis. *Hort. cliff.* 447. * *Hort. nps.* 287. *Roy. lugdb.*
 79. *Mat. med.* 429. *Dalib. parif.* 294.
 R r r 3 Casta-

Castanea sylvestris. Baub. pin. 419.

Castanea. Cam. epit. 118.

β. *Castanea sativa. Baub. pin. 418.*

Habitat in Italia & australioris Europæ montibus. h

pumila.

2. *FAGUS* foliis lanceolato-ovatis acute serratis subtus tomentosis, amentis filiformibus nodosis. *Gren. virg. 118. **

Fagus foliis ovato-lanceolatis serratis. Roy. lugdb. 79.

Castanea pumila virginiana, racemoso fructu parvo in singulis capsulis echinatis unico. Pluk alm. 90. Castesb. car. 1. p. 9. t. 9.

Castanea americana, foliis averſa parte argentea lanugine villosis. Pluk. alm. 90. t. 156. f. 2.

Habitat in America septentrionali. h

ſylvatica.

3. *FAGUS* foliis ovatis obsolete serratis. *Hort. cliff. 447. Fl. ſuec. 785. Roy. lugdb. 79. Mat. med. 428. Dalib. parif. 294.*

Fagus. Baub. pin. 419. Cam. epit. 112.

Habitat in Europa. h

CARPINUS.

Betulus.

1. *CARPINUS* squamis strobilorum planis. *Hort. cliff. 447. Fl. ſuec. 786. Roy. lugdb. 79. It. ſcan. 46. Dalib. parif. 294.*

Carpinus. Dod. pempt. 841. Cam. epit. 71.

Ostrya ulmo ſimilis, fructu in umbilicis foliaceis. Baub. pin. 427.

Habitat in Europa, Canada. h

Ostrya.

2. *CARPINUS* squamis strobilorum inflatis. *Hort. cliff. 447. Roy. lugdb. 80. Gren. virg. 118. Cold. noveb. 227.*

Ostrya ulmo ſimilis, fructu racemoso lupulo ſimili. Baub. pin. 427.

Ostrya italica, carpinifolio, fructu longiore ſ. brevioris habitore. Mich. gen. 223. t. 104. f. 1. 2.

β. *Aceris cognata Ostrya dicta flaveſcens. Pluk. alm. 7. t. 156. f. 1.*

Habitat in Italia, Virginia. h

CORYLUS.

Avellana.

1. *CORYLUS* ſtipulis ovatis obtuſis. *Hort. cliff. 448. Fl. ſuec. 787. Mat. med. 431. Hort. upſ. 286. Roy. lugdb.*

lugdb. 81. *Hall. belv.* 159. *Dalib. paris.* 294 *Gmel.*
fib. 1. p. 150.

Corylus sylvestris. *Bauh. pin.* 418.

Avellana nux sylvestris. *Fuchf. hist.* 398.

β. *Corylus sativa*, fructu albo minore, f. vulgaris. *Bauh.*
pin. 418.

γ. *Corylus sativa*, fructu rotundo maximo. *Bauh. pin.* 418.

δ. *Corylus sativa*, fructu oblongo rubente. *Bauh. pin.* 418.

ε. *Corylus nucibus in racemum congestis.* *Bauh. pin.* 418.

Habitat in Europæ sepibus. β

2. *CORYLUS stipulis linearibus acutis.* *Hort. cliff.* 448. *Columna.*
Roy. lugdb. 81.

Corylus byzantina. *Herm. lugdb.* 191.

Avellana peregrina humilis. *Bauh. pin.* 418.

Avellana pumila byzantina. *Clus. hist.* 1. p. 11.

Habitat Byzantii. β

PLATANUS, 945

1. *PLATANUS foliis palmatis.* *Hort. cliff.* 447. *Roy. orientalis.*
lugdb. 78.

Platanus. *Bauh. pin.* 431 *Clus. hist.* 1. p. 9. *Dalech.*
hist. 92. *Bell. itin.* 9.

Platanus orientalis verus. *Park. theatr.* 1427.

Habitat in Asia, Tauro, Macedonia, Atho, Lemno,
Creta, locis rignis. β

2. *PLATANUS foliis lobatis.* *Hort. cliff.* 78. *Roy. lugdb. occidentalis.*
7. *Gron. virg.* 119.

Platanus occidentalis. *Catesb. car.* 1. p. 56. t. 56.

Platanus occidentalis f. virginienfis. *Park. theatr.* 1421.

Habitat in America septentrionali. β

Confer. *Platanus orientalis.* *Pock. itin.* 2. t. 89.

LIQUIDAMBAR. 946

1. *LIQUIDAMBAR foliis palmato-angulatis.* *Styraciflua.*
Liquidambar. *Bauh. pin.* 502. *Gron. virg.* 190. *Hort.*

upf. 287. *Mat. med.* 432. *Hort. cliff.* 486. *Roy.*
lugdb. 534. *Cold. noveb.* 228.

Liquidambar arbor f. Styraciflua aceris folio. *Pluk. alm.*

224. t. 42. f. 6. *Catesb. car.* 2. p. 65. t. 65.

Styrax aceris folio. *Raj. hist.* 1681. 1799.

Habitat in Virginia, Mexico. β

2. *LIQUIDAMBAR foliis oblongis sinuatis.* *peregrina.*
Habitat in Canada.

MONADELPHIA.

PINUS.

- 947
Bluestem. 1. * *Foliis pluribus ex eadem basi vaginali.*
 1. PINUS foliis geminis: primordialibus solitariis glabris. *Hort. cliff.* 450. *Fl. suec.* 788. *Mat. med.* 434. *Roy. lugdb.* 89. *Dalib. parif.* 295. *Gmel. sib.* 1. p. 178.
Pinus sylvestris. Baub. pin. 491. *Fl. lapp.* 346. *Daléch. hist.* 45.
 2. *Pinus maritima altera. Baub. pin.* 492.
 3. *Pinaster latifolius, julis virescentibus f. pallescentibus. Baub. pin.* 492.
 4. *Pinaster tenuifolius, julo purpurascente. Baub. pin.* 492.
Habitat in Europæ borealis sylvis glareosis. h
Pinea. 2. PINUS foliis geminis: primordialibus solitariis ciliatis. *Hort. cliff.* 450. *Hort. ups.* 288. *Mat. med.* 435. *Roy. lugdb.* 89.
Pinus sativa. Baub. pin. 490.
Pinus officulis duris, foliis longis. Baub. hist. 1. p. 248.
Pinus. Cam. epit. 93.
Habitat in Italia. h
Teda. 3. PINUS foliis ternis. *Gron. virg.* 190. *
Pinus foliis longissimis ex una theca ternis. Cold. novob. 230.
Pinus virginiana tenuifolia tripilis, f. ternis plerumque ex uno folliculo setis, strobilis majoribus. Pluk. alm. 297. *Raj. dendr.* 8.
Habitat in Virginiz, Canadæ paludosis. h
Cembra. 4. PINUS foliis quinis lævibus. *It. scan.* 32. *
Pinus foliis quinis, cono erecto, nucleo eduli. Hall. belv. 150. *Gmel. sib.* 1. p. 179. 2. 39.
Pinus sativa, cortice fisso, foliis setosis subrigidis ab una vagina quinis. Amm. rutb. 178.
Pinus sylvestris montana tertia. Baub. pin. 491.
Pinus sylvestris Cembra. Cam. epit. 42.
Pinaster. Bell. conif. 19. *Mich. gen.* 223.
Larix sempervirens, foliis quinis, nucleis edulibus. Broyn. E. N. C. cent. 7. obs. 2.
Habitat in Alpibus Sibiriz, Tatariz, Helvetiz, Valle-
liæ

MONOECIA MONADELPHIA. 1001

æ, Baldi, Allobrogum, Tirolensium, Tridentinorum. *h*

5. PINUS foliis quinis scabris.

Strobil.

Pinus foliis longissimis ex uva theca quinis. Cold. noveb. 229.

Pinus virginiana, conis longis non ut in vulgari echinatis. *Pluk. alm. 297.*

Habitat in Virginia, Canada. h

Cotyledones 6 a 10. Folia quina, molliora, triquetra, scabra, angulis inconspicue serratis. Strobili spithamei: apice squamarum crassiore; hæc conjungit Pinum & Abietem Authorum.

6. PINUS foliis fasciculatis acutis.

Cedrus.

Abies foliis fasciculatis acuminatis. Hort. cliff. 449. Roy. lugdb. 89.

Cedrus conifera, foliis laticis. Baub. pin. 490. Ray. hist. 1404.

Cedrus libani. Barr. ic. 499. Edward. ornith. t. 188.

Cedrus phœnicea. Reneal. spec. 27.

Cedrus. Bell. itin. 162. Cam. epit. 57.

Habitat in Syriæ, Libani, Amani, Tauri montibus. h

7. PINUS foliis fasciculatis obtusis.

Larix.

Abies foliis fasciculatis obtusis. Hort. cliff. 450. Mas. med. 437. Roy. lugdb. 89. Gmel. fib. 1. p. 176.

Larix, folio deciduo, conifera. Baub. hist. 1. p. 265. Hort. anglic. 43. f. 11.

Larix. Baub pin. 493. Dod. pempt. 668. Cam. epit. 45. 46.

Habitat in Alpibus Helveticis, Vallesiæ, Stiriæ, Corinthiæ, Tridentinis, Sibiria. h

** Foliis solitariis & basi distinctis.*

8. PINUS foliis solitariis emarginatis.

Picea.

Abies foliis solitariis apice emarginatis. Hort. cliff. 449. Roy. lugdb. 89. Gmel. fib. 1. p. 176.

Abies laxi folio, fructu sursum spectante. Tournef. inst. 585.

Abies conis sursum spectantibus s. mas. Baub. pin. 505.

Abies femina s. Elate teleja. Baub. hist. 1. p. 221.

Habitat in Alpibus Helvetiæ, Sueviæ, Baviaræ, Scotia. h

- Balsamea.* 9. PINUS foliis solitariis subemarginatis: subtus linea duplici punctata.
Abies foliis solitariis confertis obtusis membranaceis.
Gron. virg. 191. *
Abies minor, pectinatis foliis, virginiana, conis parvis subrotundis. *Pluk. alm.* 2. t. 121. f. 1. *Raj. denar.* 8.
Habitat in Virginia, Canada. 2
Habitus antecedentis, sed Folia latiora, obtusiora, utrinque per ramos pectinatim digesta, at duplici serie, superiore sc. breviora, subtus sunt picta linea duplici glauca: singula ex 8 ordinibus longitudinalibus punctorum altorum. Apex foliorum saepe bifidus.
- Ame.* 10. PINUS foliis subulatis mucronatis laevibus bifariam versis.
Abies foliis solitariis apice acuminatis. *Hort. cliff.* 449.
Mat. med. 436. *Fl. suec.* 789. *Fl. lapp.* 347. *Roy. lugdb.* 88. *Hall. belv.* 147. *Dalib. parif.* 295. *Gmel. sib.* 1. p. 175.
Picea. *Cam. epis.* 47.
 β. *Picea* major prima f. *Abies* rubra. *Baub. pin.* 493.
 γ. *Abies* alba f. femina. *Baub. pin.* 505.
Abies minor. *Baub. pin.* 493.
Habitat in Europæ, Asiæ borealibus humidiusculis. 3

THUJA.

- 948
occidentalis. 1. THUJA strobilis laevibus: squamis obtusis. *Hort. cliff.* 449. *Hort. upf.* 289. *Roy. lugdb.* 87. *Gmel. sib.* 1. p. 182.
Thuja theophrasti. *Baub. pin.* 488.
Arbor vitæ. *Clus. hist.* 1. p. 36. *Dod. pempt* 630.
Habitat in Canada, Sibiria. 3
- orientalis.* 2. THUJA strobilis squarrosis: squamis acuminatis reflexis. *Hort. upf.* 289.
Thuja strobilis uncinatis: squamis reflexo-acuminatis. *Roy. lugdb.* 87. * *Hort. cliff.* 449.
Habitat in China. 3

CUPRESSUS.

- 949
semperverens 1. CUPRESSUS foliis imbricatis, frondibus quadrangulis.
Cupressus foliis imbricatis erectis. *Hort. cliff.* 449. *Hort. upf.* 288. *Mat. med.* 433. *Roy. lugdb.* 88. *Dalib. parif.* 295.

Cu-

MONOECIA MONADELPHIA. 1003

Cupressus meta in fastigium convoluta f. Femina. *Tournef. inst.* 587.

Cupressus. *Bauh. pin.* 488. *Cam. epit.* 52.

β. Cupressus ramos extra se spargens f. Mas. *Tournef. inst.* 587.

Habitat in Creta. b

2. CUPRESSUS foliis distichis patentibus. *Hort. cliff. disticha.* 499. *Hort. upf.* 289. *Gron. virg.* 119. *Roy. lugdb.* 88.

Cupressus americana. *Catesb. car.* 1. p. 11 t. 11.

Cupressus virginiana, foliis acaciæ deciduis. *Comm. hort.* 1. p. 113. t. 59.

Cupressus virginiana, foliis acaciæ cornigeræ paribus & deciduis. *Pluk. alm.* 125. t. 85. f. 6.

Habitat in Virginia, Carolina. b

3. CUPRESSUS foliis imbricatis, frondibus ancipitibus. *Thyoides.*

Cupressus nana mariana, fructu cæruleo parvo. *Pluk. mant.* 61. t. 345. f. 1.

Habitat in Canada. *Kalm.* b

Frons f. Rami cum foliis imbricati & compressi exacte eodem modo, quo Thuja, at fructus, magnitudine baccarum juniperi, finditur ut in Cupresso.

ACALYPHA. 950

1. ACALYPHA involucris femineis cordatis incis, fo- *virginica.* liis ovato-lanceolatis petiolo longioribus. *Hort. upf.* 290. *M. zeyl.* 342. *

Acalypha foliis ovato-lanceolatis, involucris femineis obtulis. *Hort. cliff.* 495. *Gron. virg.* 116.

Mercurialis tricoccus hermaphroditica f. ad foliorum juncturas e foliolis cristatis julifera simul & fructum gerens. *Burm. zeyl.* 248 t. 99. f. 4.

Habitat in Zeylouna, Virginia. ☉

2. ACALYPHA involucris femineis cordatis subcrena- *indica.* tis, foliis ovatis petiolo brevioribus. *Fl. zeyl.* 341. *

Mercurialis zeylanica tricoccus cum acetabulis. *Herm. lugdb.* 686. t. 687. *Raj. hist.* 1854. *suppl.* 108.

Cupameni. *Rheed. mal.* 10. p. 161. t. 81. *Raj. suppl.* 107.

β. Urtica minor iners spicata, follo subrotundo serrato, fructu tricocco. *Sloan. jam.* 38. *hist.* 1. p. 125. t. 82. f. 3.

Wellia-Cupameni. *Rheed. mal.* 10. p. 165. t. 82.

Habitat in Indiis ad fimeta. ☉

3. A-

- austrialls.* 3. ACALYPHA involucris femineis integerrimis, foliis lanceolatis obtusis.
 Ricinoides castaneæ folio. *Plum. spec.* 20?
Habitat in America meridionali.

CROTON.

- 951
castaneæ folium. 1. CROTON foliis lanceolatis obtusis serratis petiolatis glabris. *Roy. lugdb.* 201.
Habitat in America meridionali.
- palustre.* 2. CROTON foliis ovato-lanceolatis plicatis serratis scabris. *Hort. cliff.* 445. *Roy. lugdb.* 201.
 Ricinoides palustre, foliis oblongis serratis, fructu hispidulo. *Mart. cent.* 38. t. 38.
Habitat in Vera Cruce, locis paludosis. ☉
- sinzorium.* 3. CROTON foliis rhombeis repandis, capsulis pendulis, caule herbaceo. *Hort. upf.* 290. * *Mat. med.* 441.
 Croton foliis ovatis plicatis scabris, capsulis e pedunculo ramoso pendulis. *Roy. lugdb.* 532. *Sauv. monsp.* 305.
 Ricinoides ex qua paratur Tournesol gallorum. T. *Nissol. act.* 2712. p. 337. t. 17.
 Heliotropium tricoecum. *Bauh. pin.* 253. *Raj. hist.* 165.
Habitat Monspeli. ☉
- argenteum.* 4. CROTON foliis ovatis tomentosis integris serratis. *Hort. cliff.* 444. * *Roy. lugdb.* 201.
Habitat in America. ☉
- sebiferum.* 5. CROTON foliis rhombico-rotundatis utrinque mucronatis integerrimis glabris.
 Ricinus chinensis sebifera, populi nigrae folio. *Pet. gaz.* 53. t. 34. f. 3.
 Arbor sebifera chinensis kiu yeu. *Martin le Compte. Ann. herb.* 375.
Habitat in Chinæ humidis. Osbeck.
 Glandulae duae supra basin foliorum. Folia figura Populi nigrae, sed minora, teneriora.
- Tiglium.* 6. CROTON foliis ovatis glabris acuminatis serratis, caule arboreo. *Fl. zeyl.* 333. * *Mat. med.* 440.
 Ricinoides indica, folio lucido, fructu glabro Grana Tiglia officinis dicto. *Burm. zeyl.* 200. t. 90.
 Pinus indica, nucleo purgante. *Bauh. pin.* 492.
 Lignum moluccense, foliis malvæ, fructu avellanæ mino-

MONOECIA MONADELPHIA. 1005

minore cortice molliore & nigricante, Pavana incolis. *Bamb. pin.* 393.

Cadel-avanacu. *Rheed. mal.* 2. p. 61. t. 75. *Raj. bist.* 167. 1855. *suppl.* 112. 666.

Habitat in India. b

7. CROTON foliis ovatis tomentosis ferrulatis petiolar- *Lacciferum.*
tis, calycibus tomentosis. *Fl. zeyl.* 344. *

Ricinoides aromatica arbor, circææ foliis hirsutis, floribus spicatis, major. *Burm. zeyl.* 201. t. 91.

Ricinus aromaticus spicatus, folio circææ, spinis ad imum duplicatis armato, laccam in granis fundens. *Pluk. alm.* 320. *Raj. suppl.* 113.

Habitat in India. b

8. CROTON foliis cordatis scabris subserratis petiolatis, *aromaticum.*
caule arboreo. *Fl. zeyl.* 347. *

Ricinoides arbor aromatica, circææ foliis, media. *Herm. zeyl.* 202. 11.

Habitat in Zeylona.

9. CROTON foliis cordatis angulatis scabris subtus to- *moluccanum.*
mentosis. *Fl. zeyl.* 346.

Nux juglans moluccana bifida. *Burm. zeyl.* 170.

Habitat in Zeylona, Moluccis.

10. CROTON foliis trilobo-hastatis lanceolatis dentatis. *basarum.*
Ricinus asperior, alceæ venetæ foliis aliquatenus accedens.

Pluk. alm. 320. t. 220. f. 2.

Habitat in India.

11. CROTON foliis inermi-ferratis: inferioribus quin- *lobatum,*
quelobis, superioribus trilobis. *Hort. cliff.* 445. *Roy.*
Ingdb. 201.

Ricinoides herbaceum, foliis trifidis s. quinquefidis & ferratis. *Mart. cent.* 46. t. 46.

Habitat in Vera Cruce. ☉

12. CROTON foliis palmatis quinquelobis trilobisque *spinosum.*
spinoso-ferratis, floribus cauli adpersis subsessilibus. *Fl.*
zeyl. p. 238.

Ricinus maderaspatanus, quinquefidis durioribus foliis margine spinosis. *Pluk. alm.* 320. t. 108. f. 3.

Habitat in India.

13. CROTON foliis ternatis ferratis lanceolatis. *urens.*
Ricinus urens, cannabinis foliis, triphyllis. *Pluk. alm.*
320. t. 120. f. 6.

Habitat in India.

JATRO-

JATROPHA.

952
*gossypifolia.*1. JATROPHA foliis quinquepartitis: lobis ovatis integris: setis glandulosis ramosis. *Roy. lugdb.* 202.*Ricinus minor*, staphisagriae folio, flore pentapetalo purpureo. *Sloan jam.* 41. *bist.* 1. p. 129. t. 84.*Ricinus americanus perennis*, floribus purpureis, staphisagriae foliis. *Comm. hort.* 1. p. 17. t. 9.*Ricinus americanus*, folio staphisagriae. *Banh. pin.* 432-518.*Habitat in America meridionali. 5**Fructificatio Masculi floris:* Perianthium semiquinquefidum: laciniis lanceolatis, ciliatis, acutis. Corolla pentapetala, rosacea: Petalis subrotundis, patentibus, convexis, subtus concavis, hinc revolutis. Nectarium ex glandulis 5, parvis, receptaculo insertis. Filamenta 10, subulata, in medio approximata: quinque alternis corolla brevioribus, erectis; Anthera subrotunda, versatiles. Pistilli rudimentum debile. *Feminei Flores ex eadem umbella cum masculis:* Calyx 5 Corolla ut in masculo. Germen subrotundum. Styli 3. dichotomi. Stigmata simplicia. Capsula subrotunda, tricoeca, trilocularis. Semina solitaria, subrotunda. *D. Royenus.**moluccana.*2. JATROPHA foliis ovatis integerrimis subdentatis. *Fl. zeyl.* 348. **Nux moluccana* folio instar ricini, semine nigro. *Herm. zeyl.* 34.*Habitat in Moluccis, Zeylona. 5**percar.*3. JATROPHA foliis cordatis angulatis. *Hort. cliff.* 449. *Mat. med.* 439. *Roy. lugdb.* 102.*Ricinus americanus major*, semine nigro. *Banh. pin.* 432.*Ricinus ficus* folio, flore pentapetalo viridi, fructu laevi pendulo. *Sloan. jam.* 40.*Mundubignacu.* *Marcgr. bras.* 97.*Habitat in America calidiore. 5**multifida.*4. JATROPHA foliis multipartitis laevibus, stipulis setaceis multifidis. *Hort. cliff.* 445. *Roy. lugdb.* 202.*Ricinoides americanus*, tenuiter diviso folio. *Breyn. cens.* 116. t. 53. *Sloan. jam.* 40. *Raj. bist.* 167. *Morif. bist.* 3. p. 348. f. 10. t. 3. f. ult.*Avellana purgatrix.* *Banh. pin.* 418.

Ma-

MONOECIA MONADELPHIA. 1007

Manihot folio tenuiter divisō. *Dill. rleb* 217. t. 173.
f. 213.

Habitat in America meridionali. b

5. *JATROPHA* foliis palmatis: lobis lanceolatis integer- *Manihot*.
rimis lævibus.

Ricinus minor, viticis obtuso folio, caule verrucoso,
flore pentapetalo albido, ex cuius radice tuberosa, suc-
co venenato turgida americani panem conficiunt
Sloan. jam. 41. *bist.* 1. p. 130. t. 85.

Manihot inodorum f. *Yucca* foliis cannabinis. *Baub.*
pin. 90. *Pluk. alm.* 241. t. 205. f. 1.

Manihot theveti, *Yucca* & *Callavi*. *Baub. bist.* 2. p. 794.

Arbor succo venenato, radice esculenta. *Baub. sin.* 512.

Habitat in America australi. b

6. *JATROPHA* foliis palmatis dentatis aculeatis. *Hort. urcns*,
cliff. 445. *Hort. upj.* 290. *Roy. lugdb.* 202.

Ricinus tithymaloïdes americanus lactescens & *urens*,
floribus albis *Comm. hort.* 1. p. 19, t. 10.

Ricinus lactescens, fici foliis: spinulis mordacis arma-
tis. *Pluk. alm.* 320. t. 220. f. 3.

Habitat in Brasilia. b

7. *JATROPHA* aculeata, foliis trilobis, caule herbaceo. *herbacea*,
Roy. lugdb. 202.

Jussieuia herbacea spinosissima urens, foliis digitatis &
laciniatis. *Houss. Ann. herb.* 256.

Habitat in Vera Cruce. 2

RICINUS. 953

1. *RICINUS*.

Ricinus foliis peltatis ferratis: petiolis glanduliferis. *Hort.*
cliff. 450. *Hort. upj.* 289. *Fl. zeyl.* 339. *Mat. med.*
438. *Roy. lugdb.*

Ricinus vulgaris. *Baub. pin.* 439.

Ricinus gallis Palma christi. *Lob. bist.* 392.

- β. *Ricinus africanus maximus*, caule geniculato rutilante
Tournef. inst. 542.

Habitat in India-utraque, Africa, Europa australi. 3

STERCULIA. 954

1. *STERCULIA* foliis ovatis integerrimis alternis petio- *Balanghar*.
latis, floribus paniculatis. *Fl. zeyl.* 350.

Nux malabarica sulcata mucilaginosā fabacea. *Pluk.*
alm. 266.

Ca-

1008 MONOECIA MONADELPHIA.

Cavalam. *Rheed. mal.* 1. p. 89. t. 49. *Raj. hist.* 1754.
Habitat in India. ♀
Fructus nutans, quinquelobus: lobis reflexis, more Pae-
nia debiscentibus.

senda.

2. STERCULIA foliis digitatis. *Fl. zeyl.* 349.
Nux zeylanica, folio multifido digitato. Pluk. alm. 266.
t. 208. f. 3.
Kail. Rheed. mal. 4. p. 75. t. 36. *Raj. hist.* 1564.
Habitat in India. ♀

955
crepitans.

HURA.

1. HURA. *Hort. cliff.* 486. t. 34. *Roy. lugdb.* 232.
Hura americana, abutili indici folio. Comm. hort. 2. p.
131. t. 66.
Burau ex pluribus nucibus arboris Huræ. Bank. hist. 1.
p. 333. *Stoan. jam.* 214.
Arbor crepitans. Hern. mex. 88.
Habitat in Mexico, Guayava, Jamaica. ♀

SYNGENESIA.

956
Angina.

TRICHOSANTHES.

1. TRICHOSANTHES pomis teretibus oblongis in-
curvis. *Hort. cliff.* 450. *Roy. lugdb.* 262
Angina sinensis, flore albo elegantissimo, fructu ob-
longo intorto. Mich. gen. 12. t. 9.
Cucurbita sinensis, fructu longo anguino vario, flore
candido capillamentis tenuissimis ornato. Till. pis. 49.
t. 22.
Habitat in China. ☉
- nervyalla.* 2. TRICHOSANTHES pomis ovatis acutis, foliis cor-
datis oblongis trinerviis dentatis.
Tota-pivi. Rheed. mal. 8. p. 33. t. 17. *Raj. suppl.* 377.
Habitat in India.
- cucumerina.* 3. TRICHOSANTHES pomis ovatis acutis, foliis cor-
datis angulatis.
Pada valam. Rheed. mal. 8. p. 39. t. 15. *Raj. suppl.* 337.
Habitat in India.
- amara.* 4. TRICHOSANTHES pomis turbinato-ovatis.
Colocynthis flore albo fimbriato, fructu oblongo. Plum.
amer. 86. t. 101. *Raj. suppl.* 332.
Habitat in Domingo. ☉

MOMOR.

MOMORDICA. 957

1. MOMORDICA pomis angulatis tuberculatis, foliis *Balsamina*.
glabris patenti-palmatis. *Hort. cliff.* 451. *Hort. upf.*
293. *Roy. lugdb.* 262.
Balsamina rotundifolia repens f. *mas.* *Bamb. pin.* 306.
Charantia. *Dod. pempt.* 670.
Habitat in India. ☉
2. MOMORDICA pomis angulatis tuberculatis, foliis *Charantia*.
villosis longitudinaliter palmatis. *Hort. cliff.* 457. *Fl.*
zeyl. 351. *Roy. lugdb.* 262.
Balsamina cucumerina indica, fructu majore flavescen-
te. *Comm. hort.* 1. p. 103. t. 54.
Pandipavel. *Rheed. mal.* 8. p. 17. t. 9.
3. *Pavel.* *Rheed. mal.* 8. p. 18. t. 10.
Habitat in India.
Involucrum cordatum, & amplexicaule urget summam
affinitatem cum prima.
3. MOMORDICA pomis oblongis: sulcis catenulatis, fo-
liis incis. *Hort. cliff.* 451. *Hort. upf.* 293. *Fl. zeyl.*
352. *Roy. lugdb.* 262.
Cucumis ægyptius reticulatus f. *Luffa arabum.* *Vest. æ-*
gypt. 199. t. 58. 59 *Morif. hist.* 1. p. 35. f. 7. t. 7.
f. 1. 2.
Habitat in Zeylona. ☉
Semina albida. *Cirrhi apice quasi umbellati ex plurimis*
minoribus cirrbis.
4. MOMORDICA pomis cylindricis longissimis. *cylindrica.*
Pepo indicus reticulatus, seminibus nigris. *Herm. lugdb.*
482. *Pluk. alm.* 286.
Pepo reticulatus & sulcatus, semine nigro. *Herm. parad.*
app. 11. *Raj. suppl.* 332.
Habitat in Zeylona, China. ☉
Caulis quinquangulatus. *Folia cucumerina, angulata,*
basi serrato-dentata, angulis acutis: intermedia pro-
ductiore, sinubus non ut in precedente ampliatis, sed
angulis rectis instructis. *Poma pedalia, subvillosa, cy-*
lindrica. *Semina nigra.* *Flores lutei; Cirrhi ex a-*
xillis.
5. MOMORDICA pomis striatis, foliis pedatis serratis. *pedata.*
Momordica fructu striato lævi. *Few. peruv.* 1. p. 754.
t. 41.
Habitat in Peru.

- bilaterium*. 6. **MOMORDICA** pomis hispidis, cirrhis nullis.
Momordica absque cirrhis. *Hort. upf.* 294. *Mat. med.* 447.
Momordica pomis ovalibus hispidis, foliis cordatis integris plicato-dentatis. *Hort. cliff.* 451. *Roy. lugdb.* 262.
Cucumis sylvestris asininus dictus. *Bauh. pin.* 314.
Cucumis sylvestris. *Cam. epit.* 946.
Habitat in Europa australi. ☉

CUCURBITA.

- lagenaria*. 1. **CUCURBITA** foliis subangulatis tomentosis basi subtus biglandulosis, pomis lignosis.
Cucurbita feminibus obsolete bicornibus. *Hort. cliff.* 451. *Hort. upf.* 291. *Mat. med.* 442. *Roy. lugdb.* 263.
Cucurbita oblonga, flore albo, folio molli. *Bauh. pin.* 313.
Cucurbita lagenaria, flore albo. *Morif. hist.* 2. p. 23. f. 1. t. 5. f. 3.
Habitat in Americae rignis. ☉
- Pepo*. 2. **CUCURBITA** foliis lobatis, pomis lævibus.
Cucurbita seminum margine tumido. *Hort. cliff.* 452. *Hort. upf.* 291. *Roy. lugdb.* 263.
Cucurbita major rotunda, flore luteo, folio aspero. *Bauh. pin.* 213.
Cucurbita indica rotunda. *Dalech. hist.* 616.
 3. *Pepo oblongus*. *Bauh. pin.* 311.
Habitat - - - - ☉
- verrucosa*. 3. **CUCURBITA** foliis lobatis, pomis nodoso-verrucosis.
Cucurbita verrucosa. *Bauh. hist.* 2. p. 222.
Melopepo verrucosus. *Tournef. inst.* 106.
Habitat - - - - ☉
- Melopepo*. 4. **CUCURBITA** foliis lobatis, caule erecto, pomis depresso-nodosis.
Melopepo clypeiformis. *Bauh. pin.* 312.
Cucurbita clypeiformis f. *Siciliana*. *Bauh. hist.* 2. p. 224.
Habitat - - - - ☉
Cirrhi sunt in caule, quamvis non scandat, nec procumbat.
- Citrullus*. 5. **CUCURBITA** foliis multipartitis. *Hort. upf.* 3. *Mat. med.* 443.

Cu-

Cucurbita seminum margine basi dilatato. Hort. cliff.
452. *Roy. lugdb.* 264.

Anguria Citrullus dicta. Bauh. pin. 312.

Citrullus folio colocynthis secto, semine nigro. Bauh.
hist. 2. p. 235.

Habitat in Apulia, Calabria, Sicilia. ☉

CUCUMIS. 959

1. CUCUMIS foliis multifidis, pomis globosis glabris. *Colocynthis.*
Hort. nps. 293. *Mat. med.* 446. *Roy. lugdb.* 263.

Cucumis foliis multifidis. Roy. lugdb. 263.

Colocynthis, fructu rotundo, major. Bauh. pin. 313.

Colocynthis. Cam. epit. 982.

Habitat - - - ☉

2. CUCUMIS foliis palmato-sinuatis, pomis subovatis *Anguria.*
echinatis. Hort. nps. 292.

Cucumis foliis palmatis. Roy. lugdb. 263.

Cucumis sylvestris americanus, anguriae folio, fructu
ovi figura: spinosis tuberculis momordicae instar mu-
ricato. Pluk. phyt. 170. f. 3.

Cucumis anguriae folio latiore aspero, fructu minere
candido, spinulis obtusis muricato. Sloan. jam. 103.

Habitat in Jamaica. ☉

3. CUCUMIS foliis rotundato-angulatis, pomis aeu-*acutangulis.*
tangulis.

Cucumis longus indicus. Grew. mus. 229. t. 17. f. 2.

Cucumis indicus striatus operculo donatus: corticose
putamine tectus. Pluk. alm. 123. t. 172. f. 1?

Habitat in Tataria, China. ☉

Flores lutei in pedunculis multifloris. Poma oblonga:
angulis 10, compressis. Folia cordata, acutae angu-
lata, serrato-dentata. Caulis pentagonus, acutus,
scandens.

4. CUCUMIS foliorum angulis rotundatis, pomis to-*Melo.*
rulosis. Hort. cliff. 451. *Hort. nps.* 292. *Mat. med.*
444. *Roy. lugdb.* 263.

Melo vulgaris. Bauh. pin. 310.

Melo. Bauh. hist. 2. p. 242.

Habitat - - - ☉

5. CUCUMIS hirsutus, foliorum angulis rotundatis, po-*Dulain.*
mis subglobosis.

Melo variegatus, aurantii figura, odoratissimus. Dill.

elsh. 223. t. 177. f. 218.

S f f 2

Melo

Melo ægyptius minor. Tournef. inst. 105.

Melo perficus minimus odoratissimus. Raj. suppl. 333.

Cucumis orientalis, fructu citriformi. Walsb. hort. 133. t. 21?

Habitat in Ægypto, Arabia. Hasselquist. ☉

Sola pilositate differt (an sufficiens?) a Melone. Caulis decumbentes, cirrhosi, obsolete pentagoni: Pili albis, pellucidis, erectis adpersi sunt. Caules, Folia, Calyces & omnia externa, qui pili valde molles & conserti; itaque tota planta ejusmodi mollibus pilis vestita omnino est, ac germen in Melone vulgari, qui reliquis partibus hispida & scaber pilis conicis, rigidiusculis, distantibus, ab hac differt.

foliatus.

6. CUCUMIS foliorum angulis rectis, pomis oblongis scabris. *Hort. cliff. 451, Hort. ups. 292. Mat. med. 445. Roy. lugab. 263.*

Cucumis sativus vulgaris. Bauh. pin. 310.

Cucumis vulgaris. Dod. pempt. 66.

Habitat - - - ☉

maderassanum,

7. CUCUMIS foliis cordatis integris denticulatis, pomis globosis glabris. *Hort. ups. 293.*

*Bryonia foliis cordatis oblongis angulatis dentatis glabris. Fl. zeyl. 356. **

Cucumis maderaspatensis, fructu minimo, Pluk. alm. 123. t. 170. f. 2.

Habitat in India. ☉

BRYONIA.

960
alba

1. BRYONIA foliis palmatis utrinque calloso-scabris. *Hort. cliff. 453. Fl. suec. 790. Roy. lugab. 264. Mat. med. 448. Dalib. parif. 296.*

Bryonia alba, baccis nigris. Bauh. pin. 297.

Vitis alba, baccis nigris. Fuchs hist. 94. Cam. epit. 987.

Habitat in Europa ad pagos & sepes. 2

palmata.

2. BRYONIA foliis palmatis lævibus quinquepartitis: laciniis lanceolatis repando-serratis. *Fl. zeyl. 353. **

Bryonia zeylanica, folio quinquepartito. Burm. zeyl. 353.

Habitat in Zeylona. 2

sardisfolia.

3. BRYONIA foliis cordatis oblongis quinquangulatis dentatis scabris. *Fl. zeyl. 354. **

Habitat in Zeylona.

4. BRYO-

4. BRYONIA foliis palmatis scabris: lacinii lanceola-^{lacinia} tis serratis: lateralibus minimis. *Hort. cliff.* 452. *Fl. zeyl.* 355. *Roy. lugdb.* 264.
Bryonia zeylanica, foliis profunde laciniatis. *Herm. lugdb.* 95. t. 95.
Bryonia zeylanica, flore luteo, fructu rubro. *Herm. parad.* 103. t. 107.
Habitat in Zeylona. 2
5. BRYONIA foliis palmatis quinquepartitis utrinque læ-^{africana} vibus: laciniiis pinnatifidis.
Bryonia foliis palmatis quinquepartitis: subtus punctato-callosis; supra lævibus, lobis laciniatis. *Hort. cliff.* 453. * *Roy. lugdb.* 264.
Bryonia africana lacinata, tuberosa radice, floribus herbaceis. *Herm. parad.* 107. t. 108.
Habitat in Æthiopia. 2
6. BRYONIA foliis palmatis supra calloso-punctatis. ^{cretica}
Hort. cliff. 453. * *Roy. lugdb.* 264.
Bryonia cretica maculata. *Baub. pin.* 297. *prodr.* 135.
Bryonia alba maculata. *Baub. hist.* 2. p. 146.
Habitat in Creta.

SICYOS. 961

1. SICYOS foliis angulatis. *Hort. cliff.* 452. *Hort. apf.* ^{angulata} 291. *Roy. lugdb.* 265.
Bryonioides flore & fructu minore. *Dill. elsh.* 58. t. 51. f. 59.
- β. Cucumis canadensis monospermos, fructu echinato.
Herm. par. 108. t. 133.
Habitat in Canada, Mexico. ☉
2. SICYOS foliis laciniatis.
Sicioides fructu echinato, foliis laciniatis. *Plum. spec.* 3. ^{lacinata}
Habitat in America calidiori.
3. SICYOS foliis ternatis incis. *Roy. lugdb.* 265. ^{trifoliata}
Bryonia alba triphylla geniculata, foliis crassius acidis.
Sloan. jam. 106. *hist.* 1. p. 233. t. 142. f. 56.
Habitat in Jamaica. 2

FEVILLE. 962

1. FEVILLE foliis cordatis angulatis. ^{cordifolia}
Nhandiroba scandens, foliis hederaceis angulosis. *Plum. gen.* 20.
Habitat in America calidior.

S f f 3

2. FE-

trilobata.

2. FEVILLEA foliis trilobis.

Ghandiroba f. Nhandiroba. *Martez. bras.* 46. *Sloan. jam.* 84. *hist.* 1. p. 200. *Raj. hist.* 1875.
Habitat in Brasilia, Jamaica.

GYNANDRIA.

963

ANDRACHNE.

Telephioides

1. ANDRACHNE procumbens herbacea.

Andrachne. Hort. cliff. 443. *Hort. nps.* 285.

Telephioides græcum humifusum, flore albo. Tournef. cor. 50. *Dill. elib.* 377. t. 282. f. 364. *Buxb. cent.*

2. p. 20. t. 12. f. 2.

Glaux procumbens, myrti tarentini folio. Bocc. mus. 2. p. 168. t. 119.

Habitat in Italia, Græcia, Media. ☉

fruticosa.

2. ANDRACHNE erecta arborca.

Habitat in China. Osbeck.

Caules fruticosi Ramis alternis, ad ramulos compressis. Folia ovata, integerrima, petiolata, glabra. Flores axillares, conferti.

Clas.

Classis XXII.

DIOECIA.

MONANDRIA.

NAJAS. 964

1. NAJAS. *Hort. cliff.* 437. *Fl. suec.* 791. *Roy. lugdb. marina.* 10.

Fluvialis vulgaris latifolia. *Vaill. a. 1719. p. 17. t. f. 2.*

Fluvialis latifolia (& angustifolia), fructu monospermo.

Mich. gen. 11. t. 8. f. 1. 2.

Potamogeton fluviatile sargazo simile lucens, foliis margine dentatis. *Pluk. alm.* 304. t. 216. f. 4. *Raj. suppl.* 121.

- β. Fluvialis species, angusto brevique follo, undequaque spinis infecta. *Raj. suppl.* 132.

- γ. Fluvialis species, folio angusto ad margines denticulis spinosis inciso, Flagellum Christi dicta. *Raj. suppl.* 121.

Habitat in Europæ maribus.

DIANDRIA.

VALLISNERIA. 965

1. VALLISNERIA. *Fl. lapp.* 371. *Hort. cliff.* 454. *Roy. spitalis. lugdb.* 9. *Dalib. paris.* 296.

2. Vallisneria palustris, algæ folio, italica, foliis in summitate denticulatis, flore purpurascens. *Mich. gen.* 12. t. 10. f. 1.

Potamogeton, algæ folio, pisanum. *Bocc. mus.* 1. p. 29.

3. Vallisnerioides palustre, algæ folio, italicum, foliis in summitate tenuissime denticulatis, floribus albis vix conspicuis. *Mich. gen.* 3. t. 10. f. 2.

Habitat in Pisæ & Florentiæ fossis.

Folia hujus si fallor vidi in Suecia & alibi, sed nunquam nobis licuit desideratissimas intueri flores.

SALIX. 966

* Foliis glabris serratis.

1. SALIX foliis serratis glabris, floribus hermaphroditis. *hermaphroditica.*
Habitat in...

Habitas Upsaliæ. 3

Convenit cum *S. pentandra* Foliis glabris, obtuse glandulis serratis, convolutis, senis sæpius ex singula gemma, flavescentibus. Amentis villosis, magnitudine & habitu iisdem. Differt vero ab eadem Ramis pallide rufescentibus, nec purpurascensibus. Foliorum pagina superiore venis exaratis, nec venis elevatis. Glandulis ad basin petiolorum extuberantibus. Tempore prius florendi & gallis copiosioribus, quam sequenti.

criandra.

2. *SALIX* foliis serratis glabris, floribus triandris.

Salix foliis elliptico-lanceolatis utrinque glabris serratis appendiculatis. *Hall. belv.* 152. *Gmel. fib.* 1. p. 155. 2. 34. f. 3.

Salix folio auriculato splendente flexilis. *Raj. bist.* 1420.

Habitat in Helvetia, Sibiria. 3

Pæquam affinis antecedenti & sequenti.

pentandra

3. *SALIX* foliis serratis glabris, flosculis pentandris. *Hort. cliff.* 454. *Fl. succ.* 792. *Gmel. fib.* 1. p. 153. 2. 34. f. 1.

Salix pentandra. Fl. lapp. 370. 2. 8. f. 3.

Salix folio laureo f. lato glabro odorato. *Raj. bist.* 1420. alix spontanea, folio amygdalino, fragilis non auriculata. *Bauh. bist.* 1. p. 214.

Habitat in Europæ paludibus montosis duris. 3

phylicifolia.

4. *SALIX* foliis serratis glabris lanceolatis: crenis undatis. *Fl. lapp.* 358. 2. 8. f. *D. Fl. succ.* 793.

β. *Salix* foliis serratis glabris oblongo-ovatis. *Fl. lapp.* 350. 2. 8 f. C.

Habitat in Sueciæ borealibus.

vitellina.

5. *SALIX* foliis serratis ovatis acutis glabris: serraturis cartilagineis, petiolis calloso-punctatis. *Hort. apf.* 295.

Salix foliis lineari-lanceolatis acuminatis. *Guett. stamp.* 1. p. 206.

Salix fativa lutea, folio crenato. *Bauh. pin.* 473.

Habitat in Europa temperatiore.

amygdalina.

6. *SALIX* foliis serratis glabris lanceolatis petiolatis, stipulis trapeziformibus. *Roy. lugdb.* 83. *Fl. succ.* 794. *Dalib. parif.* 297.

Salix foliis serratis glabris acuminatis appendiculatis. *Fl. lapp.* 349. 2.

Salix folio amygdalino utrinque virente aurito. *Bauh. pin.* 43.

Salix

Salix folio auriculato splendente flexilis. Raj. cant. 144.
Habitat in Europæ sylvis.

7. *SALIX* foliis ferratis glabris subovatis acutis sessilibus, hastata stipulis subcordatis.

Salix foliis ferratis glabris subovatis sessilibus appendiculatis. Fl. lapp. 354. t. 8. f. G. Fl. succ. 797.

Salix foliis glabris ovatis ferratis, appendicibus latissimis. Hall. belv. 151.

Habitat in Lapponia, Helvetia. h

8. *SALIX* foliis ferratis glabris ovato-lanceolatis, petio-fragillia lis dentato-glandulosis. *Fl. lapp. 349. t. 8. f. B. Fl. succ. 795.*

Salix foliis ferratis glabris lanceolatis acuminatis appendiculatis. Roy. lugdb. 83.

*Salix, folio longo latoque splendente, fragilis. Raj. angl. 3. p. 448. cantabr. 143. It. scan. 200. **

Habitat in Europæ borealibus. h

Arbor procera Amenta ad basin pedunculi gerant foliola 2. f. 3, sepius caduca. Rami digito concussi ad genicula annosina diffiliunt.

9. *SALIX* foliis ferratis glabris linearilanceolatis, ramis 'babylonica' pendulis. *Hort. cliff. 454. Roy. lugdb. 84.*

Salix orientalis, flagellis deorsum pulchre pendentibus. Tournef. cor. 41.

Habitat in Oriente. h

10. *SALIX* foliis ferratis glabris lanceolatis: inferioribus purpurea oppositis. *It. scan. 252. **

Salix folio longo subluteo non auriculata, viminibus rubris. Raj. angl. 3. p. 450.

Habitat in Europæ australioribus. h

Rami sæpe sanguineo-corallini. Gemmæ univalves prominent Folia 3, transversim cauli applicata (contra in plerisque) ex 2 exterioribus transversim positis.

11. *SALIX* foliis ferratis glabris lanceolato-linearibus: superioribus oppositis obliquis. *Roy. lugdb. 83. n. 4. Daub. parif. 299.*

Salix foliis utrinque lævibus: inferne glaucis, squamis & capsulis tomentosis. Hall. belv. 153.

Salix humilior, foliis angustis subcæruleis ex adverso binis. Raj. hist. 1421. cant. 144. angl. 3. p. 448.

Salix tenuior, folio minore utrinque glabro, fragilis. Baab. hist. 1. p. 213. f. 2.

S f f f

Sa-

Salix Helice theophrasti. Dalecb. hist. 277.

Habitat in Europa australiori ad sepes, locis aquosis. & Rami angulosi. Folia adulta subtus glauca, superiora opposita.

myrtinities

12. *SALIX* foliis serratis glabris ovatis venosis. *Fl. lapp.* 353. t. 8. f. F. & t. 7. f. 6. *Fl. succ.* 799.

Salix myrtilli folio. spica crassissima. Hall. belv. 153.

Salix alpina pumila myrtinities. Pont. comp. 149.

Salix alpina, foliis angustioribus splendentibus serratis. Scheuch. alp. 340.

Salix alpina pumila repens, foliis oblongis exiguis: superne splendentibus, inferne cinereis creberrimis & tenuissimis crenis. Till. pis. 151.

Habitat in Alpibus Lapponiæ, Helvetiæ, Italiæ. &

arbuscula.

13. *SALIX* foliis subserratis glabris subdiaphanis subtus glaucis, caule suffruticoso. *Fl. succ.* 798. *Gmel. sib. 1. p. 166.*

Salix foliis serratis glabris obovatis. Fl. lapp. 352. t. 8. f. E.

β. *Salix* foliis integris glabris ovatis confertis pellucidis. *Fl. lapp. 356.*

γ. *Salix* foliis serratis glabris lanceolatis utrinque acutis. *Fl. lapp. 360. t. 8. f. M.*

Habitat in Lapponiæ campis arenosis. &

herbacea.

14. *SALIX* foliis serratis glabris orbiculatis. *Fl. lapp. 355. t. 8. f. N. & t. 7. f. 3. 4. Fl. succ. 800. Roy. lugdb. 82. Hall. belv. 155.*

Salix saxatilis minima. Baux. pin. 474. prodr. 159.

Salix alpina lucida repens, alni rotundæ folio. Boeck. mus. 2. p. 19. t. 11. f. ult. Raj. suppl. 12. angl. 3. p. 448.

Herba facie pyrolæ. Mart. spitsb. 47. t. G. f. B.

Habitat in Alpibus Lapponiæ, Helvetiæ. &

* *Foliis glabris integerrimis.*

reticulata.

15. *SALIX* foliis integerrimis glabris ovatis obtusis.

Salix foliis integris glabris ovatis subtus reticulatis. *Fl. lapp. 359. t. 8. f. L. & t. 7. f. 1. 2. Fl. succ. 801.*

Salix pumila, folio rotundo. Baux. hist. 1. p. 12. Scheuch. alp. 43 340. Raj. hist. 1423.

Salix pumila, folio rotundo integerrimo, julo gracili. Hall. belv. 154.

Habitat in Alpibus Lapponiæ, Helvetiæ. &

Gen

Gemmæ promant Folia 3, ovata, obtusa, petiolata, integerrima, obsolese pilosa: supra viridia rugosa; subtus glauca reticulata: venis primariis rubris demum viridibus. Spica ex eadem gemma.

16. SALIX foliis integris glabris ovatis acutis. *myrtilloides.*

Salix foliis integris glabris ovatis alternis. *Fl. lapp.*

357. t. 8. f. J. K. *Fl. succ.* 804.

Habitat in Suecia septentrionali. b

17. SALIX foliis integerrimis, subtus tenuissime villosis, *glauca.*

ovato-oblongis. *Fl. lapp.* 363. t. 7. f. 5. & t. 8. f. P.

Fl. succ. 802.

Salix alpina pyrenaica. *Bauh. pin.* 474. *prodr.* 159.

Burf. XXIV. 120.

Habitat in Alpibus Lapponicis & Pyrenaicis. *Sauv.* b

* Foliis integerrimis, villosis.

18. SALIX foliis integerrimis utrinque villosis obovatis *aurica,*

appendiculatis. *Fl. lapp.* 369. t. 8. f. T. *Fl. succ.* 810.

Roy. lugdv. 84

Salix folio rotundo minore. *Dill. app.* 37. *Raj. angl.*

3. p. 450.

Habitat in Europæ borealis sylvis. b

19. SALIX foliis utrinque lanatis subrotundis acutis. *Fl. laneta,*

lapp. 368. t. 8. f. X. & t. 7. f. 7. *Fl. succ.* 809.

Salix humilis latifolia erecta. *Bauh. pin.* 474. *prodr.* 159.

Raj. bist. 1426. *Hall. helv.* 154.

Habitat in Alpibus Lapponicis. b

Folia conferta, ovata, utrinque lanata, integerrima, subsefilia.

20. SALIX foliis integerrimis hirsutis lanceolatis. *Fl. lapp. Lapponum.*

366. t. 8. f. T. *Fl. succ.* 808. *Hall. helv.* 155. t. 5.

Habitat in Alpibus Lapponiæ ubique, Helvetiæ. b

21. SALIX foliis integris ovatis acutis supra: subvillosis, *arenaria,*

subtus tomentosis. *Fl. succ.* 806. *Is. goth.* 206.

Salix foliis integris subtus villosis ovatis acutis *Fl. lapp.*

362. t. 8. f. U. Q.

Salix foliis integris utrinque hirsutis lanceolatis. *Hall.*

helv. 155. t. 5. f. 2.

Salix pumila, foliis utrinque candicantibus & lanuginosis.

Bauh. pin. 474. *Raj. angl.* 3. p. 447. *Gmel. sib.* 1.

p. 164. t. 36. f. 1.

Habitat in Europæ paludibus. x-b.

22. SA-

- incubacea.* 22. *SALIX* foliis integerrimis lanceolatis: subtus villosis nitidis, stipulis ovatis acutis. *Fl. succ.* 807.
Salix foliis ovato-lanceolatis integerrimis, ramis decumbentibus, radice reptatrice. *Roy. lugdb.* 84. *Dalib. paris.* 299.
Salix pumila, foliis ellipticis integerrimis: subtus glaucis, spica rotundiore. *Hall. belv.* 153.
Habitat in Europæ pascuis durinsculis, uliginosis. &c.
- repens.* 23. *SALIX* foliis integerrimis lanceolatis utrinque subpilosis, caule repente. *Fl. succ.* 814.
Habitat inter montes Sueciæ, locis humidis. &c.
Caulis crassitie digiti. Rami virgati, decumbentes. Ramuli adpersi villis albis. Folia ovata s. ovato-oblonga, integerrima, utrinque glabra, subtus glauca, petiolata, stipulis nullis: infima opposita transversim & porum pilosa. Amenta lateralia Feminis capsula rufescentes.
- fusca.* 24. *SALIX* foliis integerrimis ovatis: subtus villosis nitidis. *Fl. lapp.* 364. t. 8. f. R. *Fl. succ.* 803. * *Roy. lugdb.* 84.
Salix pumila, folio utrinque glabro. *Banh. hist.* 1. p. 217.
Habitat in Europæ pascuis humidis. &c.
Gemma univalvis, bipartita: Amentum absque foliis.
- osmarinifolia.* 25. *SALIX* foliis integerrimis lanceolato-linearibus strictis sessilibus subtus tomentosis. *Roy. lugdb.* 84. n. 13.
Salix humilis angustifolia. Banh. pin. 474.
Salix homilis repens angustifolia. Lob. ic. 2. p. 137.
Banh. hist. 1. p. 214.
Habitat in Europæ campis depressis. &c.
Folia subtus nitida, sericea, villosa.
 * *Foliis subserratis, villosis.*
- Caprea.* 26. *SALIX* foliis ovatis rugosis: subtus tomentosis, undatis superne denticulatis. *Roy. lugdb.* 83. *Fl. succ.* 811. *Gmel. sib.* 1. p. 811. *Dalib. paris.* 298.
Salix foliis subcrenatis utrinque villosis ovato-oblongis. *Fl. lapp.* 365 t. 8. f. S. *Roy. lugdb.* 83. *Dalib. paris.* 298.
Salix folio rugoso obiter serrato subtus lanuginoso, juto crassissimo. *Hall. belv.* 155.
Salix latifolia rotunda. Banh. pin. 474. *Raj. angl.* 3. p. 449.
- β. *Salix* foliis integris, subtus villosis, ovato-lanceolatis utrinque acutis. *Fl. lapp.* 365. t. 8. f. N.

- Salix folio rotundo acuminato integerrimo subtus sericeo. *Hall. belv.* 154.
27. Salix foliis oblongis subtus villosis: inferioribus crenatis, superioribus integris. *Fl. lapp.* 367. t. 8. f. U.
28. Salix foliis elliptico-lanceolatis subtus sericeis appendiculatis. *Gmel. fib.* 1. p. 169.
Habitat in Europæ siccis. 3
 Gemmæ inferiores Folia, superiores Amenta non foliacea producant. 3
27. SALIX foliis subintegerrimis lanceolato-linearibus *viminatis*. longissimis acutis subtus sericeis, ramis virgatis. *Fl. succ.* 813. *Dalib. parif.* 297.
 Salix foliis undulatis lineari-lanceolatis subtus tomentosis. *Roy. lugdb.* 84.
 Salix folio longissimo angustissimo utrinque albedo. *Bauh. pin.* 474.
 Salix folio longissimo. *Raj angl.* 3. p. 450.
 Salix foliis angustis & longissimis, foliis crispis: subtus albicantibus. *Bauh. hist.* 1. p. 212.
28. Salix foliis ex lineari-lanceolatis integris subtus incanis. *Gmel. fib.* 1. p. 163.
Habitat in Europa ad pagos. 3
 Frutex junceus, altissimus. Gemmæ foliaceæ & florifera distinctæ ut in Caprea. Folia ante explicationem in hac revoluta.
28. SALIX foliis subserratis oblongo-ovatis: subtus subcinerea, villosis, stipulis dimidiato-cordatis. *Fl. succ.* 805.
 Salix foliis integris glabris lanceolato-ovatis. *Fl. lapp.* 358.
Habitat in Europæ nemoribus paludosis subhumidis. 3
 Amenta florum fusca, infra folia collocata, Pedunculo Foliolis aliquot minimis, lanceolatis. Stipulæ dimidiato-cordatæ, utrinque serratæ glandulis tribus.
29. SALIX foliis lanceolatis acuminatis serratis utrinque *alba* pubescentibus: serraturis intimis glandulosis. *Hort. cliff.* 473. *Fl. succ.* 812. *Mat. med.* 449. *Dalib. parif.* 297. *Roy. lugdb.* 83.
 Salix foliis elliptico-lanceolatis serratis subtus sericeis, vimine fragili. *Hall. belv.* 152.
 Salix vulgaris alba arborescens. *Bauh. pin.* 473.
 Salix alba. *Is. scan.* 200.
Habitat ad pagos & urbes Europæ. 3
 Est arbor maxima cum Salice fragili.

Spe

Species hujus generis difficillime extricantur. Solum palustre, arenosum, alpestre, calidum mutavit mirametamorphosi species, ut de iisdem basilarint sapius Botanici; accessit Descriptio ubique imperfecta & barbara. Incipienda itaque harum historia e novo; quod fit, si rejiciantur veterum nomina, ne confundant synonyma dubia certas plantas, & in floribus & Herbariis describantur 1. Gemmae univalves s. multivalves. 2. Foliatio intra gemmas situ & plicatione. 3. Amenta ex eadem vel distincta gemma, supra aut infra folia. 4. Folia serrata aut integerrima: paginis nudis aut villosis; 5. Stamina numerus. 6. Arbores frutices, repentes.

TRIANDRIA.

EMPETRUM.

album.

1. EMPETRUM erectum. Hort. cliff. 470. Roy. lugdb. 206.

Empetrum lusitanicum, fructu albo. Tournef. inst. 579.

Erica erecta, baccis candidis. Bauh. pin. 436.

Erica coris folio X. Clus. hist. 1. p. 45.

Habitat in Lusitania. ♀

Sequenti nimis affinis.

nigrum.

2. EMPETRUM procumbens. Hort. cliff. 470. Fl. suec. 832. Roy. lugdb. 206. Hall. helv. 162.

Empetrum. Fl. lapp. 379.

Erica baccifera procumbens nigra. Bauh. pin. 436.

Erica baccifera. Clus. pan. 29.

Habitat in Europæ frigidissimæ montosis paludosis, ♀

OSYRIS.

alba.

1. OSYRIS. Roy. lugdb. 202. Sauv. monsp. 56.

Osyris frutescens baccifera. Bauh. pin. 212.

Casia poetica monspeliensis. Cam. epit. 26. Lob. ic. 432.

Casia latinorum. Alp. exor. 41.

Habitat in Italia, Hispania, Monspeli. ♀

Femina potius hermaphrodita, cum staminibus gaudeat, tamen sine mare sterilis.

Obs. Osyris foliis obtusis. Hort. upf. 295. est similis s. Nitraria, cujus flores nondum vidi, Frutex Lycio similis, sed foliis conjugatis sessilibus.

TE-

TETRANDRIA.

VISCUM. 969

1. VISCUM foliis lanceolatis obtusis, caule dichotomo, *album*,
spicis axillaribus.
Viscum. *Hort. cliff.* 441. *Fl. suec.* 816. *Mat. med.* 450.
Roy. lugdb. 400. *Hall. herb.* 162. *Dalib. paris.* 300.
Viscum baccis albis. *Bauh. pin.* 425.
Viscum. *Cam. epit.* 555.
Habitat in Europæ arboribus, parasitica. h
2. VISCUM foliis lanceolatis obtusis, spicis lateralibus. *rubrum*.
Viscum foliis longioribus, baccis rubris. *Catesb. car.*
2. p. 81. t. 81.
Habitat in Carolina parasitica. h
3. VISCUM foliis obovatis, racemis lateralibus. *purpureum*.
Viscum foliis latioribus, baccis purpureis, pediculis in-
sidentibus. *Catesb. car.* 2. p. 95. t. 95.
Habitat in Carolina, parasitica. h
4. VISCUM caule prolifero ramulissimo aphyllis com-*opuntioideis*,
pressis.
Viscum opuntioideis ramulis compressis. *Sloan. jam.* 168.
hist. 2. p. 93. t. 201. f. 1. *Raj. dendr.* 52.
Habitat in Jamaica, parasitica. h
5. VISCUM caule verticillato. *verticilla-
tum*.
Viscum ramulis & foliis longis densissimis striatis & ra-
diatis. *Sloan. jam.* 168. *hist.* 2. p. 93. t. 201. f. 2. *Raj.*
dendr. 52.
Habitat in Jamaica, parasitica. h
6. VISCUM caule herbaceo tetragono brachiato, foliis *terrestre*,
lanceolatis.
Habitat in Philadelphie pratis subhumidis. Kalm. IX.
32.
Caulis herbaceus, pedalis, lavis, brachiatus, tetragonus.
Folia lanceolata, petiolata, levia, integerrima, atri-
na: inferiora minora. Spicæ axillares, sessiles, car-
nosæ, oblongæ. Fructificatio imperfecta: flores feminei
a me tantum visi. Fructus mihi ignotus; de genere
itaque dubium.

HIPPOPHAE. 970

1. HIPPOPHAE foliis lanceolatis.

Rhamnoides

Hip-

Hippophaë. *Fl. lapp.* 372. *Fl. suec.* 815. *Hort. cliff.* 454. *Roy. lugdb.* 207.

Rhamnus salicis folio angustiore, fructu flavescente. *Bauh. pin.* 755.

Rhamni species. *Cam. epit.* 81.

Rhamnoides florifera, salicis folio. *Tournef. cor.* 53.

Habitat in Europæ maritimis arenosis. h

equadens. 2. HIPPOPHAE foliis ovatis.

Habitat in Canada. Kalm. h

Frutex facie antecedentis, sed Folia latiora, dimidio breviora, ovata s. oblongo-ovata. Rami oppositi. Racemi simplices inter folia prima, erecti, foliis dimidio breviores. Folia supra viridia, adpersa pilis fasciculatis, divergentibus, nudo oculo vix discernendis. Subtus folia pilis & squamis argentea, adpersa squamis s. punctis ferrugineis.

MYRICA.

Gale. 1. MYRICA foliis lanceolatis subserratis, caule suffruticoso.

Myrica foliis lanceolatis, fructu sicco. *Fl. lapp.* 373. *Mat. med.* 451. *Fl. suec.* 817. *Hort. cliff.* 455. *Roy. lugdb.* 527. *Dalib. parif.* 300.

Rhus myrtifolia belgica. *Bauh. pin.* 414.

Chamaeleagnus. *Dpd. pempt.* 780. *app.*

Habitat in Europæ septentrionalis uliginosis. h

condensa. 2. MYRICA foliis lanceolatis subserratis, caule arborecente.

Myrica foliis lanceolatis subserratis, fructu baccato. *Hort. cliff.* 455. *Hort. nps.* 295. *Gron. virg.* 120. *Roy. lugdb.* 527.

Myrtus brabanticae similis carolinienfis baccifera, fructu racemoso sessili monopyreno. *Pluk. alm.* 250. t. 48. f. 9. *Catesb. car.* 1. p. 69. t. 69.

β. Myrtus brabanticae similis carolinienfis humilior, foliis latioribus & magis serratis. *Catesb. car.* 1. p. 13. t. 13. *Habitat in Carolina, Virginia, Pennsylvania. h*

asplenifolia. 3. MYRICA foliis oblongis alternatim sinuatis. *Hort. cliff.* 456. *Gron. virg.* 192. *Cold. noveb.* 224.

Gale mariana, asplenii folio. Pet. mus. 773.

Myrt brabanticae affinis americana, foliorum laciniis asplenii modo divis. *Pluk. alm.* 250. t. 100. f. 6. 7.

Habitat in America septentrionali. h

4 MY-

4. MYRICA foliis oblongis opposite sinuatis. *Hort. cliff. quercifolia.*
 456. *Roy. lugdb.* 527.
Cōriotrægematodendros africana, botryos amplioribus
 foliis densis. *Pluk. amaltb.* 65.
Laurus africana minor, folio quercus. *Comm. hort.* 2. p.
 161. t. 81. *Raj. dendr.* 85.
Habitat in Æthiopia. ♀
5. MYRICA foliis subcordatis serratis sessilibus. *Hort. cordifolia.*
cliff. 456. *Roy. lugdb.* 527.
Alaternoides ilicis folio crasso hirsuto. *Waltb. hort.* 3.
 t. 3.
- Tithymali facie planta æthiopica, ilicis aculeato folio.
Pluk. alm. 373. t. 319. f. 7.
Gale capensis, ilicis cocciferæ folio. *Pet. mus.* 774.
Habitat in Æthiopia. ♀

PISTACIA. 972

1. PISTACIA foliis subternatis. *Hort. cliff.* 456. *Roy. trifolia.*
lugdb. 78. *Sauv. meth.* 219.
Pistacium mas ficulum, folio nigricante. *Bocc. mus.* 2.
 p. 139. t. 93.
Terebinthus f. *Pistacia trifolia.* *Tournef. inst.* 580.
Habitat in Sicilia. ♀
2. PISTACIA foliis pinnatis ternatisque suborbiculatis. *narbonensis.*
Pistacia foliis sæpius quinq. orbiculatis. *Sauv. monsp.* 219.
Terebinthus peregrina, fructu majore pistaciis simili e-
 duli. *Bauh. pin.* 400. *Tournef. inst.* 579.
Terebinthus indica major, fructu rotundo. *Bauh. bist.*
 1. p. 277.
3. Terebinthus major, pistaciæ folio. *Lob. adv.* 412.
Habitat Monspeli. ♀
3. PISTACIA foliis impari pinnatis: foliolis subovatis vena
 recurvis. *Mat. med.* 454.
Pistacia peregrina, fructu racemoso f. *Terebinthina indi-*
ca. *Bauh. pin.* 401.
Pistacia. *Bauh. bist.* 1. p. 275.
Terebinthus indica theophrasti, *Pistacia dioscoridis.* *Lob.*
adv. 413. *Tournef. inst.* 380.
Habitat in Persia, Arabia, Syria, India. ♀
4. PISTACIA foliis impari-pinnatis: foliolis ovato-lan- *Terebinthus*
 ceolatis. *Hort. cliff.* 456. *Mat. med.* 452. *Hort. upf.*
 296. *Roy. lugdb.* 78. *Sauv. monsp.* 219.

T t t

Tere-

Terebinthus vulgaris. *Baub. pin.* 400. *Tournef. inst.* 579.
Terebinthus. *Clus. hist.* 1. p. 15. *Dod. pempt.* 871.
Habitat in Europa australi, Africa boreali, India. &
Confer *Therebinthus peregrina*, fructu minore cæruleo
 eduli. *C. B.* 296.

- Lentiscus.* 5. *PISTACIA* foliis abrupte pinnatis: foliolis lanceola-
 tis. *Hort. cliff.* 456. *Hort. nsp.* 996. *Mat. med.* 453.
Roy. lugdb. 79. *Sauv. monsp.* 219.
Lentiscus vulgaris. *Baub. pin.* 399. *Tournef. inst.* 580.
Lentiscus. *Cl. f. hist.* 1. p. 14. *Dod. pempt.* 871.
Habitat in Hispania, Lusitania, Italia. &
Confer *Lentiscus ex chio*, ex qua fuit mastiche. *Tournef. it.* 1. p. 285.

- Simaruba.* 6. *PISTACIA* foliis pinnatis deciduis: foliolis ovatis.
Terebinthus major, *betulæ* cortice, fructu triangulari.
Sloan. jam. hist. 2. p. 89. t. 199. *Catesb. car.* 1. p. 30.
 t. 30?
Simaruba. *Mat. med.* 535.
Habitat in Jamaica, Carolina. &

CERATONIA.

- Siliqua.* 1. *CERATONIA.* *Hort. nsp.* 296. *Mat. med.* 455. *Roy.*
lugdb. 225.
Siliqua edulis. *Baub. pin.* 400.
 2. *Siliqua.* *Cam. epis.* 139.
Habitat in Apulia, Sicilia, Creta, Cypro, Syria, Pala-
estina. &

PISONIA.

- aculeata.* 1. *PISONIA* aculeata.
Pisonia. *Hort. cliff.* 457.
Pisonia aculeis rigidissimis hamatis, floribus racemosis.
Amm. herb. 581.
Pisonia aculeata, fructu glutinoso & racemoso. *Plum.*
gen. 7.
Rhamnus f. *Lycium Fringego* jamaicensibus dictum.
Pluk. alm. 318. t. 108. f. 2.
Paliuro affinis arbor spinosa, flore herbaceo pentapeta-
 loide, fructu sicco nudo cannulato lappaceo. *Sloan.*
jam. 137. *hist.* 2. p. 25. t. 167. *Raj. dendr.* 95.
Habitat in America meridionali. &

- mitis.* 2. *PISONIA* inermis.
Pisonia malabarica non spinosa. *Amm. herb.* 582.

Katu

Karu-kara-walli. *Rheed. mal.* 7. p. 33. t. 17.
Habitat in India. ♀

ANTIDESMA. 975

1. ANTIDESMA. *Fl. zeyl.* 357. * *acniteria.*
 Arbor indica, ovalli folio, flosculis plurimis in spicis summo ramulo dispositis, acniteria. *Pluk. mant.* 22. v. 329. f. 1.
Noelietali. Rheed. mal. 4. p. 19. t. 56. *Raj. hist.* 1616.
 ♂ Baccifera maderaspatana, ribis more floribus muscosis juli instar. *Pet. mus.* 621. *Raj. dendr.* 27.
Habitat in India. ♀

SPINACIA. 976

1. SPINACIA. *Hort. cliff.* 457. *Hort. upj.* 296. *Mat. oleacea.*
med. 536. *Roy. lugdb.* 220.
Spinacia mas. Dalech. hist. 543. ♀
Spinacia semine spinoso. Moris. hist. 2. p. 598.
Lapathum hortense f. *spinacia semine spinoso. Bauh. pin.* 114.
Spinachia femina. Dalech. hist. 543. ♂
 ♀ *Lapathum hortense* f. *Spinacia semine non spinoso. Bauh. pin.* 115.
Habitat - - - - ☉

ACNIDA. 977

1. ACNIDA. *Mitch. gen.* 28. *Gen. nov.* 1091. *cannabina.*
Cannabis foliis simplicibus. Gron. virg. 192. *
Cannabis virginiana. Bauh. pin. 320.
Habitat in Virginiae paludibus salis.

CANNABIS. 978

1. CANNABIS. *fativa.*
Cannabis foliis digitatis. Hort. cliff. 457. *Hort. upj.* 297. *Mat. med.* 457. *Dalib. paris.* 300. *Roy. lugdb.* 221.
Cannabis fativa. Bauh. pin. 320. ♀
Cannabis mas. Dalech. hist. 497. ♀
Cannabis erratica. Bauh. pin. 320. ♂
Cannabis femina. Dalech. hist. 497. ♂
Habitat in India. ☉

979

HUMULUS.

- Lupulus.* 1. HUMULUS. *Hort. cliff.* 458. *Fl. suec.* 818. *Mat. med.* 458. *Dalib. parif.* 301. *Roy. lugdb.* 222.
 ♀ *Lupulus mas. Baub. pin.* 298.
Lupulus salictarius Fuchs. hist. 124.
 ♂ *Lupulus femina. Baub. pin.* 298. *Cam. epit.* 954.
Habitat in Europæ sepibus & ad radices montium. &

980

ZANONIA.

1. ZANONIA.

- Penar-valli mascula. Rheed. mal.* 8. p. 39. t. 49.
Penar-valli femina. Rheed. mal. 8. p. 31. t. 47. 48.
Habitat in Malabar.

981

HEXANDRIA.

TAMUS.

- communis.* 1. TAMUS foliis cordatis indivisis. *Hort. cliff.* 458.
Roy. lugdb. 228. *Hall. belv.* 165.
Bryonia lævis f. nigra racemosa, cujus baccæ rufescunt & nigrescunt. Baub. pin. 297.
Bryonia sylvestris baccifera. Baub. prodr. 135.
 ♂ *Bryonia lævis f. nigra racemosa. Baub. pin.* 297.
 ♀ *Bryonia lævis f. nigra baccifera. Baub. pin.* 297.
Vitis sylvestris f. Tamus. Dod. pempt. 400.
Habitat in Europa australi. &
- cretica.* 2. TAMUS foliis trilobis.
Tamnus cretica, trifido folio. Tournef. cor. 3.
Habitat in Creta. &

982

SMILAX.

* *Caule aculeato, angulato.*

- aspera.* 1. SMILAX caule aculeato angulato, foliis dentato-aculeatis cordatis.
Smilax caule angulato aculeato, foliis cordato-oblongis acutis aculeatis. Hort. cliff. 458. *Roy. lugdb.* 228.
Smilax aspera, fructu rubente. Baub. pin. 296.
Smilax aspera, rutilo fructu. Clus. hist. 1. p. 112..
 β. *Smilax aspera minus spinosa, fructu nigro. Baub. pin.* 236.
Smilax aspera, nigro fructu. Clus. hist. 1. p. 113.
 γ. *Smilax viticulis aspera, foliis longis angustis mucronatis.*

lævibus: auriculis ad basin rotundioribus. *Pluk. alm.* 348. t. 110. f. 3.

Habitat in Hispaniâ, Italiâ, Siciliâ sepibus h.

2. **SMILAX** caule aculeato angulato, foliis inermibus *excelsa*, sagittatis.

Smilax caule angulato aculeato alternatim inflexo, foliis sagittatis acutis inermibus. *Roy. lugdb.* 228.

Smilax orientalis, sarmentis aculeatis excelsâ arborescens, foliis non spinosis. *Tournef. cor.* 45. *Buxb. cent.* 1. p. 18. t. 27. *mal.*

Smilax aspera. *Alp. aegypt.* 140. t. 141?

Habitat in Oriente, Syria. h.

3. **SMILAX** caule aculeato angulato: foliis inermibus: *ceylonica*, caulinis cordatis: ramis ovaro-oblongis. *Fl. Zeyl.* 364.*

Smilax indica spinosa, folio cinnamomi, Pseudo China quibusdam. *Burm. zeyl.* 22.

Habitat in Zeylona. h.

4. **SMILAX** caule aculeato angulato, foliis inermibus *Sarsaparilla*, ovatis retuso-mucronatis.

Smilax caule angulato aculeato, foliis ovatis acutis inermibus. *Roy. lugdb.* 228.

Smilax caule angulato aculeato; foliis dilato-cordatis inermibus acutis. *Hort. cliff.* 459. *Gron. virg.* 120. *Mat. med.* 460.

Smilax viticulis asperis virginiana, folio hederaceo leni Zarza nobilissima. *Pluk. alm.* 348. t. 111. f. 2. *Raj. suppl.* 345.

Smilax aspera peruviana, f. *Sarsaparilla*. *Bauh. pin.* 296.

Habitat in Peru, Brasilia, Mexico, Virginia. h.

* *Caule aculeato, tereti.*

5. **SMILAX** caule aculeato teretiufculo, foliis inermibus ovato-cordatis.

Smilax caule aculeato, foliis orbiculato-ovatis inermibus quinquenerviis. *Mat. med.* 461.

Smilax minus spinosa, fructu rubicundo, radice virtuosa China dicta f. *Sankiva*. *Kampf. amoen.* 781. t. 782.

China radix. *Bauh. pin.* 896.

Fruticulus convolvulaceus spinosus sinicus, floribus parvis umbellatis. *Pluk. amalsh.* 101. t. 408. f. 1.

Habitat in China, Japonia. h.

Petiolibus *bidentatis*. Folia ovato-cordata, obtusa cum acumine.

- rotundifolia*. 6. **SMILAX** caule aculeato tereti, foliis inermibus reniformi-cordatis acutis.
Habitat in Canada. Kalm. 3
Caulis teres, nudus, flexuosus, aculeis rectis rarioribus.
Folia cordata, inermia, nuda, quinquenervia, latiora quam longa, acuta petiolis brevibus innixa; e petiolo Cirrhi duo, filiformes.
- aurifolia*. 7. **SMILAX** caule aculeato tereti, foliis inermibus ovato-lanceolatis trinerviis. *Gron. virg. 193.*
Smilax lævis, lauri folio, baccis nigris. Catesb. car. 1. p. 15. t. 15.
Habitat in Virginia, Carolina. 3
Folia, quoad consistentiam, crassissima in suo genere.
- tamnoides*. 8. **SMILAX** caule aculeato tereti, foliis inermibus cordatis oblongis multinerviis.
Smilax bryoniæ nigra foliis, caule spinoso, baccis nigris. Catesb. car. 1. p. 52. t. 52.
Habitat in Carolina, Virginia, Pennsylvania.
- seduca*. 9. **SMILAX** caule aculeato tereti, foliis inermibus ovatis.
Habitat in Canada. Kalm. 3
Caulis teres, nudus, flexuosus, aculeis rectis rarioribus apice nigris. Folia ovata, trinervia, annua, acuta: Petiolis brevibus e quibus Cirrhi duo, capillares. Umbellæ infra folia pedunculis vix petiolo longioribus insidentes.
- * *Caulis inermi, angulato.*
- hera nox*. 10. **SMILAX** caule inermi angulato, foliis ciliato-aculeatis.
Smilax aspera indiæ occidentalis. Bauh. pin. 296.
Smilax, foliis latis in margine spinosis, caroliniana, stipite quadrato. Pluk. alm. 348. t. 111. f. 1.
 β. *Smilax caroliniana, stipite quadrato leni, foliis angulatis asperis auriculatis ad basin angulosis. Pluk. alm. 348. t. 111. f. 3.*
Habitat in Carolina.
- herbacea*. 11. **SMILAX** caule inermi angulato, foliis inermibus ovatis. *Gron. virg. 193.*
Smilax lævis marilandica, foliis hederæ nervosis prælongis pediculis insidentibus, flosculis minimis in umbellam parvam congestis. Ray. suppl. 345.

Smi-

Smilax claviculata, hederæ folio, tota lævis, e terra mariana. *Pluk. alm.* 348. t. 110. f. 4?

Habitat in Virginia, Marilandia. 2

* *Caule inermi, tereti.*

12. **SMILAX** caule inermi tereti, foliis inermibus lam-^{lanceolata.} ceolatis. *Hort. cliff.* 459. *Gron. virg.* 120.

Smilax, viticulis asperis, virginiana, foliis angustis lævibus nullis auriculis prædita. *Pluk. alm.* 349. t. 110. f. 4? *Raj. suppl.* 344?

Habitat in Virginia.

Differt nostra a Plukenetiana aculeorum absentia.

13. **SMILAX** caule inermi tereti, foliis inermibus: cau-^{Pseudo Chi-} linis cordatis, racemis ovato-oblongis. na.

Smilax caule tereti inermi, foliis inermibus: caulinis cordatis, ramorum lanceolatis, pedunculis longissimis. *Gron. virg.* 193.

Smilax caule tereti inermi, foliis cordato-ovatis acutis inermibus, petiolis bidentatis. *Hort. cliff.* 459. *Gron. virg.* 120.

Smilax virginiana, spinis innocuis armata, latis canellæ foliis, radice arundinacea crassa nodosa & carnosæ. *Pluk. alm.* 349. t. 110. f. 5.

Smilax aspera, fructu nigro, radice nodosa magna lævi farinacea. *Sloan. jam.* 105. *bist.* 1. p. 231. t. 143. f. 1.

China spuria nodosa. *Banh. pin.* 297

Habitat in Virginia, Jamaica.

CISSAMPELOS. 983

1. **CISSAMPELOS** foliis peltatis corœatis emarginatis. *Peruvia.*

Mat. med. 459.

Clematis baccifera glabra & villosa, rotundo & umbilicato folio. *Plum. amer.* 78. t. 93. *fil. t.* 183. *Sloan. jam.* 85. *bist.* 1. p. 200.

Convolvulus brasiliensis, flore octopetalo, monococcos. *Raj. bist.* 1331.

Hedera monophyllos virginiana, umbilicatis foliis, baccata, officulo compresso lunato singulari. *Pluk. mant.* 99.

Caapeba folio orbiculari & umbilicato lævi. *Plum. gen.* 33.

Caapeba brasiliensis. *Marcgr. bras.* 24. *Pis. bras.* 94.

3. *Caespeba* folio orbiculari umbilicato & tomentoso.
Plum. gen. 33.
Habitat in America meridionali.
- Caespeba.* 2. *CISSAMPELOS* foliis margine petiolatis integris.
Caespeba folio orbiculari non umbilicato. *Plum. gen. 33.*
Habitat in America meridionali.
- Smilacina.* 3. *CISSAMPELOS* foliis cordatis acutis angulatis.
Smilax ? lenis folio anguloso hederateo. *Catesb. car. 1*
p. 51. t. 51.
Habitat in Carolina.
Fructificationem perfectam non vidi.

984

RAJANIA.

- hastata.* 1. *RAJANIA* foliis hastato-cordatis. *Hort. cliff. 458.*
Jan-Raja scandens, folio oblongo angusto auriculato.
Plum. gen. 33.
Bryonia fructu alato, foliis auriculatis. *Plum. amer. 84.*
t. 98. fil. 1. 178.
Habitat in Domingo.
- cordata.* 2. *RAJANIA* foliis cordatis septinerviis.
Jan-Raja scandens, folio tamni. *Plum. gen. 33.*
Habitat in America meridionali.
Folia cordata, oblonga, acuminata, reticulata. Flores
racemosi.
- quinquefolia* 3. *RAJANIA* foliis quinis.
Jan-Raja scandens quinquefolia. *Plum. gen. 33.*
Habitat in America.
Folia ad singula genicula quina, ovato-lanceolata, obtu-
siuscula, petiolata. Racemi ex medio internodiorum,
non autem ex axillis.

185

DIOSCOREA.

- pentaphylla.* 1. *DIOSCOREA* foliis digitatis. *Hort. cliff. 459. Fl.*
zeyl. 363.
Ricophora pentaphyllos, caule spinoso, fructu oblongo
 triquetro, malabaræa. *Pluk. alm. 321.*
Nureni-kelengu. Rheed. mal. 7. p. 67. t. 35.
Habitat in India.
- triphylla.* 2. *DIOSCOREA* foliis ternatis. *Hort. cliff. 459.*
Tsiageri-nuren. Rheed. mal. 7. p. 63. t. 33.
Habitat in Malabaræa.
Præcedenti maxime affinis, an satis distincta?

3. DIO-

3. **DIOSCOREA** foliis cordatis, caule aculeato *Bulbifer-aculeata*.
Hort. cliff. 459.
Dioscorea ind. orient. folio tamni longiore, floribus spicatis, spicis plurimis ex uno pedunculo exeuntibus, scapo eorum medio geniculato. *Amm. herb.* 257.
Kattu-kalengu. Rheed. mal. 7. p. 71. t. 37.
Cara. Francisc. gart. 1. 25. f. 8. 8
Habitat in Malabaria.
4. **DIOSCOREA** foliis cordatis, caule alato bulbifero. *alata*.
Fl. zeyl. 360. *
Volubilis rubra, caule membranulis exstantibus alato, folio cordato nervoso. *Sloan. jam.* 46. *bist.* 1. p. 139.
Katfil-kekengu. Rheed. mal. 7. p. 71. t. 38. *Raf. suppl.* 134.
Habitat in Indiis.
5. **DIOSCOREA** foliis cordatis, caule lævi bulbifero. *bulbifera*.
Hort. cliff. 459. *Fl. zeyl.* 359. *
Rhizophora zeylanica, scammonii folio singulari, radice rotunda. *Herm. par.* 217. t. 217.
Rhizophora indica, bryoniæ nigræ similis, ad foliorum ortum verrucosa. *Pluk. alm.* 321. t. 220. f. 6.
Katu-katfil. Rheed. mal. 7. p. 69. t. 36.
Habitat in Indiis.
6. **DIOSCOREA** foliis cordatis alternis, caule lævi. *sativa*.
Hort. cliff. 459. t. 28. *Fl. zeyl.* 358. *Roy. lugdb.* 527.
Dioscorea scandens, foliis tamni, fructu racemoso. *Plum. gen.* 9.
Volubilis nigra, folio cordato nervoso. *Sloan. jam.* 46. *bist.* 1. p. 140.
Mu-kelengu. Rheed. mal. 8. p. 97. t. 51.
Habitat in Indiis. 2
7. **DIOSCOREA** foliis cordatis alternis oppositisque, *villosa*.
caule lævi.
Dioscorea foliis cordatis acuminatis: nervis lateralibus ad medium folii terminatis. *Groz. virg.* 121.
Bryoniæ similis floridana, muscotis floribus quernis, foliis subtus lanugine villosis: medio nervo in spinulam abeunte. *Pluk. alm.* 46. t. 375. f. 5.
Habitat in Virginia, Florida.
8. **DIOSCOREA** foliis oppositis ovatis acuminatis. *Fl. oppositifolia*.
zeyl. 361. *

Inhame maderas. foliis binis pulchre venosis. *Presgaz.*
50. t. 31. f. 6.
Habitat in India. 2

OCTANDRIA.

POPULUS.

- 786
- alba.* 1. **POPULUS** foliis subrotundis dentato-angulatis subtus tomentosis. *Hort. cliff.* 460. *Fl. suec.* 820. *Roy. lugdb.* 82. *Hall. belv.* 156. *Gmel. sib.* 1. p. 151. *Dalib. parif.* 302.
Populus alba, majoribus foliis. *Bauh. pin.* 429.
Populus alba. *Dod. pempt.* 835.
Habitat in Europa temperatiori. 5
- tremula.* 1. **POPULUS** foliis subrotundis dentato-angulatis utrinque glabris. *Hort. cliff.* 460. *Fl. suec.* 819. *Roy. lugdb.* 82. *Hall. belv.* 156. *Gmel. sib.* 1. p. 151. *Dalib. parif.* 301.
Populus tremula. *Bauh. pin.* 429.
Populus lybica. *Dod. pempt.* 836.
Habitat in Europæ frigidioribus. 5
Folia involuta basi internæ glandula duplici. 5
- nigra.* 3. **POPULUS** foliis deltoideis acuminatis serratis. *Hort. cliff.* 460. *Fl. suec.* 821. *Mat. med.* 462. *Roy. lugdb.* 82. *Gron. virg.* 194. *Hall. belv.* 156. *Gmel. sib.* 1. p. 151. *Dalib. parif.* 301.
Populus nigra. *Bauh. pin.* 429. *Dod. pempt.* 836.
Habitat in Europa temperatiore. 5
Folia involuta absque glandulis ad basin, hinc diversa a precedenti.
- balsamifera.* 4. **POPULUS** foliis subcordatis oblongis crenatis. *Hort. cliff.* 460. *Roy. lugdb.* 82.
Populus foliis cordatis crenatis basi nudis, petiolis teretibus. *Wach. altr.* 294.
Populus nigra, folio maximo, gemmis balsamum odoratissimum fundentibus. *Catesb. car.* 1. p. 34. t. 34.
Populus foliis ovatis acutis serratis. *Gmel. sib.* 1. p. 152. t. 33.
Habitat in America septentrionali. 5
- betamophylla.* 5. **POPULUS** foliis cordatis subrotundisque: primoribus villosis.

Po-

DIOECIA ENNEANDRIA. 1035

Populus foliis cordatis obsolete serratis: infimis serraturis glandulosis, petiolis lateraliter utrinque planis. Wacb. utr. 294?

Populus magna, foliis amplis: aliis cordiformibus; aliis subrotundis: primoribus tomentosis. Gron. virg. 194. Habitat in Virginia. 5

RHODIOLA. 487

1. RHODIOLA. *Fl. lapp. 378. Fl. succ. 831. Mat. med. Rosen. 477. Hort. cliff. 470. Roy. lugdb. 457.*

Rhodia radix. Bauh. pin. 286. Clus. hist. 2. p. 65. Raj. hist. 690.

Telephium luteum minus, radice rosam redolente. Moris. hist. 3. p. 468. f. 12. t. 10. f. 8.

Habitat in Alpibus Laponiæ, Austriæ, Helvetiæ, Britanniæ. 2

ENNEANDRIA. 988

MERCURIALIS.

1. MERCURIALIS caule simplicissimo, foliis scabris. *perennis. Hort. cliff. 461. Fl. succ. 823. Roy. lugdb. 203. Dalib. paris. 302*

♀ *Mercurialis montana testiculata. Bauh. pin. 122.*

Cynocrambe mas. Cam. epit. 999.

♂ *Mercurialis montana spicata. Bauh. pin. 122.*

Cynocrambe femina. Cam. epit. 999.

Habitat in Europæ nemoribus. 2

2. MERCURIALIS caule brachiato, foliis glabris. *Hort. annua. cliff. 461. Hort. upf. 298. Mat. med. 464. Roy. lugdb. 263. Dalib. paris. 302.*

Mercurialis testiculata f. mas. Bauh. pin. 121. 2

Mercurialis mas. Dod. pempt. 658. 2

♂ *Mercurialis spicata f. femina, Dod. pempt. 658*

Habitat in Europæ temperate umiosis. 3

3. MERCURIALIS caule subfruticoso, foliis tomentosis. *Hort. upf. 461. Roy. lugdh. 203. Saxo. monsp. 128.*

Phyllon testiculatum. Bauh. pin. 122. 2

Phyllum marificum. Clus. hist. 2. p. 48. 2

♂ *Phyllon spicatum Bauh. pin. 122.*

Phyl-

Phyllon feminificum. *Clas. hist.* 2. p. 48.

Habitat in G. Narbonensi, Hispania. *h*

procumbens. 4. MERCURIALIS androgyna. *Virid. clif.* 98. *Roy. lugdb.* 203.

Mercurialis procumbens dicoctos africana, foliis violæ tricoloris. *Herm. app.* 10.

Ricinocarpus afra. *Boerb. lugdb.* 1. p. 254.

Habitat in Africa.

989

HYDROCHARIS.

Morsus ranae. 1. HYDROCHARIS. *Hort. clif.* 460. *Fl. succ.* 822.

Roy. lugdb. 10. *Hall. belv.* 301. *Dalib. paris.* 303.

Nymphæa alba minima. *Bauh. pin.* 193.

Rana morsus. *Dod. pempt.* 583. f. 2. *3*; 1. *2*

β. Morsus rana: flore pleno odoratissimo. *Raj. cens.* 101. *3*

2. Nymphæa alba minor. *Bauh. pin.* 193.

Habitat in Europæ fossis limosis. *☉* *2*

Gemmisera f. bulbisera est.

990

DECANDRIA.

CARICA.

Papaya. 1. CARICA foliorum lobis sinuatis. *Hort. clif.* 461.

Fl. zeyl. 365. *Roy. lugdb.* 225.

Arbor platani folio, fructu peponis magnitudine eduli. *Bauh. pin.* 131.

Ficus arbor utriusque indiæ, platani foliis, monopeteches. *Pluk. alm.* 145. t. 278. f. 1.

Papaya-maram. *Rheed. mal.* 1. p. 23. t. 15. f. 1. *2*

3 Ambapaya. *Rheed. mal.* 1. p. 21. t. 15. f. 2.

Papaya. *Ebret. t.* 3. f. 1.

Habitat in Indiis. *h*

Posoposa. 2. CARICA foliorum lobis integris. *Hort. clif.* 462. *2*

Roy. lugdb. 225.

Papaya ramosa, fructu pyriformi. *Few. peruv.* 2. p. 52. t. 39. f. 1.

Ficus arbor papaje sylvestris nomine missa. *Pluk. alm.* 146. t. 278. f. 2^p

Platani folio arbor Posoposo philippensis. *Pet. gaz.* 68. t. 43. f. 2^p

Habitat Surinami. *h*

KIG-

DIOECIA DECANDRIA.

1037

KIGGELARIA. 1001 751

1. KIGGELARIA. *Hort. cliff.* 462. t. 29. * *Roy. lugdb. africana.* 478.

Laurus non odorata, fructu globofo, africana. Sterb citr. 246. t. 12. f. B. C. D.

Evonymo affinis æthiopica sempervirens, fructu globofo scabro, foliis falicis rigidis ferratis. Herm. lugdb. 139. *Pluk. alm.* 139. t. 176. f. 3.

Habitat in Æthiopia. b

CORIARIA. 1002 752

1. CORIARIA foliis ovato-oblongis. *Hort. apf.* 299. *myrtifolia. Sauv. monsp.* 151.

Coriaria. Hort. cliff. 462. *Roy. lugdb.* 222.

Rhus myrtifolia monspeliaca. Bauh. pin. 414. *Angl. hortul.* 64. t. 20. f. 1.

Rhus plinii myrtifolia monspeliensium. Lob. ic. 2. p. 98.

Habitat Monspeli b

2. CORIARIA foliis cordato-ovatis sessilibus.

Coriaria ruscifolia, vulgo Deu. Few. peruv. 3. p. 17. t. 12.

ruscifolia.

Habitat in Peru. b

DATISCA. 1003 753

1. DATISCA caule lævi.

Cannabina.

Cannabis foliis pinnatis. Hort. cliff. 457. *Roy. lugdb.* 221.

Cannabis lutea fertilis. Alp. exot. 300. t. 298. *Morif. bist.* 3. p. 432. f. 8. t. 25. f. 4.

Luteola herba sterilis. Bauh. pin. 100.

Cannabis lutea cretica. Alp. exot. 296. t. 295.

Cannabis lutea sterilis. Alp. exot. 301. t. 300.

Luteola herba foliis canuabinis. Bauh. pin. 100.

Habitat in Creta. 2

2. DATISCA caule hirsuto.

hirta.

Habitat in Philadelphia. Kalm.

Hæc species major; Caulis exstantibus pilis undique hirsutus. Folia pinnata, precedentis quodammodo similia, at Foliola majora, magis alterna & magis basi decurrentia ac confluentia.

PO-

POLYANDRIA.

CLIFFORTIA.

- ilicifolia.* 3. CLIFFORTIA foliis subcordatis dentatis.
 Cliffortia foliis dentatis. *Hort. cliff.* 463. t. 30. *Roy. lugdb.* 252.
 Arbuscula atra, folio acuto ilicis caulem amplexo rigido. *Dill. elrh.* 36. t. 31. f. 35.
Habitat in Æthiopia. ♀
- versifolia.* 2. CLIFFORTIA foliis lanceolatis integerrimis. *Hort. cliff.* 463. t. 31. *Roy. lugdb.* 252.
 Frutex æthiopicus conifer, fructu parvo sparsim intra folia rusci, seminibus cylindræis. *Pluk. alm.* 159. t. 297. f. 2.
Habitat in Æthiopia. ♀
- polygonifolia.* 3. CLIFFORTIA foliis linearibus pilosis. *Hort. cliff.* 501. t. 32
Habitat in Æthiopia. ♀
- trifoliata.* 4. CLIFFORTIA foliis ternatis: Intermedio tridentato. *Roy. lugdb.* 253.
 Myrica foliis ternatis: intermediis cuneiformibus tridentatis. *Hort. cliff.* 456.
 Thymelææ? affinis æthiopica, foliis tridentatis & ex omni parte hirsutæ pubescentibus. *Pluk. alm.* 367. t. 317. f. 4.
 Arbuscula atra, foliolis trifoliatis sine pedunculo ad caulem natis, semine papposo. *Boerb. lugdb.* 2. p. 262.
Habitat in Æthiopia. ♀

MONADELPHIA.

JUNIPERUS.

- Oxycedrus.* 1. JUNIPERUS foliis quaternis patentibus subulatis mucronatis. *Sauv. ad. monsp.* 1743. p. 52.
 Juniperus foliis quaternis patentibus subulatis pungentibus. *Roy. lugdb.* 90.
 Juniperus foliis ternis patentibus subulatis pungentibus trinerviis. *Sauv. monsp.* 169.
 Juniperus major, bacca rufescente. *Barb. pin.* 489.
Oxycedrus. Clus. hist. 1. p. 39.
Habitat in Hispania, G. Narbonensi. ♀

2. JU-

2. JUNIPERUS foliis quadrifariam imbricatis acutis *tinvisera*.
Roy. lugdb. 90.
Cedrus hispanica procerior, fructu maximo nigro
Tournef. inst. 588.
Habitat in Hispania. ♀
3. JUNIPERUS foliis undique imbricatis ovatis obtusis. *Roy. lugdb.* 90. *Mat. med.* 466. *Sauv. mousp.* 169. *Gmel. sib.* 1. p. 182.
Cedrus, folio cupressi, media, majoribus baccis. *Bauh. pin.* 488.
Cedrus phœnicea altera plinii & theophrasti. *Lob. ic.* 221.
Habitat in Gallia, Sibiria, ♀
4. JUNIPERUS foliis omnibus quadrifariam imbricatis: junioribus ovatis, senioribus acutis. *Roy. lugdb.* 90.
Juniperus barbadensis, cupressi foliis, ramulis quadratis.
Pluk. alen. 201. t. 197. f. 4.
Juniperus bermudiana. *Hort. angl.* 42. t. 1. f. 1. ex figura.
Habitat in America. ♀
5. JUNIPERUS foliis inferioribus ternis: superioribus *bermudiana*, binis decurrentibus subulatis patulis acutis. *Roy. lugdb.* 90.
Juniperus bermudiana. *Herm. lugdb.* 345. t. 347.
Habitat in America. ♀
6. JUNIPERUS foliis oppositis erectis decurrentibus: *Sabina*, oppositionibus pyxidatis.
Juniperus foliis interne adnatis: oppositionibus concatenatis. *Hort. cliff.* 464. *Hort. xps.* 299. *Mat. med.* 467. *Roy. lugdb.* 90. *Gmel. sib.* 1. p. 183.
Sabina folio cupressi. *Bauh. pin.* 487.
Sabina. *Dod. pempt.* 854.
8. *Sabina* folio tamarisci dioscoridis. *Bauh. pin.* 487.
Habitat in Italia, Sibiria, Olympo, Ararat, Lusitania. ♀
7. JUNIPERUS foliis ternis basi adnatis: junioribus *virginiana*, imbricatis, senioribus patulis. *Hort. cliff.* 464. *Roy. lugdb.* 90. *Gron. virg.* 194.
Juniperus major americana. *Raj. hist.* 1413. 1414.
Juniperus maxima, cupressi folio minimo, cortice exteriore in tenues philyras spirales ductili. *Sloax. jam.* 128. *hist.* 2. p. 2. t. 157. f. 3. *Raj. dendr.* 12.
Habitat in Virginia, Carolina. ♀

8. JU.

- communis.* 8. JUNIPERUS foliis ternis patentibus mucronatis.
 Juniperus foliis patentibus. *Hort. cliff.* 465. *Fl. suec.* 824.
Mat. med. 465. *Roy. lugdb.* 89. *Hall. belv.* 146.
Gmel. fib. 1. p. 182. *Dalib. paris.* 303. binerviis. *Sauv.*
monsp. 169.
 Juniperus vulgaris fruticosa. *Bauh. pin.* 488.
 B. Juniperus vulgaris arbor, *Bauh. pin.* 488.
 C. Juniperus minor montana, folio latiore, fructu longiore. *Bauh. pin.* 489.
Habitat in Europæ septentrionalis sylvis. h
- phœnicea.* 9. JUNIPERUS foliis ternis oblitteratis imbricatis obtusis.
 Cedrus, foliis cupressi, major, fructu flavescente *Bauh.*
pin. 487.
 Juniperus major. *Clus. hist.* 1. f. 38.
Habitat in Europa australi, Monspelii. ‡

TAXUS.

- 996*
~~1006~~
taccato. 1. TAXUS foliis approximatis.
 Taxus *Hort. cliff.* 464. *Fl. suec.* 825. *Roy. lugdb.* 87.
Hall. belv. 146. *Dalib. paris.* 303.
 Taxus. *Bauh. pin.* 505. *Cam. epit.* 840. *Dod pempt.*
 859.
Habitat in Europa, Canada. h
- nucifera.* 2. TAXUS foliis distantibus. †
 Taxus nucifera. *Kempf. amœn.* 814. t. 815.
Habitat in Japonia. h

EPHEDRA.

- 1007*
diffachya. 1. EPHEDRA pedunculis oppositis: amentis geminis.
Hort. cliff. 465.
 Polygonum bacciferum maritimum minus. *Bauh. pin.*
 15.
 Triagum. *Cam. hort.* 171. t. 46.
Habitat in G. Narbonensis, Hispaniæ saxosis collibus
marinis. h
- monosperma* 2. EPHEDRA pedunculis pluribus: amentis solitariis.
Gmel. fib. 1. p. 171. t. 38. f. 1.
 Ephedra minima, flagellis brevioribus & tenuioribus.
Amm. ruth. 254. t. 26.
 Ephedra monospermos. *Amm. ruth.* 255.
Habitat in Sibiria montibus apricis, sterilissimis. h

STN-

SYNGENESIA.

RUSCUS. *1068 998*

1. RUSCUS foliis supra floriferis nudis. *Hort. cliff. aculeatus.*
463. Hort. upf. 300. Mat. med. 768. Dalib. parif.
304. Roy. lugdb. 227.
Ruscus myrtioli aculeatus. Tournef. inst. 79.
Ruscus. Baub. pin. 470.
Ruscum. Dod. pempt. 474.
Habitat in Galliæ, Italiæ nemorosis asperis. 3
2. RUSCUS foliis subtus floriferis nudis. *Hort. cliff. Hypophyl-*
465. Hort. upf. 300. Roy. lugdb. 227. lum.
Laurus alexandrina, fructu folio insidente. Baub. pin.
305.
Laurus alexandrina, Chamædaphne. Col. ecphr. 1. p.
164. t. 165. f. 1.
3. RUSCUS latifolius, fructu ex medio foliorum extra pen-
dente. Dill. elth. 333. t. 251. f. 303.
Habitat in Italia ad latera collium. 2
3. RUSCUS foliis subtus floriferis sub foliolo. *Hort. Hypoglossum*
cliff. 466. Hort. upf. 300. Mat. med. 469. Roy. lugdb.
227.
Laurus alexandrina, fructu pediculo insidente. Baub.
pin. 304.
Hypoglossum dioscoridis; Laurotaxa plinii. Col. ecphr.
1. p. 166. t. 165. f. 2.
Habitat in Hungariæ, Liguriæ, Italiæ montibus um-
brofis. 2
4. RUSCUS foliis margine floriferis. *Hort. cliff. 464. Androgynus*
Ruscus latifolius e foliorum tinu florifer & baccifer.
Dill. elth. 332. t. 250. f. 332.
Habitat in Canariis. 3
5. RUSCUS racemo terminali hermaphroditico. *Hort. racemosus.*
cliff. 496. Roy. lugdb. 227.
Laurus alexandrina ramosa, fructu e summitate cauli-
um prodennte. Herm. lugdb. 360. 379. t. 681.
Laurus alexandrina, fructu longis pediculis caulibus
allegato. Moris. bles. 279.
Habitat - - -
Calyx & Corolla coalita in unum corpus carnosum,
connuentia.

GYNANDRIA.

99099

CLUTIA.

- alaternoides* 1. **CLUTIA** foliis sessilibus lanceolatis. *Hort. cliff.* 509.
Roy. lugdb. 203.
 Croton foliis linearilanceolatis. *Hort. cliff.* 444. 8
 Chamælea foliis oblongis. *Burm. afr.* 116. t. 43. f. 1.
 Chamælea foliis latis oblongis, floribus ex alis in ipi-
 cam erectis. *Burm. afric.* 118. t. 43. f. 3. 8.
Alaternoides africana, telephii legitimi imperati folio.
Comm. hort. 2. p. 3. t. 9.
Habitat in Æthiopia. 5
- pulchella*, 2. **CLUTIA** foliis ovatis integerrimis, floribus laterali-
 bus.
Clutia foliis petiolatis. *Hort. cliff.* 500. 431. *Roy. lugdb.*
 202.
Frutex æthiopicus, portulacæ folio, flore ex albo vire-
 scente. *Comm. hort.* 1. p. 177. t. 91.
Habitat in Æthiopia. 5
- velutæ.* 3. **CLUTIA** foliis ovalibus retusis, floribus racemosis
 axillaribus. *Fl. zeyl.* 967.
 Corni f. Sorbi species. *Bont. jav.* 103.
Scherunam-cottani. *Rheed. mal.* 2. p. 23. t. 18. male.
Raj. hist. 1623.
Habitat in India. 5
- Eluteria*. 4. **CLUTIA** foliis cordato-lanceolatis. *Fl. zeyl.* 366.
Mat. med. 470.
Eluteria. *Hort. cliff.* 486.
Ricinus dulcis arborescens americanas, populnea fron-
 de argentea. *Pluk. alm.* 321. t. 220. f. 5. *Seb. thes.*
 1. p. 56. t. 35. f. 3.
Habitat in Indiis. 5
- Castarilla*. 5. **CLUTIA** foliis lanceolatis acutis integerrimis petio-
 latis. †
Ricinoides æleagni folio. *Catesb. car.* 2. p. 46. t. 46.
Habitat in Carolina. 5

Clas-

Classis XXIII.

POLYGAMIA.

MONOECIA. 1000

MUSA.

1. MUSA spadice nutante. *paradisac.*
 Musa racemo simplicissimo. *Hort. cliff. 467. Hort. ups.*
 301. *Fl. zeyl. 368. Roy. lugdb. 10.*
 Musa cliffortiana. *Linn. mus. 1 t. 1.*
 Musa. *Clas. exot. 229. Sterb. citr. t. 4.*
 Ficus indica, fructu racemoso, folio oblongo. *Baub.*
pin. 508.
 Palma humilis, longis latisque foliis. *Baub. pin. 107.*
Habitat in India. &

2. MUSA spadice erecto. *Bihai.*
 Bihai amplissimis foliis, florum vasculis variegatis.
Plum. gen. 50.
 Bihai. *Ovid. l. 8. c. 9.*
 3. Bihai amplissimis foliis, florum vasculis coccineis.
Plum. gen. 50
 7. Bihai amplissimis foliis, florum vasculis subnigris. *Plum.*
gen. 50.
Habitat in America calidiore.

OPHIOXYLON. 1001

1. OPHIOXYLON. *Fl. zeyl. p. 239.* *serpentinum*
 Ophioxylon foliis quaternis. *Fl. zeyl. 398. * Mas. med.*
 474.
 Ligustrum foliis ad singula genicula ternis. *Burm. zeyl.*
 141. t. 64.
 Clematis indica, persicæ foliis, fructu periclymeni. *Baub.*
pin. 304.
Habitat in Zeylona. &

CELTIS. 1002

1. CELTIS foliis ovato-lanceolatis. *Hort. cliff. 39. Roy. australis.*
lugdb. 207. Dalib. parif. 304.
 Lotus fructu cerasi. *Baub. pin. 447.*
 Lotus f. Celtis. *Cam. epit. 155.*

U u u 1

Lotus

Lotus arbor. *Lob. ic.* 186.

Habitat in Europa australi & Africa citeriore. ♀

orientalis. 2. **GELTIS** foliis oblique cordatis serratis: subtus villosis. *Fl. zeyl.* 369. *

Ulmus fructu baccato. *Hort. cliff.* 83.

Salvifolia arbor orientalis, foliis tenuissime crenatis.

Pluk. alm. 329. t. 221. f. 4.

Mallam-toddali. *Rheed. mal.* 4. p. 83. t. 40.

Habitat in Indiis. ♀

occidentalis. 3. **CELTIS** foliis oblique ovatis serratis acuminatis.

Celtis procera, foliis ovato-lanceolatis serratis, fructu pullo. *Gron. virg.* 195.

Lotus arbor virginiana, fructu rubro. *Raj. hist.* 1917.

Habitat in Virginia. ♀

Folia tenera, ovato-lanceolata, parum pubescentia; adulta lato-ovata, acuminata, acumine & basi integerrima, ceterum serrata, nuda, nervoso-venosa, latere postico duplo minore. Maxime affinis prima.

VERATRUM.

album.

1. **VERATRUM** racemo supradecomposito, corollis erectis.

Veratrum caule ramoso. *Mat. med.* 471.

Veratrum pedunculis corolla erecta patente brevioribus.

Gmel. fib. 1. p. 75.

Helleborus albus, flore subviridi. *Bauh. pin.* 186.

Elleborus albus, exalbido flore. *Clus. hist.* 1. p. 274.

Habitat in Russiæ, Sibiricæ, Austriæ, Helvetiæ, Italiæ,

Græciæ montosis. &

ulgerum. 2. **VERATRUM** racemo composito, corollis patentissimis.

Veratrum pedunculis corolla patentissima longioribus.

Gmel. fib. 1. p. 76.

Helleborus albus, flore atro rubente. *Bauh. pin.* 186.

Morisi. hist. 3. p. 485. f. 12. t. 4. f. 1.

Habitat in Hungariæ, Sibiricæ apricis siccis. &

Præcedenti maxime affinis; diversa loco, colore, pedunculis villosis, corolla patente nec erecta, racemo composito, nec subdiviso in paniculam.

interum.

3. **VERATRUM** racemo simplicissimo, foliis sessilibus. Veratrum caule simplicissimo. *Gron. virg.* 195. *Gen. nov.* p. 16. f. 2.

Reseda

Refeda foliis lanceolatis, caule simplicissimo. *Gron. virg.* 59.

Habitat in Virginia, Canada.

Radix bulbosa. Folia radicalia plura, lato-lanceolata, nervosa, firma, glabra. Caulis pedalis, simplicissimus. Folia caulina nonnulla, inaequaliter posita, angusta, lanceolata, sessilia, nec ut in reliquis vaginantia, parva. Racemus terminalis, simplicissimus, primum ovatus, deum longitudine caulis, densus, flosculis parvis, flavis.

ANDROPOGON. 1004

1. ANDROPOGON spica solitaria, floribus inferiori-
bus muticis. contortam.

Aegilops maderaspatana, glumis pilosis aristatis. *P. Schench. gram.* 92.

Gramen secalinum indicum, spica gracili tomentosa: longioribus aristis ad se invicem intortis. *Pluk. aim.* 173. t. 191. f. 5. *Morif. hist.* 3. p. 180. f. 2. t. 4. f. ult.

Habitat in India.

2. ANDROPOGON spica oblonga, floribus lanatis re-
motis divaricatis: arista flexuosa nuda. *divaricatum*

Lagurus humilior, panicula conica laxa nutante cul-
mum terminante. *Gron. virg.* 135.

Habitat in Virginia.

3. ANDROPOGON panicula nutante, aristis tortuosis
laevibus, glumis calycinis hirsutis. *nutans*

*Andropogon folio superiore spathaceo, pedunculis late-
ralibus oppositis unifloris: aristis flexuosis. Gron. virg.*
133.

Gramen avenaceum, panicula minus sparsa, glumis al-
ba sericea lanugine obductis. *Sloan. jam.* 35. *hist.* 1.
p. 43. t. 14. f. 2.

Habitat in Virginia, Jamaica.

4. ANDROPOGON panicula laxa, aristis tortuosis. *alopecuroides*
Andropogon culmo paniculato. Gron. virg. 133.

Gramen *Dactylon alopecuroidis* facie, panicula longis-
sima e spicis plurimis tomentosis constante. *Sloan,
jam.* 3. *hist.* 1. p. 113. t. 14. f. 70. f. 1.

Habitat in America septentrionaliore.

- distachyon.* 5. ANDROPOGON spicis binis terminalibus, culmo indiviso.
Habitat in Helvetia. Burser.
 Culmi plus quam pedales. Folia plana. Rami nulli. Spicæ duæ, æquales, oblongæ, erectiusculæ, culmum terminantes: singulæ ex flosculis viginti pluribus. Pedicelli arcti, brevissimi, planiusculi, basi pilosi, superne latiores. Flores solitarii. Calyces uniflori: gluma exterior flore fere longior latiorque apice bifida: aristâ longa, articulata, patente. Gluma calycina interior minima.
- Schænanthus.* 6. ANDROPOGON paniculæ spicis conjugatis ovatis.
Fl. zeyl. 465. (sub Laguro) Mat. med. 31.
Juncus odoratus aromaticus. Baub. pin. 11. theatr. 163.
 Gramen ad junceum accedens aromaticum majus syriacum. *Morif. hist. 3. p. 229. f. 8. t. 9. f. 25.*
 Gramen Dactylon aromaticum, multiplici panicula, spicis brevibus tomento candicantibus ex eodem pediculo binis, *Pluk. alm. 175. t. 190. f. 1.*
Habitat in India, Arabia.
- virgineum.* 7. ANDROPOGON paniculæ spicis conjugatis, pedunculis medio villosis, spicis lana brevioribus.
*Roy. lugdb. 53 * Gron. virg. 132.*
Habitat in America.
- bicorne.* 8. ANDROPOGON paniculæ spicis conjugatis, flosculis basi lanatis folio spataneo obvolutis.
Lagurus spicis oblongis pedunculatis e singula ala pluribus. Hort. cliff. 25. Gron. virg. 135.
 Gramen Dactylon bicornè tomentosum maximum, spicis numerosis. *Sloan. hist. 1. p. 42.*
Habitat in Brasilia, Jamaica.
- hirtum.* 9. ANDROPOGON paniculæ spicis conjugatis, calycibus hirsutis. *Roy. lugdb. 53. Gron. virg. 133.*
Festuca junceo folio, spica gemina. Baub. pin. 9. prodr. 19. theatr. 145.
 Gramen Dactylon spica gemina. *T. Scheuchb. gram. 95.*
 Gramen Dactylon siculum, multiplici panicula, spicis ab eodem exortu geminis. *Pluk. alm. 175. t. 92. f. 1.*
Habitat in Lusitania, Sicilia, Smyrniæ.
- Nardus.* 10. ANDROPOGON paniculæ ramis supradecompositis proliferis. *Fl. zeyl. 45.* (sub Laguro) Mat. med. 32.*
Calamus odoratus mathioli. Baub. pin. 17. theatr. 263.
Habitat in India.

1. **ANDROPOGON** spicis digitatis plurimis, flosculis sessilibus: aristato muticoque, pedicellis lanatis. *Ischaemum*.
Ischaemum panicula digitata, spiculis oppositis disjunctis-
 ve, florum serie duplici, culmo tereti. *Guett. stamp.*
 187. *Dalib. parif.* 288.
Gramen dactylon spicis villosis. *Bauh. pin.* 8. *theatr.*
 116. *Schenck. gram.* 94.
Gramen Dactylon spicis aristatis geniculatis. *Barr. ic.*
 753. f. 2
Habitat in Europa austriacis aridis.
Flosculi in singula spica dense gemini: altero aristato,
altero mutico muticula.
2. **ANDROPOGON** spicis digitatis plurimis erectiusculis articu-
 latis levibus: flosculis utrinque aristatis. *susculum*
Gramen Dactylon indicum, spicis villosis subrubescen-
 tibus. *Morif. hist.* 3. p. 185. f. 8. t. 3. f. 15.
Gramen Dactylon elatius, spicis plurimis tomentosis.
Sloan. jam. 33. *hist.* 1. p. 111. t. 65. f. 2?
Habitat in Indiis
Spica in meis specimenibus glabra sunt.

HOLCUS. 1005

1. **HOLCUS** glumis villosis, seminibus aristatis. *Hort. Sorghum*.
upf. 301.
Holcus glumis glabris. *Hort. cliff.* 468.
Milium arundinaceum, subrotundo semine, *Sorgho* no-
 minatum. *Bauh. pin.* 26. *Morif. hist.* 3. p. 196. f.
 8. t. 5. f. 7.
Sorghi. *Bauh. hist.* 2. p. 447.
Habitat in India. ☉
Arista inferne fusca, superne lavis albida.
2. **HOLCUS** glumis glabris, seminibus muticis. *Saccharatus.*
Milium indicum, arundinaceo caule, granis flavescen-
 tibus. *Hern. lugd.* 425. *
Milium indicum sacchariferum altissimum, semine fer-
 rugineo. *Breyn. prodr.* 2.
Frumentum indicum quod *Milium indicum* vocant.
Bauh. theatr. 489.
Habitat in India. ♂
Altitudo humana. Folia facie *Sacchari* linea longitudi-
 nali alba. Radices ex infimis canlis geniculis descen-
 dunt in terram. Flores masculi vix se aperiant.
3. **HOLCUS** glumis glabris, floribus hermaphroditis mu-
 ticis, semineo aristato. *balepensis.*

Gramen arundinaceum halepense, tragopoginis folio, panicula miliacea. *Pluk. alm.* 176. t. 32. f. 1.

Gramen paniculatum arundinaceum syriacum Holliann dictum. *Morif. bist.* 3. p. 201.

Habitat in Syria, Mauritania. &

Singulo flori femineo adstant flores duo masculi pedicellati.

lanatum.

4. HOLCUS glumis bifloris villosis: hermaphrodito mutico, masculo arista recurva.

Aira foliis pubescentibus, panicula contracta: flosculo hermaphrodito mutico, masculo arista uncinata, calyce brevior. *Fl. succ.* 67. *It. æl.* 122.

Aira flosculo masculo aristato, femineo mutico. *Hort. cliff.* 27. *Roy. lugdb.* 66.

Gramen pratense paniculatum molle. *Baub. pin.* 2. *prodr.* 3. *theatr.* 27. *Schench. gram.* 234.

Gramen lanatum. *Dalech. bist.* 425.

Habitat in Europæ pascuis arenosis. &

odoratus.

5. HOLCUS glumis trifloris muticis glabris: hermaphrodito diandro.

Hierocloe. *Gmel. fib.* 1. p. 101.

Aira foliis planis, floribus trifloris aristatis: flosculo intermedio mutico diandro. *Fl. succ.* 70.

Poa spiculis bifloris cordatis. *Fl. lapp.* 53.

Gramen paniculatum odoratum. *Baub. pin.* 7. *prodr.* 7. *Schench. gram.* 236.

Gramen maræ borussorum. *Laf. pruss.* 111. t. 26.

Habitat in Europæ frigidioris pascuis humentibus. &

Flosculi duo laterales masculi triandri: intermedius s. tertius hermaphroditus diandrus.

laxa.

6. HOLCUS glumis glabris bifloris muticis acuminatis, panicula filiformi debili.

Aira calycibus trivalvibus trifloris. *Gron. virg.* 136.

Habitat in Virginia, Canada.

Habitus *Aira ceruleæ*. Culmi bipedales, angusti, subnutantes. Folia plura, levia, margine scabra, ore piloso. Panicula filiformis, parum admodum ramosa, subnutans, debilis; pedunculi sepius biflori. Flosculo masculo ovato, crasso.

striatus.

7. HOLCUS glumis striatis bifloris muticis acuminatis, panicula conferta oblonga.

Aira panicula oblonga, floribus muticis: hermaphrodito masculoque, calycibus diphyllis. *Gron. virg.* 135.

Habi-

Habitat in Virginie paludibus.

Folia plana, longa, vaginis crassiusculis striatis. Panicula attenuata: Pedunculis brevissimis, confertis, ramosis. Glumæ oblongæ, striatæ, acuminatæ. Calyx monophyllus, minimus.

ISCHÆMUM. 1006

1. ISCHÆMUM seminibus muticis.

muticum.

Tagadi. Rheed. mal. 12. p. 91. t. 49.

Habitat in India. 2

Spica bipartita, imbricata: pedunculis alternis, simplicibus, bifloris, angulatis, culmo adpressis. Flos alter singulum pedunculum terminat Hermaphroditus ex Calyce biglumi unifloro; alter vero s. inferior Flos androgynus, basi exteriori unicuique pedunculo sessilis insertus, ex Calyce bivalvi, duriusculo, longitudine flosculorum, continens flosculos duos: alterum Masculinum; alterum Femininum. Pistillorum stigmata crassa, barbata. Semina mutica. Pedunculus spicæ foliis vestitus.

2. ISCHÆMUM seminibus aristatis.

aristatum.

Habitat in China. Osbeck.

Structura spicæ & florum omnino eadem cum precedente, at Culmus altior, Spica longior & Pedunculus spicæ nudus. Semina arista intorta flosculis longiore armata.

CENCHRUS. 1007

1. CENCHRUS panicula spicata, glumis muricatis setaceis ciliaribus. Guett. stamp. 2. p. 188. Dalib. paris. 305. Sauv. monsp. 40.

Gramen caninum maritimum, spica echinata. Bauh. pin.

2. Scheuchb. gram. 76.

Gramen caninum maritimum asperum. Bauh. prodr. 2. t. 2.

Gramen caninum maritimum spicatum, echinatis glumis. Barr. ic. 718.

Habitat in Europæ australioris maritimis.

2. CENCHRUS spica ovata simplici. Roy. lugdb. 71. capitatus. Sauv. monsp. 40.

Gramen spica subrotunda echinata. Bauh. pin. 7. Scheuchb. gram. 74.

U u u s

Gra-

- Gramen montanum echinatum tribuloides capitatum.
Col. eschr. 1. p. 340 t. 338. f. 1.
 Gramen minimum, spica globosa echinata. *Barr. rar.*
 1170. t. 28. f. 1. & t. 862. f. 2.
Habitat in G. Narbonensi, Italia
- echinatus.* 3. CENCHRUS spica oblonga conglomerata. *Roy.*
lugdb. 72.
 Panicastrella americana major annua, spica laxa purpureascente. *Mich. gen.* 36.
 Gramen locustis tumidioribus echinatis. *Scheuch. gram.*
 77.
 Gramen americanum, spica echinata, majoribus locustis.
Pluk. alm. 177. t. 92. f. 3.
 Gramen echinatum maximum, spica rubra f. alba. *Sloan. jam.* 30. *hist.* 1. p. 108. *Raj. suppl.* 602.
 Gramen americanum, spica echinata, majoribus glutinis.
T. Morisf. hist. 3. p. 195.
Habitat in Jamaica, Curassao. ☉
- tribuloides* 4. CENCHRUS glutinis femineis globosis muricato-spinosis hirsutis.
 Cenchrus capitulis spinosis tomentosis. *Gron. virg.* 122.
 Panicastrella americana minor annua, spica angusta densa albicante. *Mich. gen.* 37.
 Gramen maritimum echinatum procumbens, culmo longiori, spicis strigosioribus. *Sloan. jam.* 30. *hist.* 1. p. 108. t. 65. f. 1. *Raj. suppl.* 602.
 Gramen tribuloides spicatum maximum virginianum.
Pluk. alm. 177.
 Gramen echinatum spicatum, locustis crassioribus, tribuloides virginianum. *Morisf. hist.* 3. p. 195. f. 8. s. 3. f. 4.
Habitat in Virginie maritimis. ☉
- frutescens.* 5. CENCHRUS capitulis lateralibus sessilibus, foliis mucronatis, caule fruticoso.
 Arundo graminea aculeata. *Alp. exot.* 105. t. 104.
 Gramen orientale spicatum fruticosum spinosum, spicis echinatis in capitulum congestis. *Tournef. cor.* 39.
Habitat in America. ☉

ÆGILOPS.

- ovata.* 1. ÆGILOPS spica ovata aristis brevioribus. *Hort. ups.*
 301. *Dalib. parif.* 304. *Sauv. monsp.* 40.

Ægi-

POLYGAMIA MONOECIA, 1051

Ægilops. Roy. lugdb. 72. Dod. pempt. 73. Cam. epist. 928. f. 2.

Festuca altera, capitulis duris. Bauh. pin. 10. theatr. 151.

Gramen spicatum, durioribus & crassioribus locustis, spica brevi. Schenck. gram. 11.

Habitat in Europa australi. ♂

3. *ÆGILOPS* spica subulata scabra, flosculo ultimo caudata, tantum aristato.

Gramen creticum, spica gracili in duas aristas longissimas & asperas abeunte. Tournef. cor. 29.

Habitat in Creta.

Glumæ omnes, excepta ultima, bidentata sunt.

4. *ÆGILOPS* spica subulata aristas longiore. *Squarrosa.*

Habitat in Oriente. Tournefort.

5. *ÆGILOPS* spica subulata scabra, aristas longitudine triuncialis, spicæ.

Festuca altera, capitulis duris, spica triuncialis. Bauh. pin. 10.

Gramen spicatum, durioribus & crassioribus locustis, spica longissima. T. Schenck. gram. 12.

Habitat in Montpelii, Massiliæ, Smyrnæ aridis.

6. *ÆGILOPS* spica subulata mutica incurva. *incurva.*

Habitat in Oriente. Tournefort.

VALANTIA. 1059

1. *VALANTIA* floribus masculis trifidis hermaphroditi germi insidentibus. *Hort. ups. 302. Sauv. monsp. 162.*

- α *Valantia* procumbens glabra. *Hort. cliff 468. Roy. lugdb. 258.*

Valantia annua quadrifolia verticillata, floribus ex viridi-pallescentibus, fructu echinato. Mich. gen. 13. t. 7.

Rubeola echinata saxatilis. Bauh. pin. 334.

Cruciata muralis minima romana. Col. expbr. 1. p. 298. t. 297. Moris. bist. 3. p. 328. f. 9. t. 21. f. 2.

- β. *Valantia* erecta hispida. *Hort. cliff. 460. * Roy. lugdb. 258.*

Habitat in G. Narbonensis, Italiæ muris. ☉

2. *VALANTIA* floribus masculis trifidis hermaphroditis pedunculo insidentibus. *Hort. ups. 302. * Dalib. parif. 305. Sauv. monsp. 162. Apa-*

Aparine semine lævi. *Vaill. parif.* 18. t. 4. f. 3.
Habitat inter Germaniæ, Galliæ, Siciliæ segetes. ☉

- articulata.* 3. **VALANTIA** floribus masculis quadrifidis, pedunculis nudis. *Hort. upf.* 303. *. *Dalib. parif.* 305.
 Cruciata orientalis latifolia erecta glabra. *Tournef. cor.* 4.
Habitat in Ægypto, Syria. ☉

- Cruciata.* 4. **VALANTIA** floribus masculis quadrifidis, pedunculis diphyllis. *Hort. upf.* 303. *Sauv. monsp.* 163. *Dalib. parif.* 305.
 Galium foliis quaternis, flosculis in alis confertis. *Hort. cliff.* 34. *Hall. helv.* 462.
 Cruciata hirsuta. *Bauh. pin.* 335.
 Cruciata. *Dod. pempt.* 257.
Habitat in Germania, Helvetia, Gallia. 2

PARIETARIA.

- officinalis.* 1. **PARIETARIA** foliis lanceolato-ovatis alternis. *Hort. cliff.* 469. *Hort. upf.* 302. *Mat. med.* 472. *Roy. lugdb.* 210. *Dalib. parif.* 305. *Gort. gelr.* 554.
 Parietaria officinarum & dioscoridis. *Bauh. pin.* 121.
 Helxine. *Cam. epis.* 849.
Habitat in Europæ temperationis ruderatis. 2
- lusitanica.* 2. **PARIETARIA** foliis ovatis obtusis, caulibus filiformibus procumbentibus.
 Parietaria lusitanica annua minima. *Tournef. inst.* 509.
Raj. suppl. 129.
 Parietaria ficula, alfine folio. *Bocc. sic.* 47. t. 24.
Habitat in Lusitania, Hispania. ☉
- cretica.* 3. **PARIETARIA** foliis subovatis, involucris fructiferis quinquefidis compressis: laciniis lateralibus majoribus.
 Parietaria cretica minor, capsulis seminum alatis. *Tournef. cor.* 38.
Habitat in Creta.
- zeylanica.* 4. **PARIETARIA** foliis oppositis. *Fl. zeyl.* 371. *
Habitat in Zeylona.

ATRIPLEX.

- Halimius.* 1. **ATRIPLEX** caule fruticoso, foliis deltoideis integris. *Hort. cliff.* 469. *Gren. virg.* 195. *Roy. lugdb.* 218.
 Hali-

Halimus latifolius f. fruticosus, *Bauh. pin.* 120.

Halimus. Clus. hist. 1. p. 53.

Habitat in Hispaniæ, Lusitaniæ, Virginæ sepibus maritimis. ☉

2. ATRIPLEX caule fruticoso, foliis obovatis. *Fl. succ. Portulacoides.*
829.

Atriplex caule fruticoso, foliis lanceolatis obtusis. *Roy. lugdb.* 218.

Halimus f. *Portulaca marina. Bauh. pin.* 120.

Portulaca marina. Dod. pempt. 771.

Habitat in Oceani: Europæ septentrionalis littoribus. ☉

3. ATRIPLEX caule herbaceo, foliis deltoideis sinuatis undulatis alternis. *Hort. nps.* 303. * *tatarica.*
Habitat in Tataria. ☉

4. ATRIPLEX caule erecto herbaceo, foliis triangularibus. *Hort. cliff.* 469. *Hort. nps.* 303. *Mat. med.* 473. *hortensis.*
Roy. lugdb. 217.

Atriplex hortensis alba f. pallide virens. *Bauh. pin.* 119.

Atriplex hortensis. Dod. pempt. 615.

- β. *Atriplex hortensis rubra. Bauh. pin.* 119.
Habitat in Tataria. ☉

5. ATRIPLEX caule herbaceo, foliis deltoideo-lanceolatis obtuse-dentatis subtus farinaceis. *Hort. cliff.* 469. *Fl. succ.* 826. *Gron. virg.* 120. *Roy. lugdb.* 218. *laciniata.*

Atriplex maritima laciniata. Bauh. pin. 120.

Atriplex marina. Dod. pempt. 615.

Habitat ad Europæ & Virginæ septentrionalis littora marina. ☉

6. ATRIPLEX caule herbaceo: valvulis femineis magnis deltoideis sinuatis. *Fl. succ.* 361. *lt. æl.* 88. *basata.*
Dalib. parif. 305.

Atriplex sylvestris, folio hastato f. deltoide. *Morif. blef.* 237. *Raj. hist.* 192.

Atriplex sylvestris annua, folio deltoide triangulari sinuato & mucronato hastæ cuspidis simili. *Morif. hist.* 2. p. 607. f. 5. t. 32. f. 14.

Habitat in Europa frigidiori. ☉

7. ATRIPLEX caule herbaceo patulo, foliis subdeltoideo-lanceolatis, calycibus seminum disco dentatis. *patula.*
Atriplex caule herbaceo, valvulis femineis seminum disco dentatis. *Fl. succ.* 828. *Dalib. parif.* 306.

Atri-

Atriplex caule herbaceo, foliis superioribus lanceolatis integerrimis. *Hort. cliff.* 469. *Roy. lugdb.* 218.

Atriplex angusto oblongo folio. *Baub. pin.* 219.

Atriplex sylvestris, polygoni f. helxines foliis. *Lob. ic.* 257.

Habitat in Europæ cultis, ruderatis. ☉

- littoralis. 8. *ATRIPLEX* caule herbaceo erecto, foliis omnibus linearibus integerrimis. *Roy. lugdb.* 218. *It. scav.* 199.*
Atriplex angustissimo & longissimo folio. *Herm. lugdb.* 79.

Habitat in Europæ septentrionalis littoribus maris. ☉

- 16 1 2
frondosa. 1. *DALECHAMPIA.* *Hort. cliff.* 485.

Dalechampia scandens, lupuli foliis, fructu hispido tricocco. *Plum. gen.* 17.

Lupulus folio trifido, fructu tricocco hispido. *Plum. amer.* 89. t. 101.

Convolvulo-Tithymalus. *Boerb. lugdb.* 2. p. 268.*

Habitat in America meridionali.

Conf. *DALECHAMPIA* scandens, aristolochiæ foliis, fructu prurigineo. *Barr. aquin.* 47. *hujus an Plukenetia?*

Qui viderit vivam plantam utrum sufficienter distinguam ab Euphorbia determinet.

- 10 1 3
tataricum. 2. *ACER* foliis cordatis indivisis serratis: lobis obsoletis, floribus racemosis.

Acer foliis oblonge cordatis inæqualiter serratis. *Krausen; alt. petr.* 1749. p. 285. t. 13.

Habitat in Tataria. ♀

Frutex altus. Folia carpini, lobis vix manifestis. Flores racemosi, ut in sequente, sed petaloides.

- Pseudo-Platanus.* 2. *ACER* foliis quinquelobis inæqualiter serratis, floribus racemosis.

Acer foliis quinquelobis acutis obtuse serratis, petiolis canaliculatis. *Hort. ups.* 93. *Sæuv. monsp.* 212.

Acer foliis palmatis obtuse serratis, floribus subapetalis racemosis. *Hort. cliff.* 143. *Roy. lugdb.* 457. *Gort. gelr.* 78.

Acer montanum candidum. *Baub. pin.* 430.

Acer majus. *Dod. pempt.* 840.

Habitat in Helvetiæ, Austriæ montanis.

3. ACER foliis quinquelobis subdentatis subtus glaucis, ^{rubrum.}
pedunculis simplicissimis aggregatis.
Acer foliis quinquelobis acuminatis acute serratis, peti-
olis teretibus. *Hort. upf.* 94.
Acer foliis palmato-angulatis, floribus subapetalis sessilibus,
fructu pedunculato corymboso. *Gron. virg.* 41.
Cold. noveb. 85.
Acer virginianum, folio subtus incano, flosculis viridi
rubentibus. *Herm. par.* 1. t. 1.
Acer virginianum, folio majore: subtus argenteo supra
viridi splendente. *Pluk. alm.* 7. t. 2. f. 2. *Catesb. car.*
1. p. 62. t. 62.
Habitat in Virginia, Pennsylvania.
4. ACER foliis quinquepartito-palmatis acuminato-dentatis. ^{saccharinum.}
Habitat in Pennsylvania. Kalm. 5
Folia in hoc profundius & versus basin divisa, acumi-
nata apicibus & densibus, venula etiam & subtus
glauca, a precedente distinctissima. Flores mihi i-
gnosi.
5. ACER foliis quinquelobis acuminatis acute dentatis ^{Platanoides.}
glabris, floribus corymbosis. *Fl. succ.* 303.
Acer foliis palmatis acute dentatis, floribus corollatis
dichotomo-corymbosis. *Hort. cliff.* 143.
Acer montanum, tenuissimis & acutissimis foliis. *Bauh.*
pin. 431.
Acer major. *Cam. epit.* 63.
Acer montanum, orientalis platani foliis atrovirentibus.
Pluk. alm. 7. t. 252. f. 1.
Habitat in Europa boreali.
6. ACER foliis trilobis acuminatis serrulatis, floribus ^{pa-} ^{penfylvani-}
ceinosis. ^{cam.}
Habitat in Pennsylvania. Kalm.
Folia in tres lobos integros maxime acuminatos divisa,
margine tenuissime inaequaliter & argute serrata.
7. ACER foliis lobatis obtusis emarginatis. ^{campestra.}
Acer foliis tripartito-palmatis: laciniiis utrinque emargi-
natis obtusis cortice sulcato. *Hort. cliff.* 147. *Roy.*
lugdb. 459. *Gort. gelr.* 78.
Acer foliis semitrilobis, lobis obtuse incisis. *Hall. helv.*
212,
Acer campestre & minus. *Bauh. pin.* 431.

Acer

Acer minus. Dod. pempt. 840.

β. *Acer campestre & minus mas f. sterile. Vaill. paris. Habitat in Scania & australiori Europa.*

mon/pessula- 8. *ACER foliis trilobis integerrimis. Roy. lugdb. 459.*
num. *Acer trifolium. Bamb. pin. 431. Pluk. alm. 7. t. 251. f. 3.*

β. *Acer orientalis, hederæ folio. Tournef. cor. 43. Pock. orient. 191. t. 85.*

Acer cretica. Alp. exot. 9. t. 8.

Habitat Montpelii & in Creta. h

Negunda. 9. *ACER foliis compositis, floribus racemosis. Hort. cliff. 144. Gron. virg. 154. Roy. lugdb. 460.*

Acer maximum, foliis trifidis & quinquesfidis, virginianum. Pluk. alm. 7. t. 123. f. 45.

Habitat in Virginia. h

1074

BEGONIA.

obliqua.

1. BEGONIA.

Begonia nivea maxima, folio aurito. Plum. spec. 21.

Aceris fructu herba anomala, flore tetrapetalo albo. Sloan. jam. 83. hist. 1. p. 199. t. 127. f. 1. 2.

Begonia purpurea maxima, folio aurito. Plum. spec. 20.

β. *Begonia, roseo flore, folio aurito, minor & glabra. Plum. spec. 20.*

γ. *Begonia, roseo flore, folio aurito, minor & hirsuta. Plum. spec. 20.*

δ. *Begonia roseo flore, folio orbiculari. Plum. spec. 20.*

ε. *Begonia roseo flore, foliis acutioribus auritis & late crenatis. Plum. spec. 20.*

ζ. *Begonia hirsuta, flore albo, folio aurito, fructu crenato. Barr. aquin. 21.*

Habitat in America meridionali.

Folia cordata altero latere oblitterato.

DIOECIA.

GLEDITSIA.

eriazanthos. 1. *GLEDITSIA. Gron. virg. 183. Hort. upf. 298.*

Melilobus. Mich. gen. 15.

Cæsalpinoides foliis pinnatis ac duplicato-pinnatis. Hort. cliff. 489.

Aca-

Acacia americana, abruæ folio, triacanthos. *Pluk. mant.*

1. *Hort. angl.* t. 21.

β. *Acacia*, abruæ folio, triacanthos, capsula ovali unicum
semen claudente. *Catesb. car.* 1. p. 43. t. 43.

Habitat in Virginia. β

FRAXINUS. 1026

1. FRAXINUS foliolis serratis, floribus apetalis. *excelsior.*

Fraxinus floribus nudis. *Hort. cliff.* 469. *Fl. suec.* 830.

Mat. med. 475. *Roy. lugdb.* 396. *Dalib. paris.* 306.

Fraxinus excelsior. *Bauh. pin.* 416.

Fraxinus. *Dod. pempt.* 771.

Habitat in Europa septentrionalibus. β

2. FRAXINUS foliolis serratis, floribus corollatis. *Ornus.*

Fraxinus floribus completis. *Hort. ups.* 304. *Hort. cliff.*

476. *Roy. lugdb.* 396. *Mat. med.* 476.

Fraxinus humilior s. altera theophrasti, minore & tenuiore folio. *Bauh. pin.* 416.

Fraxinus floriteria botryoides. *Morif. prel.* 265. *Hort. angl.* 33. t. 9

Fraxinus tenuiore & minore folio. *Bauh. hist.* 1. p. 177.

Habitat in Europa australi. β

3. FRAXINUS foliolis integerrimis, petiolis teretibus. *americana.*

Gron. virg. 122. *Roy. lugdb.* 533.

Fraxinus carolinensis, foliis angustioribus utrinque acuminatis pendulis. *Catesb. car.* 1. p. 80. t. 80.

Habitat in Carolina Virginia. β

DIOSPYROS. 1077

1. DIOSPYROS foliorum paginis discoloribus. *Lotus.*

Diospyros foliis utrinque bicoloribus. *Roy. lugdb.* 441.

Diospyros foliis utrinque diverse coloratis. *Hort. cliff.* 149.

Lotus africana latifolia. *Bauh. pin.* 447

Pseudo-Lotus. *Cam. epit.* 156.

β. *Lotus africana angustifolia* s. femina. *Bauh. pin.* 447

Lotus africana altera. *Cam. epit.* 157. *bon.*

Habitat in G. Narbonensi, Italia, Mauritania. β

Confr. Si s. Kaki. *Kampf. amœn.* 85. t. 86.

2. DIOSPYROS foliorum paginis concoloribus. *virginiana.*

Diospyros foliis utrinque concoloribus. *Hort. cliff.*

149. *Roy. lugdb.* 441.

Diospyros floribus dioicis. *Gron. virg.* 156.

X x x

Gua-

Guajacana loco arbori affinis virginiana Pishamia dicta.

Pluk. alm. 180. t. 244. f. 5. *Raj. bist.* 1918.

Guajacana. *Catesb. car.* 2. p. 76. t. 76.

Loti africanæ similis indica. *Bauh. pin.* 448.

Habitas in America septentrionali. 3

1088
aquatica.

NYSSA.

1. NYSSA.

Nyssa foliis integerrimis. Hort. cliff. 462.

Nyssa pedunculis multifloris. Gron. virg. 121.

Arbor in aqua nascens, foliis latis acuminatis & dentatis, fructu elæagni majore. *Catesb. car.* 1. p. 60. t. 63.

Arbor in aqua nascens, foliis latis acuminatis & non dentatis, fructu elæagni minore. *Catesb. car.* 1. p. 41. t. 41.

2 Nyssa pedunculis unifloris. Gron. virg. 121.

Cynoxylon americanum, folio crassiusculo molli & tenaci. Pluk. alm. 127. t. 172. f. 6.

Habitas in America septentrionalis aquosus.

1089
æthiopickm.

ANTHOSPERMUM.

1. ANTHOSPERMUM. Hort. cliff. 455. t. 27.* Roy. lugdb. 208.

Frutex africanus ambaram spirans. *Pluk. alm.* 159. t. 183. f. 1. *Walsh. hors.* 24. t. 9.

Habitas in Æthiopia 3

1090
echinatus

ARCTOPUS.

1. ARCTOPUS. Hort. cliff. 495.*

Valerianoides cortusæ matthioli facie planta æthiopica, foliorum ad lacinias superna parte spicis stelliformiter echinata & ad oras pilis longioribus fimbriata. *Pluk. mant.* 185. t. 271. f. 5.

2 *Arctopus foliis superne spinis stelliformibus echinatis lacinatis & in cilia diductis, floribus umbellatis. Burm. afr.* 1. p. 1. t. 1.

Habitas in Æthiopia. 2.

21
ovinanefolium.

PANAX.

1. PANAX foliis ternis quinatis. Gron. virg. 147. Mat. med. 116.

Aureliana canadensis. *Last. gins.* 51. t. 1. *Catesb. car.* 3. p. 16. t. 16.

Arb-

Araliastrum, quinquifolii folio, majus Ninfin vocatum.

Vaill. sex. 43.

Habitat in Canada, Pennsylvania, Virginia. 2

Semina producant folium ternatum absque cotyledonibus
uti monocotyledones.

PANAX foliis ternis ternatis. *Gron. virg.* 35.

trifolium.

Araliastrum, fragariæ folio, minus. *Vaill. sex.* 43.

Nasturtium marianum, anemones sylvaticæ foliis, enneaphyllon, floribus exiguis. *Pluk. mant.* 135. t. 435.
f. 7.

Habitat in Virginia. 2

Precedenti maxime affinis & forte varietas, mari prioris facie & parvitate.

POLYOECIA.

FICUS.

1022

FICUS foliis palmatis. *Hort. cliff.* 471. *Hort. upf.* 305. *Corica.*

Mat. med. 478. *Amæn. acad.* 1. p. 24. *Roy. lugdb.* 211.

Ficus communis. *Baub. pin.* 457.

Ficus. *Dod. pempt.* 812.

Caprificus. *Baub. hist.* 1. p. 124.

Ficus humilis. *Baub. pin.* 457.

Habitat in Europa australi, Asia. 3

FICUS foliis cordatis subrotundis integerrimis. *Hort. Sycomorus*
cliff. 471. * *Roy. lugdb.* 211. *Amæn. acad.* 1. p. 26.

Ficus folio mori, fructum in caudice ferens. *Baub. pin.* 459

Moro-Ficus. *Baub. hist.* 1. p. 124. f. 1. 2.

SycoMorus, Ficus pharaonis. *Cam. math.* 103. f. 3.

Habitat in Ægypto. 3

Conser. Ficus indica, tiliz folio subtus albo & villosa,
polyrhizos. *Pluk. phys.* 178. f. 3.

FICUS foliis cordatis integerrimis acuminatis. *Hort. religiosa.*
cliff. 471. *Fl. zeyl.* 372. *Amæn. acad.* 1. p. 30.

Ficus malabariensis, folio cuspidato, fructu rotundo
paavo gemino. *Pluk. alm.* 144. t. 178. f. 2.

Arealu. Rhed mal. 1. p. 47. t. 27.

Habitat in India.

FICUS foliis ovatis integerrimis obtusis, caule infer- *benghalensis.*
ne radicato. *Hort. cliff.* 471. *Amæn. acad.* 1. p. 29.
Roy. lugdb. 212.

X x x 2

Ficus

Ficus benghalensis, folio subrotundo, fructu orbiculato.

Comm. hort. 1. p. 119. t. 62.

Peralu. Rheed. mal. 1. p. 49. t. 28.

Habitat in India. ♀

indica.

5. *FICUS* foliis lanceolatis petiolatis, pedunculis aggregatis, ramis radican-
tibus.

Ficus indica, foliis mali cotoneæ similibus, fructu sic-
bus simili. *Bauh. pin.* 457.

Ficus indica theophrasti. *Tabern. hist.* 1370. *Amæn. acad.* 1. p. 27. *

Kalon-alou. Rheed. mal. 3. p. 73. t. 57. *Raj. hist.* 1437.

- β. *Ficus* foliis lanceolatis integerrimis. *Hort. cliff.* 471.
Roy. lugdb. 212.

Ficus indica maxima, folio oblongo, funiculis e sum-
mis ramis dimissis radices agentibus se propagans, fru-
ctu minori sphaerico sanguineo. *Steen. jam.* 189. *hist.*
2. p. 140. t. 223. *Raj. dendr.* 16.

Ficus americana, arbuti foliis non serratis, fructu pili
magnitudine. *Pluk. alm.* 144. t. 1788.

Ficus citri folio, fructu parvo purpureo. *Catesb. ex.*
3. p. 18. t. 18.

Tiela. Rheed. mal. 3. p. 85. t. 63.

Habitat in Indiis. ♀

- β. ejusdem esse speciei cum α docuit ramentum proprie-
tas radicandi, flores rubri, *Catesb.* & anteptu-
rum nota.

racemosa.

6. *FICUS* foliis ovatis acutis integerrimis, caule arbo-
reo, fructu racemoso. *Amæn. acad.* 1. p. 30.

Alty-alu. Rheed. mal. 1. p. 43. t. 25. *Raj. hist.* 1434.

Habitat in India. ♀

pumila.

7. *FICUS* foliis ovatis acutis integerrimis, caule repente.
Amæn. acad. 1. p. 30.

Ficus sylvestris procumbens, folio simplici. *Kämpf. a-
mæn.* 803. t. 804.

Habitat in China, Japonia. 2

*Ficum a Cryptogamia ad Polygamsiam amandavi, sus-
cendente fructificationis structura, receptaculi umbilico
in quibusdam patulo, diffinitate sequentium; jussu il-
lustriss. D. O. Munchausen. L. B. Botanici a-
cutissimi.*

2
asymonia
wa. ovata
hence tota
21115 hinc
1024

Clas-

Classis XXIV.

CRYPTOGAMIA.

FILICES.

EQUISETUM. 1

1. EQUISETUM caule spicato, frondibus compositis. *sylvaticum.*

Fl. succ. 834.

Equisetum setis ramosis internodio multoties longioribus. *Roy. lugdb.* 496.

Equisetum sylvaticum. *Fl. lapp.* 391. *Tabern. hist.* 562.

Equisetum sylvaticum, tenuissimis setis. *Bauh. pin.* 16. *theatr.* 145.

Habitat in Europæ septentrionalis pratis sylvaticis. &

2. EQUISETUM scapo fructificante nudo, sterili frondoso.

Equisetum scapo fructificante nudo, caule sterili ramis compositis. *Fl. succ.* 833. *Mat. med.* 479. *Dalib. paris.* 308.

Equisetum setis quadrangularibus internodio longioribus, caulibus reptantibus. *Gnest. stamp.* 1. p. 201.

Equisetum setis simplicibus internodio multoties longioribus. *Roy. lugdb.* 496.

3. EQUISETUM arvense. *Fl. lapp.* 390. *Hort. cliff.* 471. *Gren. virg.* 123.

Equisetum arvense, longioribus setis. *Bauh. pin.* 16.

Equisetum minus. *Fuchs. hist.* 323.

Habitat in Europæ agris, pratis. &

4. EQUISETUM caule angulato, frondibus simplicibus.

Equisetum setis simplicibus internodia vix superantibus.

Roy. lugdb. 496. *Fl. succ.* 835. *Dalib. paris.* 307.

Equisetum palustre, *Fl. lapp.* 392.

Equisetum palustre, brevioribus setis. *Bauh. pin.* 16. *theatr.* 242.

5. EQUISETUM palustre minus polystachyon. *Bauh. pin.* 16, *prodr.* 24. *Raj angl.* 3. p. 131. t. 5. f. 3.

Habitat in Europæ aquis. 24.

X x x 3

4 E-

fluviatile.

4. **EQUISSETUM** caule striato, frondibus subsimpli-
cibus.

Equisetum caule non sulcato latissimo, verticillis den-
tissimis. *Hall. belv.* 144. *Fl. suec.* 836. *Dalib. parif.* 307.

Equisetum fluviatile. *Fl. lapp.* 393.

Equisetum palustre longioribus setis. *Banh. pin.* 15.
ibent. 242.

Habitat in Europa ad ripas lacuum, fluviorum. &

terrestre.

5. **EQUISSETUM** caule subnudo lævi. *Fl. suec.* 837.
Dalib. parif. 308.

Equisetum scapo nudo simplicissimo. *Roy. Ingd.* 496.

Equisetum nudum lævius nostras. *Raj. angl.* 3. p. 131.
t. 5. f. 2.

Habitat in Europæ paludibus turfosis, profundis. &

hyemale.

6. **EQUISSETUM** caule nudo scabro basi subramoso.
Fl. suec. 838. *Dalib. parif.* 308.

Equisetum scapo nudo simplicissimo. *Raj. Ingd.* 496.
Gron. virg. 196.

Equisetum hyemale. *Fl. lapp.* 394.

Equisetum foliis nudum ramosum. *Banh. pin.* 16.

Equisetum nudum minus variegatum basilienfe. *Banh.*
pin. 16. *prodr.* 25.

Equisetum. *Cam. epit.* 770. f. A.

Habitat in Europæ sylvis, asperis, uliginosis. &

ONOCLEA.

sensibilis.

1. **ONOCLEA.** *Gen. nov.* 1093.

Angiopteris. *Mitch. gen.* 29.

Osmunda frondibus pinnatis: foliolis superioribus basi
coadunatis: omnibus lanceolatis pinnato subulatis.
Hort. cliff. 272. *Gron. virg.* 196.

Filix mariana, *osmundæ facie*, racemifera. *Pluk. mant.*
80. t. 404. f. 2.

Polypodium virginianum majus, *osmundæ facie*, tene-
rius. *Morif. bist.* 3. p. 563. f. 14. t. 2. f. 10.

Filix indica, *polypodii facie.* *Mentz. pag.* 6. t. 10.

Habitat in Virginia. &

OPHIOGLOSSUM.

algatum.

1. **OPHIOGLOSSUM** fronde ovata. *Fl. suec.* 839.
Dalib. parif. 309,

Ophioglossum folio ovato, spica disticha. *Hort. cliff.*
472.

Ophio.

- Ophioglossum vulgatum. *Baub. pin. 354. Plum. fil. XXXVI. t. B. f. 5.*
 Ophioglossum. *Baub. pin. 364.*
 3. Ophioglossum minus, subrotundo folio. *Baub. pin. 354. prodr. 150.*
Habitat in Europæ pratis sylvaticis.
2. OPHIOGLOSSUM fronde lanceolata. *Lusitanicum,*
 Ophioglossum pumilum autumnale. *Grisl. last.*
 Ophioglossum angustifolium minimum. *Barr. rar. 1280. t. 252. f. 2.*
Habitat in Lusitania.
3. OPHIOGLOSSUM fronde cordata. *reticulatum,*
 Ophioglossum cordatum & reticulatum. *Plum. fil. 141. t. 164.*
 Ophioglossum vulgaris facie. *Pet. fil. 179. t. 10. f. 4.*
Habitat in America meridionali.
4. OPHIOGLOSSUM fronde palmata basi spicifera. *palmatum.*
 Ophioglossum palmatum. *Plum. fil. 139. t. 163. Pet. fil. 178. t. 11. f. 2.*
Habitat in America meridionali.
5. OPHIOGLOSSUM caule flexuoso tereti, frondibus scandens.
 oppositis pinnatis: foliolis utrinque spiciferis. *Hort. cliff. 473. Fl. zeyl. 374. **
 Filix scandens pulchra brasiliiana. *Breyn. cent. 185. t. 96.*
 Lonchitis scandens brasiliiana, pinnulis eleganter laciniatis. *Morif. hist. 3. p. 568. f. 14. t. 3. f. 15.*
 Tlieru-valli-panna. *Rheed. mal. 12. p. 65. t. 33.*
Habitat in India. b.
6. OPHIOGLOSSUM caule flexuoso angulato, frondibus diphyllis: pinnis trifido-palmatis. *flexuosum.*
*Fl. zeyl. 375. **
 Valli-panna. *Rheed. mal. 12. p. 6. t. 32.*
Habitat in India. b.
Præcedenti nimis affinis.

OSMUNDA. 4

* *Scapis insidentibus cauli ad basin folii.*

1. OSMUNDA scapo caulino solitario, frondibus verticillatis lanceolatis indivisis. *zeylanica.*
 Osmunda frondibus lanceolatis indivisis verticillato-terminalibus. *Fl. zeyl. 373. **
Habitat in Zeylona.

X x x 4

2. OS-

- Lunaria.* 2. OSMUNDA scapo caulino solitario, fronde pinnata solitaria. *Fl. suec.* 842. *Dalib. paris.* 309.
 Osmunda fronde pinnata caulina: pinnis lunulatis. *Hort. cliff.* 472.
 Osmunda folio pinnatifido: pinnis lunulatis. *Fl. lapp.* 389.
 Lunaria racemosa minor & vulgaris. *Bauh. pin.* 354.
 Lunaria minor. *Cam. epit.* 643.
 β. Lunaria racemosa ramosa major. *Bauh. pin.* 355.
 Lunariæ rarior species. *Cam. epit.* 644.
 γ. Lunaria racemosa minor, adianti folio. *Breyn. cent.* 183. t. 93.
 δ. Lunaria minor, rutaceo folio. *Bauh. pin.* 355.
Habitat in Europa.
- virginiana.* 3. OSMUNDA scapo caulino solitario, fronde supradecomposito.
 Osmunda fronde pinnata caulina, fructificationibus spicatis. *Gron. ving.* 196.
 Osmunda asphodeli radice. *Plum. fil.* 136. t. 159. *Pet. fil.* 168. t. 9. f. 2.
Habitat in America.
- phyllitidis.* 4. OSMUNDA scapis caulinis geminis, fronde pinnata, caule lævi.
 Osmunda lanceolata & subtiliter serrata. *Plum. fil.* 133. t. 156.
 Osmunda racemifera, phyllitidis folio vix crenato. *Pet. fil.* 163. t. 8. f. 15.
Habitat in America meridionali.
- hirta.* 5. OSMUNDA scapis caulinis geminis, fronde pinnata, caule hirta.
 Osmunda spicis geminis. *Pet. fil.* 164. t. 14. f. 5.
 Osmunda hirsuta, lonchitidis folio. *Plum. fil.* 114. t. 157.
 Lunaria hirsuta, lonchitidis folio, spica gemella. *Raj. suppl.* 102.
 Lunaria elatior americana hirsuta, pinnis lonchitidis integris, duplici spica. *Morif. bist.* 3. p. 598.
 Lonchitis hirsuta florida. *Plum. amer.* 18. t. 26.
Habitat in Martinica.
- hirsuta.* 6. OSMUNDA scapis caulinis geminis, fronde bipinnata hirsuta.
 Osmunda, spicis geminis, villosa. *Pet. fil.* 165. t. 8. f. 16.

Osmunda molliter hirsuta & profunde laciniata. Plum. fil. 138. t. 162.

Lunaria elatior, matricariæ folio, spica duplici. Sloan. jam. 14. bist. 1. p. 71, t. 25. f. 6. Moris. bist. 3. p. 595. Raj. suppl. 101.

Habitat in Americæ meridionalis rupibus.

7. OSMUNDA scapis caulinis geminis, fronde supra-*adiantifolia* decomposita.

Osmunda, filiculæ folio, major. Plum. fil. 135. t. 158.

Osmunda adianti nigri facie. Pet. fil. 167. t. 9. f. 1.

Lunaria elatior, adianti albi folio, duplici spica. Sloan. jam. 14. bist. 1. p. 71. Moris. bist. 3. p. 595.

Adiantum saxosum floridum. Plum. amer. 29. t. 43. Raj. suppl. 101.

Habitat in Dominicæ, Jamaicæ rupibus.

* *Scapis nudis, e radice enatis.*

8. OSMUNDA scapis radicatis: racemis verticillatis, *verticillata*, fronde supradecomposita.

Osmunda verticillata. Plum. fil. 137. t. 160. Pet. fil. 171. t. 12. f. 4.

Habitat in America meridionali.

9. OSMUNDA scapo radicato, fronde pinnata: pinnis *cervina* integerrimis.

Osmunda linguæ cervinæ folio. Plum. fil. 132. t. 154.

Osmunda racemifera, phyllidis folio. Pet. fil. 162. t. 8. f. 3.

Habitat in America meridionali.

10. OSMUNDA scapo radicato, fronde pinnata: pinnis *bipinnata*, pinnatifidis.

Osmunda latis crenis incisa. Plum. fil. 133. t. 155.

Habitat in America meridionali.

11. OSMUNDA scapo radicato paniculato, fronde *subfiliculifolia* pradecomposita.

Osmunda frondibus radicatis quater alternatim pinnatis, scapo nudo, pedunculis alternis alternatim spicatis.

Hort. cliff. 172.

Osmunda, filiculæ folio, altera. Plum. fil. 138. t. 161.

Osmunda cicutæ folio. Pet. fil. 170. t. 9. f. 3.

Habitat in America meridionali.

* *Fronde ipsa fructificationes ferente.*

12. OSMUNDA fronde bipinnata apice racemifera. *regalis.*

X x x 5

Of-

Osmunda scapo paniculato polyphyllis. *Fl. suec.* 840.

Dalib. parif. 308.

Osmunda frondibus caulinis simpliciter pinnatis: pinnis lanceolatis. *Hort. cliff.* 472.

Osmunda regalis f. *Filix florida*. *Plum fil.* 35. c. B. f. 4.

Filix ramosa non dentata florida. *Banb. pin.* 357.

Filix palustris. *Dod. pempt.* 473.

β. *Filix florida* f. *Osmunda regalis*, foliis alternis, surculis seminiferis. *Gron. virg.* 123.

Filix non dentata florida, foliis alternis & in summo caule feminibus occultatis. *Pluk. alm.* 156. t. 181. f. 1.

Filix botrytes f. *floridana major virginiana*, pinnulis non dentatis alternatim positis. *Morif. hist.* 3. p. 593.

Habitat in Europa, Virginia ad fluvios.

claytoniana. 13. *OSMUNDA* fronde pinnata: pinnis pinnatifidis apice coarctato-fructificantibus.

Osmunda frondibus pinnatis ex adverso binis: foliolis oppositis integris decurrentibus, spicis seminiferis e superiore parte caulis ad modum & formam frondium binatim egressis. *Clayt*

Habitat in Virginia.

Conjungit O. *Regalis* *precedentes cum hac planta & hæc sequentes cum precedenti ut separari nequeant.*

* *Frondibus aliis foliaceis, aliis fructificantibus.*

cinereum. 14. *OSMUNDA* fronde pinnata: pinnis pinnatifidis: scapis hirsutis, racemis oppositis compositis.

Osmunda mariana, dryopteris folio, in medio caulis florifera. *Pet. mnsf.* 86.

Osmunda non ramosa, tassis floriferis imo folii scapo infra foliosos adnascentibus. *Kaj. suppl.* 86.

Habitat in Marilandia,

Struthiopteris. 15. *OSMUNDA* frondibus pinnatis: pinnis pinnatifidis, scapo fructificante disticho. *Fl. suec.* 841.

Filix palustris altera, fusco pulvere hirsuta. *Banb. pin.* 358.

Filicestrum septentrionale & palustre. *Amm. ruth.* 175.

Habitat in Suecia, Helvetia, Russia, Norvegia. &

Spirant. 16. *OSMUNDA* frondibus lanceolatis pinnatifidis: laciniis confluentibus integerrimis parallelis.

Pteris fronde pinnata lanceolata, laciniis parallelis integerrimis. *Fl. suec.* 844. *Dalib. parif.* 310.

Pteris

Pteris fronde pinnata: foliolis linearibus parallelis. Roy. Lugdb. 497.

Lonchitis minor. Baub. pin. 359.

Lonchitis aspera minor. Cam. epis. 665.

Habitat in Europa.

17. **OSMUNDA** frondibus supradecompositis: pinnis alternis subrotundis incis. *crispa*.

Adiantum foliis minutim in oblongum scissis, pediculo viridi, Baub. pin. 755.

Adiantum album tenuifolium, rutæ murariæ accedens. Baub. bist. 3. p. 743. Raj. bist. 147.

Adiantum album alpinum crispum. Raj. bist. 47. angl. 3. p. 126.

Adiantum album floridum f. Filicula petræa crispa. Pluk. alm. 9. t. 3. f. 2.

Filix botryoides f. Filicula petræa florida anglica, foliis plurisariam divis. Moris. bist. 3. p. 593. f. 14. t. 4. f. 4.

Habitat in Anglia, Helvetia.

Planta quasi media inter Osmundas ultimas & Acrosticha.

ACROSTICHUM. 5

* *Fronde simplici indivisa.*

1. **ACROSTICHUM** frondibus lineari-lanceolatis acutis, caule scandente. *Fl. zeyl. 380. * Amœn. acad. 1. p. 268. **

Tiri-panna. Rheed. mal. 12. p. 141. t. 33.

Habitat in India.

Frustrificatio non dum visa nobis fuit.

2. **ACROSTICHUM** frondibus lanceolato-ovatis integerrimis, caule scandente. *Amœn. acad. 1. p. 268.*

Lingua cervina scandens, citrei foliis, minor. Plum fil. 107. t. 116.

Habitat in America meridionali.

3. **ACROSTICHUM** frondibus integerrimis glabris petiolatis: sterilibus subrotundis: fertilibus linearibus. *Fl. zeyl. 378. * Amœn. acad. 1. p. 268. t. 2. **

Phyllitis repens scandensve, foliis duplicibus: aliis latis & subrotundis; aliis longis & angustis. Raj. suppl. 13.

Filix malabarica. Pet. gaz. 3. t. 53. f. 12.

Matet a-mala-maravara. Rheed. mal. 12. p. 87. t. 29.

Habitat in Malabar. Zeylona, Africa.

4. A.

- crinitum*. 4. **ACROSTICHUM** fronde ovata obtusa hirsuta supra crinita.
Phyllis crinita, latissimo folio. *Pet. fil.* 145. t. 13. f. 14.
Habitat - - - - -
- * *Fronde simplici divisa.*
- septentrionale*. 5. **ACROSTICHUM** fronde nuda lineari laciniata. *Fl. lapp.* 380. *Fl. succ.* 856. *Amoen. acad.* 1. p. 269. *Roy. lugdb.* 507. *Hall belv.* 134. *Dalib. paris.* 314.
Filicula saxatilis corniculata. *Baub. pin.* 358.
Filix saxatilis tragi. *Lob. ic.* 47.
Habitat in Europæ fissuris rupium. &
- pectinatum*. 6. **ACROSTICHUM** nudum simplicissimum, spica secunda adscendente reflexa compressa. *Amoen. acad.* 1. p. 270. f. 4. *
Adiantum pectinatum, e capite bonæ spei. *Pet. masf.* 743.
Juncus elegantissimus, capitulis pectinatis. *Pluk. alm.* 200. s. 95 f. 7. *Raj. suppl.* 629.
Juncus africanus, pectinato capite. *Morif. bist.* 3 p. 233. f. 8. s. 9. f. 30.
Habitat in Æthiopia.
- dichotomum*. 7. **ACROSTICHUM** nudum dichotomum, spicis secundis adscendentibus reflexis compressis.
Filix cochine. *Pet. gaz.* t. 70. f. 12.
Habitat in China.
Refert A. pectinatum caule & spica, at Caulis est pluries dichotomus totidem spicis.
- digitatum*. 8. **ACROSTICHUM** caulibus nudis triquetris, fronde digitata lineari integerrima æquali. *Fl. zeyl.* 379. * *Amoen. acad.* 1. p. 269. f. 1. *
Planta pluribus foliis triquetris instar graminis cyperini summitate foliorum in plurima folia biuncialia divisus.
Herm. zeyl. 27.
Habitat in Zeylona.
- polypodioides*. 9. **ACROSTICHUM** fronde pinnatifida, pinnis linearibus integerrimis patentibus connatis. *Amoen. acad.* 1. p. 271. *
Acrostichum fronde pinnata: foliolis linearibus alternis sessilibus, terminali plerumque trifida. *Gron. virg.* 198.
Polypodium minus virginianum, foliis brevibus subtus argenteis. *Morif. bist.* 3. p. 563. f. 14. t. 3. f. 5.
Pe-

Polypodium minus, pinnulis raris subtus cinereis. *Sloan. jam.* 16. *bist.* 1. p. 79.

Filix polypodium dicta minima jamaicensis, foliis aversa parte pulvere asplenii ritu circumquaque resparsis. *Pluk. alm.* 153. t. 289. f. 1.

Habitat in Virginia, Jamaica.

* *Fronde pinnata.*

10. **ACROSTICHUM** fronde pinnata: pinnis alternis *aureum*, linguiformibus integerrimis glabris. *Hors. cliff.* 477. *Amæn.* 1. p. 273.

Phyllitis ramosa aurea. *Pet. fil.* 142. t. 8. f. 5.

Lonchitis palustris maxima. *Sloan. jam.* 15. *bist.* 1. p. 76.

Lingua Cervina aurea. *Plum. fil.* 87. t. 104.

Filix palustris aurea, foliis linguae cervinae. *Plum. amer.* 5. t. 7. *Raj. suppl.* 63.

Filix simpliciter pinnatis foliis s. *Phyllitis ramosa jamaicensis maxima*, aversa parte ferruginea lanugine circumquaque obducta. *Pluk. alm.* 154. t. 288. f. 1.

Habitat in Jamaica, Dominicae humentibus.

11. **ACROSTICHUM** frondibus pinnatis; pinnis ob-*orbisfolium*, longis integris serratis acutis, stipitibus squamosis.

Lonchitis calamifera, pinnis serratis. *Pet. fil.* 153. t. 9. f. 8.

Filix scandens latifolia serrata. *Plum. amer.* 8. t. 12. *Raj. suppl.* 76.

Filix major, in pinnas tantum divisa raras oblongas latasque crenatas. *Sloan. jam.* 18. *bist.* 1. p. 84. t. 38.

Filix jamaicensis hirsuta, sorbi aucupariae folia quodammodo referens. *Morif. bist.* 3. p. 576.

Filix jamaicensis, pinnatis fraxini foliis tenuissime crenatis mucronatis. *Pluk. alm.* 153. t. 286. f. 3.

Habitat in Jamaica, Domingo.

12. **ACROSTICHUM** fronde pinnata: pinnis alternis *arcolatum*, linearibus apice serratis. *Gron. virg.* 124. *Amæn. acad.*

1. p. 274.

Filix mariana, pinnulis semiuiferis angustissimis. *Pet. act.* 246. p. 398.

Habitat in Virginia, Marilandia.

13. **ACROSTICHUM** frondibus pinnatis: pinnis alter-*plaigneures*, nis ovatis crenatis sessilibus sursum arcuatis. *Gron. virg.* 123. *Amæn. acad.* 1. p. 272. *

Filix

- Filix polypodium dicta minima virginiana platyneuros.
Pluk. alm. 153. t. 289. f. 2. *Raj. suppl.* 58.
 Asplenium virginianum, polypodii facile. *Raj. suppl.* 59.
 Polypodium minus virginianum, foliis brevibus subtus
 argenteis. *Murif. hist.* 3. p. 563. f. 14. t. 2. f. 5.
Habitat in Virginia.
- nodosum.* 14. ACROSTICHUM fronde pinnata: pinnis lanceola-
 tis oppositis, stipitis articulis nodosis.
 Phyllitis pulverulenta nodosa major. *Pet. fil.* 143. t. 6.
 f. 16.
 Filix latifolia nodosa. *Plum. amer.* 4. t. 6. *Raj. suppl.*
 63.
 Lingua cervina nodosa major. *Plum. fil.* 108.
Habitat in Dominica.
- ovifoliatum.* 15. ACROSTICHUM fronde pinnata: foliolis ternatis lanceolatis. *Amoen. acad.* 1. p. 274. *
 Acrostichum fronde pinnata; foliolis digitatis: pinnis linearibus: intermedia longiore. *Hort. upf.* 476.
 Phyllitis ramosa trifida. *Sloan. jam.* 88. *hist.* 1. p. 88.
 t. 45. f. 2.
 Phyllitis pulverulenta, foliis angustis trifidis. *Pet. fil.*
 141. t. 8. f. 9.
 Lingua cervina triphylla angusta & leviter serrata. *Plum. fil.* 12. t. 144.
 Filix jamaicensis non ramosa trifolia angustifolia, caule lævi. *Murif. hist.* 3. p. 572. *Raj. suppl.* 73.
Habitat in Jamaica.
- * *Fronde juv-bipinnata.*
- siliquosum.* 16. ACROSTICHUM fronde pinnata: pinnis alternis
 sursum pinnulatis linearibus: inferioribus bipartitis. *Fl. zeyl.* 376. * *Amoen. acad.* 1. p. 270. f. 3.
 Acrostichum maderaspatanum, foliis laciniatis. *Pet. mus.* 741.
 Millefolium aquaticum. *Rumpf. amb.* 5. p. 17. t. 74. f. 1.
 Planta siliquosa bivalvis unicapularis, siliquis in plurimos ramulos foliatis diffusis, crithmi maritimi segmenta quodammodo mentientibus. *Pluk. alm.* 298.
 t. 215. f. 3.
Habitat in Zeylona.
- obtusifolides* 17. ACROSTICHUM fronde pinnata: pinnis alternis
 utrinque pinnatifidis: sterilibus latoribus. *Fl. zeyl.* 377. t. 4. * *Amoen. acad.* 1. p. 275.
Habitat in Zeylonæ aquis.
18. A-

18. ACROSTICHUM frondibus sub-bipinnatis: pinnis *marantæ*. opposito-coadunatis subtus hirsutissimis: basi subdentatis.

Lonchitis folio ceterach. Bauh. pin. 259. Barr. ic. 858. 857.

Lonchitis aspera marantæ. Cam. epit. 666.

Filicula crispa, lanugine hepatici coloris vestita. Pluk. alm. 150. t. 281. f. 4.

Habitat in Europa australi.

Stipites spississimi, atropurpurei. Frons ferme bipinnata, lanceolata, subtus dense imbricata squamis paleaceis, croceis, efficcata contrahitur & occultas paginam anticum. Foliola opposita, fere pinnata. Pinnæ ovate, obtusæ, crassæ, integerrimæ; at inferiores basi sæpe utrinque dentatæ. Fructificationes sparsæ inter squamas foliaceas per totam paginam posticam.

19. ACROSTICHUM fronde sub-bipinnata: pinnis opposito-coadunatis obtusis subtus hirsutis: basi integerrimis.

Polypodium fronde duplicato-pinnata: pinnulis coadunatis obtusis subtus hispidis. Fl. suec. 850.

Polypodium duplicato-pinnatum: pinnulis obrutis remotis subtus villosum. Fl. lapp. 383.

Polypodium lonchitidis marantæ species cambrobritannica. Pluk. alm. 150. t. 89 f. 5.

Filicula alpina tenerior, alis latiusculis brevioribus integris profunde dentatis. Moris. hist. 3. p. 576. f. 14. t. 3. f. 23.

Habitat in Europæ frigidissimæ rupibus.

Maxime affinis precedenti, at vix digito longior, nec spissimanum; stipite virescente, nec atropurpureo ut illa.

20. ACROSTICHUM fronde pinnata: pinnis sessilibus *ebenæ*, oblongis sinuatis: summis brevissimis integerrimis. *Ammon. acad. 1. p. 272. **

Filix non ramosa minor (& minima), caule nigro, furculis raris, pinnulis angustis raris brevibus acutis subtus niveis. Sloan. jam. 20. hist. 1. p. 92. t. 53. f. 1. Raj. suppl. 83.

Habitat in Jamaica sepibus humidiusculis.

21. ACROSTICHUM fronde pinnata: pinnis pinnatifidis integerrimis. *Thelypteris.*

Filix mollis f. glabra, vulgari mari non ramosæ accedens. Bauh. hist. 730. t. 731. f. 1. The-

Thelypteris palustris non ramosa. *Rupp. jen.* 277.

Habitat in Europæ septentrionalioris paludibus.

Habitus Polyp. Filicis maris. Stipes glaber. Folla sterilia, latiora, obtusiora.

* *Fronde bipinnata.*

- cruciatum*, 22. ACROSTICHUM fronde bipinnata: foliolis oppositis lanceolatis: infimis cruciatim appendiculatis. *Amoen. acad.* 1. p. 275. *

Filix pulverulenta, pinnulis obtuse dentatis. *Plum. fil.* 26. t. 38.

Adiantum pulverulentum, pinnis infra auritis. *Pet. fil.* 161. t. 9. f. 10.

Habitat in America meridionali.

- barbatum*, 23. ACROSTICHUM fronde bipinnata: foliolis oppositis: pinnulis lanceolatis obtusis serratis sessilibus alternis. *Hort. cliff.* 476. *Amoen. acad.* 1. p. 274. *

Filix africana floridæ similis, in ambitu foliorum argute denticulata. *Pluk. alm.* 156. t. 181. f. 5.

Habitat in Africa

- calomelanos*, 24. ACROSTICHUM frondibus bipinnatis: pinnis alternis lanceolaris acuminatis pinnatifidis.

Acrostichum fronde alternatim duplicato-pinnatum: pinnulis lanceolatis sessilibus: inferioribus serratis incisis.

Hort. cliff. 476. *Amoen. acad.* 1. p. 274.

Filix non ramosa major, caule nigro, furculis raris, pinnulis angustis raris longis dentatis. *Sloan. jam.* 20. *bist.* 1. p. 92. t. 30. f. 2.

Filix albissimo pulvere conspersa. *Plum. fil.* 30. t. 40.

Pet. fil. 156. t. 9. f. 11.

Adiantum nigro simile, albissimo pulvere conspersum.

Plum. amer. 30. t. 40.

Adiantum calomelanos americanum f. *Adiantum nigrum*, foliis prona parte candidissimis. *Pluk. alm.* 11. t. 124. f. 3

Habitat in America meridionali.

Stipes ita ater, at folia subsus nivea.

- pulchrum*, 25. ACROSTICHUM bipinnatum: pinnulis indivisis. *Sauv. monsp.* 246. †

Adiantum pulcherrimum. *Bauh. bist.* 3. p. 737.

Adiantum foliis longioribus pulverulentis, pediculo nigro. *Bauh. pin.* 355.

Habitat Monspellii, inque Harcynia.

Pollen est omnia crocei coloris. *Sauv.*

PTERIS. *C** *Frondibus simplicissimis.*

1. PTERIS fronde simplici lanceolata subangulata glabra: *lanacolata*.
apice fructificante.

Lingua Cervina foliis acutis & ad oras summitatum pulverulentis. *Plum. amer.* 28. t. 40. *fil.* 116. t. 132.

Morif. bist. 3. p. 558. *Raj. suppl.* 52.

Phyllitis lineata hinc inde dentata. *Pet. fil.* 123. t. 6. f. 5.

Habitat in Domingo.

2. PTERIS fronde simplici lineari integerrima longitudi-*lineata*.
naliter fructificante.

Lingua Cervina longissimis & angustissimis foliis. *Plum. amer.* 28. t. 41. *fil.* 123. t. 143. *Morif. bist.* 3. p. 558. *Raj. suppl.* 52.

Phyllitis lineata, graminis folio longissimo. *Pet. fil.* 126.

t. 14. f. 3.

Habitat in Domingo.

3. PTERIS fronde simplici lineari apice trifido. *tricuspidata*.

Lingua Cervina angusto trifidoque folio. *Plum. fil.* 121.

t. 140.

Phyllitis lineata apice trifido. *Pet. fil.* 124. t. 10. f. 6.

Habitat in Domingo.

4. PTERIS fronde simplici dichotoma subtus hispida: *a-furcata*.
picibus fructificantibus.

Lingua Cervina furcata. *Plum. fil.* 122. t. 141.

Phyllitis aspera, furcis lineatis. *Pet. fil.* 125. t. 6. f. 6.

Habitat in America meridionali.

* *Frondibus simpliciter pinnatis, compositis, oblongis.*

5. PTERIS frondibus pinnatis: foliolis pinnatifidis, cau-*arboreo*.
dice arboreo aculeato.

Filix ramosa arborescens & aculeata. *Plum. fil.* 6. t. 5.

Filix arborescens latifolia aculeata. *Plum. amer.* 3. t. 3.

Filix lineata arborea, caudice aculeato. *Pet. fil.* 132. t.

4. f. 2.

Habitat in Martinica.

6. PTERIS frondibus pinnatis: pinnis oppositis ovato-*grandifolia*.
linearibus acuminatis integerrimis.

Pteris fronde pinnata: pinnis lanceolato-linearibus integerrimis sessilibus erectiusculis. *Hort. cliff.* 473.

Phyllitis ramosa ad margines pulverulenta. *Pet. fil.* 127.

t. 6. f. 15.

Y y y

Lingua

Lingua Cervina latifolia membrana tenui marginata.
Plum. fil. 88. t. 106.

Filix latifolia ad margines pulverulenta. *Plum. amer.*
6. t. 8.

Filix pinnata latifolia ad margines. pulverulenta. *Raj.*
suppl. 64.

Filix americana maxima: lobis oppositis acutioribus ad
margines pulverulentis. *Morif. hist.* 3. p. 571.

Habitat in Dominicæ, Martinicæ *uliginosis.*

longifolia. 7. PTERIS frondibus pinnatis: pinnis linearibus repandis: basi cordatis. *Hort. cliff.* 473.

Lonchitis lineata, pinnis longissimis. *Pet. fil.* 129. t. 6.
f. 12.

Lonchitis americana major, alis longissimis non serratis ad basin auriculatis oppositis ad oras pulverulentis. *Morif. hist.* 3. p. 568.

Lonchitis non ramosa, longis angustis & ad basin auriculatis foliis. *Plum. fil.* 52. t. 69. *Raj. suppl.* 65.

Filix non ramosa, longissimis angustis & ad basin auriculatis foliis. *Plum. amer.* 12. t. 18.

Habitat in Dominicæ *sylvis juxta rivulos.*

vinosa. 8. PTERIS frondibus pinuatis: pinnis linearibus rectis basi rotundatis.

Habitat in China *Osbeck.*

Præcedenti valde similis.

stipularis. 9. PTERIS fronde pinnata: pinnis linearibus sessilibus, stipulis lanceolatis.

Filix altera longissimis angustis & ad basin foliosis foliis.

Plum. amer. 12. t. 19. *fil.* t. 70.

Filix lineata ad basin foliolosa. *Pet. fil.* 130. t. 12. f. 1.

Habitat in America *meridionali.*

trichomanoides. 10. PTERIS fronde pinnata: pinnis subovatis obtusis repandis: subtus hirsutis. *Hort. cliff.* 473.

Trichomanes argenteum ad oras nigrum. *Plum. fil.* 57.
t. 75. *Pet. fil.* 137. t. 9. f. 16.

Trichomanes majus, pinnis sinuatis subtus niveis. *Sloan. jam.* 17. *hist.* t. p. 80. t. 35. f. 1. *Raj. suppl.* 60.

Habitat in Jamaica, Dominicæ *rupibus.*

raspa. 11. PTERIS fronde pinnata: pinnis oblongo-ovatis obtusis: subtus lanatis.

Acrostichum fronde pinnata: foliolis oblongo-ovatis vix cordatis hirsutis obsolete serratis. *Hort. cliff.* 475.

Filix

Filix minor, rufa lanugine tota obducta, in pinnas tantum divisa raras non crenatas subrotundas. *Sloan. jam.*

19. *hist.* 1. p. 87. t. 45. f. 1.

Habitat in America septentrionaliore.

* *Frondibus sub-bipinnatis f. ramosis.*

12. PTERIS frondibus quinquangulis trifoliatis: pinnis *pedata*, pinnatifidis: lateralibus bipartitis.

Pteris fronde simplici quinquangula producta tripartita: intermedia trifida: lateralibus bifidis. *Hort. cliff.* 473.

Hemionitis profunde laciniata ad oras pulverulenta. *Plum. amer.* 24. t. 34. *fil.* 130 t. 152.

Hemionitis foliis atro-virentibus maxime dissectis; f. Filix geranii robertiani folio. *Sloan. jam.* 15. *hist.* 1. v. 73.

Filix marginifera, geranii folio. *Pet. fil.* 176. t. 8. f. 12.

Filix Hemionitis americana, petroselini foliis profunde laciniatis. *Pluk. alm.* 155. t. 286. f. 5. *Raj. suppl.* 55.

Adiantum monophyllum americanum, foliis profunde laciniatis ad oras pulverulentum. *Morif. hist.* 3. p. 592.

- β. Adiantum monophyllum antegoanum, ad oras pulverulentum platani in modum divisum: lacinia media longius producta. *Morif. hist.* 3. p. 392.

Habitat in Jamaica, Dominica, Sibirica rupibus.

Planta sibirica minor & subsus nivea: margine ferrugineo.

13. PTERIS fronde supradecomposita: foliolis pinnatis: *aquilina*, pinnis lanceolatis: infimis pinnatifidis: superioribus minoribus. *Hort. cliff.* 473. *Fl. suec.* 343. *Valerb. par.* 309.

Filix ramosa major, pinnulis obtusis non dentatis. *Bauh. pin.* 357.

Filix femina. *Fuchs. hist.* 596. *Cam. epit.* 992.

Habitat in Europae sylvis, praesertim caduis.

14. PTERIS fronde supradecomposita, pinnis linearibus: *caudata*, infimis basi pinnato dentatis; terminalibus longissimis. *Hort. cliff.* 473.

Filix ramosa, pinnulis longiusculis: partim auriculatis. *Plum. amer.* 14. t. 22. *fil.* 23. t. 29.

Filix femina vulgaris facie. *Pet. fil.* 135. t. 3. f. 12.

Filix femina ramosissima jamaicensis, pinnula alas claudente longissima. *Pluk. alm.* 156.

Filix femina f. ramosa major, pinnulis angustis obtusis

• Y y y 2

non

non dentatis: impari furculum terminante longissima.
Sloan. jam. 24. bist. 1. p. 101. Moris. bist. 3. p. 583.

- β. Filix femina f. ramosa major, pinnulis angustissimis rarissimisque. *Sloan. jam. 24. bist. 1. p. 101. t. 63.*
Habitat in Jamaica, Dominicæ saxosis.

- acrilata.* 15. PTERIS fronde decomposita: foliolis pinnatis: infimis semipinnatifidis, terminalibus baseosque longissimis.

Lonchitis ramosa, limbo pulverulento. Plum. amer. 21 t. 30. fil. t. 51. Pet. fil. 131. t. 3. f. 10.

Habitat in Jamaica, Dominica.

Folia lateralibus infima longiora ad basin, latere tantum inferiore pinnata.

- atropurpurea.* 16. PTERIS fronde decomposita pinnata: pinnis lanceolatis: terminalibus longioribus.

Pteris adianti facie, caule ramulis petiolisque politiore nitore nigricantibus. Gron. virg. 197.

Habitat in Virginia.

Stripes spishamensis, atropurpureus, nitidus. Frons pinnata: pinnis lanceolatis, lato margine revolutis, quæ magis inferius & etiam magis petiolata: inferioribus petiolis uno alterove pari pinnarum minorum sessilium auctis.

- transita.* 17. PTERIS fronde pinnata: pinnis pinnatifidis: infima bipartita.

Filix pinnalis lonchitidis obtusis non dentatis ad oras pulverulentis. Plum. amer. 10. t. 14. fil. t. 15.

Filix lonchitidis pinnis. Pet. fil. 134. t. 3. f. 11.

Habitat in Domingo, Martinica, Jamaica.

- semipinnata.* 18. PTERIS fronde subbipinnata: foliolis lateralibus loboque infimo semipinnatifidis.

Habitat in China. Osbeck.

Scapus levis. Frondes lanceolata, pinnata. Foliola opposita, remota, pinnatifida: apicibus terminalibus elongatis; margo foliorum terram spectans sapius & unice pinnatifidus, non vero superior, hinc quasi semipinnata evadunt foliola, infima demum lacina pinnarum cæteris longior est & similiter semipinnatifida; reliquis omnibus integris, obtusis.

- dichotoma.* 19. PTERIS frondibus dichotomis: pinnatis integerrimis. Filix furcata, pinnulis longiusculis non dentatis. *Plum. fil. 22. t. 28. amer. 13. t. 20. Raj. suppl. 96.*

Filix

Filix furcata, taxi foliis. *Pet. fil.* 51. t. 5. f. 4.

Filix femina f. *ramosa major* dichotoma, pinnulis lonchitidis longis angustis non dentatis. *Sloan. jam.* 24. *bist.* 1. p. 102.

Filix jamaicensis dichotomos f. ramis bifidis, feminae nostratis, pinnulis ramosissima. *Pluk. alm.* 156.

Habitat in Jamaica, Martinicae sylvis.

Fructificatio a me visa non fuit.

BLECHNUM. 7

1. BLECHNUM fronde pinnata: pinnis lanceolatis oppositis basi emarginatis. *orientalis.*

Lonchitis juxta nervum pulverulenta. *Pet. fil.* 128. t. 3.

f. 9. *Morif. bist.* 3 p. 568. f. 14. t. 2. f. 16.

Lonchitis f. *Hermionitis* pinnata juxta nervum pulverulenta. *Plum. Raj. suppl.* 66?

Habitat in America meridionali. Pinnæ lanceolatae plurimæ.

2. BLECHNUM fronde pinnata: pinnis linearibus alternis. *occidentalis.*

Habitat in China *Osbeck.*

Frons tripedalis, stipite basi antice setis griseis magnis vestito; latus anterius stipitis sulcis 3 longitudinaliter exaratus. Folia lineari-lanceolata, sessilia, glabra, integerrima, ad angulum acutum striata, digiti longitudine.

HEMIONITIS. 6

1. HEMIONITIS frondibus lanceolatis integerrimis. *lanceolata.*

Phyllitis angustifolia lineis catenatis. *Pet. fil.* 122. t.

6. f. 4. quæ *Plum. fil.* 127. f. 6.

Habitat in America meridionali.

2. HEMIONITIS frondibus palmatis hirsutis. *palmata.*

Hemionitis fronde palmata, lineis seminalibus reticulatis. *Hort. cliff.* 474.

Hemionitis aurea hirsuta. *Plum. amer.* 23. t. 33. f. 1. t. 51.

Hemionitis folio hirsuto & magis dissecto f. *ranunculi* folio. *Sloan. jam.* 14. *bist.* 1. p. 72. *Morif. bist.* 3. p. 560. f. 14. t. 1. f. 5.

Filix, ranunculi folio, prolifera *Pet. fil.* 177. t. 8. f. 11.

Filix Hemionitis dicta, saniculæ foliis, villosa. *Pluk. alm.* 155. t. 291. f. 4.

Habitat in America meridionali.

LONCHITIS. 9

- livida.* 1. LONCHITIS frondibus pinnatifidis obtusis integerrimis, furculis ramosis hirsutis.
 Filix ramosa, pinnulis quercinis. *Plum. fil.* 18. t. 20.
 Adiantum pinnis quercinis, ramis villosis. *Pet. fil.* 82. t. 4. f. 5.
Habitat in America meridionali.
- ovata.* 2. LONCHITIS frondibus pinnatis: pinnis infimis bipartitis, furculis indivisis aculeatis.
 Filix latifolia spinulis mollibus & nigris aculeata. *Plum. fil.* 14. t. 17.
 Filix spinosa, seminibus junulatis. *Pet. fil.* 172. t. 4. f. 4.
Habitat in America meridionali.
- repens.* 3. LONCHITIS frondibus pinnatis: pinnis alternis sinuatis, furculis ramosis aculeatis.
 Filix aculeata repens. *Plum. fil.* 11. t. 12.
 Filix ramosa aculeata, repens. *Pet. fil.* 39. t. 4. f. 6.
Habitat in America meridionali.

ASPLENIUM. 10

* *Fronde simplici.*

- rhizophylla.* 1. ASPLENIUM frondibus cordato-ensiformibus indivisis: apice filiformi radicante. *Amoen. acad.* 2 p. 337.
 Phyllitis filicifolia parva saxatilis virginiana per summitates foliorum radicata. *Pluk. alm.* 154. t. 105. f. 3.
 Phyllitis non sinuata minor, apice folii radices agente
Sloan. jam. 14. *hist.* 1. p. 71. t. 26. f. 1.
 Phyllitis saxatilis virginiana per summitates foliorum prolifera. *Morif. hist.* 3. p. 557. f. 14. t. 1. f. 14.
Habitat in Jamaica, Virginia, Canada, Sibiria.
- Hemionitis.* 2. ASPLENIUM frondibus simplicibus cordato-hastatis quinquelobis integerrimis, stipitibus lævibus.
 Hemionitis vulgaris. *Bauh. pin.* 353.
 Hemionitis vera. *Clus. hist.* 2. p. 214.
Habitat in Italia, Hispania.

Ma.

Maxime affinis sequenti, at diameter frondis longitudinalis vix superat transversalem.

3. ASPLENIUM frondibus simplicibus cordato-lingu- *Scolopendri-*
latis integerrimis, stipitibus hirsutis. *um.*

Asplenium frondibus ensiformibus integris: basi cordatis inflexis, petiolis hirsutis. *Vir. cliff.* 103. *Hort. cliff.* 474. *Fl. suec.* 843. *Mat. med.* 481. *Roy. lugdb.* 498. *Hall. belv.* 134. *Dalib. parif.* 310. *Sauv. monsp.* 27.

Lingua Cervina officinarum. *Bauh. pin.* 353. *Plum. fil.* 34. t. A. f. 4.

Phyllitis. *Cam. epit.* 579.

- B. Phyllitis crispa. *Bauh. hist.* 3. p. 757.

γ. Phyllitis f. Lingua Cervina maxima, undulato folio auriculato per basin. *Pluk. phyt.* 248. f. 1.

δ. Lingua Cervina medio folii nervo in aculeum abeunte. *Morif. hist.* 3. p. 557. f. 14. t. 1. f. 9.

ε. Lingua Cervina multifido folio. *Bauh. pin.* 353.

ζ. Phyllitis f. Lingua Cervina crispa, folio multifido, ramosa. *Pluk. phyt.* 248. f. 2.

Habitat in Europæ umbrosis, nemorosis, saxosis. &

4. ASPLENIUM frondibus simplicibus lanceolatis integerrimis glabris. *indus.*

Phyllitis indica maxima, latissimis foliis planis musæ facie. *Morif. hist.* 3. p. 558. f. 14. t. 1. f. 15.

Scolopendria indiæ orientalis, musæ facie. *Brœyn. cent.* 129. t. 99.

Habitat in Javæ summis arboribus. Osbeck.

Folia bipedalia, latitudine spithamæ, robusta & consistentia, glabra, ad angulum acutum striata. Lineis fructificantes inter singulos nervos s. strias parallelas, breves, utpote vix $\frac{1}{4}$ latitudinis folii attingentes. Radix insidet summis arboribus, unde folia exeunt in orbem digesta, erecta, instar umbellæ, intra cujus centrum Aves plerumque nidos suos conficiunt.

5. ASPLENIUM fronde simplici lanceolata serrata. *serratum.*

Lingua Cervina longo lato serratoque folio. *Plum. amer.* 27. t. 39.

Phyllitis major margine crenato. *Pet. fil.* 106. t. 6. f. 7.

Phyllitis non sinuata, foliorum limbis leviter serratis.

Sloan. jam. 14. *hist.* 1. p. 72.

Habitat in America calidiore.

6. ASPLENIUM fronde binata: foliolis lanceolatis subsinuatis connatis. *bifolium.*

Lingua Cervina geminato folio. *Plum. fil.* 116. t. 133.
Phyllitis folio geminato dentato. *Pet. fil.* 107. t. 13.
f. 11.

Habitat in America meridionali.

* *Fronde pinnatifida.*

Ceterach. 7. **ASPLENIUM** frondibus pinnatifidis: lobis alternis
confluentibus. *Hort. cliff.* 474. *Mat. med.* 482. *Hall.*
belv. 135. *Dalib. parif.* 310. *Sauv. monsp.* 270.

Asplenium f. *Ceterach.* *Bauh. hist.* 3. p. 479. *Morif. hist.*
3. p. 561. f. 14. t. 2. f. 1. *Plum. fil.* XXXIII. t. B.
f. 3. *Barr. ic.* 1043. 1044.

Ceterach officinarum. *Bauh. pin.* 354.

Habitat in Walliæ, Italiæ fissuris rupium.

obtusifolium. 8. **ASPLENIUM** fronde subpinnata: pinnis obtusis
sinuatis decurrentibus alternis.

Adiantum alis latioribus. *Pet. fil.* 117. t. 2. f. 4.

Habitat in America.

* *Fronde pinnata.*

falcifolium. 9. **ASPLENIUM** fronde pinnata: pinnis falcato-lanceo-
latis crenatis: basi sursum angulatis.

Lonchitis glabra major. *Plum. amer.* 18. t. 27. *fil.* t.
60. *Raj. suppl.* 67.

Lonchitis striata. *Pet. fil.* 110. t. 3. f. 2.

Lonchitis major, pinnulis latioribus leviter denticulatis:
superiore latere auriculatis. *Sloan. jam.* 16. *hist.* 1. p.
78.

Habitat in Antillis.

Trichoma- 10. **ASPLENIUM** frondibus pinnatis: pinnis subrotun-
ner. dis crenatis. *Fl. lapp.* 388. *Fl. suec.* 854. *Hort. cliff.*
474. *Mat. med.* 483. *Roy. lagdb.* 498. *Hall. belv.*
135. *Dalib. parif.* 311.

Trichomanes f. *Polytrichum officinarum.* *Bauh. pin.*
356. *Plum. fil.* 26. t. V. f. 1.

Trichomanes. *Fuchf. hist.* 796. *Cam. epit.* 925.

β. *Trichomanes* foliis eleganter incilis. *Tournef. inst.*
539. t. 315. f. 7. C.

γ. *Trichomanes minus & tenerius.* *Bauh. pin.* 356.

Habitat in Europæ fissuris rupium.

Trich. den- 11. **ASPLENIUM** frondibus pinnatis: pinnis cuneifor-
tatum. mibus obtusis crenato-emarginatis.

Polytrichum saxatile dentatum. *Plum. amer.* 35. t. 50.

Tri-

Trichomanes latifolium dentatum. Plum. fil. 2. 101.
Trichomanes striatum vulgaris facie. Pet. fil. 120. 2.
 2. f. 15.

Habitat in America meridionali.

12. **ASPLENIUM** fronde pinnata: pinnis ovatis super-^{marinum.}
 nie basi gibbis serratis obtusis. *Hort. cliff. 474. Hall.*
helv. 136.

Chamae Filix marina anglica. Moris. hist. 3. p. 573. f.
 14. 2. 3. f. 25. bon.

Filicula maritima ex insulis stœchadis. Bauh. pin. 359.
Burf. XX: 39.

Habitat in Anglia, insulis Stœchadum. Burf.

13. **ASPLENIUM** fronde pinnata: pinnis falcato-lan-^{cultrifolium}
 ceolatis inciso serratis: basi decursum angulatis. *Hort.*
cliff. 474.

Lonchitis latifolia, pediculis lucidis & glabris. Plum.
fil. 45. 2. 59.

Habitat in Martinica.

* *Fronde decomposita.*

14. **ASPLENIUM** frondibus duplicato-pinnatis. pinnis ^{Adiant. ni-}
 obovatis superne crenatis; foliolis inferioribus majo-^{gram.}
 ribus.

Adiantum foliis longioribus pulverulentis, pediculo ni-
gro. Bauh. pin. 355.

Adiantum nigrum officinarum. Bauh. hist. 3. p. 742.

Dryopteris nigra. Dod. pempt. 466.

Habitat in Europa australiore.

15. **ASPLENIUM** frondibus alternatim decompositis: ^{Ruta mur.}
 foliolis cuneiformibus crenulatis. *Roy. lugdb. 498. via.*

Fl. succ. 855. Mat. med. 484. Dalib. parif. 311.

Ruta muraria. Bauh. pin. 356.

Paronychia. Cam. epis. 785.

Habitat in Europa ex rupium fissuris. 2

16. **ASPLENIUM** frondibus tripinnatis: foliolis alter-^{Omopteris.}
 nis: pinnis lanceolatis inciso-serratis.

Asplenium frondibus duplicato-pinnatis, foliis inferiori-
buss majoribus: foliolis ovatis superne crenatis. Roy.
lugdb. 498. Dalib. parif. 311.

Adiantum foliis longioribus pulverulentis, pediculo ni-
gro. Bauh. pin. 355.

Adiantum nigrum officinarum. Bauh. hist. 3. p. 471.

Habitat in Italia, Gallia.

Y y y 5

17. A-

- Teleb ramosum.* 17. **ASPLENIUM** frondibus duplicato-pinnatis: pinnis obovatis crenatis: foliolis inferioribus minoribus. *Roy. lugd.* 498.
Trichomanes ramosum majus & minus. Bauh. pin. 256.
Trichomanes ramosum. Bauh. hist. 3. p. 747. *Raj. anglic.* 3. p. 119.
Habitat in Arvernâ rupibus.
- marginatum* 18. **ASPLENIUM** fronde composita: pinnis oppositis lanceolatis serratis.
Phyllitis ramosa margine membranaceo. Pet. fil. 108. t. 12. f. 2.
Habitat in America.
- squamosum.* 19. **ASPLENIUM** fronde composita: pinnis acuminatis incis, stipite squamolo.
Lonchitis ramosa, caule squamoso. Pet. fil. 112. t. 5. f. 2.
Habitat in America.
- striatum.* 20. **ASPLENIUM** fronde composita: pinnis pinnatifidis obtusis crenatis: terminali acuminata.
Filix striata, pinnis crenatis, major & minor. Pet. fil. 113. 114. t. 3. f. 3. 4.
Habitat in America.

POLYPODIUM.

* *Frondē indivisa.*

- lanceolatum.* 1. **POLYPODIUM** frondibus lanceolatis integerrimis glabris, fructificantibus solitariis, radice nuda.
Phyllitis, folio longo, angustifolia, maculis majoribus. Pet. fil. 8. t. 6. f. 2.
Habitat in America meridionali.
Dicimus fructificationes solitarias, quæ serie longitudinali tantum ab utroque latere disponuntur; seriales, quæ transversim, simplici serie, inter singulos nervos transversales digeruntur; sparsas, quæ inter singulos nervos transversales duobus pluribusque ordinibus digestæ sunt.
- Lycopodioides.* 2. **POLYPODIUM** frondibus lanceolatis integerrimis glabris, fructificationibus solitariis, caule squamoso repente. *Hort. cliff.* 474.
Lingua Cervina scandens, caulibus squamosis. Plin. fil. 104. t. 119.

Phyl-

- Phyllitis scandens*, caulibus squamosis. *Plum. amer.* 29.
t. 42. *Raj. suppl.* 53.
- Phyllitis minor scandens*, foliis angustis. *Sloan. flor.* 15.
hist. 1. p. 73.
- Filix maculata scandens*, caulibus squamosis. *Pet. fil.*
12. t. 4. f. 15.
- Filix sarmentosa bifrons* f. *Dryopteris scandens jamaicensis* inter filicem & lycopodium media. *Pluk. alm.*
156. t. 290. f. 3.
- Habitat in America*: Martinica, Domingo, Jamaica.
- Confer.* *Filix maculata scandens*, caulibus squamosis angustifolia. *Pet. fil.* 13. t. 4. f. 16.
- Filix scandens quercus folio*, caulibus lycopodii. *Pet. fil.* 21. t. 15. f. 2.
3. **POLYPODIUM** frondibus lanceolatis integerrimis piloselloides. hirsutis: sterilibus ovatis, fertilibus lanceolatis, fructificationibus solitariis.
- Lingua cervina minima repens & hirsuta*. *Plum. fil.* 103. t. 118.
- Filix foliis ovalibus & longis pilosis*. *Pet. fil.* 175. t. 10. f. 5.
- Habitat in America meridionali.*
4. **POLYPODIUM** frondibus crenatis glabris: sterilibus heterophyllis subrotundis sessilibus; fertilibus lanceolatis, fructificationibus solitariis.
- Lingua cervina heterophylla scandens & repens*. *Plum. fil.* 105. t. 120.
- Filix maculosa scandens*, foliis rotundis & longis glabris. *Pet. fil.* 174. t. 15. f. 4.
- Habitat in America meridionali.*
5. **POLYPODIUM** frondibus lanceolatis glabris integerrimis, fructificationibus serialibus.
- Phyllitis maculata*, amplissimo folio. *Pet. fil.* 1. p. 6. t. 8.
6. *Phyllitis maculata*, breviori & crassiori folio. *Pet. fil.* 2. p. 6. t. 1.
- Habitat in America meridionali.*
6. **POLYPODIUM** frondibus lanceolatis glabris integerrimis, fructificationibus sparsis.
- Polypodium*, linguae cervinae folio, majus. *Plum. amer.* 26. t. 38. *Raj. suppl.* 52.
- Phyllitis folio longo & marginato*. *Pet. fil.* 5. t. 6. f. 10.
- 7 *Phyl-*

Phyllitis folio non sinuato tenuiori, rotundis pulverulentis maculis averſa parte punctato. *Sloan. jam.* 14. *biſt.* 1. p. 72.

Habitat in America meridionali.

Conſer. Phyllitis folio longo leviter undulato. *Pet. fil.* 4. t. 6. f. 9.

Phyllitis, folio longo, anguſtiſolia. *Pet. fil.* 7. t. 6. f. 3..

compoſitum.

7. POLYPODIUM frondibus lanceolatis glabris integerrimis; apice multifido, fructificationibus ſparſis.

Lingva cervina, multifido cacumine, laciniata. *Plum. fil.* 115. 2. 131.

Phyllitis folio, longo apice palmato. *Plum. fil.* 6. t. 6. f. 11.

Habitat in America meridionali.

Suſpecta affinitatis eſt cum precedenti.

trifurcatum.

8. POLYPODIUM frondibus lanceolatis glabris repandoſinuatis: apice trilobis.

Lingva Cervina ſinuosa in ſummitate trifulca. *Plum. fil.* 120. t. 138.

Polypodium vulgare, ſummo trifido. *Pet. fil.* 34. t. 15. f. 3.

Habitat in America meridionali.

Suſpecta etiam eſt ob affinitatem cum proxime precedentibus duabus.

* *Fronde pinnatifida lobis coadunatis.*

triſpatum.

9. POLYPODIUM frondibus pinnatifidis glabris: lobis ſemiorbiculatis crenatis.

Polypodium, aſplenii folio criſpo, pendulum. *Pet. fil.* 25. t. 13. f. 12.

Habitat in America meridionali.

ſuſpenſum.

10. POLYPODIUM frondibus pinnatifidis glabris: lobis ſemiovatis acutis.

Polypodium pendulum glabrum. *Plum. fil.* 67. t. 87.

Polypodium, aſplenii folio brevi, pendulum. *Pet. fil.* 24. t. 1. f. 15.

Habitat in America meridionali.

aſplenioſiſſimum.

11. POLYPODIUM frondibus pinnatifidis pilosis: lobis ſemiovatis acutis.

Polypodium aſplenii folio villoſo. *Pet. fil.* 26. t. 7. f. 16.

β. Poly-

- β. Polypodium pendulum** subtus villosum. *Pet. fil.* 35.
t. 14. f. 1.
Polypodium pendulum hirsutie rufa pubescens. *Plum.*
fil. 68. *t. 88.*
Habitat in America meridionali.
Affinitas hujus cum antecedenti suspecta est.
- 12. POLYPODIUM** frondibus serrato-pinnatifidis lan- *scolopendri-*
ceolatis. oides.
Polypodium incisuris asplenii. *Plum. fil.* *t. 91.*
Polypodium minus, asplenii folio. *Pet. fil.* 23. *t. 1. f.*
14.
Filix non ramosa scolopendrioides. *Plum. amer.* 7. *t. 11.*
Habitat in Jamaica.
- 13. POLYPODIUM** frondibus pinnatifidis: pinnis ob- *vulgare.*
longis subserratis obtusis, radice squamata.
Polypodium fronde pinnata: pinnis lanceolatis indivisis
serruatis alternis connato-lesilibus. Hort. cliff. 475.
Fl. succ. 845. *Roy. lugdb.* 499. *Dalib. parisi.* 311.
Mat. med. 485.
Polypodium pinnatum: pinnis lanceolatis integris. *Fl.*
lapp. 380.
Polypodium vulgare. *Banh. pin.* 359. *fil.* 27. *t. A. f.*
2.
Polypodium majus. *Dod. pempt.* 464.
β. Polypodium minus. *Banh. pin.* 359. *Dod. pempt.* 464.
Habitat in Europæ rimis rupium.
- 14. POLYPODIUM** frondibus pinnatifidis: pinnis ob- *virginia-*
longis subserratis obtusis, radice lævi. num.
Polypodium minus vulgaris facie. *Pet. fil.* 22. *t. 7. f.*
9.
Polypodium radice tenui & repente. *Plum. amer.* 25.
t. 36. fil. 60. *t. 77. Raj. suppl.* 56.
Polypodium virginense minus, foliis obtusioribus. *Mo-*
rif. hist. 3. *p.* 563. *f.* 14. *t. 2. f.* 3.
Habitat in Virginia.
Antecedenti simillima, sed minor, & subtus glabra.
- 15. POLYPODIUM** frondibus pinnatifidis: lobis lance- *Ontar.*
olatis alternis obtusis distantibus.
Polypodium, lonchitidis folio, minus. *Pet. fil.* 32. *t. 1.*
f. 16.
Habitat in America.
- 17. POLYPODIUM** frondibus pinnatis lanceolatis: lo- *postinatum*
bis

bis approximatis ensiformibus parallelis acutis horizontalibus, radice nuda.

Polypodium lonchitidis folio, *Pet. fil.* 31. t. 7. f. 14.

Polypodium nigrum tenerius sectum. *Plum. amer.* 26. t. 37.

Habitat in Jamaica.

taxifolium. 17. **POLYPODIUM** frondibus pinnatis: lobis approximatis ensiformibus parallelis acutis adscendentibus, radice hirta.

Polypodium taxi folio pendulum. *Pet. fil.* 35. t. 14. f. 1.

Polypodium tenue & pendulum. *Plum. amer.* 69. t. 89.

Habitat in America meridionali.

struthionis. 18. **POLYPODIUM** frondibus pinnatis: lobis approximatis ensiformibus repandis horizontalibus.

Polypodium fronde pinnata: foliolis lanceolato-linearibus undulis acuminatis oppositis connatis. *Hort. cliff.* 475.

Polypodium crispum struthionis pennis. *Pet. fil.* 30. t. 3. f. 8.

Polypodium radice subcærulea & punctata. *Plum. fil.* 60. t. 78.

Habitat in America.

squamatum. 19. **POLYPODIUM** frondibus pinnatifidis scabris: pinnis lanceolatis distantibus horizontalibus integerrimis.

Polypodium longifolium, squamulis argenteis. *Pet. fil.* 29. t. 7. f. 11.

Habitat - - - - -

loricatum. 20. **POLYPODIUM** frondibus pinnatifidis lævibus: pinnis lanceolatis distantibus horizontalibus repandis.

Polypodium longifolium, radice dulci. *Pet. fil.* 27. t. 7. f. 10. 12.

Habitat in America.

aiatum. 21. **POLYPODIUM** frondibus pinnatifidis lævibus: pinnis oblongis distantibus dentatis.

Polypodium serratum majus cotta aiata. *Pet. fil.* 37. t. 7. f. 13.

Habitat in America.

cambicum. 22. **POLYPODIUM** frondibus pinnatifidis: pinnis lanceolatis lacero-pinnatifidis serratis.

Poly-

- Polypodium fronde pinnatifida:** foliolis lanceolatis sinuato-pinnatifidis. *Hors. cliff.* 475. *Rov. lugdb.* 499.
- Polypodium cambrobritannicum,** lobis foliorum profunde dentatis. *Morif. hist.* 3. p. 565. f. 14. t. 2. f. 8. bene.
- Polypodium cambro-britannicum,** pinnulis ad margines laciniatis. *Raj. hist.* 137. *angl.* 3. p. 117.
- Filix amplissima,** lobis foliorum laciniatis, cambrica. *Pluk. alm.* 153. t. 30. f. 1.
- Habitat in Anglia.*
23. **POLYPODIUM** frondibus pinnatifidis lævibus: pin- *aureum.*
nis oblongis distantibus: intimis patulis; terminali
maxima.
- Polypodium fronde pinnata:** pinnis lanceolatis integer-
rimis basi connatis patulis: terminali maxima. *Hors.*
cliff. 475.
- Polypodium majus aureum.** *Plum. amer.* 25. t. 25.
fil. 59. t. 76. *Raj. suppl.* 56.
- Polypodium maximum.** *Pet. fil.* 15. t. 7. f. 8.
- Polypodium altissimum.** *Sloan. jam.* 15. *hist.* 1. p. 75.
- Polypodium jamaicense** majus & elatius, alis longiori-
bus: punctis aureis averſa parte notatis. *Morif. hist.* 3. p.
563.
- Habitat in America ad caudices veterum arborum.*
24. **POLYPODIUM** frondibus sterilibus brevioribus *quercifol-*
obtusis sinuatis; fructificantibus alternis pinnatis lan- *um.*
ceolatis. *Fl. zeyl.* 382.
- Polypodium exoticum,** folio quercus. *Baub. pin.* 359.
Morif. hist. 3. p. 564. f. 14. t. 1. f. 15.
- Polypodium malabaricum,** foliis biformibus. *Pet. mus.*
794. *Raj. suppl.* 57.
- Polypodium indicum.** *Clus. exot.* 88. t. 89. *Pis. mont.*
195.
- Panna-kelengo-marava.** *Rheed. mal.* 12. p. 23. t. 11.
- Habitat in India.*
- * *Fronde trifoliata: pedunculo foliolis tribus.*
25. **POLYPODIUM** fronde ternata sinuato-lobata: in- *trifoliatum.*
termedia majora.
- Polypodium fronde ternata sinuata:** foliolis lateralibus
bifidis; intermedio petiolato trifido majore. *Hors. cliff.*
475.
- Polypodium trifoliatum.** *Pet. fil.* 18. t. 7. f. 3
- Lonchitis maxima trifolia americana.** *Raj. suppl.* 66.
- Hemic-

Hemionitis maxima trifolia. *Plum. amer.* 22. t. 32. fil. 127. t. 148.

Hemionitidi affinis Filix major trifida auriculata: pinnulis latissimis sinuatis. *Sloan. jam.* 18. *bist.* 1. p. 85. t. 42.

Filix f. Hemionitis dicta caribæorum, amplissimis foliis, trifoliata. *Pluk. alm.* 155. t. 291. f. 3.

Habitat in Caribæis insulis.

Confer. Hemionitis maxima quinquefolia. *Plum. Raj. suppl.* 67. *Pet. fil.* 148. t. 7. f. 2.

Polypodium auriculatum. *Pet. fil.* 16. t. 7. f. 5.

Polypodium auriculatum pentaphylloides. *Pet. fil.* 17. t. 7. f. 6.

Polypodium auritum majus: laciniis crispis. *Pet. fil.* 19. t. 7. f. 7.

Polypodium auritum tricuspidatum: oris pilosis. *Pet. fil.* 20. t. 7. f. 4.

* *Fronde pinnata simpliciter.*

Len. hick. 26. **POLYPODIUM** fronde pinnata: pinnis lunulatis ciliato-ferratis declinatis, petiolis strigosis. *Horr. cliff.* 475. *Gron. virg.* 123.

Lonchitis aspera. *Baub. pin.* 359.

Lonchitis aspera major. *Morif. bist.* 3. p. 566. f. 14. t. 2. f. 1.

Habitat in Alpibus Helvetiæ, Baldi, Arvonix, Montpelii, Virginix.

Virginica plantæ pinnae magis elongata & minus incurvata, serraturis minus ciliato-spinosis.

auriculatum 27. **POLYPODIUM** fronde pinnata: pinnis falcato-lanceolatis ferratis: basi truncatis sursum auritis. *Fl. zeyl.* 383.

Filix zeylanica, lonchitidis facie. *Breyn. cent.* 189. t. 44. f. 2. *Raj. suppl.* 69. *Burm. zeyl.* 98. t. 44. f. 2.

Filix f. Lonchitis aspera maxima indorum. *Pluk. alm.* 152. t. 30. f. 4.

Lonchitis aspera zeylanica, alis falcatis levissime denticulatis & superiore latere auriculatis. *Morif. bist.* 3. p. 566.

Habitat in India.

triangulum 28. **POLYPODIUM** frondibus pinnatis: pinnis triangularibus dentatis.

Trichomanes folio triangulo dentato. *Pet. fil.* 76. t. 1. f. 10.

Habitat in America.

29. **PO-**

29. **POLYPODIUM** frondibus pinnatis: pinnis corda- *cordifolium*.
tis obtusis integerrimis repandis.
Trichomanes folio cordato. Pet. fil. 75. t. 1. f. 11.
Habitat in America.
- * *Fronde sub-bipinnata, cujus pinnae confluent basi, ut semi-bipinnata potius, quam perfecte duplicato-pinnata sit.*
30. **POLYPODIUM** frondibus sub-bipinnatis: foliolis in- *Phegopteris*.
fimis reflexis: paribus pinnula quadrangulati coaduna-
tis. *Hors. cliff. 475. Fl. suec. 851.*
Polypodium pinnatum: pinnis lanceolatis pinnatifidis in-
tegris: inferioribus nutantibus. Fl. lapp. 367. Hall. helv.
138.
Filix non ramosa minor & sylvestris. Pet. epist. 56 t.
50.
Filix minor britannica, pediculo pallidiore, alis inferioribus deorsum spectantibus. Moris. hist. 3. p. 575. f.
14. t. 4. f. 17.
Filix virginiana non ramosa, foliis planis. Pluk. alm.
151. t. 284. f. 2.
Habitat in Europae sagetis & in Virginia.
31. **POLYPODIUM** frondibus sub-bipinnatis: foliolis *retroflexum*.
plurimis reflexis: pinnis laceris.
Filicula alis inferioribus pendulis. Pet. fil. 72. t. 1.
f. 9.
Habitat in America.
32. **POLYPODIUM** frondibus sub-bipinnatis lanceola- *fragrans*.
tis: foliolis confertis: lobis obtusis serratis, stipite paleaceo.
Dryopteris rubrum idæum spirans. Amm. ruth. 251.
Habitat in Sibiria.
Habitus P. F. maris, at longe minor. Foliola densius congesta, lobis lateralibus obtusis, profundius serratis.
33. **POLYPODIUM** fronde sub-bipinnata lanceolata: *fontanum*.
foliolis subrotundis, stipite lævi.
Filicula fontana minor. Baux. pin. 358.
Habitat in Sibiria, Galloprovincia.
Habitus P. fragilis, at foliola ætiora, minus profunde subdivisa, punctis floriferis majoribus, prodeuntibus non ex squama subrotunda, sed ex rima f. valvula lineari, oblonga, alba.

- parafitum.* 34. **POLYPODIUM** fronde semi-bipinnata lanceolata: lobulis rotundatis integerrimis striatis.
Kari-welli-panna. Rheed. mal. 12. p. 35. t. 17.
Habitat in India supra arbores. Osbeck. 41.
Stipes laevis. Foliola lanceolata ad medium pinnatifida, lobis obtusis non serratis substriatis, punctis fronsificantibus sparsis.
- varium.* 35. **POLYPODIUM** fronde lateribus bipinnata: foliolo infimo pinnatifido.
Habitat in China. Osbeck.
Stipes hirtus pilis vagis. Frons pinnata: Foliolis lateralibus ad basin pinnatifidis: Pinnæ omnes sessiles argute & tenuissime serrata. Pinnulae baseos deorsum spectantes juxta stipitem, longiores & semipinnatifida, qua nota facillime innotescit. Sic frons apice simpliciter pinnata, lateribus bipinnata, pinnula baseos elongata, pinnatifida.
- crissatum.* 36. **POLYPODIUM** frondibus sub-bipinnatis: foliis ovato-oblongis: pinnis obtusiusculis apice acute serratis. *Polypodium* fronde duplicato-pinnata: pinnulis obtusis coadunatis crenatis. *Fl. succ. 848.*
Filix non ramosa major pinnulis latiusculis longioribus in profundos denticulos divisilis. Moris. hist. 3. f. 14. t. 3. f. 11.
Habitat in Europa septentrionali.
Florent hinc foliola superiora, non inferiora.
- F. mas.* 37. **POLYPODIUM** fronde bipinnata: pinnis obtusis crenulatis, stipite paleaceo. *Hort. cliff. 475. Fl. succ. 846. Mat. med. 486. Roy. lugdb. 499. Hall. belv. 139. Dalib. parif. 313.*
Filix mas non ramosa dentata. Baub. pin. 358.
Filix mas. Fuchf. hist. 594.
Habitat in Europæ sylvis.
- F. femina.* 38. **POLYPODIUM** fronde bipinnata: pinnulis lanceolatis pinnatifidis acutis. *Fl. lapp. 386. Fl. succ. 847. Roy. lugdb. 499. Hall. belv. 140. Dalib. parif. 313.*
Filix mas non ramosa, pinnulis angustis rarioribus profunde dentatis. Moris. hist. 3. p. 579. J. 14. t. 3. f. 3. Pluk. phyt. 180. f. 4.
Habitat in Europæ frigidioris subhumidis.
- aculeatum.* 39. **POLYPODIUM** fronde bipinnata: pinnis lunulatis dentatis, stipite trigoso. *Roy. lugdb. 500. Dalib. parif. 314.*
Filix

CRYPTOGAMIA FILICES. 1091

Filix aculeata major. *Baub. pin.* 358. *prodr.* 151.

Habitat in Europa.

40. POLYPODIUM fronde bipinnata: foliolis pinnis-*rhæticum*.
que remotis lanceolatis: serraturis acuminatis.

Filix rhætica tenuissime denticulata. *Baub. hist.* 3. p. 740.

Filicula fontana major f. *Adiantum album*, filicis folio.

Baub. pin. 151.

Habitat in Gallia, Helvetia.

41. POLYPODIUM fronde bipinnata: pinnis oblongis *novaboracense*.
integerrimis parallelis, stipite lævi.

Habitat in Canada. *Kalm.*

Habitus & structura omnino P. Filicis maris, at pinna magis approximata, integerrima & punctis floris minimis copiosissimis; stipites dein laevis est.

42. POLYPODIUM fronde bipinnata: pinnis basi sinu-*marginale*.
ato-repandis, fructificationibus marginalibus.

Habitat in Canada. *Kalm.*

Habitus P. F. maris, sed pinna oblonga non crenata, at versus basin obtuse serrata; manifeste distinguitur pinnis ab utroque margine subtus punctis 2. f. 3. majoribus, distinctis, nec pluribus in disco uti F. maris.

43. POLYPODIUM frondibus bipinnatis: foliolis re-*bulbiferum*.
motis: pinnis oblongis obtusis serratis subtus bulbiferis.

Polypodium fronde duplicato-pinnata, foliolis oblongis obtusis incisis dorso bulbiferis. *Roy. lugdb.* 500.

Filix saxatilis canadensium globulifera. *Pluk. alm.* 150.

Filix baccifera. *Corn. canad.* 5. *Morif. hist.* 3. p. 579.

f. 14. t. 3. f. 10.

Habitat in Canada.

44. POLYPODIUM fronde bipinnata: foliolis remotis: *F. fragile*.
pinnis subrotundis incisis. *Fl. lapp.* 384. *Fl. suec.* 849.

Roy. lugdb. 500. *Dalib. parif.* 312.

Filix saxatilis, cauliculo tenui fragili. *Pluk. alm.* 150. t. 180. f. 5.

Filix pumila saxatilis. 2. *Clus. pen.* 706.

Filix saxatilis non ramosa nigris maculis punctata. *Baub. pin.* 358?

Habitat in collibus Europæ frigidioris.

45. POLYPODIUM fronde bipinnata: foliolis suboppo-*regium*.
sitis: *Z z z z*

sitis : pinnis alternis laciniatis. *Hort. cliff.* 475. *Roy. lugdb.* 500. *Dalib. parif.* 312.

Filix regia, fumarie pinnulis. *Vaill. parif.* 52. t. 9. f. 1.
Habitat in Gallia.

Fructificationum pulvilli in hoc majores, quam in affinis.

- leptophyllum.* 46. **POLYPODIUM?** frondibus bipinnatis : sterilibus brevissimis : pinnis cuneiformibus lobatis. *Magn. monsp.* 5. t. 5.

Adiantum minimum, folio vario Magn. monsp. 5. t. 5.

Adiantum filicinum leptophyllum elatius hispanicum Barr. rar. 1270. t. 431. *ops.*

Habitat in Hispania, Lusitania, Galloprovincia.

- Savoneta.* 47. **POLYPODIUM?** frondibus bipinnatis : pinnis pinatifidis lanceolatis serratis, radicibus lanatis.

Habitat in China, unde habui cum radice.

Radix decumbens, crassa, vellere molliissimo. densissimo intense flavo obvestita. Stipites pedales f. sesquipedales, virides, laeves, obsolete angulati. Frons bipinnata: Foliolis oblongis infimis oppositis; reliquis alternis, axillis patulis. Pinnæ lanceolatae, acutae, pinnatifide lobis acutis apice serratis. Sunt hæ frondes glabrae, nec fructificationem obtinui, sed ex facie huc resuli.

* *Spinis sparsis aculeata aut arborescentia.*

- arborescens.* 48. **POLYPODIUM** frondibus bipinnatis ferratis, caudice arboreo inermi.

Filix arborescens, pinnulis dentatis. Plum. fil. 1. t. 1. *amer.* 1. t. 1. 2.

Filix arborescens, pinnis dentatis. Pet. fil. 41. t. 1. f. 1. 2.

Habitat in America meridionali.

- spinosum.* 49. **POLYPODIUM** frondibus bipinnatis ferratis, caudice arboreo aculeato.

Filix arborescens latifolia aculeata. Plum. amer. 3. t. 3. *fil.* 3. t. 3.

Filix arborescens, pinnis latis, caudice spinoso. Pet. fil. 42. t. 4. f. 1.

Habitat in America meridionali.

- horridum.* 50. **POLYPODIUM** frondibus supradecompositis integerrimis semi-fagittatis, caudice aculeato.

Filix ramosa pinnis latis gladiatis, caudice spinoso. Pet. fil. 50. t. 5. f. 1.

Filix latifolia ramosa, cauliculis nigris & spinosis. Plum. amer. 3. t. 4. *Kaj. suppl.* 92.

Habitat in America meridionali.

51. PO-

51. **POLYPODIUM** fronde supradecomposita: pinnis *pyramidale*, terminalibus lanceolatis longissimis serratis, stipite inferne aculeato.
 Filix ramosa pyramidalis, pinnis parvis. *Pet. fil.* 40. t. 4. f. 2.
Habitat in America.
52. **POLYPODIUM** fronde supradecomposita: pinnis *asperum*, obtusis apice serratis: terminalibus acuminatis, caudice arboreo aculeato.
 Filix arborescens, caudice & caule spinosis. *Pet. fil.* 47. t. 4. f. 7.
Habitat in America.
53. **POLYPODIUM** fronde bipinnata: pinnis ovatis *innervatum*, denticulato-spinosis.
 Filix pinnis aculeatis. *Pet. fil.* 53. t. 1. f. 6.
Habitat in America.
54. **POLYPODIUM** fronde bipinnata hirsuta: pinnis ob- *villosum*, longis obtusis: terminalibus acuminatis.
 Filix dentata, caule & costa villosis. *Pet. fil.* 52. t. 4. f. 10.
 Filix ramosa villosa major, crenis rotundis dentata. *Plum. amer.* 15. t. 23. *fil.* 21. t. 27. *Raj. suppl.* 97.
 Filix ramosa major hirsuta, ramulis raris, pinnulis asplenii crebris latis brevibus subrotundis non dentatis. *Sloan. jam.* 23. *bist.* 1. p. 100.
Habitat in America.
55. **POLYPODIUM** fronde bipinnata: pinnis horizon- *detussum*, talibus integerrimis obtusis: terminalibus lanceolatis.
 Filix, taxifoliis, major & minor. *Pet. fil.* 61. 62. t. 2. f. 5. 6.
Habitat in India.
- * *Fronde supradecomposita.*
56. **POLYPODIUM** fronde supradecomposita: foliolis *Dryopteris*, ternis bipinnatis. *Fl. suec.* 852. *Dalib. parif.* 314.
 Polypodium trifidum, ramis pinnatis: pinnis pinnatifidis. *Fl. lopp.* 387.
 Filix ramosa minor, pinnulis dentatis. *Bauh. pin.* 358.
 Filix pumila saxatilis. *Clus. bist.* 2. p. 212.
Habitat in Europæ nemoribus.
57. **POLYPODIUM** fronde supradecomposita pilosa: *spolunear*, foliis lanceolatis pinnatis: pinnis oppositis pinnatifidis.
Fl. zeyl. 384. Z z z 3 Filix

Filix bermudensis elegans ramosa, pinnis rarioribus dentatis, cauliculis mucosa lanugine obductis. Pluk. alm.

155. t. 244. f. 2.

Habitat in Indiis.

Lusitanicum. 58. **POLYPODIUM** fronde supradecomposita: foliolis alternis: pinnis oblongis longitudinaliter pinnatifidis.

Roy. lugd. 500.

Filix lusitanica, polypodii radice. Magn. hors. 79. t. 79.

Habitat in Lusitanica.

12 ADIANTUM.

* *Fronde simplici.*

reniforme. 1. **ADIANTUM** frondibus reniformibus simplicibus stipitatis multifloris.

Adiantum monophyllum, foliis asari rotundis. Moris. hist. 3. p. 592.

Filix Hemionitis dicta maderensis, foliis asari rotundioribus, crenarum segmentis oblongo quadratis, per ambitum circumcirca reflexis. Pluk. alm. 155. t. 287. f. 5.

Hemionitis asari folio. Sloan. jam. 14. hist. 1. p. 15.

Habitat in Madera.

philippense. 2. **ADIANTUM** frondibus reniformibus simplicibus alternis petiolatis lobatis multifloris.

Adiantum philippense, folio rotundo laciniato. Pet. gaz. 8. t. 4. f. 4.

Habitat in Philippinis.

* *Fronde composita.*

radiatum. 3. **ADIANTUM** fronde digitata: foliolis pinnatis: pinnis unifloris.

Adiantum alis radiatis. Pet. fil. 93. t. 14. f. 3.

Adiantum ramosum radiatum. Plum. amer. 33. t. 49.

Adiantum nigrum majus non ramosum, surculis e pedunculi summitate tanquam e centro stellæ in modum radiatis. Sloan. jam. 22. hist. 1. p. 95.

Adiantum jamaicense, pinnulis auriculatis, ramosum quinque ramulis ex eodem caulis puncto expansis. Pluk. alm. 11. t. 253. f. 3.

Lonchitis radiata, polytrichi facie. Plum. fil. 53. t. 100.

Trichomanes americanum radiatum. Moris. hist. 3. p. 590. f. 14. t. 4. f. 9.

Habitat in Jamaica, Domingo.

4. ADI-

4. **ADIANTUM** fronde pedata: foliolis pinnatis: pin- *pedatum.*
nis antice gibbis incisīs fructificantibus.
Adiantum fronde bifida; foliolis alternis pinnatis: pinnis
obtusis recurvis superne incisīs multifloris. *Roy. lugdb.*
503.
Adiantum fronde supradecomposita bipartita, foliis par-
tialibus alternis: foliolis trapeziformibus. *Gron. virg.*
123.
Adiantum fruticosum americanum, summis ramulis re-
flexis & in orbem expansis. *Pluk. alm.* 10. t. 124. f.
2. *Morif. hist.* 3. p. 588. f. 14. 1. 5. f. 12.
Adiantum fruticosum brasilianum. *Bank. pin.* 355.
prodr. 150.
Adiantum americanum. *Corn. canad.* 7. t. 6. *Raj. hist.* 148.
Habitat in Canada, Virginia. &
5. **ADIANTUM** fronde composita: pinnis tripartitis ob- *trilobum.*
tusis incisīs multifloris.
Adiantum triphyllum repens. *Pet. fil.* 100. t. 11. f. 9.
Habitat in America.
6. **ADIANTUM** fronde pinnata: pinnis obtusis obova- *serrulatum.*
tis serratis unifloris.
Trichomanes majus nigrum, pinnis leviter dentatis tra-
pezii figura. *Sloan. jam.* 17. t. 35. f. 2.
Habitat in Jamaica.
* *Fronde decomposita.*
7. **ADIANTUM** frondibus decompositis: pinnis alter- *flabellu-*
nis rhombeo rotundatis multifloris, pedunculis supra *tum.*
pubescentibus.
Adiantum chinense perelegans ramosum, folio flabel-
liformi cum rubedine perfuso. *Pluk. alm.* 11. t. 4.
f. 3.
Adiantum, foliis coriandri, ramosum. *Raj. hist.* 1853.
Habitat in China. *Osbeck.*
Stipes rami & pedicelli supra canaliculati & ferrugi-
neo villosi.
8. **ADIANTUM** fronde decomposita, foliis alternis ter- *trifoliatum.*
natis linearibus unifloris.
Adiantum clavatum triphyllum *Pet. fil.* 99. t. 11. f. 4.
Habitat in America.
9. **ADIANTUM** fronde decomposita: pinnis alternatis *obusatum*
pinnatifidis: lobis inæqualibus.
Habitat in China.

Z z z 4 .

Frons

Fons lanceolata, glabra. Foliola alterna, lanceolata; alternatim pinnata. Pinnæ alternatim pinnatifide: lobis linearis-cuneiformibus, incis.

- Capillus ve-* 10. **ADIANTUM** frondibus decompositis: foliis alternis: pinnis cupeiformibus lobatis pedicellatis.
neris. Adiantum ramis pinnatis, foliis circuli sectorem referentibus. *Sauv. monsp.* 245.
 Adiantum foliis coriandri. *Baub. pin.* 356. *Mat. med.* 480.
 Adiantum. *Cam. epit.* 924.
Habitat in Europa australi.

- pulverulentum.* 11. **ADIANTUM** frondibus bipinnatis: pinnis ovalibus antice truncatis unifloris, caule hirtio.
 Adiantum mas caule squamoso & pediculo pulverulentum. *Pet. fil.* 84. *t.* 5. *f.* 7.
 Adiantum nigrum ramosum pulverulentum & falcatum. *Plum. fil.* 32. *t.* 47. *fil.* *t.* 55. *Raj. suppl.* 100.
Habitat in America.

* *Fronde supradecomposita.*

- clavatum.* 12. **ADIANTUM** fronde supradecomposita: foliolis alternis: pinnis cuneiformibus integerrimis alternis unifloris.
 Adiantum fronde ramosissima divisa: foliolis linearibus retulis inferne attenuatis. *Hort. cliff.* 474.
 Adiantum clavis simplicibus. *Pet. fil.* 95. *t.* 14. *f.* 7.
 Adiantum minus, foliis in summitate retulis. *Plum. amer.* 34. *t.* 50. *f.* B.
 Filicula pinnis obtulis. *Plum. fil.* 101. *f.* B.
Habitat in Dominica.

- aculeatum.* 13. **ADIANTUM** frondibus supradecompositis: pinnis palmatis multifloris, caule aculeato.
 Adiantum caule fruticoso supra-decomposito aculeato, frondibus palmatis. *Hort. cliff.* 473.
 Adiantum frutescens spinosum & repens. *Plum. fil.* 77. *t.* 94.
 Adiantum repens spinosum. *Pet. fil.* 78. *t.* 11. *f.* 6.
 Filix ramosa major, caule spinoso, pinnulis rotundis profunde laciniatis, cerefolii foliis. *Sloan. jam.* 23. *biff.* 1. *p.* 99. *t.* 61.
Habitat in Jamaica, Dominica. 5

14. ADI-

14. **ADIANTUM** frondibus supradecompositis: foliis alternis: pinnis rhombeis incisus utrinque fructificationibus. *trapeziforme.*

Adiantum fronde alternatim supradecomposita: foliolis trapeziformibus acuminatis inciso-crenatis. *Hort. cliff.* 473. *

Adiantum ramosius, pediculis lucidis & nigris. *Plum. fil.* 78. t. 95.

Adiantum nigrum ramosum maximum, foliis magnis trapezii in modum figuratis. *Sloan. jam.* 23. *bist.* 1. p. 98. t. 59. *bona.*

Adiantum fruticosum, coriandri, folio, jamaicense, pediculis foliorum politiore nitore nigricantibus. *Pluk. alm.* 10. t. 254. f. 1.

Habitat in America meridionali.

15. **ADIANTUM** frondibus supradecompositis: pinnis hexagonis. sexangularibus emarginatis integerrimis utrinque unifloris.

Adiantum pinnis hexagonis furcatis. *Pet. fil.* 94. t. 10. f. 2.

Habitat in America.

TRICHOMANES.

* *Fonde simplici.*

1. **TRICHOMANES** fronde simplici oblonga lacera. *membrana-
centra.*

Darea americana lichenoides. *Pet. mus.* 763.

Phyllitis scandens minima, musci facie, foliis membranaceis subrotundis. *Sloan. jam.* 15. *bist.* 1. p. 74. t. 27. f. 1. *Morif. bist.* 13. p. 558. *Raj. suppl.* 53.

Habitat in America.

Confer *Adiantum* membranaceum lichenoides. *Pet. fil.* 101. t. 13. f. 5. *Filix* *Hemionitis* *Lichenoides americana* auricularis *cæsalpini* æmula, radice repente. *Pluk. phyt.* 285. f. 3.

2. **TRICHOMANES** frondibus pinnatifidis lanceolatis: cristis pinnis parallelis subserratis.

Trichomanes fronde lanceolato-pinnata: pinnis sessilibus parallelis subserratis. *Hort. cliff.* 476.

Polypodium cristatum calyciferum. *Plum. fil.* 67. t. 86.

Darea filicis maris facie. *Pet. fil.* 103. t. 11. f. 8.

Habitat in Martinica.

Z z z 5

3. TRI-

- polypodiell-* 3. TRICHOMANES fronde lanceolata pinnatifida repanda, floribus solitariis terminalibus.
Habitat in India.
 Frons digiti longitudine & latitudine, pinnatifida ferme ad costam, lobis lanceolatis, obtuse serratis, apice acutis, & unico flore in singulis lobis apice instructis.
 Frons insuper pellucida venis alternis.
- * *Fronde composita.*
- hirsutum.* 4. TRICHOMANES frondibus pinnatis: pinnis alternis pinnatifidis pilosis.
 Filicula digitata. *Plum. spec.* 13. *fil.* 1. 50. t. B.
 Darea americana major pubescens. *Pet. mus.* 762.
 Darea atkinsii digitata. *Pet. fil.* 105. t. 15. f. 5.
Habitat in America.
- pyxidifera* 5. TRICHOMANES frondibus sub-bipinnatis: pinnis alternis confertis lobatis linearibus.
 Filix pyxidifera. *Plum. fil.* 74. t. 50.
 Darea tunbrigensis facie. *Pet. fil.* 104. t. 13. f. 13.
Habitat in America.
- tunbrigensis.* 6. TRICHOMANES frondibus pinnatis: pinnis lobatis oblongis crenulatis.
 Adiantum petraeum perpallidum anglicum, foliis bifidis trifidisque. *Raj. angl.* 3. p. 123. *suppl.* 77.
 Adiantum radicosum humifparsum f. Filicula pellucida nostras, coriandri foliis mollicellis, globuliferum. *Pluk. alm.* 10. t. 3. f. 5. & 6.
 Darea tunbrigensis minor. *Pet. mus.* 761.
 Adiantum montanum italicum, adianti foliis. *Bocc. mus.* 2. p. 24. t. 2. f. 1.
Habitat in Anglia, Italia.
- adiantoides.* 7. TRICHOMANES frondibus pinnatis: pinnis ensiformibus acuminatis inciso-serratis: serraturis bifidis.
Fl. zeyl. 385.
 Adiantum africanum rutæ murariæ æmulum, segmentis longioribus acutis. *Pluk. alm.* 10. t. 123. f. 6.
 Filix non ramosa zeylanica, foliis adianti in modum serratis. *Burm. zeyl.* 97. t. 43.
Habitat in India, Africa.
- * *Fronde supradecomposita.*
- stans.* 8. TRICHOMANES frondibus supradecompositis: foliolis alternis: pinnis alternis oblongis serratis. *Adi-*

Adiantum scandens ramosissimum, laciniis retusis dissectum. *Plum. fil.* 76. t. 93.

Darcia ramosissima scandens. *Pet. fil.* 102. t. 22. f. 5
Habitat in America.

9. TRICHOMANES fronde supra-decomposita, foliis ^{chinensis} pinnisque alternis lanceolatis: foliolis laciniis cuneiformibus.

Habitat in China. Osbeck.

Radix vix villosa. Stipes *levis*, antice subcanaliculatus sed margine obtuso, nec marginato ut in sequenti. Frons lanceolata, pedalis ex foliis alternis, lanceolatis. Folia alterna, alternatim pinnata. Pinnæ alternatim pinnatifidæ segmentis cuneiformibus. alternis, sapinus bifloris.

10. TRICHOMANES fronde supradecomposita tripar-^{canariensis} tita: foliolis alternis: pinnis alternis pinnatifidis.

Frons ramosa canariensis, rutæ murariæ pinnulis angustis altius incisus mediæ costæ alternatim alligatis. *Pluk. alm.* 156. t. 291. f. 2.

Habitat in Canariis, Lusitania, ad latera montium.

11. TRICHOMANES fronde supradecomposita: pin-^{capillaceum} nis filiformibus linearibus unifloris.

Adiantum capillaceum. *Plum. fil.* 83. t. 99. *Pet. fil.* 96. t. 10. f. 7.

Habitat in America.

MARSILEA.

1. MARSILEA foliis oppositis simplicibus.

Salvinia vulgaris aquis innatans, foliis subrotundis punctatis læte virentibus. *Mich. gen.* 107. t. 58.

Lenticula palustris latifolia punctata. *Bauh. pin.* 362.

Lens palustris patavina. *Bauh. hist.* 3. p. 785.

Habitat in Italiæ fossis paludosis stagnantibus lente fluentibus.

2. MARSILEA foliis quaternatis.

Lemma. *Juss. art.* 1740. p. 263. *Guss. stamp.* 1. p. 62.

Lenticula palustris quadrifolia. *Bauh. pin.* 362. *Mapp. albat.* 166. t. 166.

Lens palustris quadrifolia. *Bauh. hist.* 3. p. 789. *Morif. hist.* 3. p. 619. f. 15. t. 4. f. 5.

Lens

Lens palustris altera. Cam. epit. 853.

Habitat in Indiæ, Sibiriz, Galliæ, Alsatiz fossis.

Genus distinguam, dum precedens (1) fueris examinata.

PILULARIA.

globulifera. 1. PILULARIA.

Pilularia palustris juncifolia. Vail. parif. 159. t. 15. f. 6. Dill. musc. 538. t. 79. f. 1. Juss. act. 1739. p. 240. t. 11.

Graminifolia palustris repens, vasculis granorum piperis æmulis. Ray. hist. 1325. angl. 3. p. 136. Moris. hist. 3. p. 608. f. 15. t. 7. f. 49.

Gramen piperinum. Pet. herb. t. 19. f. 8.

Muscus aureus capillaris palustris, inter folia folliculis rotundis quadripartitis. Pluk. alm. 246. t. 48. f. 1.

Habitat in Europæ inundatis.

ISOËTES.

lacustris. 2. ISOËTES. It. jean. 417. t. 419. Nov. gen. 1109.

Marsilea foliis subulatis semicylindricis articulatis. Fl. suec. 996.

Calamaria folio longiore & graciliore. Dill. musc. 541. t. 80. f. 2.

Subularia lacustris f. Calamistrum herba aquatica alpina.

Ray. angl. 1. p. 210. t. 210.

Habitat in Europæ frigida fundo lacuum.

MUSCI.

LYCOPODIUM.

linifolium. 1. LYCOPODIUM foliis alternis remotis lanceolatis, floribus axillaribus.

Selago linearis foliis. Dill. musc. 440. t. 57. f. 5.

Muscus maximus, linearis foliis. Plum. fil. 144. t. 166. f. C. b.

Habitat in America meridionali.

nudum. 2. LYCOPODIUM foliis oppositis, spicis terminalibus; floribus distantibus.

Lycopodioides frutescens, spicis laxis nudis. Dill. musc. 468. t. 64. f. 4.

Muscus frutescens, fructu tricapsulari. Plum. fil. 145. t. 170. f. A. A.

Habitat in Indiis.

Cau-

Caulis erectus, fere lignosus, tetragonus. Folia minima, acuta, remotissima, opposita. Rami terminales, dichotomi, triquetri, erecti. Pedunculi brevissimi, alterni, laterales, uniflori, apice bidentati, in eujus bifurcatura Anthera sessilis, rotundo-tricocca, trilocularis, polline foeta.

3. **LYCOPODIUM** foliis verticillato-quaternis, spicis *Phlegmaria*. terminalibus dichotomis.

*Lycopodium erectum dichotomum, foliis cruciatis, spicis gracilloribus. Dill. musc. 450. t. 61. f. 5. Fl. zeyl. 386. **

Selago indiarum orientalis f. Phlegmaria admirabilis zeylanica. Breyn. cent. 186, t. 92. Raj. hist. 1582. suppl. 655.

Tana-pouel paatsia maravara. Rheed. mal. 12. p. 27. t. 14.

Habitat in Malabar, Zeylona.

4. **LYCOPODIUM** foliis sparsis filamentosis, spicis *clavatum*. teretibus pedunculatis geminis.

Lycopodium caule repente, foliis patulis, pedunculis ipica gemina terminatis. Fl. lapp. 418. Fl. succ. 859.

Hort. cliff. 476. Mat. med. 488. Roy. lugdb. 506.

Hall. belv. 92. Dalib. parif. 339.

Lycopodium vulgare pilosum anfragosum & repens. Dill. musc. 441. t. 58. f. 1.

Muscus terrestris clavatus. Baub. pin. 360.

Habitat in Europæ sylvis muscosis.

5. **LYCOPODIUM** foliis sparsis filamentosis, spicis *rupestre*. terminalibus tetragonis.

Lycopodium spica sessili tetragona, squamis carinatis mucronatis. Aman. acad. 2. p. 337.

Lycopodium rupestre pilosum & incanum, spicis acute quadrangulis. Dill. musc. 457. t. 65. f. 11.

Muscus rupestris repens virginianus, clavis foliolis erectis quadratis. Pluk. alm. 248. t. 100. f. 2. Morif.

hist. 3. p. 624. f. 15. t. 5 f. 4. Raj. suppl. 32.

Habitat in Virginia, Canada, Sibiria.

5. **LYCOPODIUM** foliis sparsis ciliatis lanceolatis, spicis *Selaginoides* solitariis terminalibus foliosis.

*Lycopodium spicis solitariis sessilibus, foliis aculeato-ciliatis, capsulis tetraspermis. Fl. succ. 863. **

Ly-

Lycopodium caule repente, ramis spicæ foliis patula in-
arctis. Fl. lapp. 421.

Selaginoides foliis spinosis. Dill. musc. 460. t. 88. f.
1.

Habitas in Europæ pascuis muscosis.

alopecuroides.

7. **LYCOPODIUM** foliis sparsis ciliatis linearibus, spicis terminalibus foliosis.

Lycopodium ramis reflexis, apicibus radicatis: foliolis subulatis basi ciliatis. Hort. cliff. 476

Lycopodium terrestre repens, caule rotundo foliis densè obdito, radices ad summam emittens & se late propagans. Gron. virg. 126.

Lycopodium alopecuroides, flagellorum extremitatibus radicosis. Dill. musc. 464. t. 62. f. 8.

Muscus terrestris repens virginianus humi diffusus, venticulis longioribus, foliolis tenuibus vestitis. Moris. hist. 3. p. 624. f. 15. t. 5. f. 12. Raj. suppl. 32.

Habitat in Virginia, Canada.

- inundatum.* 8. **LYCOPODIUM** foliis sparsis integerrimis, spicis terminalibus foliosis.

Lycopodium caule repente, spicis solitariis sessilibus foliolis, foliolis lævibus. Fl. suec. 862. Dalib. parif. 340.

Lycopodium palustre repens, clava singulari. Vaill. parif. 123. t. 16. f. 11. Dill. musc. 452. t. 62. f. 7.

Habitat in Europæ inundatis.

Selago.

9. **LYCOPODIUM** foliis sparsis octifariis, caule dichotomo erecto fastigiato, floribus sparsis.

Lycopodium caule erecto dichotomo. Fl. lapp. 420. Fl. suec. 857. Mat. med. 487. Hort. cliff. 476. Roy. lugdb. 506.

Selago vulgaris abietis rubræ facie. Dill. musc. 435. t. 56. f. 1.

Muscus erectus ramosus saturate viridis. Bauh. pin. 360.

Muscus erectus abietiformis. Moris. hist. 3. p. 624. f. 5. t. 5. f. 9.

Habitat in Europæ borealis sylvis acerosis.

obscurum.

10. **LYCOPODIUM** foliis sparsis decurrentibus, farnentis repentibus, surculis erectis dichotomis.

Lycopodioides radiatum dichotomum. Dill. musc. 274. t. 67.

Habit-

Habitat in Philadelphia.

Sarmenta erecta, dichotoma at non fastigiata: ramis lateralibus scilicet brevioribus. Folia sparsa attamen rariora; situ indeterminato, glabra, rigidiuscula, linearia, acuta, basi decurrentia s. adnata cauti, deimpunctula. Fructificatio neque a me visa fuit.

11. LYCOPODIUM foliis sparsis quinquefariis subser-^{annuotinum.} ratis, surculis annotino-articulatis, spicis terminalibus glabris erectis.

Lycopodium caule erecto, spica glabra terminali. Fl. lapp. 419. Fl. succ. 858. Hall. belv. 93.

Lycopodium elatius juniperinum, clavis singularibus sine pediculis. Dill. musc. 455. t. 63. f. 9.

Muscus clavatus, juniperinis foliis reflexis. clavis singularibus sine pediculis. Moris. hist. 3. p. 624. f. 15. t. 5. f. 3.

Muscus terrestris repens, clavis singularibus erectis.

Pluk. alm. 248. t. 258. f. 5.

Habitat in Europæ memoribus.

12. LYCOPODIUM foliis sparsis curvatis, caule ramo-^{cernuum.} sissimo, spicis nutantibus.

Lycopodium erectum ramosissimum, spicis ovatis sessilibus nutantibus. Fl. zeyl. 387.

Lycopodium frutescens capillaceum crispum, spicis brevibus nutantibus. Dill. musc. 456. t. 63. f. 10.

Lycopodium zeylanicum erectum ramosissimum. Burm. zeyl. 144. t. 66.

Muscus zeylanicus erectus perpetuo virens, in arboris proceritatem excrescens. Pluk. alm. 257. t. 47. f. 9.

Muscus zeylanicus terrestris clavatus erectus. Moris. hist. 3. p. 624. f. 15. t. 5. f. 6.

Muscus clavatus erectus, crispatis foliolis. Pluk. alm. 149. t. 431. f. 3.

Muscus maximus, coridis folio, viticulis late ramosis.

Plum. fil. 144. t. 165.

Bellon-patia. Rheed. mal. 12. p. 73. t. 39.

Habitat in Indiis.

13. LYCOPODIUM foliis sparsis imbricatis, ramulis *Bryopteris*, convolutis.

Lycopodioides squamosum hispidum: extremitatibus convolutis. Dill. musc. 472. t. 66. f. 11.

Muscus erectus indicus platyceros filicem referens.

Moris. hist. 3. p. 628. f. 15. t. 7. f. 51.

Mu-

Muscus filicinus elegans f. *Bryopteris*. *Pluk. alm.* 247.
z. 100. *f.* 3. *Raj. suppl.* 32.
Habitat in India.

sanguinolentum. 14. *LYCOPODIUM* foliis quadrifariis imbricatis, caulis repentibus dichotomis, spicis sessilibus tetragonis. *Aman. acad.* 2. p. 363. *f.* 26. *
Habitat in Camtschatca.

alpinum. 15. *LYCOPODIUM* foliis quadrifariam imbricatis acutis, surculis erectis dichotomis, spicis sessilibus teretibus.
Lycopodium caule repente, ramis tetragonis. *Fl. lapp.* 417. *z.* 91. *f.* 6. † *Fl. succ.* 861. * *Hort. cliff.* 476.
Roy. lugdb. 506. *Hall. helv.* 93.
Lycopodium sabinae facie. *Dill. musc.* *z.* 58. *f.* 2.
Muscus clavatus, foliis cupressi. *Raj. hist.* 121.
Habitat in Alpibus Lapponiae, Helvetiae,

complanatum. 16. *LYCOPODIUM* foliis bifariis connatis: superficialibus solitariis, spicis geminis pedunculatis.
Lycopodium caule repente, ramis triquetro-planis. *Fl. lapp.* 416 *Fl. succ.* 860. *
Lycopodium sabinae facie. *Dill. musc.* 445. *synonyma.*
Muscus clavatus, foliis cupressi. *Baub. pin.* 360.
Muscus terrestris ramosus pulcher. *Baub. hist.* 3. p. 759.
Lycopodium digitatum, foliis arboris vitae, spicis bigemellis teretibus. *Dill. musc.* 448. *z.* 59. *f.* 3.
Habitat in Europæ & Americæ septentrionalis sylvis acerosis.
Confer. *Lycopodium altissimum, spicis bigemellis quadrangulis.* *Dill. musc.* 449. *z.* 60. *f.* 4.

carolinense. 17. *LYCOPODIUM* foliis bifariis patulis: superficialibus solitariis, scapis longissimis monostachyis.
Lycopodium pinnatum repens, spicis & pediculis singularibus longis. *Dill. musc.* 452. *z.* 62. *f.* 6.
Habitat in Carolina.

helveticum. 18. *LYCOPODIUM* foliis bifariis patulis: superficialibus distichis, spicis geminis pedunculatis.
Lycopodioides denticulatum pulchrum repens, spicis pediculis insidentibus. *Dill. musc.* 465. *z.* 64. *f.* 2.
Lycopodioides foliis ellipticis alterne majoribus. *Hall. helv.* 93.
Muscus denticulatus major. *Baub. pin.* 360.

Mu-

Muscus repens, foliis alternis subrotundis per siccitatem falcatis. *Morif. hist.* 3. p. 626. f. 15. t. 6. f. 34.
Habitat in Alpibus Helvetiæ.

19. **LYCOPODIUM** foliis bifariis: alternis minoribus, *apodami*
 caule repente, spicis sessilibus.

Lycopodioides denticulatum pulchrum repens, spicis apodibus. *Dill. musc.* 467. t. 64. f. 3.
Habitat in Carolina, Virginia, Pennsylvania.

20. **LYCOPODIUM** foliis bifariis: superficialibus *di-stichis*
stichis, caule erecto tereti.

Lycopodioides dentatum erectum filicinum, caule tereti ramosissimo. *Dill. musc.* 468. t. 65. f. 5.

Muscus squamosus erectus. *Plum. amer.* 35. t. 24. fil. t. 43. *Raj. suppl.* 31.
Habitat in America calidiore.

21. **LYCOPODIUM** foliis bifariis: superficialibus *di-canaliculatis*
stichis, caule erecto canaliculato.

Lycopodioides erectum filicinum, pinnulis acaciæ, caule sulcato. *Dill. musc.* 469. t. 65. f. 6.

Muscus denticulatus major ex Ceram. *Pluk. amalth. app.* t. 453. f. 8.
Habitat in Amboina.

22. **LYCOPODIUM** foliis bifariis patentibus: *superfi-plumosis*
cialibus distichis, surculis erectiusculis, spicis terminalibus tetragonis.

Lycopodioides dentatum dichotomum rigidum minus.
Dill. musc. 471. t. 66. f. 10.

Muscus squamosus repens. *Plum. amer.* 36. t. 24. fil. 43. *Raj. suppl.* 31.

Planta muscosa & plumosa guyanensis perelegans minor.
Breyn. cent. 1. p. 194. t. 101
Habitat in Indiis.

Confer. *Lycopodioides dichotomum taxiforme*. *Dill. musc.* 471. t. 66. f. 9.

Lycopodioides dentatum dichotomum rigidum minus.
Dill. musc. 471. t. 66. f. 11.

Lycopodioides Lonchitidis auriculatis foliis *Dill. musc.* 470. t. 66. f. 8.

23. **LYCOPODIUM** foliis bifariis patentibus: *superfi-ornithopodi-*
cialibus distichis, surculis repentibus, spicis sessilibus. *oides.*

Lycopodium procumbens repens, ramis subspicatis, foliolis

A a a a

lis

lis distichis alternis ovatis appendiculatis. *Fl. zeyl.* 388.

Lycopodioides repens, pinnulis ornithopodii. *Dill. musc.* 464. t. 66. f. 1. *B.*

Muscus lycopodioides denticulatus scandens repensve, in extremitatibus ramulorum polyspermos. *Raj. suppl.* 32.

Habitat in India.

denticulatum.

24. *LYCOPODIUM* foliis bifariis; superficialibus imbricatis, surculis repentibus, floribus sparsis.

Lycopodioides imbricatum repens. *Dill. musc.* 462. t. 66. f. 1. *A.*

Muscus denticulatus minor. *Bamb. pin.* 360.

Muscus terrestris lusitanicus. *Clas. hist.* 2. p. 249.

Habitat in Lusitania, Hispania, Iberia.

PORELLA.

pinnata.

1. *PORELLA.* †

Porella pinnis obtusis. *Dill. musc.* 459. t. 68. f. 1.

Habitat in Pennsylvania.

Hanc neque ego vidi, nec eam in solo natali investigare potuit D. Kalm.

SPHAGNUM.

palustre.

1. *SPHAGNUM* ramis deflexis. *Fl. lapp.* 415. *Fl. succ.* 864.

Sphagnum palustre molle deflexum, squamis cymbiformibus. *Dill. musc.* 240. t. 32. f. 1.

Muscus palustris in ericetis nascens. *Pluk. phyt.* 101. f. 1. *Vaill. parif.* 139. t. 23. f. 3.

2. *Sphagnum palustre molle deflexum*, squamis capilla-ceis. *Dill. musc.* 243. t. 32. f. 2.

Habitat in Europæ paludibus profundis sylvaticis.

alpinum.

2. *SPHAGNUM* subramosum erectum.

Sphagnum subulatum viridissimum, capsulis ovatis. *Dill. musc.* 245. t. 32. f. 3.

Habitat in Angliæ summis alpinæ paludibus.

PHASCUM.

acaulon.

1. *PHASCUM* acaule, anthera sessili.

Sphagnum acaulon bulbiforme majus. *Dill. musc.* 251. t. 32. f. 11.

Mu-

Muscus trichodes acaulos minor latifolius. *Pet. gaz.* 126. t. 95. f. 14.

β. *Sphagnum acaulon* bulbiforme minus. *Dill. musc.* 252. t. 32. f. 12.

Muscus trichodes acaulos minor latifolius. *Vaill. parif.* 128. t. 27. f. 2.

Habitat in Europæ agris, hortis & ad fossas.

Confer Sphagnum acaulon maximum, foliis in centro ciliaribus. Hall. itin. alp. 83. t. 1. *Dill. musc.* 253. t. 32. f. 13.

Sphagnum acaulon trichodes. Dill. musc. 251. t. 32. f. 10.

Muscus trichodes minor acaulos, capillaceis foliis. *Vaill. parif.* 128. t. 29. f. 4.

2. PHASCUM caulescens erectum.

caulescens.

Sphagnum foliis tenuibus gramineis pellucidis. Dill. musc. 550. t. 85. f. 15.

Habitat in Pennsylvania.

3. PHASCUM caulescens repens.

repens.

Sphagnum vermiculare repens, capsulis intra foliorum squamosorum alas minimis. Dill. musc. 550. t. 85. f. 16.

Habitat in arborum truncis.

FONTINALIS.

1. FONTINALIS foliis complicato-carinatis trifariis a-antipyretica cutis, antheris lateralibus. *Fl. suec.* 866.

Fontinalis capitulis lateralibus. Roy. lugdb. 506.

Fontinalis triangularis major complicata e foliorum alis capsulifera. Dill. musc. 254. t. 33. f. 1.

Muscus squamosus, foliis acutissimis, in aquis nascens. *Vaill. parif.* 140. t. 33. f. 5.

Muscus aquaticus, terrestri vulgari similis, major. *Buxb. cent.* 3. p. 39. t. 69. f. 2.

Muscus aquaticus, viticulis longis minus ramens lucidis. *Morif. bist.* 3. p. 626. f. 15. t. 6. f. 32.

Muscus aquaticus denticulatus. *Laf. pruss.* 173. t. 52.

Habitat in Europæ fluviis.

2. FONTINALIS foliis ovatis concavis trifariis acutis, minor, antheris terminalibus.

Fontinalis triangularis minor carinata, e cymis capsulifera. Dill. musc. 257. t. 33. f. 2.

Fontinalis capitulis terminalibus. Roy. lugdb. 506.

A a a a a

Habi-

Habitat in Europæ fluviiis.

Præcedenti nimis affinis.

- squamosa* 3. FONTINALIS foliis imbricatis subulato-lanceolatis, capitulis lateralibus.
Fontinalis squamosa tenuis sericea atrovirens. *Dill. musc.* 259. t. 33. f. 3.
Fontinalis minor lucens. *Bamb. hist.* 3. p. 778.
Habitat in Britannia, Gallia.

- pennatum* 4. FONTINALIS foliis bifariis patentibus, capsulis lateralibus.
Sphagnum pennatum undulatum, vagina squamosa. *Dill. musc.* 250. t. 32. f. 9.
Muscus terrestris major, ramulis compressis, foliis superficie crispis. *Vaill. parif.* 129. t. 27. f. 4.
Habitat in Helvetiæ arboribus.
Confer SPHAGNUM pennatum undulatum, vagina squamosa. *Dill. musc.* 250. t. 32. f. 8.
SPHAGNUM pennatum undulatum, vagina pilosa. *Dill. musc.* 249. t. 32. f. 7.



SPLACHNUM.

- rubrum* 1. SPLACHNUM umbraculo convexo. *Amæn. acad.* 2. p. 272. t. 3. f. 2. *
Muscus norvegicus, umbraculo ruberrimo insignitus. *Pet. musc.* 70. t. 1. f. 10. *Dill. musc.* 547. t. 83. f. 9.
Muscus coronatus norvegicus, pediculo longissimo, umbraculo amplo. *Morif. hist.* 3. p. 630. f. 15. t. 7. f. 10.
Habitat in Norvegia, Finlandia, Russia, Sibiria.
- lucum* 2. SPLACHNUM umbraculo plano. *Amæn. acad.* 2. p. 277. t. 3. f. 1. *
Habitat in Sueciæ paludibus sylvaticis.
- ampullaceum* 3. SPLACHNUM umbraculo ampullaceo clauso. *Amæn. acad.* 2. p. 280.
Bryum ampullaceum, foliis thymi pellucidis, collo striatiore. *Dill. musc.* 343. t. 44. f. 3.
Muscus capillaceus minor, capitulis geminatis. *Vaill. parif.* 130. t. 26. f. 4.
Muscus aureus capillaris minor, capitulis geminatis crebrius mutuo incubitu adnatis. *Pluk. alm.* 246. *Morif. hist.* 3. p. 629. f. 15. t. 16. f. 10.

Mu-

Muscus capillaceus, capitulis pyriformibus tubulosis.

Buxb. cent. 2. p. 1. t. 1. f. 1.

Habitat in Europæ frigidioris paludibus sylvaticis.

POLYTRICHUM. 6

2. POLYTRICHUM caule simplici (*Fl. lapp. 395.*) an- continued.
thera parallelepipedæ. *Fl. suec. 868. Dalib. parif. 316.*

Polytrichum quadrangulare vulgare, yuccæ foliis serratis. *Dill. musc. 420. t. 54. f. 1.*

Polytrichum aureum majus. *Baub. pin. 356.*

Muscus juniperifolius, capitulo quadrangulo. *Vaill. parif. 131. t. 23. f. 8.*

3. Polytrichum quadrangulare, juniperi foliis brevioribus & rigidioribus. *Dill. musc. 424. t. 54. f. 2.*

Polytrichum aureum medium. *Baub. pin. 356.*

Muscus coronatus humilis. *Morif. bist. 3. p. 630. f. 15. t. 7. f. 6. 8.*

Muscus erectus, juniperi folio glauco. *Vaill. parif. 131. t. 23. f. 6.*

7. Polytrichum quadrangulare minus, juniperi foliis pilosis. *Dill. musc. 426. t. 54. f. 3.*

Habitat in Europæ α. uliginosis, β. pascuis, γ sterilissimis. 2

2. POLYTRICHUM caule ramosissimo, pedunculis alpinum, terminalibus.

Polytrichum alpinum ramosum, capsulis e summitate ellipticis. *Dill. musc. 427. t. 55. f. 4.*

Habitat in Alpibus Helvetiæ, Snowdon.

3. POLYTRICHUM caule ramosissimo, pedunculis a- urnigerum.
xillaribus.

Polytrichum caule ramoso, capsulis basi destitutis apophysi. *Fl. suec. 870. Dalib. parif. 315.*

Polytrichum ramosum setis ex alis urnigeris. *Dill. musc. 427. t. 55. f. 5.*

Muscus ramosus erectus, calyptra villosa. *Vaill. parif. 131. t. 28. f. 13.*

Habitat in Europa, Jamaica.

MNIUM. 7

1. MNIUM caule simplici, foliis ovatis. *Fl. suec. 911. pellucidum*

Mnium serpilli foliis tenuibus pellucidis. *Dill. musc. 232. t. 31. f. 2.*

Bryum alternans calyptra variegata & scyphulis foliosis.

Hall. belv. 118. t. 4. f. 8.

A a a a 3

Mu-

Muscus coronatus minimus, capillaceis foliis, capitulis oblongis. *Vaill. parif.* 130. t. 24. f. 7.

Habitat in Europæ pascuis succulentis umbrosis.

fontanum. 2. *MNIUM* caule simplici geniculis inflexo. *Fl. Lapp.* 414. *Fl. Suec.* 913.

Bryum palustre, scapis teretibus stellatis, capsulis magnis subrotundis. *Dill. musc.* 340. t. 44. f. 2.

Muscus capillaceus tenuissimus, pediculo longissimo purpurascente, capitulo rotundiore. T. *Vaill. parif.* 134. t. 24. f. 10.

Muscus trichodes palustris erectus coronatus, capitulis sphaericis amplioribus. *Morif. hist.* 3. p. 628. f. 15. t. 6. f. 8.

Muscus parvus stellaris. *Pink. a.l.m.* 255. t. 47. f. 6.

Habitat in Europa ad fontes paludosos frigidos.

palustre. 3. *MNIUM* caule dichotomo, foliis subulatis. *Fl. Lapp.* 413. *Fl. Suec.* 912.

Mnium majus, ramis longioribus bifurcatis. *Dill. musc.* 233. t. 31. f. 3.

Muscus palustris erectus flavescens, capillaceo folio. *Vaill. parif.* 135. t. 24. f. 1.

Muscus palustris, flagellis erectis luteolis raro divisis, capitulis oblongis adiant. *Morif. hist.* 3. p. 629. f. 15. t. 6. f. 9.

Muscus stellaris ramosus palustris, pediculo aureo erecto, capitulo magno sphaerico. *Buxb. cent.* 1. p. 45. t. 65. f. 1. 2. 3. 4.

Habitat in Europæ paludibus.

androgynum 4. *MNIUM*, caule ramoso, androgynum. *Fl. Suec.* 910. *Mnium perangustis & brevibus foliis.* *Dill. musc.* 230. t. 31. f. 1.

Bryum angustiori folio, capsula longa pyriformi inclinata. *Hall. belv.* 114.

Muscoides muscus capillaris minimus, capitulo minimo pulverulento. *Vaill. parif.* t. 29. f. 6.

Muscus trichodes parvus, capitulo glomerato s. botryoide. *Mich. gen.* 108. t. 59. f. 8. H. K.

Muscus coronatus humilis minimus, foliolis pediculis & capitulis minimis erectis. *Morif. hist.* 3. p. 631. f. 15. t. 7. f. 20.

Habitat in Europæ sylvis.

hygrometricum. 5. *MNIUM* acule, anthera nutante, calyptra reflexa tetragona.

Bry-

Bryum capsula nutante turbinato-ovata, calyptra reflexa quadrangulari. *Fl. suec.* 903. *Dalib. parif.* 324.

Bryum bulbiforme aureum, calyptra quadrangulari, capsulis pyriformibus nutantibus. *Dill. musc.* 407. t. 52. f. 75.

Muscus foliis scutellatis, capitulo pyriformi nutante. *Vaill. parif.* 135. t. 26. f. 16.

Muscus coronatus minor, foliolis latiusculis ad caulem convolutis, capitulis cernuis & obtusis aureis. *Morif. bist.* 3. p. 631. f. 15. t. 7. f. 17.

Habitat in Europæ sylvis apricis sterilibus.

6. MNIMUM caule dichotomo: axillis pedunculiferis, antheris erectis, foliis carinatis.

Bryum capsula erectiuscula oblonga, caulium divisuris pedunculiferis: summitatibus stellatis, foliis carinatis. *Fl. suec.* 896. *Dalib. parif.* 325.

Bryum tenue stellatum, setis purpureis. *Dill. musc.* 386. t. 49. f. 51.

Muscus trichodes parvus, foliis musci vulgaris, capitulis longis acutis. *Morif. bist.* 3. p. 628. f. 15. t. 6. f. 4. 16.

Habitat in Europæ pascuis.

7. MNIMUM antheris erectis, operculis filiformibus longitudine capitulorum.

Bryum stellare nitidum pallidum, capsulis tenuissimis. *Dill. musc.* 381. t. 48. f. 44.

Habitat in Europæ muris, aggeribus, sepibus.

Caulis erectus, brevissimus. Pedunculi atropurpurei, flexuosi, Anthera cylindrica, livida. Opercula & Calyptrae filiformes, longiores & angustiores, quam in reliquis.

8. MNIMUM foliis arefactione revolutis.

cirratum.

Bryum cirratum & stellatum, tenuioribus foliis. *Dill. musc.* 379. t. 48. f. 42.

Muscus muralis minimus roseus stellaris. *Vaill. parif.* 130. t. 24. f. 8.

Habitat in Europæ sylvis & ad sepes.

9. MNIMUM foliis ovatis acuminatis pellucidis, pedunculis subradicalibus, anthera nutante.

Bryum annonium lanceolatum pellucidum, capsulis oblongis pendulis. *Dill. musc.* 399. t. 50. f. 68.

Habitat in Europæ humidiusculis.

horum.

10. MNIUM antheris pendulis, pedunculo curvato, furculo simplici, foliolis margine scabris.

Bryum stellare horum sylvarum, capsulis magnis nutantibus. *Dill. musc.* 402. t. 51. f. 71.

Muscus capillaceus major stellatus. *Vaill. parif.* t. 24. f. 4. v.

Muscus polytrichoides humilior, foliis brevioribus raris pallide viridantibus et vix pellucidis. *Morif. hist.* 3. p. 630. f. 15. t. 6. f. 3. 4.

Habitat in Europæ sylvis, aggeribus.

radum.

11. MNIUM antheris pendulis, calyptra recurvata, foliis pellucidis.

Bryum capsulis pendulis, calyptra recurvata, foliis subulatis acutis integerrimis, caule simplicissimo. *Fl. suec.* 907.

Bryum pendulum horum molle, foliis & lanceolatis & gramineis. *Dill. musc.* 401. t. 51. f. 7.

Habitat in sylvis & declivitatibus.

styriforme.

12. MNIUM antheris pendulis turbinatis, stipite filiformi, floribus femineis setiferis.

Bryum antheris nutantibus turbinatis, foliis ensiformibus nitidis. *Fl. suec.* 902.

Bryum trichodes aureum, capitulis pyriformibus nutantibus. *Dill. musc.* 391. t. 50. f. 60.

Habitat in Europæ rupibus.

Polytrichoides.

13. MNIUM calyptra villosa.

Polytrichum nanum, capsulis subrotundis galeritis, aloes folio non serrato. *Dill. musc.* 428. t. 55. f. 6. *Dalib. parif.* 315.

Muscus capillaceus minor, calyptra tomentosa. *Vaill. parif.* 131. t. 26. f. 15.

Muscus coronatus rigidus minor & humilior, capitulis villosis brevioribus. *Morif. hist.* 3. p. 630. f. 15. t. 7. f. 7.

- β. Polytrichum parvum, aloes folio serrato, capsulis oblongis. *Dill. musc.* 429. t. 55. f. 7.

Adiantum aureum medium, in ericetis proveniens. *Vaill. parif.* 131. t. 29. f. 11.

Muscus, adianti aurei facie, humilis, capitulo rotundo turgido, calyptra villosa incana. *Buxb. cent.* 1. p. 43. t. 63. f. 1.

- γ. Polytrichum acaulon capillaceum, capsulis cylindricis. *Dill. musc.* 434. t. 55. f. 12.

Habitat in Europæ ericetis subnatis.

14.

34. **MNIUM** pedunculis aggregatis, foliis patentibus pellucidis. *serpyllifolium*.
- α.** **Mnium** pedunculis aggregatis foliis obovatis integerri-punctatum. mis obtusis punctatis.
- Bryum** capsulis nutantibus, foliis subovatis obtusis pellucidis punctatis, caulibus sterilibus procumbentibus, pedunculis aggregatis. *Fl. succ.* 901. *Dalib. parif.* 323.
- Bryum** caule procumbente, foliis alternis ovatis pellucidis. *Fl. lapp.* 404.
- Bryum** pendulum, serpylli folio rotundiore pellucido, capsulis ovatis. *Dill. musc.* 416. t. 53. f. 81.
- Muscus** folio lato subrotundo, capitulo singulari nutante, pediculo longo subrubenti insidente. *Vaill. parif.* 136. t. 26. f. 5.
- Muscus** polytrichoides palustris major, serpylli latioris folio pellucido. *Morif. bist.* 3. p. 627. f. 15. t. 6. f. 39. 40.
- β.** **Mnium** pedunculis aggregatis, foliis alternis acutis sercuspidatum ratis.
- Bryum** pendulum, foliis variis pellucidis, capsulis ovatis. *Dill. musc.* 413. t. 53. f. 79.
- Muscus** palustris, foliis subrotundis. *Vaill. parif.* 26. f. 18.
- Muscus** polytrichoides pellucidus fere, foliis denticulatis ad margines veluti crispis. *Pluk. alm.* 257. t. 98. f. 8.
- γ.** **Mnium** pedunculis aggregatis, foliis rosaceo-congestis proliiferum, lanceolatis acutis.
- Bryum** stellare roseum majus, capsulis ovatis pendulis. *Dill. musc.* 411. t. 52. f. 77.
- Muscus** stellaris roseus. *Bauh. pin.* 361. *prodr.* 151. *Buxb. cent.* 2. p. 2. t. 1. f. 3.
- Muscus** coronatus humilis, foliolis latioribus stellatim nascentibus donatus. *Morif. bist.* 3. p. 630. f. 15. t. 7. f. 9.
- δ.** **Mnium** pedunculis aggregatis, foliis oblongis undulato-undulatum. tis.
- Bryum** caule adscendente, foliis ovato-oblongis patulis pellucidis. *Fl. lapp.* 399. *Roy. lugdb.* 502. *Guett. stamp.* 90. *Dalib. parif.* 324.
- Bryum** dendroides polycephalon, phyllitidis folio undulato pellucido, capsulis ovatis pendulis. *Dill. musc.* 410. t. 52. f. 76.
- Muscus** roseus polycephalus, linearis foliis undulatis. *Vaill. parif.* 135. t. 24. f. 3. *Mich. gen.* 108. t. 59. f. 5.

A a a a 5

Ha-

Habitat in Europa fere ubique.

Varietates has pro distinctis speciebus sumunt varii; ego ob suas causas ita conjunxi, ut qui distinguendus velit, facile distinctas tradat.

triquetrum. 15. **MNIUM** foliis trifariis patulis subulato-lanceolatis carinatis.

Bryum annotinum palustre, capsulis ventricosis pendulis. Dill. musc. 404. t. 51. f. 72?

Muscus denticulatus lucens fluviatilis maximus, ad ramorum apices adianti capitulis ornatus. Vail. parij. 135. t. 24. f. 2. 2?

Habitat in Sueciæ paludibus tarsofis.

Caules plurimi, in cespitem congesti, digitales & longiores, tomentosi. Folia trifariam posita, unde ramuli triquetri patuli, recentia pallide viridia, antiqua fusca, parva, distantia, carinata, acuminata. Pedunculi solitarii, terminales, purpurei. Capsulæ erectæ, oblique s. arcuatae, obtusissimæ, flavæ. Stellæ semina ramos terminant in eadem planta cum capsula, sed distinctis ramis.

trichomanis 16. **MNIUM** foliis distichis integerrimis.

Mnium trichomanis facie, foliolis integris. Dill. musc. 236. t. 31. f. 5.

Habitat in Sueciæ, Angliæ & adis.

sum.

17. **MNIUM** foliis distichis bifidis.

Mnium trichomanis facie, foliolis bifidis. Dill. musc. 237. t. 31. f. 6.

Jungermannia terrestris repens, foliis ex rotunditate acuminatis bifidis: apertura pene visibili. Mich. gen. 8. s. 5. f. 14.

Habitat in Europæ vasis.

jungermannia, 18. **MNIUM** foliis distichis, pinnis subtus auriculatis.

*Mnium caule procumbente simplici, foliis imbricatis integerrimis alternis antice appendiculatis. Fl. suec. 914. **

Lichemastrum alpinum purpureum, foliis auritis & cochleariformibus. Dill. musc. 479. t. 69. f. 1.

Jungermannia alpina palustris purpurea cambrica, foliis rotundioribus auritis tenuissime denticulatis. Mich. gen. 6. t. 5. f. 16.

Habitat in Europæ subhumidis.

BRY-

BRYUM. 8

* *Antheris sessilibus*

1. BRYUM antheris sessilibus terminalibus, calyptra apocarpum, minima.

Sphagnum caulibus ramosis, foliis undique imbricatis capsulas obtegentibus. *Fl. suec.* 864.

Sphagnum foliis pilo terminales. *Ray. Lugdb.* 500.

Sphagnum subhirsutum obscure virens, capsulis rubellis. *Dill. musc.* 245. t. 32. f. 4.

Muscus apocarpus hirsutus saxi adnascens, capitulis obscure rubris. *Vaill. paris.* 129. t. 31. f. 15.

2. Sphagnum nodosum hirsutum incanum. *Dill. musc.* 246. t. 32. f. 5.

Muscus squamosus saxatilis tortuosus ac nodosus. *Vaill. paris.* 123. t. 27. f. 18.

Habitat in Europæ saxi, arboribus.

2. BRYUM antheris sessilibus sparsis, calyptris striatis striatum, sursumve pilosis.

Polytrichum calyptra striata sursumve pilosa. *Fl. suec.* 869. *Dalib. paris.* 314.

Polytrichum bryi ruralis facie, capsulis sessilibus majus. *Dill. musc.* 430. t. 55. f. 8.

Muscus apocarpus arboreus ramosus. *Vaill. paris.* 129. t. 25. f. 5. 6.

3. Polytrichum bryi ruralis facie, capsulis sessilibus minus. *Dill. musc.* 431. t. 55. f. 9.

Muscus capillaceus minimus acaulos, calyptra striata. *Vaill. paris.* 30. t. 27. f. 10.

Muscus humilis tectorum subfuscus, capitulis brevibus, pileolis erectis, pedunculis curtis. *Morif. hist.* 3. p. 629. f. 15. t. 6. f. 13.

4. Polytrichum capsulis sessilibus, foliis brevibus rectis carinatis. *Dill. musc.* 432. t. 55. f. 10.

5. Polytrichum capillaceum crispum, calyptris acutis pilosissimis. *Dill. musc.* 433. t. 55. f. 11.

Muscus capillaceus minimus, calyptra villosa. *Vaill. paris.* 130. t. 26. f. 9.

Habitat in Europæ a. arboribus b. saxi, c. rupibus, d. sylvis.

* *Antheris pedunculatis erectis.*

3. BRYUM antheris erectis sphericis. *Fl. suec.* 888. *Dalib. nomiforma.* *parif.* 317.

Bryum caule erecto, foliis setaceis, capitulis globosis. *Fl. lapp.* 400. Bry-

Bryum capillaceum, capsulis sphaericis. *Fl. lapp.* 400
Dill. musc. 339. t. 44. f. 1.

Muscus trichodes minimus sericeus capillaceus, capitulis sphaericis. *Morif. hist.* 3. p. 628. f. 15. t. 6. f. 6.
Vaill. parif. 129. t. 24. f. 9. 12.

Habitat in Europæ scopulis humosis.

pyriforme. 4. *BRYUM* Antheris erectis obovatis, calyptra subulata, surculis acaulibus, foliis ovatis muticis.

Bryum antheris erectis turbinato-ovatis: operculo obtusiusculo. *Fl. suec.* 890. *Dalib. parif.* 317,

Bryum serpillifolium pellucidum, capsulis pyriformibus. *Dill. musc.* 345. t. 44. f. 6.

Muscus capillaceus minimus, capitulis pyriformibus turgidis. *Vaill. parif.* 129. t. 29. f. 3.

Muscus minimus latifolius, capitulis erectis turgidis. *Buxb. cent* 1. p. 44. t. 64. f. 1.

Muscus coronatus humilis stellaris, foliis latiusculis, capitulis pyriformibus erectis turgidiusculis. *Morif. hist.* 3. p. 631. f. 15. t. 7. f. 16.

Habitat in Europæ pratis ad aggeras.

extinctorium 5. *BRYUM* anthera erecta oblonga minori calyptris laxis æqualibus. *Fl. suec.* 891. *Dalib. parif.* 325.

Bryum calyptra extinctorii forma, minus. *Dill. musc.* 349. t. 45. f. 8.

Muscus capillaceus minimus, calyptra longa conoide nitida. *T. Vaill. parif.* 137. t. 26. f. 1.

Muscus capillaris minimus cucullatus. *Magn. hort.* 139. t. 82.

Muscus coronatus humilis tenuifolius, pilcolo magno acuto infra aperto, extinctorium referente. *Morif. hist.* 3. p. 630. f. 15. t. 7. f. 12.

β. *Bryum calyptra extinctorii figura*, majus & ramosum. *Dill. musc.* 350. t. 45. f. 9.

Habitat in Europæ arenosis.

subulatum. 6. *BRYUM* antheris erectis subulatis surculis acaulibus. *Bryum capsulis longis subulatis.* *Dill. musc.* 350. t. 45. f. 10.

Muscus capillaris, corniculis longissimis incurvis R. *Vaill. parif.* 133. t. 25. f. 8.

Habitat in Europæ aggeribus humidiusculis.

ravale. 7. *BRYUM* antheris erectiusculis, foliis piliferis recurvis. *Fl. suec.* 900. *Dalib. parif.* 320. *It. scan.* 118.

Bryum caule erecto, foliis reflexis seta terminatis, capitulis talcatis. *Fl. lapp.* 397. *Bryum*

Bryum rurale ungviculatum hirsutum elatius & ramosius. Dill. musc. 352. t. 45. f. 12.

Muscus capillaris tectorum, densis cespitibus nascens, capitulis oblongis, foliis in pilum oblongum desinentibus. Vaill. parif. 133. t. 25. f. 3.

Muscus erectus hirsutus, capitulis longis acutis ungviculis avicularum similibus, vulgarissimus. Morif. hist. 3. p. 628. f. 15. t. 6. f. 12.

Habitat in Europæ muris, rusticorum tectis, arborum truncis. &

8. BRYUM antheris erectis, foliis piliferis rectiusculis, ^{murale.} sarculis simplicibus cespitosis.

Bryum tegulare humile pilosum & incanum. Dill. musc. 355. t. 45. f. 14.

Muscus capillaris minor, capitulis erectis, vulgarissimus, foliis in pilum desinentibus. Vaill. parif. 133. t. 24. f. 15.

Muscus vulgaris omnium vulgarissimus non villosus. Vaill. parif. 133. t. 24. f. 14.

Muscus capillaris minor, capitulis erectis vulgarissimus non villosus. Vaill. parif. t. 25. f. 4.

Habitat in Europæ saxis, muris, tectis, tegulis.

9. BRYUM antheris erectiusculis, pedunculis aggrega-^{scoparium.} tis, foliis secundis recurvatis, caule declinato. *Fl. succ. 898. Dalib. parif. 319.*

Bryum caule inclinato, foliis arrectis subulatis, capitulis erectiusculis. Fl. lapp. 398.

Bryum reclinatum, foliis falcatis, scoparum effigie. Dill. musc. 357. t. 46. f. 16.

Muscus capillaceus major, pediculo & capitulo tenuioribus. Vaill. parif. 132. t. 28. f. 12.

Muscus capillaceus, foliis pallidis, pediculis pluribus ex communi theca egredientibus. Buxb. cens. 2. p. 8. t. 4. f. 1.

Habitat in Europæ sylvis arenosis, in truncis putridis. &

10. BRYUM antheris erectiusculis, pedunculis subfoli-^{undulatum.} tariis, foliis lanceolatis carinatis undulatis patentibus serratis. *Fl. succ. 899. Dalib. parif. 320.*

Bryum phyllitidis folio rugoso acuto, capsulis incurvis. Dill. musc. 360. t. 46. f. 18.

Muscus erectus, linariæ folio, major. Vaill. parif. 132. t. 26. f. 17.

Habitat in Europæ sylvis, aggeribus, pratis umbrosis.

11. BRY-

- glaucum.* 11. BRYUM antheris erectiusculis: operculo arcuato, foliis erectis imbricatis, furculis ramosis.
 Bryum albidum & glaucum fragile majus, foliis erectis, setis brevibus. *Dill. musc.* 362. t. 46. f. 20.
 Muscus trichodes montanus albidus fragilis. *Morif. hist.* 3. p. 630. f. 15. t. 6. f. 22.
 Muscus erectus capillaceus densissimus, glauco folio. *Vaill. paris.* 131. t. 26. f. 13.
 Muscus saxatilis ericoides. *Banb. pin.* 362. *prodr.* 151.
Habitat in Europæ ericetis apricis.
- albidum.* 12. BRYUM antheris erectis foliis lingulatis obtusis patulis †
 Bryum nanum, lariginis foliis albis. *Dill. musc.* 364. t. 46. f. 21.
Habitat in Insula Providentia.
- pellucidum.* 13. BRYUM antheris erectiusculis, foliis recurvis acutis, caule hirsuto,
 Bryum palustre pellucidum, capsulis & foliis brevibus recurvis. *Dill. musc.* 364. t. 46. f. 23.
 Muscus polytrichodes elatior, foliis angustis pellucidis & fere membranaceis. *Pluk. alm.* 247. t. 44. f. 7.
 Muscus polytrichodes angustifolius pellucidus ramosus. *Pluk. alm.* 247.
 β. Bryum erectis capitulis brevibus, foliis reflexis. *Dill. musc.* 365. t. 46. f. 24.
Habitat in Europæ paludibus.
- aciculare.* 14. BRYUM antheris erectis: operculo aciformi, foliis erectis subsecundis.
 Bryum montanum hemiheterophyllum, operculis acutis. *Dill. musc.* 366. t. 46. f. 25.
Habitat in Anglia.
- flexuosum.* 15. BRYUM antheris erectis, foliis setaceis, pedunculis flexuosis. *Fl. suec.* 892.
 Bryum pilosum molle, setis intortis. *Dill. musc.* 373. t. 47. f. 33.
Habitat in Europæ sylvis.
- heteromallum.* 16. BRYUM antheris erectis, foliis setaceis secundis. *Fl. suec.* 895.
 Bryum heteromallum. *Dill. musc.* 375. t. 47. f. 37.
 Muscus capillaceus, foliis unam partem spectantibus. *Vaill. paris.* 132. t. 27. f. 7.
 Muscus trichodes, foliis capillaceis, capitulis minoribus. *Morif. hist.* 3 p. 628. f. 15. t. 6. f. 5. 7.
Habitat in Europæ ericetis, juniperetis.

17. BRY-

17. **BRYUM** antheris erectis, foliis setaceis imberbibus: *tertiofirmum*.
 siccitate retorquentibus. *Fl. suec.* 497.
Bryum alpinum, crinium crispum simile, capsula barbata. *Hall. belv.* 110. t. 4 f. 2.
Bryum cirratum, setis & capsulis longioribus. *Dill. musc.* 377. t. 48. f. 4.
Muscus alpinus cirrhosus s. crinium tortorum instar crispatus. *Scheuch. alp.* 65. t. 19. f. 15.
Muscus helveticus trichodes crispatus. *Pet. gaz.* 159. t. 65. f. 8.
Habitat in montibus Helvetiæ, Sveciæ, Cambriæ.
18. **BRYUM** antheris erectis subrotundis: operculo *mu-truncatulum*
 cronato. *Fl. suec.* 889. *Dalib. paris.* 313.
Bryum exiguum, creberrimis capsulis rufis. *Dill. musc.* 347. t. 45. f. 7.
Muscus capillaceus, omnium minimus. *Vail. paris.* 130 t. 26. f. 2.
Muscus stellaris minimus, pediculo brevissimo, capitulis turgidis. *Buxb. cent.* 2. p. 4. t. 2. f. 2.
Muscus coronatus minimus, foliis & capitulis oblongis in pediculis brevissimis. *Morif. hist.* 3. p. 631. f. 15. t. 7. f. 15.
Habitat in Europa ad agros, fossas, sepes.
Capsulæ amisso operculo apparent valde truncatæ, obovatae & rufæ, quibus manifeste differt a sequente.
19. **BRYUM** antheris erectis ovatis, foliis lanceolatis *a-viridulum*.
 cumminatis imbricato-patulis. *Fl. suec.* 893.
Bryum capillaceum breve, pallide & læte virens, capsulis ovatis. *Dill. musc.* 380. t. 48. f. 43.
Muscus capillaceus omnium minimus, foliolis longioribus & angustioribus. *Vaill. paris.* 130. t. 29. f. 5.
Habitat in Europæ agris, aggeribus.
Folia viridissima. Capsulæ flavescences, ciliatæ.
20. **BRYUM** acaulon, foliolis setaceis, antheris obtu-*paludosum*.
 sissimis patulis.
Bryum trichodes acaulon palustre minimum, setis & capsulis brevissimis *Dill. musc.* 387. t. 49. f. 53.
Habitat in Europæ humidis.
Differt a Viridulo, capsulis fuscis & foliis siccitate minime crispis.
21. **BRYUM** antheris erectis, caule erectiusculo, ramis la-*Hypnoides*.
 teralibus brevibus fertilibus. *Fl. lapp.* 396. *Fl. suec.* 894. *Dalib. paris.* 326.

Bry-

- oryum* hypnoides polycephalon lanuginosum montanum. *Dill. musc.* 372. t. 47. f. 32.
Muscus alpinus ramosior erectus, flagellis brevioribus, lanuginosus. Pluk. alm. 255. t. 47. f. 5.
Muscus terrestris vulgari similis lanuginosus. Moris. hist. 3. p. 625. f. 15. t. 5. f. 7.
 β. *Bryum hypnoides hirsutie virescens, fasciculare alpinum. Dill. musc.* 370. t. 47. f. 28.
 γ. *Bryum hypnoides, ericæ facie, capsulis barbatis, alpinum. Dill. musc.* 371. t. 47. f. 31.
Habitat in Europæ frigida rupibus, saxis, sabulosis.
- verticillatum.* 22. *BRYUM* antheris erectis, pedunculis siccitate contortis, foliis piliferis, furculis fastigiatis.
Bryum pilosum verticillatum. Dill. musc. 374. t. 47. f. 35.
Habitat ad latera collium.
- celsi.* 23. *BRYUM* antheris erectiusculis, pedunculis longissimis, foliis setaceis, furculis acaulibus.
Bryum trichodes acaulon minimum, setis & capsulis oblongis. Dill. musc. 388. t. 49 f. 54.
Habitat Upsaliæ.
- trichodes.* 24. *BRYUM* antheris erectiusculis: ore ciliato absque annulo, pedunculo longissimo.
Bryum trichodes aureum, capsulis incurvis obtusis in setis longis. Dill. musc. 389. t. 49. f. 58.
Habitat in Suecia, Germania.
- squarrosum.* 25. *BRYUM* antheris obliquis, foliis quinquesariis imbricatis recurvis, caule ferrugineo-tomentoso.
Muscus palustris, foliis reflexis cauliculos quinquangulos formantibus. Buxb. cent. 4. p. 36. t. 65. f. 1.
Habitat in Europæ borealis paludibus apricis. Læf.
Muscus distinctissimus ab Hypno squarroso. Surculi plures, conferti, erectiusculi, undique fusco pubescentes: Ramo uno alterove brevi. Folia minima, sessilia, ovata, acuminata, subcarinata, late viridia, imbricata, revoluta, quinquesariam exstantia. Pedunculus terminalis purpureus, longus. Capsula oblonga, obliqua.
- * *Antheris nutantibus.*
- argentum.* 26. *BRYUM* antheris pendulis, furculis cylindricis imbricatis lævibus *Fl. suec.* 909. *Dalib. parif.* 321. *Bry-*

- Bryum pendulum* julaceum argenteum & sericeum.
Dill. musc. 392. t. 50. f. 62.
- Muscus squamulosus* argenteus, ericæ folio. *Vaill. paris.* 134. t. 26. f. 3.
- Muscus minimus* e viridi argenteus, capitulis oblongis cernuis. *Morif. hist.* 3. p. 627. f. 15. t. 6. f. 17.
- β. *Bryum pendulum*, surculis teretibus viridibus. *Dill. musc.* 394. t. 50. f. 63.
- Habitat in Europæ sectis, muris, rupibus.*
27. **BRYUM** antheris subrotundis, pedunculis reflexis, *pulvinatum*. foliis piliferis. *Fl. suec.* 904. *Dalib. paris.* 321.
- Bryum* foliis setaceis, pedunculo reflexo, capitulis subrotundis. *Fl. lapp.* 401.
- Bryum* orbiculare pulvinatum hirsutiae canescens, capsulis immersis. *Dill. musc.* 395. t. 50. f. 65.
- Muscus* capillaceus lanuginosus minimus. T. *Vaill. paris.* 133. t. 29. f. 2.
- Muscus*, capitulo inflexo, pene acaulos. *Pet. gaz.* 127. t. 95. f. 15.
- Habitat in Europæ muris, rupibus.*
28. **BRYUM** antheris pendulis, foliis lanceolatis acuminato-setaceis, pedunculis longissimis. *Fl. suec.* 905. *Dalib. paris.* 322.
- Bryum* foliis subulatis, pedunculo longissimo, capitulo reflexo. *Fl. lapp.* 402.
- Bryum pendulum* ovatum caespitium & pilosum, seta bicolori. *Dill. musc.* 396. t. 50. f. 66.
- Muscus* capillaceus minimus, capitulo nutante, pediculo purpureo. *Vaill. paris.* 134. t. 29. f. 7.
- Muscus* trichodes, capitulo parvo reflexo, pediculo ima medietate rubro, summa luteo-viridi. *Morif. hist.* 3. p. 629. f. 15. t. 6. f. 15.
- Habitat in Europæ muris, sectis, glareosis.*
29. **BRYUM** antheris pendulis, foliis ovatis setiferis caespitium, pedunculis longissimis. *Fl. suec.* 906. *Dalib. paris.* 322.
- Bryum* foliis latiusculis congestis, capsulis longis nutantibus. *Dill. musc.* 398. t. 50. f. 67.
- Muscus* capillaceus major, capitulis crassioribus cylindraceis nutantibus. *Vaill. paris.* 134. t. 24. f. 6.
- Muscus* capillaris, foliis latiusculis congestis, capitulis longis reflexis. *Morif. hist.* 3. p. 629. f. 15. t. 6. f. 19.

B b b b

Mu-

Muscus capillaris, folio longiore. *Magn. port.* 138. t. 138.

Habitat in Europæ siccis, muris, glareosis.

Precedenti valde affinis.

carneum. 30. *BRYUM* antheris pendulis subglobosis, foliis acutis alternis. *Fl. suec.* 908.

Bryum lanceolatum pellucidum, capsulis subrotundis, pedunculis carnis. *Dill. musc.* 400. t. 50. f. 69.

Muscus stellaris humilis, capitulis pendulis. *Buxb. cent.* 2. p. 5. t. f. 6.

Habitat in Europæ aridis.

HYPNUM.

9

* *Frondibus pennatis.*

spinaforme. 1. *HYPNUM* fronde simplicissima: foliolis patentibus subulatis, pedunculis radicalibus.

Hypnum erectum non ramosum spinarum halecumarum. *Dill. musc.* 332. t. 43. f. 68.

Adiantum s. *Polytrichum aureum medium*, capitulo proprio pediculo insidente. *Sloan. jam.* 12. *bist.* 1. p. 68. t. 25. f. 4.

Habitat in Jamaica.

taxifolium. 2. *HYPNUM* fronde simplicissima pinnata lanceolata basi pedunculifera.

Hypnum foliis pinnatis contiguis linearibus, pedunculis radicalibus, capitulo subrotundo erecto. *Roy Lugdb.* 504.

Hypnum taxiforme minus: basi capsulifera. *Dill. musc.* 263. t. 34. f. 2.

Muscus pennatus, capitulis adianti. *Vaill. paris.* 136. t. 24. f. 11.

Habitat in Europæ umbrosis.

denticulatum. 3. *HYPNUM* fronde pinnata simplici: pinnis duplicatis basi pedunculifera.

Hypnum denticulatum pennatum, pinnulis duplicatis recurvis. *Dill. musc.* 266. t. 34. f. 5.

Muscus pennatus denticulatus minor. *Vaill. paris.* 137. t. 29. f. 8.

Muscus pennatus, capitulis adianti. *Morif. bist.* 3. p. 626. f. 15. t. 6. f. 36.

Habitat in Europa supra terram umbratam.

4 HYP-

4. HYPNUM fronde simplicissima pinnata lanceolata a-*bryoides*.
pice pedunculifera.
Hypnum taxiforme exiguum versus summitatem capsu-
liferum. *Dill. musc.* 262. t. 34. f. 1.
Muscus polytrichoides, exiguis capitulis in summis sur-
culis, f. foliis subrotundis erectis. *Vaill. parif.* 136. t.
24. f. 13. *Morif. hist.* 3. p. 629. f. 15. t. 6. f. 11.
Muscus terrestris minimus, foliis pennatim dispositis.
Buxb. cent. 1. p. 44. t. 64. f. 3.
Habitat in Europæ umbrosis.
Muscus minimus capituli ora magna ciliis rubris spe-
ctabilis.
5. HYPNUM fronde pinnata ramosa apice pedunculi-*acacioides*.
fera.
Hypnum pinnulis acaciæ, setis brevissimis. *Dill. musc.*
265. t. 34. f. 4.
Habitat in Patagonia.
6. HYPNUM fronde pinnata ramosa erecta medio pe-*adiantoides*.
dunculifera.
Hypnum taxiforme palustre ramosum majus & e-
rectum. *Dill. musc.* 264. t. 34. f. 3.
Muscus taxiformis ramosus. *Vaill. parif.* 136. t. 28. f. 5.
Muscus filicifolius f. pennatus minor, pinnulis plurimis
ad mediam costam adnexis. *Buxb. cent.* 2. p. 3. t.
1. f. 4.
Habitat in Europæ paludosis. &
7. HYPNUM fronde pinnata ramosa: foliolis imbrica-*complan-*
tis acutis complicatis compressis. *-atum.*
Hypnum pennatum compressum & splendens, capsulis
ovatis. *Dill. musc.* 268. t. 34. f. 7.
Muscus trichomanoides filicifolius splendens. *Vaill.*
parif. 139. t. 23. f. 4.
Muscus terrestris repens primæ speciei similis. *Morif.*
hist. 3. p. 625. f. 15. t. 5. f. 5.
Habitat in Europæ ad truncos arborum.
Confer Hypnum pennatum trichomanoides splendens ra-
mosum. *Dill. musc.* 269. t. 34. f. 8.
8. HYPNUM fronde pinnata ramosa, foliolis patentis-*ornithopodi-*
simis ovatis carinatis mucronatis. *-oides.*
Hypnum pennatum, ornithopodii pinnulis pellucidis.
Dill. musc. 270. t. 34. f. 9.
Habitat in Patagonia.

B b b b 2

* Sar-

* *Surculis vagis.*

- lucens.* 9. HYPNUM surculis ramosis, frondibus subpinnatis: foliolis punctatis.
Hypnum pennatum aquaticum lucens, longis latisque foliis. *Dill. musc.* 270. t. 34. f. 10.
Habitat in Europæ paludosis.
- undulatum.* 10. HYPNUM surculis ramosis, frondibus subpinnatis: foliolis undulatis complicatis.
Hypnum pennatum undulatum, lycopodii instar sparsum. *Dill. musc.* 271. t. 36. f. 11.
Hypnum repens crispum, ramulis compressis filicinorum more dispositis. *D. Hall. herb.* 105. t. 3. f. 5.
Habitat in Anglia, Hibernia, Helvetia.
Calyptra apice fusco distinctum. Dill.
- crispum.* 11. HYPNUM surculis ramosis, frondibus subpinnatis: foliolis undulatis planis.
Hypnum pennatum undulatum crispum, setis & capsulis brevibus. *Dill. elth.* 273. t. 36. f. 12.
Muscus terrestris major, ramulis compressis, foliis superficie crispis. *Morif. hist.* 3. p. 625. f. 15. t. 5. f. 10.
Habitat in Europæ locis saxosis.
- triquetrum.* 12. HYPNUM ramis vagis recurvis, foliis ovatis recurvatis patulis. *Fl. suæc.* 881.
Hypnum ramis inæqualibus, foliis laxè imbricatis triangularibus acutis. *Fl. lapp.* 409.
Hypnum vulgare triangulum maximum & pallidum. *Dill. musc.* 293. t. 38. f. 28.
Muscus ramosus major spermatophorus. *Vaill. parif.* 137. t. 28. f. 9.
Habitat in Europæ pratis, sylvis, sepibus.
- reticulatum.* 13. HYPNUM ramis vagis subrepentibus, foliolis ovatis mucronatis imbricatis.
Hypnum dentatum vulgatissimum operculis obtusis. *Dill. musc.* 295. t. 38. f. 29.
Muscus myosuroides rutabuli fructu. *Vaill. parif.* - t. 27. f. 8. & t. 23. f. 2.
Muscus denticulatus minor sericeus nostras, capitulis adianti. *Morif. hist.* 3. p. 626. f. 15. t. 6. f. 35.
Habitat in Europæ sepibus, sylvis ad arborum radices.

* *Saxo*

* *Surculis pennatis.*

14. HYPNUM furculis pennatis: ramulis distantibus, *filicinum*.
 foliolis imbricatis incurvis acutis secundis. *Fl. suec.*
 875.
 Hypnum repens filicinum crispum. *Dill. musc.* 282. t.
 36. f. 19.
 Muscus filicinus palustris. *Vaill. parif.* 138. t. 29. f. 9.
Habitat in Europæ humidiusculis.
- 15 HYPNUM furculis proliferis plano-pinnatis, pedun- *proliferum*.
 culis aggregatis. *Fl. lapp.* 405. *Fl. suec.* 872.
 Hypnum filicinum, tamarisci foliis minoribus non splen-
 dentibus. *Dill. musc.* 276. t. 35. f. 14.
 Muscus filicinus major. *Morif. bist.* 3. p. 625. f. 15. t.
 5. f. 20. *Tournef. inst.* 556. t. 326. f. C. *Vaill. pa-*
ris. 140. t. 25. f. 1.
Habitat in Europæ pratis, sylvis.
16. HYPNUM furculis subproliferis plano-pinnatis cu- *delicatulum*.
 spidatis, pedunculis aggregatis.
 Hypnum filicinum, tamarisci foliis minimis non splen-
 dentibus, setis capsulis brevioribus. *Dill. musc.* 546.
 t. 83. f. 6.
Habitat in America, Europa septentrionali.
Præcedenti simillima, sed longe tenerior & forte sola
varietas; antheras non vidi.
17. HYPNUM furculis plano-pinnatis continuatis, pe- *parietinum*.
 dunculis aggregatis. *Fl. lapp.* 406. *Fl. suec.* 873.
 Hypnum filicinum, tamarisci foliis majoribus splenden-
 tibus. *Dill. musc.* 274. t. 35. f. 13.
 Muscus filicinus major. *Buxb. cent.* 2. p. 2. t. 1. f. 2. &
 p. 3. t. 2. f. 1. & 4. t. 28. f. 1. & t. 29. f. 1.
Habitat in Europa frequens.
18. HYPNUM furculis subbipinnatis decumbentibus: ra- *prælongum*.
 mulis remotis, foliolis ovatis, antheris cernuis.
 Hypnum filicinum, triangularibus parvis foliis, præ-
 longum. *Dill. musc.* 278. t. 35. f. 15.
 Muscus terrestris minor, foliis crispis. *Buxb. cent.* 4.
 p. 37. t. 63. f. 3.
Habitat in Europæ terris, truncis, lignis.
Hæc species difficiliter determinatur, quam dignosci-
tur.
19. HYPNUM furculis pinnatis: ramulis approximatis: crista, ca-
 apicibus recurvis. *Fl. suec.* 871. *frons-*

B b b b 3

Hyp-

Hypnum ramulis plano-pinnatis: pinnis contiguis apicibus deflexis. *Fl. lapp.* 407.

Hypnum ilicinum, cristam castrensem repræsentans. *Dill. musc.* 284. t. 36. f. 20.

Muscus cristam castrensem repræsentans flavescens nemorosus ramosus cassubicus. *Morif. bist.* 3. p. 625. f. 15. t. 5. f. 8.

Muscus terrestris repens subflavus, foliolis crispis minoribus, ramulisque densius confertis. *Vaill. parif.* 141. t. 27. f. 14.

Muscus pennatus major, caulibus ramosis in summitate veluti pinnatus. *Loef. pruss.* 167. t. 42.

Habitat in Europa, Pennsylvania, ad radices Abietis.

abietinus.

20. HYPNUM furculis pinnatis teretiusculis remotis inæqualibus. *Fl. lapp.* 408. *Fl. suec.* 874.

Hypnum lutescens, alis subulatis tenacibus. *Dill. musc.* 280. t. 35. f. 17.

Muscus pennatus minor, caulibus ramosis, in summitate veluti spicatus. *Loef. pruss.* 167. t. 43. *Vaill. parif.* t. 29. f. 12.

Muscus terrestris, furculis filamentosis tenacibus abietinis semel tantum divisis. *Morif. bist.* 3. p. 625. f. 15. t. 5. f. 22.

Habitat in Europæ sylvis abietinis.

Pedunculi e media rachi furculi, solitarii, purpurei, recti, longitudine furculi, Calyptra erecta, subulata, pallida. Anthera rassa, recurvata magis quam in ullo alio, margine oris integro, intra quem Cilia brevia, patula.

* *Foliis reflexis.*

cupressiforme 21. HYPNUM furculo subpinnato, foliis secundis recurvis apice subulatis. *Fl. suec.* 880.

Hypnum crispum cupressiforme, foliis aduncis. *Dill. musc.* 287. t. 37. f. 23.

Muscus squamosus ramosus minor & crispus. *Vaill. parif.* 139. t. 27. f. 13.

Muscus terrestris aureus minor, flagellis clavisque brevioribus. *Morif. bist.* 3. p. 624. f. 15. t. 5. f. 8.

Habitat in Europæ sylvis ad arborum radices.

aduncum.

22. HYPNUM furculo erectiusculo subramoso, foliis secundis recurvatis subulatis, ramis recurvatis. *Fl. suec.* 879.

Hyp-

Hypnum palustre erectum summitatibus aduncis. Dill. musc. 292. t. 37. f. 26.

Habitat in Europæ uliginosis.

23. HYPNUM ramis vagis procumbentibus recurvis, foliis secundis acuminatis.

Hypnum scorpioides palustre magnum, lycopodii instar sparsum Dill. musc. 290. t. 37. f. 25.

Habitat in Anglia, Sueciæ paludibus profundis.

Muscus procumbens, ramis vagis fuscis, apicibus aduncis flavis.

24. HYPNUM furculis repentibus, ramis vagis tereti-viticulosum, foliis patulis acuminatis.

Hypnum subhirtutum, viticulis gracilibus erectis, capsulis teretibus. Dill. musc. 307. t. 39. f. 43.

Muscus squamosus, viticulis longioribus glabris. Vaill. parif. 137. t. 23. f. 1.

Muscus montanus, flagellis longis tenuibus cupressinis ut plurimum indivisis. Moris. hist. 3. p. 624. f. 15. t. 5. f. 7.

Muscus montanus gracilis ramosus, viticulis longioribus glabris. Pluk. alm. 255. t. 47. f. 4.

Muscus dens foliis juniperinis in cespitem aggestis. Loef. pruss. 49. t. 49.

Habitat in Europæ montosis aridis ad arbores.

25. HYPNUM ramis vagis, foliis ovatis apice subulatis undique reflexis. *Fl. succ. 878. **

Hypnum repens, triangularibus reflexis foliis, majus. Dill. musc. 303. t. 39. f. 38.

Muscus terrestris erectior, foliis reflexis. Moris. hist. 3. p. 524. f. 15. t. 6. f. 2.

26. Hypnum repens, triangularibus reflexis foliis, minus. *Dill. musc. 304. t. 39. f. 39.*

Muscus erectus, foliis reflexis. Vaill. parif. 139. t. 27. f. 5.

Habitat in Europæ subhumidis.

27. HYPNUM furculis reptantibus, ramis vagis erectis, loreum, foliis secundis, antheris subrotundis.

Hypnum loreum montanum, capsulis subrotundis. Dill. musc. 305. t. 39. f. 40.

Muscus squamosus major, foliis angustioribus acutissimis. Vaill. parif. 138. t. 25. f. 2.

Muscus repens major, foliis & flagellis longis & tenuibus donatus. Moris. hist. 3. p. 626. f. 15. t. 5. f. 24.

B b b b 4

Mu-

Muscus montanus. Tabern. ic. 809.

Musco denticulato similis. Bamb. pin. 360.

Habitat in Europæ montosis.

* *Surculis dendroidibus f. fasciculatis.*

dendroides. 27. HYPNUM furculo erecto, ramis fasciculatis terminalibus simpliciusculis, antheris erectis. *Fl. suec. 886.*
Hypnum caule erecto superne undique ramoso. *Fl. lapp. 411.*

Hypnum dendroides sericeum, setis & capsulis longioribus erectis. *Dill. musc. 313. t. 40. f. 48.*

Muscus squamosus ramulosus erectus alopecuroides. *Tournef. inst. 554. t. 326. f. B. Vaill. paris. 137. t. 26. f. 6.*

Muscus dendroides elatior, ramulis crebris minus furculosis, capitulis pedunculis brevibus infidentibus. *Morif. bist. 3. p. 626. f. 15. t. 5. f. 31.*

Habitat in Europæ pratis sylvisque subhumidis.

alopecuroides. 28. HYPNUM furculo erecto, ramis fasciculatis terminalibus subdivisis, antheris subnutantibus. *Fl. suec. 887.**

Hypnum dendroides obscurius, setis & capsulis brevioribus subnutantibus. *Dill. musc. 315. t. 41. f. 49.*

Muscus dendroides sylvarum erectus, ramulis kali æmulis, radice repente. *Morif. bist. 3. p. 626. f. 15. t. 5. f. 30.*

Muscus squamosus alopecuroides, flagellis recurvis. *Vaill. paris. 137. t. 23. f. 5.*

Habitat in Europæ sylvis humidiusculis.

* *Surculis teretiusculis.*

curtipendulum. 29. HYPNUM furculis vagis teretibus, foliis ovatis acutis patulis, antheris pendulis. *Fl. suec. 877.*

Hypnum dentatum curtipendulum, viticulis rigidis. *Dill. musc. 333. t. 43. f. 69.*

Habitat in Europæ, America, ad radices arborum, saxorum.

pinum. 30. HYPNUM furculis pinnato-sparsis subulatis, foliis ovatis obtusis conniventibus. *Fl. suec. 876.*

Hypnum ramis inæqualibus sparsis, foliis ovatis imbricatis, capitulis obliquis. *Fl. lapp. 410.*

Hypnum cupressiforme vulgare, foliis obtusis. *Dill. musc. 309. t. 40. f. 45.*

Muscus squamosus cupressiformis. *Vaill. paris. 138. t. 28. f. 3.*

Habitat in Europæ pascuis, sylvis.

31.

31. HYPNUM furculis ramisque vagis teretibus erectiis- illecebrum.
 iculis obtusis.

Bryum caule erecto ramoso, foliis ovatis undique imbricatis. *Fl. lapp.* 317.

Hypnum cupressiforme rotundius vel illecebræ æmulum. *Dill. musc.* 311. t. 40. f. 46.

Muscus terrestris, furculis kali geniculati aut illecebræ æmulis, foliis subrotundis squamatim incumben- tibus.

Vaill. parif. 137. t. 25. f. 7.

Habitat in Europa & Americæ septentrionalis pascuis.

32. HYPNUM furculis teretibus ramosis, foliolis acutis patulis distantibus.

Hypnum aquaticum, flagellis & teretibus & pennatis.

Dill. musc. 308. t. 40. f. 44.

Hypnum ramosum fluitans pennatum. *Raj. angl.* 3. p. 81.

Habitat in Europa, ad fluviorum ripas.

33. HYPNUM furculis vagis: apice foliis convolutis cuspidatum, acuminato. *Fl. succ.* 884.

Hypnum palustre extremitatibus cuspidatis & pungentibus. *Dill. musc.* 300. t. 39. f. 34.

Muscus saxatilis repens, coma sparsa. *Buxb. cent.* 2. p. 6. t. 3. f. 1. 2. 3. 4.

Habitat in Europæ paludibus muscosis, quas replet.

* *Surculis confertis.*

34. HYPNUM furculo repente, ramis confertis erectis, sericeum, foliis subulatis, antheris erectis. *Fl. lapp.* 412. *Fl. succ.* 882. *Dalib. parif.* 326.

Hypnum vulgare sericeum recurvum, capsulis erectis cuspidatis. *Dill. musc.* 323. t. 42. f. 59.

Muscus terrestris luteo-viridens sericeus repens. *Morif. hist.* 3. p. 626. f. 15. t. 5. f. 25.

Muscus arboreus splendens sericeus. *Vaill. parif.* 132. t. 27. f. 3.

Habitat in Europa, ad truncos, muros, inque campis.

35. HYPNUM furculo repente, ramis confertis erectis, velutinum, foliis subulatis, antheris submutantibus. *Fl. succ.* 883.

Hypnum velutinum, capsulis ovatis cernuis. *Dill. musc.* 326. t. 42. f. 61.

Muscus squamosus ramosus tenuior, capitulis incurvis. *Vaill. parif.* 138. t. 26. f. 9.

B b b b s

Mu-

- Muscus terrestris minor omnium vulgatissimus. *Buxb. cent. 4. p. 35. t. 62. f. 23.*
Habitat in Europa ad radices arborum umbrosas.
- serpens.* 36. HYPNUM furculis repentibus, ramis filiformibus, foliis oblitterat s.
 Hypnum trichodes serpens, setis & capsulis longis erectis. *Dill. musc. 329. t. 42. f. 64.*
 Muscus terrestris omnium minimus, capitulis majusculis oblongis erectis. *Morif. bist. 3. p. 625. f. 15. t. 5. f. 14. 21. Vaill. parif. 138. t. 28. f. 2. 6. 7. 8.*
Habitat in Europa, Virginia, ad candices, ligna, saxa.
- sciuroides.* 37. HYPNUM furculis erectis ramosis incurvatis.
 Hypnum arboreum sciuroides. *Dill. musc. 319. t. 41. f. 54.*
 Muscus arboreus splendens myosuroides. *Vaill. parif. t. 27. f. 12.*
 Muscus repens, ferici modo lucens, viticulis longioribus erectis. *Morif. bist. 3. p. 626. f. 15. t. 5. f. 27.*
Habitat in Europæ truncis arborum.
 Pedunculi maturi semis instar torti; femina furculi distincti crescunt.
- myosuroides.* 38. HYPNUM furculis ramosissimis: ramis subulatis utrinque attenuatis teretibus.
 Hypnum myosuroides tenuius, capsulis nutantibus. *Dill. musc. 317. t. 41. f. 51.*
 Muscus trichodes humilis ramosus, capitulis oblongis tumidiusculis donatus. *Morif. bist. 3. p. 628. f. 15. t. 6. f. 3.*
 Muscus squamosus minor myosuroides, capitulis incurvis. *Vaill. parif. 137. t. 27. f. 6.*
Habitat in Europa supra lapides arborumque radices.
- clavellatum.* 39. HYPNUM repens, ramulis erectis confertissimis, antheris incurvis: operculis inflexis.
 Hypnum clavellatum parvum repens, setis brevibus. *Dill. musc. 551. t. 81. f. 17.*
Habitat in Europæ lignis putridis.
- julaceum.* 40. HYPNUM ramis erectis teretibus imbricatis obtusis, perichætiis longitudine fere pedunculorum.
 Hypnum julaceum, perichætio setas æquante. *Dill. musc. 321. t. 41. f. 5. 6.*
 Hypnum fabinæ foliis, capitulis atro-fuscis, pediculis brevibus. *Gron. virg. 125.*
Habitat in America septentrionali.

ALGÆ.

JUNGERMANNIA. / 0

* *Frondibus pinnatis secundis.*

1. JUNGERMANNIA frondibus simpliciter pinnatis: *asplenoides*.
foliolis ovatis subciliatis.

Jungermannia major, foliis brevioribus & obtusioribus
non dentatis. *Mich. gen.* 8. t. 5. f. 3. o. o.

Lichenastrum asplenii facie, pinnis laxioribus. *Dill.*
musc. 482. t. 69. f. 5.

Muscus bifolius procumbens, foliis subrotundis. *Laf.*
pruss. 167. t. 47.

2. Jungermannia caule ramoso repente, foliis imbricatis
ovato-crenulatis: alternis minoribus. *Roy. lugdb.* 508.

Jungermannia major, foliis subrotundis tenuissime den-
ticulatis. *Mich. gen.* 7. t. 5. f. 1. 2.

Lichenastrum asplenii facie, pinnis confertioribus. *Dill.*
musc. 483. t. 69. f. 6.

Hepaticoides polytriche facie. *Vaill. parif.* 99. t. 19. f. 7.

Muscus trichomanoides, foliis rotundioribus pellucidis
squamatim conjuncte sibi incumbentibus. *Morif. bist.*
3. p. 627. f. 15. t. 6. f. 42.

Habitat in Europæ Indiæ udis umbrosis.

2. JUNGERMANNIA frondibus simpliciter pinnatis: *viticniosa*.
foliolis subulatis.

Jungermannia terrestris, viticulis longis, foliis perexi-
guis densissimis ex rotunditate acuminatis. *Mich. gen.*
8. t. 5. f. 4.

Lichenastrum trichomanis facie e basi & medio florens.
Dill. musc. 484. t. 69. f. 7.

Habitat in Europæ udis umbrosis sylvis.

3. JUNGERMANNIA frondibus simpliciter pinnatis: *polyanthes*.
foliolis integerrimis imbricatis convexis.

Jungermannia palustris repens, foliis densissimis ex ro-
tunditate acuminatis. *Mich. gen.* 8. t. 5. f. 5.

Lichenastrum trichomanes facie, polyanthemum breve
& repens. *Gill. musc.* 486. t. 70. f. 5.

Habitat in Europæ palustribus.

4. JUNGERMANNIA frondibus simpliciter pinnatis *lanceolata*.
lanceolatis apice floriferis, foliolis integerrimis.

Jungermannia palustris minima repens, foliis subrotun-
dis densissimis late virentibus. *Mich. gen.* 8. t. 5. f.
67. Li

Lichenastrum trichomanis facie minus, ab extremitate
florens. *Dill. musc.* 486. t. 70. f. 10.
Habitat in Europæ humidis umbrosis.

- bidentata.* 5. **JUNGERMANNIA** frondibus simpliciter pinnatis
apice floriferis, foliolis bidentatis.
Jungermannia major repens, foliis bifidis. *Mich. gen.*
5. t. 5. f. 12.
Lichenastrum, pinnulis acutioribus & concavis bifidis, ma-
jus. *Dill. musc.* 487. t. 70. f. 11.
Hepaticoides, polytrichi facie, foliis bifidis, major. *Vaill.*
parif. 99. t. 19. f. 8.
Muscus lichenoides, foliis pennatis bifidis, minor. *Mo-*
rif. bist. 3. p. 627. f. 15. t. 6. f. 47.
Habitat in Europæ ericetis umbrosis.

- bicuspidata.* 6. **JUNGERMANNIA** frondibus simpliciter pinnatis
medio floriferis, foliolis bidentatis.
Jungermannia minima repens, foliis bifidis, vagina flo-
rum cylindracea. *Mich. gen.* 9. t. 6. f. 17.
Lichenastrum, pinnulis acutissime bifidis, minimum. *Dill.*
musc. 288. t. 70. f. 13.
Habitat in Europæ umbrosis humidis.

* *Frondibus pinnatis: pinnulis auriculatis superne
tectis.*

- undulata.* 7. **JUNGERMANNIA** frondibus supra bipinnatis a-
pice floriferis; foliolis subrotundis integerrimis.
Lichenastrum pinnulis auriculatis majoribus & non cre-
natis. *Dill. musc.* 490. t. 71. f. 17.
Hepatica saxatilis undulata seminifera. *Vaill. parif.* 98.
t. 19. f. 6.
Habitat in Europa.

8. **JUNGERMANNIA** frondibus supra bipinnatis apice
floriferis, foliolis ciliatis.
Jungermannia nemorosa, foliis acutioribus auritis tenu-
issime denticulatis, flore insidente pediculo brevior.
Mich. gen. 7. t. 5. f. 8.
Lichenastrum auriculatum, pinnis minoribus crenatis.
Dill. musc. 490. t. 7. f. 18.
Habitat in Europæ sylvis.

- resupinata.* 9. **JUNGERMANNIA** frondibus supra bipinnatis in-
ferne floriferis: foliolis crenulatis imbricatis rotundis.
Jungermannia furculosa resupinata, foliolis pinnato-im-
bricatis

bricatis deflexis rotundis integris: auricula laterali. *Fl. suec.* 926.

Lichenastrum imbricatum, pinnulis rotundis, crispum.
Dill. musc. 491. t. 71. f. 19.

Habitat in Europæ rupibus.

10. *JUNGERMANNIA* frondibus supra bipinnatis *a- albicans*, pice floriferis: foliolis linearibus recurvatis.

Jungermannia furculis pinnatis: foliolis recurvis planis auriculatis. *Fl. suec.* 916.

Lichenastrum auriculatum: pinnulis angustis planis recurvis. *Dill. musc.* 492. t. 71. f. 20.

Hepaticoides albescent, foliis pinnatis *Vaill. parif.* 100. t. 19. f. 5.

Habitat in Europæ umbrosis.

11. *JUNGERMANNIA* frondibus subtus bipinnatis, *trilobato*, foliolis quadratis obsolete trilobis.

Jungermannia furculis planis pinnatis: foliolis obsolete trilobis auriculatis. *Fl. suec.* 917.

Jungermannia foliis latiusculis obtusis undulatis & veluti angulosis. *Mich. gen.* 7. t. 5. f. 10.

Lichenastrum pinnulis obtusè trifidis nervo geniculato.
Dill. musc. 493. t. 71. f. 22.

Habitat in Svecia, Anglia, Italia.

12. *JUNGERMANNIA* frondibus subtus bipinnatis *a- reptans*, pice reptantibus, foliolis quadridentatis.

Lichenastrum multistidum exiguum ad basin florens per siccitatem imbricatum. *Dill. musc.* 494. t. 71. f. 24.

Habitat in Europæ Ericetis adis.

* *Frondibus imbricatis.*

13. *JUNGERMANNIA* furculis repentibus: foliolis *complanata*, inferne auriculatis duplicato-imbricatis, ramis æqualibus. *Fl. suec.* 922.

Jungermannia foliis circinatis imbricatis dispositis ex viridi flavescens. *Mich. gen.* 7. t. 5. f. 21.

Lichenastrum imbricatum majus, squamis compressis & planis. *Dill. musc.* 496. t. 72. f. 26.

Habitat in Europa ad arborum truncos.

14. *JUNGERMANNIA* furculis repentibus, foliis inferne auriculatis duplicato-imbricatis, ramis apice latioribus. *Fl. suec.* 919.

Liche-

Lichenastrum imbricatum minus, squamis convexo-concavis. *Dill. musc.* 497. t. 72. f. 27.

Hepaticoides, furculis & foliolis thuyæ instar compressis, minor. *Vaill. parif.* 99. t. 19. f. 10.

Muscoides minimum, foliis alternis: superioribus circinatis, inferioribus cucullatis, floræ pediculo ferme carente. *Mich. gen.* 10. t. 6. f. 6.

Habitat in Europa, America, ad arborum truncos.

- tamarisci.* 15. **JUNGERMANNIA** foliis imbricatis serie duplici: superioribus subrotundis convexis obtusis quadruplo-majoribus. *Fl. lapp.* 425. *Fl. suec.* 923. *Roy. lugd.* 505

Lichenastrum imbricatum, tamarisci narbonensis facie. *Dill. musc.* 499. t. 72. f. 31.

Hepaticoides quæ *Muscus trichomanoides* terrestris minor floridus. *Vaill. parif.* 100. t. 23. f. 10.

Muscoides squamosum saxatile nigro-purpureum, furculis angustioribus, foliis circinatis minoribus. *Mich. gen.* 10. t. 6. f. 5.

Habitat in Europa ad truncos arborum, rupes.

- platyphylla.* 16. **JUNGERMANNIA** furculis procumbentibus subtus imbricatis, foliolis cordatis acutis.

Lichenastrum arboris vitæ facie, foliis minus rotundis. *Dill. musc.* 501. t. 72. f. 32.

Hepaticoides, furculis & foliolis thuyæ instar compressis, major. *Vaill. parif.* t. 19. f. 9.

Muscoides squamosum atrovirens, foliis subrotundis. *Mich. gen.* 9. t. 6. f. 3. 4.

Muscus muralis floridus, foliis subrotundis creberrime imbricatis dispositis. *Morif. hist.* 3. p. 627. f. 15. t. 6. f. 44.

Habitat in Europæ & Americæ septentrionalis sylvis.

- ciliaris.* 17. **JUNGERMANNIA** furculis repentibus, foliolis duplicato-imbricatis: interne auriculatis ciliatis. *Fl. suec.* 918. *

Jungermannia caule ramoso, furculis compressis, foliis imbricatis serie duplici ovato-acutis ciliatis. *Fl. lapp.* 426.

Lichenastrum filicinum pulchrum villosum. *Dill. musc.* 503. t. 73. f. 35.

Muscus palustris, absinthii folio. *Vaill. parif.* 140. t. 26. f. 11?

Habitat in Europa passim.

18. JUN-

18. JUNGERMANNIA furculis erectiusculis bifari- *varia*.
 am imbricatis, foliis bipartitis.
 Jungermannia repens, foliis cordatis carinatis. *Mich.*
gen. 8. t. 5. f. 9.
 Lichenastrum foliis variis. *Dill. musc.* 505. t. 73. f.
 36.
Habitat in Europæ sylvis ericetis.
 * *Surculis undique imbricatis foliolis sparsis.*
19. JUNGERMANNIA furculis teretibus, foliolis un- *gulacea*.
 dique imbricatis, floribus pedunculatis.
 Lichenastrum alpinum, bryjulacei argentei facie. *Dill.*
musc. 506. t. 73. f. 38.
Habitat in Alpibus Britannicæ.
20. JUNGERMANNIA furculis teretibus, foliolis *rupestris*.
 subulatis secundis.
 Jungermannia furculosa erectiuscula, foliolis undique
 imbricatis acuminatis hinc reflexis. *Fl. suec.* 920. *
 Lichenastrum alpinum nigricans, foliis capillaceis reflexis. *Dill. musc.* 507. t. 73. f. 40.
Habitat in Europæ frigidis rupibus humentibus.
21. JUNGERMANNIA furculis teretibus, foliolis ca- *trichophylla*.
 pillaceis æqualibus.
 Jungermannia furculis erectiusculis, foliolis imbricatis
 acuminatis undique erectis. *Fl. suec.* 921.
 Lichenastrum trichodes minimum in extremitate florens.
Dill. musc. 505. t. 73. f. 37.
Habitat in Europæ frigidis rupibus.
22. JUNGERMANNIA furculis teretibus, foliolis o- *alpina*.
 vatīs patentibus, calycibus imbricatis.
 Lichenastrum alpinum atrorubens teres, calycibus squa-
 mosis. *Dill. musc.* 506. t. 73. f. 39.
Habitat in Alpibus Britannicæ, inque rupibus Sueciæ.
 * *Acaules frondibus simplicibus.*
23. JUNGERMANNIA acaulis, foliolo frondis innato. *epiphylla*.
Fl. suec. 930. *
 Jungermannia hypophylla. *It. wigth.* 213.
 Marilca major atrovirens, floribus albicantibus e folio-
 rum medio egredientibus. *Mich. gen.* 5. t. 4. f. 1.
 Lichenastrum capitulis rotundis e foliorum medio ena-
 scentibus. *Dill. musc.* 508. t. 74. f. 41.
 Lichen petræus, caliculo calceato. *Barrb. pin.* 362.
 Lichen

Lichen alter minor, folio calceato. *Col. ecphr.* 1. p. 331. t. 332.

Habitat in Europæ ripis elatioribus ambrosiæ adis.

viridis.

24. JUNGERMANNIA acaulis, fronde oblonga sinuata pingui. *Fl. succ.* 929.

Martileia media pinguis pallide virens, floribus majoribus nigricantibus ad foliorum latera egredientibus.

Mich. gen. 5. t. 4. f. 2.

Lichenastrum capitulis oblongis juxta foliorum divissuras nascentibus. *Dill. musc.* 509. t. 74. f. 42.

Lichen parvus erectus, foliis profunde laciniatis. *Pluk. alm.* 216. t. 42. f. 2.

Habitat in Europæ paludibus.

multifida.

25. JUNGERMANNIA acaulis, fronde bipinnatifida. Lichenastrum ambrosiæ divissura. *Dill. musc.* 511. t. 74. f. 43.

Habitat in Angliæ ericetis.

furcatum.

26. JUNGERMANNIA acaulis, fronde lineari ramosa: extremitatibus furcatis obtusiusculis. *Fl. succ.* 928.

Martilia minima angustifolia, floribus nigricantibus ex inferiori foliorum parte e subhirsuta & turbinata vagina. *Mich. gen.* 5. t. 4. f. 4.

Lichenastrum tenuifolium furcatum, thecis globosis pilosis. *Dill. musc.* 512. t. 74. f. 45.

Hepatica arborea globulifera. *Vaill. parif.* 98. t. 23. f. 11.

Habitat in Europa ad truncos, rupes & e terra.

passila.

27. JUNGERMANNIA acaulis, fronde subbipinnatifida: lobis imbricatis, perichætio plicato.

Lichenastrum exiguum, capitulis nigris lucidis e cotylis parvis nascentibus. *Dill. musc.* 513. t. 74. f. 46.

Habitat in Europæ rupibus, ericetis humentibus.

TARGIONIA.

hypophylla.

1. TARGIONIA.

Targionia minima & vulgaris. *Mich. gen.* 3. t. 3.

Lichen petræus minimus, tractu orobi *Bamb. pin.* 362. *Dill. musc.* 532. t. 78. f. 9.

Lichen alter acaulos hypophyllocarpus. *Col. ecphr.* 1. p. 331. f. 333.

Lichen terrestris minor seminifer. *Buxb. cent.* 1. p. 41. t. 61. t. 4.

Habitat in Italia, Hispania, Constantinopoli.

MAR.

MARCHANTIA.

/ 2

1. MARCHANTIA calyce communi decemfido. *Fl. polymorpha.*
Swec. 931. *Fl. lapp.* 422. *Roy. lugdb.* 507.

Marchantia major, capitulo stellato: radiis teretibus.

Mich. gen. 2. t. 1. f. 1.

Lichen fontanus major stellatus æque ac umbellatus & cyathophorus. *Dill. musc.* 523. t. 76. f. 6.

Lichen petræus latifolius, f. Hepatica fontana. *Bauh. pin.* 362.

Lichen. *Fuchf. hist.* 179. *Cam. epit.* 782.

2. Marchantia stellata. *March. act. parif.* 1713. p. 307. t. 5.

Marchantia capitulo stellato: radiis teretibus. *Mich. gen.* 2. t. 1. f. 2.

Lichen domesticus minor stellatus æque ac umbellatus & cyathophorus. *Dill. musc.* 527. t. 77. f. 7.

Lichen petræus stellatus. *Bauh. pin.* 362.

Hepatica secunda. *Tabern. ic.* 815.

3. Marchantia calyce communi octopartito: laciniis e plano convexis. *Fl. lapp.* 423. *Roy. lugdb.* 507.

Marchantia capitulo non dissecto. *Mich. gen.* 1. t. 1. f. 5.

Lichen petræus umbellatus. *Bauh. pin.* 362.

Hepatica tertia. *Tabern. ic.* 815.

Habitat in Europa juxta aquas, locis umbrosis.

4. MARCHANTIA calyce communi dimidiato palma-chenopoda. to quadrifido.

Lichen anapodocarpus. *Plum. fil.* 143. t. 142. *Dill. musc.* 531. t. 77. f. 8.

Habitat in Martinica.

5. MARCHANTIA calyce communi quadripartito: laciniis tubulosis. *Hort. cliff.* 477.

Lichen seminifer lunulatus, florifer pileatus tandem cruciatus. *Dill. musc.* 521. t. 75. f. 5.

Lichen hepaticus, capitulis quadrifidis. *Buxb. cent.* 1. t. 6.

Lunaria vulgaris. *Mich. gen.* 4. t. 4.

Habitat in Europæ umbrosis.

6. MARCHANTIA calyce communi hemispherico a-tenella. piculato: margine radiato.

Lichen pileatus parvus carinatus, capitulis fimbriatis.

Dill. musc. 521. t. 75. f. 4.

Habitat in Virginia. D. Gronovius.

C c c c

5. Mar-

hemisphærica. 5. MARCHANTIA calyce communi quinquefido hemisphærico, perichætio nullo. *Fl. succ.* 932. *Fl. lapp.* 424. *Roy. lugdb.* 507.

Hepatica media, capitulo hemisphærico. *Mich. gen.* 3. t. 2. f. 2.

Lichen pileatus parvus, foliis crenatus. *Dill. musc.* 519. t. 75. f. 2.

Lichen petreus, capitulis stellatis crassis, pediculis brevibus insidentibus *Buxb. cent.* 2. p. 5. t. 5. f. 1.

Habitat in Europæ paludosis.

Perichætia in hac specie nulla, nec Calyces seminiferi.

conica. 6. MARCHANTIA calyce communi quinqueloculati subovato.

Lichen vulgaris major pileatus & verrucosus. *Dill. musc.* 516. t. 75. f. 1.

Lichen petreus, caliculo pileolum sustinente. *Bauh. pin.* 362.

Lichen plinii primus & pileatus. *Col. ecphr.* 1. p. 330. t. 331.

Hepatica vulgaris maior s. officinarum italiz. *Mich. gen.* 3. t. 2. f. 1.

Habitat in Europæ locis umbrosis, ripis elevatis.

androgyna. 7. MARCHANTIA calyce communi integro hemisphærico.

Lichen pileatus angustifolius dichotomus. *Dill. musc.* 520. t. 75. f. 3.

Hepatica minor angustifolia, capitulo hemisphærico. *Mich. gen.* 3. t. 2. f. 3.

Habitat in Italia, Jamaica inter saxorum scopulones. Monoica s. Androgyna est hæc species.

BLASIA.

pusilla.

1. BLASIA. *Fl. succ.* 933.

Blasia pusilla, lichenis pyxidati facie. *Mich. gen.* 14. t. 7.

Mnium lichenis facie. *Dill. musc.* 237. t. 31. f. 7.

Habitat in Europa ad latera fossarum, solo ex arena sterili.

RICCIA.

crystallina.

1. RICCIA frondibus superficie papillofis.

Riccia foliis aspergine crystallina pertusis, margine incrassatis. *Fl. succ.* 935. *It. æl.* 154.

Riccia minima & minor pinguis aspergine crystallina pertusa. *Mich. gen.* 107. t. 57. f. 7. 3.

Hepa-

Hepatica palustris, lobis inflatis. *Vaill. parif.* 98. t. 19. f. 2.

Lichen palustris, rutæ folio. *Gelf. apf.* 28. *Dill. musc.* 355. t. 78. b. 12.

Habitat in Europæ locis subhumidis.

2. **RICCIA** frondibus glabris bipartitis acutis. *Fl. suec. minima.* 934.

Riccia minima nitida, segmentis angustioribus acutis.

Mich. gen. 107. t. 57. f. 6.

Lichen omnium minimus, foliolis fissis super terram expansis. *Dill. musc.* 534. t. 78. f. 11.

Habitat in Europæ inundatis.

3. **RICCIA** frondibus glabris canaliculatis bilobis obtusis.

Riccia minima pingvis, foliis latiusculis ample sulcatis e glauco virescentibus. *Mich. gen.* 107. t. 57. f. 4.

Hepatica palustris bifurca, locis brevioribus carinatis. *Vaill. parif.* 98. t. 19. f. 1.

Lichen minimus, foliis venosis bifariam vel trifariam se dividendo progredientibus. *B. Dill. musc.* 533. t. 78. f. 10.

Lichen terrestris minor, foliis rutæ. *Buxb. cent.* 2. p. 10. t. 5. f. 5.

Habitat in Anglia, Italia, Gallia.

4. **RICCIA** frondibus dichotomis lineari-filiformibus. *It. fluitans.* *Scan.* 241.

Lichenastrum aquaticum fluitans tenuifolium furcatum.

Dill. musc. 213. t. 74. f. 47.

Hepatica palustris dichotoma: segmentis angustioribus. *Vaill. parif.* 98. t. 10. t. 3.

Habitat in Europæ fossis, piscinis, ad ripas supra aquam extensa.

ANTHOCEROS. 15

1. **ANTHOCEROS** frondibus indivisis sinuatis puncta-punctatis.

Anthoceros foliis minoribus magis claciniatis. *Dill. musc.* 476. t. 68. f. 1.

Anthoceros minor, foliis magis carinatis atque eleganter crenatis, subtus incurvatis. *Mich. gen.* 11. t. 7. f. 2.

Habitat in Angliæ, Italiæ uliginosis, umbrosis.

2. **ANTHOCEROS** frondibus indivisis sinuatis lævibus. *lævis.*

Anthoceros. *Hort. cliff.* 477. *Roy. Lugdb.* 507.

C c c 2

Antho-

Anthoceros foliis majoribus minus laciniatis. Dill. musc.
476. t. 68. f. 2.

Anthoceros major. Mich. gen. 11. t. 7. f. 1.

Lichen hepaticus, pediculis gramineis. Buxb. cens. 1
p. 40. t. 61. f. 1.

Habitat in Europa & America boreali.

- multifidus.* 3. **ANTHOCEROS** frondibus bipinnatifidis linearibus
Anthoceros folio tenuissimo multifido. Dill. musc. 477
t. 88. f. 4.
Habitat in Germania.

LICHEN.

* *Leprosi tuberculati.*

- scriptus.* 1. **LICHEN** leprosus albicans, lineolis nigris ramosis
characteriformibus. *Fl. suec.* 941.
Lichenoides crusta tenuissima, peregrinis veluti literis
inscripta. Dill. musc. 128. t. 18. f. 1.
Habitat in Europæ corticibus arborum.
- geographicus* 2. **LICHEN** leprosus flavescens: lineolis nigris mappam
referens. *Fl. suec.* 940.
Lichenoides nigro-flavum tabulæ geographicæ instar pi-
ctum. Dill. musc. 126. t. 18. f. 5.
Habitat in Europæ rupibus altis.
- cucurbitum.* 3. **LICHEN** leprosus albicans, lineolis simplicibus pun-
ctisque nigris confertis.
Lichenoides punctatum & rugosum nigrum. Dill. giff
125. t. 18. f. 2.
Habitat in Europæ sylvis supra arborum trunco.
- sanguinarius.* 4. **LICHEN** leprosus cinereo-virescens, tuberculis atris.
Fl. suec. 938.
Lichenoides leprosum, crusta cinereo-virescente, tuber-
culis integerrimis. Dill. musc. 126. t. 18. f. 3.
Habitat in Europæ rupibus truncisque arborum.
Tubercula majuscula, gibba, atra, at diffracta intus
rubra sunt.
- fusco-ater.* 5. **LICHEN** leprosus fuscus, tuberculis atris.
Habitat in Europæ rupibus.
- calcareus.* 6. **LICHEN** leprosus candidus, tuberculis atris. *Fl. suec.*
937. *It. wigsh.* 29. 211. *It. gotl.* 183.
Lichenoides tartareum tinctorium candidum, tuberculis
atris. Dill. musc. 128. t. 18. f. 8.
Habitat in Europæ rupibus marmoreis.

7. LICHEN leprosus ater, tuberculis viridibus confertis *atro-virens.*
Fl. succ. 939.
Habitat in Europæ rupibus.
Tubercula parva, flavo-virescentia, conferta, ut tota a-
rea flavo-virens conspiciatur, quam cingit margo ni-
ger.
8. LICHEN leprosus niger, tuberculis albis. *atro-albus,*
Habitat in Alpium rupibus.
9. LICHEN leprosus flavus, tuberculis rubris. *ventosus,*
Habitat in Alpium rupibus.
10. LICHEN leprosus albus, tuberculis albis farinaceis, *fragilis,*
It. scan. 59.
Habitat in Europæ, vestiens truncos Fagi.
11. LICHEN leprosus cinereus, tuberculis albidis rugo-
sis.
Lichen geographicus. It. scan. 48.
Habitat in Carpini truncis, ramis.
12. LICHEN leprosus candidus, tuberculis incarnatis. *erictorum,*
Fl. succ. 936. It. scan. 34.
Lichen crustaceus, calyce convexo foliis majore. Fl.
lapp. 455.
Coralloides fungiforme carneum, basi leprosa. Dill.
musc. 76. t. 14. f. 1.
Fungus omnium minimus turbinatus croceus. Bocc.
mus. 2. p. 150. t. 10.
Fungilli incarnati coloris minuti musco insidentes.
Mentz. pag. 1. 6.
Habitat in Europæ sylvis sterilissimis, glareosis subnudis,
aliginosis.

* *Leprosi scutellati.*

13. LICHEN crustaceus flavus, scutellis luteis. *candelarius.*
Byssus farinacea flava. It. al. 30. It. w:goth. 159.
Habitat in Europæ parietibus, muris, truncis arborum,
presertim Quercus.
14. LICHEN crustaceus ex albido virefcens, scutellis *tartareis.*
flavescentibus: margine albo.
Lichen leprosus candidus, scutellis fuscis: margine al-
bo. Fl. succ. 942.
Lichen tinctorius ruber. It. al. 29. It. w:goth. 146.

C c c c 3

Li

Lichenoides tartareum farinaceum. scutellarum umbone fusco. *Dill. musc.* 131. t. 18. f. 12.

Lichen crustaceus saxatilis farinaceus verrucosus candidus omnium crassissimus, receptaculis florum nigris. *Mich. gen.* 96. t. 52. f. 6.

Habitat in Europa ad parietes rupium.

pallidus. 15. LICHEN crustaceus albicans, scutellis pallidis. *Fl. succ.* 944.

Lichenoides crustosum orbiculare incanum. *Dill. musc.* 135. t. 18. f. 17.

Habitat in Europæ corticibus arborum.

subfuscus. 16. LICHEN crustaceus albicans, scutellis subfuscis: margine cinereo subcrenato. *Fl. succ.* 943.

Lichenoides crustaceum & leprosum, scutellis subfuscis. *Dill. musc.* 134. t. 18. f. 16.

Habitat in Europa, arboribus & rupibus innascens.

appaliensis. 17. LICHEN crustaceus, foliis subulatis stratis.

Habitat in summis Jugis supra terram sterilissimam campi polonici, Upsalia, nec alibi nobis visus.

Magnitudo Lichens saxatilis, cinereo-cibus, constans Setis rectis, albis, emarcidis, prostratis, inequalibus, confusis, simplicissimis, longitudine unguis, fragilissimis, sæpe pluribus basi coalitis. Scutella alba, margine obtuso, majuscula, radicalia, nec setis insidentia.

* *Imbricatus.*

centrifugus. 18. LICHEN imbricatus, foliolis multifidis obsolete laticibus albidis centrifugis, scutellis rufo-fuscis. *Fl. succ.* 945.

Lichen foliis planis multifidis obtusis: laciniis linearibus, calycibus concavis. *Fl. lapp.* 448. t. 11. f. 2.

Lichen imbricatum viridans, scutellis badiiis. *Dill. musc.* 180. t. 24. f. 75.

Habitat in Europæ frigida rupibus. rupestris.

saxatilis. 19. LICHEN imbricatus, foliolis sinuatis scabris lacunosus, scutellis folio concoloribus. *Fl. succ.* 946. *Mat. med.* 494.

Lichen foliis planis acute laciniatis rubiginoso-albidis. *Fl. lapp.* 447. *Roy. lugdb.* 507.

Lichen tinctorius. *It. scan.* 409. *

Lichen opere phrygio ornatus. *Vaill. parif.* t. 21. f. 5. Liche-

Lichenoides vulgatissimum cinereo-glaucum lacunosum & cirrhosum. *Dill. musc.* 188. t. 24. f. 83.

Musco-fungus Lichenoides minor cinereus vulgatissimus. *Morif. hist.* 3. p. 634 f. 15. t. 7. f. 6.

Habitat in Europæ rupibus. rupestris.

20. *LICHEN imbricatus*, foliolis multifidis glabris ob-*omphalodes*.
tusis incanis, punctis vagis eminentibus. *Fl. succ.* 947.

Lichen nigricans omphalodes. *Vaill. parif.* 116. t. 20. f. 10.

Lichenoides saxatile tinctorium, foliis pilosis purpureis.

Dill. musc. 185. t. 24. f. 80.

Habitat in Europæ rupibus. rupestris & arboreus.

21. *LICHEN imbricatus*, foliolis lobatis nitidis lividis *olivaceus*.
Fl. succ. 948.

Lichen foliis planis multifidis obtusis: laciniis apice latioribus. *Fl. lapp.* 449. *Ray. lugdb.* 511.

Lichen crustæ modo arboribus adnascens, olivaceus.
Vaill. parif. t. 20. f. 8.

Lichenoides olivaceum, scutellis amplioribus verrucosis.

Dill. musc. 184. t. 24. f. 78.

Habitat in Europæ rupibus. arboreus & rupestris.

22. *LICHEN imbricatus*: foliolis linearibus dichotomis *sahlbomense*.
planiusculis acutis nigris, scutellis atris. *Fl. succ.* 1140.

Lichenoides tinctorium atrum, foliis minimis crispis.

Dill. musc. 188. t. 24. f. 81.

Habitat in Europæ rupibus nudis. rupestris.

23. *LICHEN imbricatus*, foliolis palmatis incurvis atris. *flygins*.
Fl. succ. 949.

Habitat in Suecia, imprimis in Insula Balthici Blå-kulla. rupestris.

24. *LICHEN* foliis imbricatis dentato-ciliatis, scutellis *cristatus*.
folio majoribus.

Lichenoides gelatinosum, foliis imbricatis & cristatis.

Dill. musc. 140. t. 19. f. 26.

Habitat in Europa australi.

25. *LICHEN imbricatus*, foliis crispis-fulvis, peltis con-*parietinus*.
coloribus fulvis.

Lichenoides vulgare sinuosum, foliis & scutellis luteis.

Dill. musc. 180. t. 24. f. 76.

Habitat in Europæ parietibus, rupibus, lignis. arboreus f. rupestris.

phyodes.

26. **LICHEN** imbricatus: laciniis obtusis subinflatis.
Lichen foliaceus subdepressus lævis, laciniis integerrimis truncatis subtus concavis lacunosis atris. *Fl. suec.* 951.

Lichen saxeus. *It. scan.* 409.

Lichen crustæ modo arboribus adnascens tenuiter divi-
sus. *Buxb. cent* 3. p. 272. t. 14. f. 1.

Lichen pulmonarius arboribus adnascens: desuper cinereus, subtus anthracinus: segmentis teretibus tubulosis.
Mich. gen. 91. t. 50. f. 1. 2.

Lichenoides ceratophyllum obtusius & minus ramosum.
Dill. musc. 154. t. 20. f. 49.

*Habitat in Europæ corticibus Betulæ, & Saxis ærapi-
bus.*

*Diagnoscitur a reliquis apicibus foliorum tumidis, &
si transversim secantur, inflatis.*

stellatis.

27. **LICHEN** imbricatus: foliolis oblongis laciniatis angustis cinereis, scutellis pullis. *Fl. suec.* 950.

Lichen foliis laciniatis cinereis: laciniis obtusiusculis sæpius apice trifidis. *Roy. lugdb.* 510.

Lichen pulmonarius vulgarissimus superne albo-cinereus, inferne nigricans, segmentis angustis, receptaculis nigricantibus. *Mich. gen.* 91. t. 43. f. 2.

Lichenoides cinereum, segmentis argutis stellatis, scutellis nigris. *Dill. musc.* 176. t. 24. f. 70.

Habitat in Europæ ramis arborum.

arbores.

Affinis plurimum L. rupestri.

* *Foliacei.*

ciliaris.

28. **LICHEN** foliaceus erectiusculus: laciniis linearibus ciliatis, scutellis pedunculatis crenatis. *Fl. suec.* 952.

Lichen foliis laciniatis cinereis: laciniis linearibus apice pilosis. *Roy. lugdb.* 509.

Lichen cinereus arboreus, marginibus pilosis, major.
Vaill. parisi. 115. t. 20. f. 4.

Lichenoides hispidum majus & rigidius, scutellis nigris.
Dill. musc. 150. t. 20. f. 45.

Musco-fungus arboreus cinereus scutellatus; marginibus pilosis. *Morif. bist.* 3. p. 634. f. 15. t. 7. f. 6.

Musculus arboreus pyxoides pilosus. *Loef. pruss.* 171. t. 50.

Habitat in Europæ arboribus.

*arbores.**cylindricus.*

29. **LICHEN** foliaceus ciliatus, peltis cylindricis rectis ipatis perforatis truncatis: duplici circulo. *Armen. acad.* 2. p. 264.

Liche.

Lichenoides foliorum laciniis crinitis. Dill. musc. 149.

t. 20. f. 42.

Habitat in Pensylvania, Lapponia. D. Montin.

30. **LICHEN** foliaceus adscendens laciniatus; margini-*islandicus.*

bus elevatis ciliatis. *Fl. suec. 959. Mat. med. 493.*

Lichen foliis oblongis laciniatis: marginibus conniven-

tibus ciliatis. Fl. lapp. 455. Roy. lugdb. 508. Hall.

helv. 75.

Lichen terrestris, foliis eryngii. Buxb. cent. 2. p. 11.

t. 6. f. 1. 2.

Lichenoides rigidum, eryngii folia referens. Dill. musc.

209. t. 28. f. 111.

Muscus islandicus purgans. Borrich. act. Haffn. 1674.

p. 126.

31. **Lichenoides** eryngii folia referens, tenuioribus & crispioribus foliis. *Dill. musc. 212. t. 28. f. 112..*

32. **Corallioides** tenuissimum nigricans, mundi muliebris in-tenuissimus, star textum. *Dill. musc. 113. t. 17. f. 32.*

Habitat in Europæ sylvis sterilissimis, Pinetis. terrestris.

Varietas 32. habitat frequens in sterilissimis collibus Suecia, Ramulis omnino teretibus, intus cavis, nec ciliatis adeoque diversissimus; at color luridus, basis rubra, varia specimina & gradus, unam, eandemque speciem esse evincunt.

33. **LICHEN** foliaceus adscendens laciniatus crispus glaberrimus.

ber lacunosus albus margine elevato. *Fl. suec. 958.*

Lichen niveus sinibus dædaleis laciniatus, ramis erectis,

calyce orbiculato. Fl. lapp. 446. t. 11. f. 1.

Lichenoides lacunosum candidum glabrum, endiviæ

crispæ facie. Dill. musc. 162. t. 21. f. 56.

Habitat in Lapponiæ, Upsaliæ, Grönlandiæ alpinis, apricis, siccis, glareosis. terrestris.

34. **LICHEN** foliaceus laciniatus obtusus glaber: supra *pulmonarius*

lacunosus; subtus tomentosus. *Fl. suec. 960. Mat.*

med. 490. St. w. gerb. 215.

Lichen foliis oblongis divisis, superficie reticulato-pun-

ctata. Fl. lapp. 444. Roy. lugdb. 509.

Lichenoides pulmonicum reticulatum vulgare, margini-

tus peltiferis. Dill. musc. 212. t. 29. f. 113.

Muscus pulmonarius. Baub. pin. 361.

Pulmonaria. Fuchs. hist. 631. Cam. epit. 783.

C c c c c

Ha-

*Habitat in Europæ sylvis æmbrosiis super arbores anti-
quas, præsertim in Fagis & Quercubus. arboreus.*

furfuraceus. 33. LICHEN foliaceus decumbens furfuraceus: laciniis
acutis: subtus lacunosis atris. *Fl. succ.* 953.

Lichen foliis planis laxis flaccidis inæqualiter laciniatis:
supra albis subtus atris. *Fl. lapp.* 450.

Lichen arboreus leucomelanus: ramulis altera parte
antracinis, altera candidissimis. *Buxb. cent.* 2. p. 12.
e. 7. f. 1.

Lichenoides cornutum amarum: desuper cinereum, in-
ferne nigrum. *Dill. musc.* 157. t. 21. f. 52.

Mucos amarus, absinthii folio. *Bauh. hist.* 3. p. 764.

Habitat in Europæ arboribus. arboreus.

ampullaceus. 34. LICHEN foliaceus planiusculus lobatus crenatus, pel-
tis globosis inflatis.

Lichen foliis laciniatis, marginibus convolutis in ves-
iculam abeuntibus. *Roy. lugdb.* 509.

Lichenoides tinctorum glabrum vesiculosum. *Dill.*
musc. 188. t. 24. f. 82.

Habitat in Lancastria Angliæ.

farinaceus. 35. LICHEN foliaceus erectus compressus ramosus fa-
rinaceus: lateribus verrucosis. *Fl. succ.* 957.

Lichen foliis multifidis, calycibus in margine foliorum
sessilibus. *Roy. lugdb.* 510.

Lichen cinereus angustior, scutis in marginibus segmen-
torum. *Vaill. parif.* 115. t. 20. f. 14. 15. 13.

Lichenoides segmentis angustioribus ad margines verru-
cosis & pulverulentis. *Dill. musc.* 172. t. 23. f. 63.

*Habitat in Europæ arboribus, præsertim Quercis
Fraxinetis.*

calicaris. 36. LICHEN foliaceus erectus linearis ramosus lacuno-
sus convexus mucronatus. *Fl. succ.* 956.

Lichen cinereus latifolius ramosus. *Vaill. parif.* 115. t.
20. f. 6.

Lichenoides coralliforme rostratum & canaliculatum.
Dill. musc. 170. t. 23. f. 62.

Musco-fungus arboreus, capitulis rostratis. *Morij. hist.*
3. p. 634. f. 15. t. 7. f. 5.

Habitat in Europæ arboribus, rapibus.

fraxineus. 37. LICHEN foliaceus erectus oblongus lanceolatus sub-
laciniatus lacunosus glaber, scutellis subpedunculatis.
Fl. succ. 955.

Li-

- Lichen foliis oblongis planis rugosis sinuatis, calycibus orbiculatis petiolatis. *Roy. lugdb.* 509.
- Lichen pulmonarius cinereus mollior in amplas laciniis divisus. *Tournef. inst.* 549. f. 325. f. A. B.
- Lichenoides longifolium rugosum rigidum. *Dill. musc.* 165. t. 22. f. 59.
- Musco Fungus quernus latifolius cinereus. *Morif. hist.* 3. p. 634. f. 15. t. 7. f. 14. 3. 4.
- Habitat in Europæ Arboribus, præsertim Faxinis. arboreus.*
38. LICHEN foliaceus rectiusculus lævis subtomentosus *fuciformis*. ramosus, laciniis lanceolatis.
- Lichenoides fuciforme tinctorium, corniculis longioribus & acutioribus. *Dill. musc.* 168. t. 22. f. 61.
- Fucus verrucosus tinctorius Roccella. *Bauh. pin.* 365.
- Bauh. hist.* 3. p. 797.
- Fucus marinus, Roccella tinctorum. *Bauh. pin.* 365.
- Habitat in India, Canariis.*
39. LICHEN foliaceus erectiusculus lacunofus: subtus *prunosus*. tomentosus albus. *Fl. suec.* 954.
- Lichen foliis mollibus candicantibus dichotomis, calycibus orbiculatis sessilibus. *Roy. lugdb.* 570.
- Lichen cinereus vulgarissimus cornua damæ referens. *Vaill. parif.* 115. t. 20. f. 11. 12.
- Lichenoides cornutum, bronchiale molle subtus incanum. *Dill. musc.* 160. t. 21. f. 55.
- Habitat in Europæ arboribus, præsertim in Prunospinosa. arboreus.*
40. LICHEN foliaceus laciniatus crispus fulvus, peltis *li-juniperinus*. vidis. *Fl. suec.* 967.
- Lichen fulvus sinubus dædaleis laciniatus. *Fl. lapp.* 450.
- Roy. lugdb.* 510.
- Habitat in Europæ juniperetis. arboreus.*
- Differt a L. parietino colore pallide flavo; foliis laxi, erectiusculis; peltis disco brunneis.*
41. LICHEN pallide viridis rugosus margine undulatus. *caperatus*. *Roy. lugdb.* 510. *Guett. stamp.* 1. p. 31.
- Lichenoides caperatum rosaceæ expentum. *Dill. musc.* 193. t. 25. f. 97.
- Musco-fungus lichenoides, crustæ modo adnascens major cinereus. *Morif. hist.* 3. p. 633. f. 15. t. 7. f. 1.
- Habitat in Europa & America ad saxa & arboreas.*
42. LI-

glauus.

42. LICHEN foliaceus depressus lobatus glaber: margine crispato farinaceo. *Fl. succ.* 966.
 Lichenoides endiviæ foliis crispis splendentibus subtus nigricantibus. *Dill. musc.* 192. t. 25. f. 96.
 Musco-Fungus lichenoides arborum crispus cinereus subtus nigricans. *Morif. bist.* 3. p. 633. f. 15 t. 7. f. 4.
Habitat in Europæ frigida, præsertim Sueciæ, truncis Betulinis. arboreus.

* *Coriacei.**aquaticus.*

43. LICHEN foliaceus repens lobatus obtusus, peltis hemisphæricis maximis.
 Lichenoides leutellis amplis. *Dill. musc.* 150. t. 20. f. 44?
Habitat in Suecia sub aqua, in paludibus.
Planta diversa a Lichene crinito. Dill. musc. t. 84. f. 8.
 Folia lobata, pallide viridia, margine integerrima. Peltis subjeffiles, primum plana, deum dilatata, folio majores, incarnato fusca, margine reflexo, supra discum folii posita.

resupinatus.

44. LICHEN foliaceus repens lobatus, peltis marginalibus posticis.
 Lichen pulmonarius major s. minor ex obscuro cinereus: inferne ex albo rufescens, receptaculis florum rubris ad latera oblongis. *Misch. gen.* 86. t. 44. f. 1. 2.
 Lichenoides fuscum, peltis posticis ferrugineis. *Dill. musc.* 206. t. 28. f. 105.
Habitat in Europæ sylvis. terrestris.

venosus.

45. LICHEN foliaceus repens ovatus planus subtus venosus villosus, peltis marginalibus horizontalibus. *Fl. succ.* 964.
 Lichenoides parvum virescens, peltis marginalibus planis. *Dill. musc.* 208. t. 28. f. 109.
Habitat in Europa ad margines scrobicularum in sylvis. terrestris.

apthosus.

46. LICHEN foliaceus repens lobatus obtusus planus, verrucis sparsis, pelta marginali adscendente. *Fl. succ.* 963. *Mat. med.* 422. *Amœn. acad.* 2. p. 69.
 Lichenoides digitatum læte virens, verrucis nigris notatum. *Dill. musc.* 207. t. 28. f. 106.
Habitat in Europæ Sylvis acerosis sterilibus sub juniperis. terrestris.

arcticus.

47. LICHEN foliaceus repens lobatus obtusus planus lævis subtus aversus villosus. *Fl. succ.* 962.

Li-

Lichen foliis planis subrotundis lobatis obtusis, calyce plano ovali lacinula propria adnato, niveus *Fl. lapp.* 442.

Habitat in Suecia boreali, sub juniperis. terrestris.
Folia lata, integra, albida, subtus nivea. Pelta subtus versus marginem, magna, livido-incarnata, magnitudine pollicis, adeoque inter omnes Lichenes latissima. Supra læve est folium.

48. LICHEN foliaceus repens lobatus obtusus planus *cinereus*, subtus venosus villosus, pelta marginali adscendente. *Fl. suec.* 961. *Mat. med.* 491.

Lichen foliis planis subrotundis lobatis obtusis calyce plano lacinia propria adnato. *Fl. lapp.* 441. *Roy. lagdb.* 508.

Lichen pulmonarius saxatilis digitatus. *Vaill. parif.* 116. t. 21. f. 16.

Lichenoides digitatum cinereum, lactucæ foliis sinuosis. *Dill. musc.* 200. t. 27. f. 102.

Musco-Fungus terrestris latifolius cinereus, hepaticæ facie. *Morif. bist.* 3. p. 632. f. 15. t. 7. f. 1.

Habitat in Europæ sylvis, juxta lapides, in terra. terrestris.

49. LICHEN foliaceus repens subrotundus planus sub- *croceus*, tus venosus villosus croceus, peltis sparsis adnatis. *Fl. suec.* 965.

Lichen foliis subrotundis planis levissime incis, calycibus orbiculatis disco folii adnatis. *Fl. lapp.* 443. t. 11. f. 3. *Roy. lagdb.* 509.

Lichen alpinus viridis subtus aurantius, scutis magnis nigris planissimis. *Hall. helv.* 74.

Lichenoides subtus croceum, peltis appressis. *Dill. musc.* 221. t. 30. f. 120.

Habitat in Lapponia, Helvetia, Grœnlandia. terrestris.

* *Umbilicati, squalentes quasi fuligine.*

50. LICHEN foliaceus gibbus punctatus subtus fulvus. *miniatus*.

Lichen pulmonarius alpinus cinereus punctatus. *Hall. helv.* 75. t. 2. f. 2.

Lichenoides coriaceum nebulosum cinereum punctatum subtus fulvum. *Dill. musc.* 223. t. 30. f. 127.

Habitat in Angliæ, Helvetiæ, Italiæ, rupibus alpinis.

- vellens.* 51. LICHEN foliaceus umbilicatus subrus hirsutissimus.
Fl. succ. 968.
 Lichen folio subrotundo peltato: margine fere integro, subtus maxime hirsutus. *Fl. lapp.* 454. *Roy. lugdb.* 511.
 Lichenoides coriaceum, latissimo folio umbilicato & verrucoso. *Dill. musc.* 545. *t.* 82. *f.* 5.
Habitat in alpinis Lapponiæ, Sueciæ, Angliæ. *rupestris.*
- pustulatus.* 52. LICHEN foliaceus umbilicatus: subtus lacunofus.
Fl. succ. 969,
 Lichen folio orbiculato margine integriusculo, superficie subtus puculata. *Fl. lapp.* 453. *Roy. lugdb.* 511.
 Lichenoides crustæ modo saxi adnascens verrucosus cinereus & veluti deustus. *Vaill. parif.* 116. *t.* 20 *f.* 9.
 Lichenoides pustulatum cinereum & veluti amustum *Dill. musc.* 227. *t.* 30. *f.* 131.
Habitat in Europæ rupibus apricis. *rupestris.*
- probo/cidens* 53. LICHEN foliaceus umbilicatus, peltis turbinatis truncatis perforatis.
Habitat in Suecia.
 Folium magnitudine pollicis, subrotundum, orbiculatum, margine inæqualiter & obtuse lobatum, centro radicatum, subtus læve, hinc inde & rarius fibræ radicantem exserens; Pagina superior planiuscula vel nonnihil contorta, cinerea, punctis fuscis elevatis exasperata. Peltæ per discum sparsæ, utræ, minimæ, turbinatæ, truncatæ, poro pertusæ usque ad folium, margine lævi plano.
- densus.* 54. LICHEN foliaceus umbilicatus undique lævis *Fl. succ.* 970. *It. w:goth.* 217.
 Lichen folio orbiculato peltato margine integriusculo undique glaber. *Fl. lapp.* 452. *Roy. lugdb.* 511.
 Lichen pulmonarius saxatilis e cinereo fulcis minimus. *Vaill. parif.* 116. *t.* 21. *f.* 14.
 Lichenoides coriaceum cinereum, peltis atris compressis. *Vaill. parif.* 219. *t.* 20. *f.* 117.
Habitat in Sueciæ, Galiiæ rupibus s. cantibus elatis apricis. *rupestris.*
- polyphyllus.* 55. LICHEN foliaceus polyphyllus utrinque lævis cre-natus.
 Lichenoides tenue pullum, foliis utrinque glabris. *Dill. musc.* 225. *t.* 30. *f.* 129.
Habitat in Europæ rupibus elatis apricis. *rupestris.*
56. LI-

56. LICHEN foliaceus polyphyllus utrinque lævis poly-*polyrhizos*.
rhizos.

Lichenoides pullum superne & glabrum, inferne nigrum
& cirrhosum. *Dill. musc.* 226. t. 30. f. 130.

*Habitas in Arvoniz, Sveciz rupibus apricis elatis. ru-
pestris.*

* *Scyphiferi.*

57. LICHEN scyphifer simplex integerrimus, stipite cy-*cocciferus*.
lindrico, tuberculis coccineis. *Fl. suec.* 972. *Mat.*
med. 496.

Lichen caule simplici, calyce turbinato margine carno-
so colorato prolifero. *Fl. lapp.* 429. *Roy. lugdb.* 512.

Lichen pyxidatus, oris coccineis & tumentibus. *Vaill. pa-
ris.* 115. t. 21. f. 4. *Mich. gen.* 82. t. 41. f. 3.

Lichenoides scyphiforme, tuberculis coccineis. *Dill.*
musc. 82. t. 14. f. 7.

Habitas in Europæ sylvis sterilibus, ericetis, rupibus.

58. LICHEN scyphifer simplex folio brevior, tubercu-*cornucopiai-*
lis coccineis. *Fl. suec.* 974. *des.*

Lichen caule simplici, calyce turbinato: centro mul-
tipliciter prolifero. *Fl. lapp.* 432. *Gron. virg.* 204.
128.

Coralloides scyphiforme: marginibus radiatis & foliatis.
Dill. musc. 85. t. 14. f. 9.

Muscus pyxoides. *Barr. rar.* 1284. t. 1268. f. 2.

Habitas in Lapponiz, Sueciz, Angliz sylvis glareosis.

59. LICHEN scyphifer simplex crenulatus, tuberculis *pyxidatus*.
fuscis. *Fl. suec.* 971.

Lichen caule simplici, calyce turbinato: margine tenui.
Fl. lapp. 428. *Gron. virg.* 203. *Roy. lugdb.* 512.

Lichen pyxidatus major. *Tournef. inst.* 549. t. 325. f.
D. Vaill. paris. 115. t. 21. f. 8. *Mich. gen.* 82. t.
41. f. 1.

Coralloides scyphiforme, tuberculis fuscis. *Dill. musc.*
79. t. 14. f. 6.

Fungus terrestris pyxidatus. *Magn. hort.* 83. t. 83.

β. Lichen caule simplici, calyce turbinato: centro simpli-
citer prolifero. *Fl. lapp.* 431. *Roy. lugdb.* 512.

Lichen pyxidatus prolifer. *T. Vaill. paris.* 115. t. 21.
f. 5. 9. 11.

Lichen pyxidatus major alpinus, receptaculis copiosiori-
bus & rufescentibus. *Mich. gen.* 82. t. 41. f. 2. 7. 9.

Habitas in Europæ sylvis.

- fimbriatus.* 60. LICHEN scyphifer simplex denticulatus, stipite cylindrico. *Fl. succ.* 973.
Lichen caule simplici, calyce turbinato: margine acuto prolifero. *Fl. lapp.* 430.
Lichen pyxidatus minor. *Vaill. parif.* 115. t. 21. f. 6.
Mich. gen. 83. t. 41. f. 45.
Coralloides scyphiforme gracile: marginibus serratis. *Dill. musc.* 84. t. 14. f. 8.
Habitat in Europæ sylvis sterilibus.
- gracilis.* 61. LICHEN scyphifer ramosus denticulatus filiformis. *Fl. succ.* 975.
Lichen pyxidatus & corniculatus ramosus alpinus e fusco cinereus, pyxidulis crenatis. *Mich. gen.* 81. t. 41. f. 5.
Coralloides scyphiforme ferratum elatius, caulibus gracilibus glabris. *Dill. musc.* 88. t. 14. f. 13.
Musco-fungus pyxidatus gracilior ramosus, calycibus serratis. *Morif. hist.* 3. p. 362. f. 15. t. 7. f. 6.
Habitat in Europæ ericetis, montosis, sylvaticis.
- digitatus.* 62. LICHEN scyphifer ramosissimus: ramis cylindricis, calycibus integris nodosis. *Fl. succ.* 978.
Lichen caule inordinate ramoso, ramis in calyces turbinatos margine carnosos desinentibus. *Fl. lapp.* 436.
Coralloides ramulosum, tuberculis coccineis. *Dill. musc.* 96. t. 15. f. 19.
Habitat in Europæ sylvis sterilibus.
- cornutus.* 63. LICHEN scyphifer simpliciusculus subventricosus, calycibus integris. *Fl. succ.* 976.
Lichen caule simplici, subulato rarius bifido. *Fl. lapp.* 434. *Roy. lugdb.* 512.
Coralloides vix ramosum, scyphis obscuris. *Dill. musc.* 90. t. 15. f. 14.
Muscus fistulosus corniculatus. *Barr. rar.* 1286. t. 1277. f. 1. *Bocc. mus.* 2. p. 149. t. 107.
Habitat in Europæ ericetis.
- deformis.* 64. LICHEN scyphifer simpliciusculus subventricosus, calycibus dentatis. *Fl. succ.* 977.
Lichen caule simplici, apice acuto s. calyce turbinato terminato. *Fl. lapp.* 433. t. 11. f. 3.
Lichen pyxoides teres, acetabulis minoribus repandis. *Mich. gen.* 80. t. 41. f. 1.

Coral-

Coralloides crassius subincanum, calycibus dentatis.

Dill. musc. 95. t. 15. f. 18. mala.

Habitat in Europæ ericetis.

* *Fruticulosi.*

5. LICHEN fruticulosus perforatus ramosissimus, ramulis nutantibus. *Fl. suec. 980.*

Lichen caule erecto tereti ramosissimo, alis perforatis. alpestris,

Fl. lapp. 437. Gron. virg. 203. 127. Roy. lugdb. 511.

Coralloides montanum, fruticuli specie ubique candicans. Dill. musc. 107. t. 16. f. 29.

Musco fungus coralloides montanus ramosissimus cinereus vulgaris. *Morif. hist. 3. p. 633. f. 15. t. 7. f. 9.*

Muscus coralloides s. cornutus montanus. *Bauh. pin. 361.*

Muscus corallinus s. Corallina montana. *Bauh. hist. 1198.*

Lichen caule ramoso, alis perforatis, ramis capitatis. sylvaticus. *Roy. lugdb. 511.*

Coralloides fruticuli specie candicans, corniculis rufescentibus. Dill. musc. 110. t. 16. f. 30.

Muscus terrestris coralloides erectus, corniculis rufescentibus. *Bauh. pin. 361. prodr. 152.*

Habitat in Alpibus, Europæ frigida sylvis sterilissimis, Alpestris differt a sylvatico, ut flos plenus a simplici.

6. LICHEN fruticulosus perforatus, ramulis brevissimis acutis. *Fl. suec. 979.*

Lichen caule ramoso, alis perforatis, ramis brevissimis acutis. *Fl. lapp. 438. Roy. lugdb. 511.*

Coralloides perforatum minus molle & tenue. Dill. musc. 99. t. 16. f. 22.

Habitat in Europæ ericetis

7. LICHEN fruticulosus subdichotomus, ramis simplicibus subulatis. *Fl. suec. 981.*

Lichen caule erecto dichotomo, ramis subulatis. *Fl. lapp. 435. Roy. lugdb. 512.*

Coralloides corniculis longioribus & rarioribus. Dill. musc. 102. t. 16. f. 26.

Muscus corniculatus. *Tabern. ic. 809. Bauh. hist. 3. p. 767.*

Habitat in Europæ sylvis ericetis.

8. LICHEN fruticulosus solidus tectus foliolis crustaceis. *Fl. suec. 982.*

D d d d

Li-

Lichen caule ramoso solido foliolis crustaceis testis. *A. lapp.* 439. *Roy. lugdb.* 512.

Lichen alpinus glaucus ramosus botryoides. *Schenck alp.* 137. t. 19. f. 4.

Coralloides crispum & botryoides alpinum. *Dill. musc.* 114. t. 17. f. 33.

Corallina alpina valde crispa. *Pet. musc.* t. 65. f. 7.

Musco-fungus coralloides terrestris dense ramificatus cinereus & veluti incrustatus norvegicus. *Moriz. bist.* 3. p. 633. f. 15. t. 7. f. 12.

Muscus cupressiformis ramosus. *Las. pruss.* 168. t. 4.

Habitat in Helvetiæ, Italiæ, Cambriæ, Lappocia, Scaniæ, Grœnlandiæ, Penſylvaniæ alpeſtribus.

fragilis. 69. LICHEN fruticulosus solidus, ramulis teretibus obtusis. *Fl. suec.* 983.

Lichen erectus ramosissimus, ramulis teretibus nodis filiformibus obtusis. *Fl. lapp.* 440. t. 11. f. 4.

Coralloides alpinum, corallinæ minoris facie. *Dill. musc.* 116. t. 17. f. 34.

Habitat in Europæ alpibus alpinisque, per Sueciam in rupibus.

Roccella. 70. LICHEN fruticulosus solidus aphyllus subramosus tuberculis alternis.

Coralloides corniculatum fasciculare tinctorium fuscescente teretis facie. *Dill. musc.* 120.

Muscus ceranoides palmentis, comis digitatis orbiculatis. *Pet. gaz.* t. 7. f. 12.

Muscus ceranoides albus fungosus apicibus nigris. *Planch. alm.* 255. t. 205. f. 6.

Fucus marinus Roccella tinctorum. *Bauh. pin.* 365.

Fucus capillaceus Roccella. *Bauh. pin.* 365.
Habitat in insulis Archipelagi, Canariis, ad rupes marinas.

* *Filamentosi.*

plicatus. 71. LICHEN filamentosus pendulus, ramis implexis scutellis radiatis. *Fl. suec.* 984. *Mat. med.* 495.

Lichen ramis filiformibus ramosis pendulis confertis. *Fl. lapp.* 457. *Gron. virg.* 203. 127.

Usnea vulgaris, loris longis implexis. *Dill. musc.* 56. t. 11. f. 1.

Muscus arboreus, Usnea officinarum. *Bauh. pin.* 365.

Habitat in Europæ & America borealis sylvis densis umbrosisque fagis.

72. L.

LICHEN filamentosus pendulus subarticulatus, ra-*barbatus*.
mis patentibus. *Fl. suec.* 985.

Lichen caule tereti coriaceo, ramulis tenuissimis cine-
reis. *Roy. lugdb.* 513.

Usnea barbata, loris tenuibus fibrosis. *Dill. musc.* 63.
t. 12. f. 6.

Muscus capillaceus longissimus *Baub. pin.* 361.

Habitat in Europæ & Americæ septentrionalis sylvis fa-
getis.

LICHEN filamentosus pendulus: axillis compressis. *puber.*
Fl. suec. 986.

Lichen ramis filiformibus ramosis pendulis, alis com-
pressis. *Fl. lapp.* 456.

Usnea jubata nigricans. *Dill. musc.* 64. t. 12. f. 7.

Muscus arborum niger & tenuior. *Dod. pempt.* 471.

Habitat in Europæ sylvis & rupibus.

LICHEN filamentosus ramosissimus decumbens im-*lanatus*.
plicatus opacus. *Fl. suec.* 987.

Usnea lanæ nigræ instar saxi adhærens. *Dill. musc.* 66.

t. 13. f. 8.

Habitat in Europæ frigida rupibus.

LICHEN filamentosus ramosissimus decumbens im-*pubescens*.
plexus nitidus

Usnea cæspitosa exilis capillacea atra. *Dill. musc.* 66.

t. 13. f. 9.

Habitat in Europa septentrionali, Lapponia, Suecia.

LICHEN filamentosus subramosus decumbens im-*chalybeiformis*.
plicato-flexuosus. *Fl. suec.* 988.

Usnea rigida horsum vorsum *extensa*. *Dill. musc.* 66.

t. 13. f. 10.

Muscus caule rigido instar fili chalybei. *Raj. anglie.* 3.

p. 65.

Habitat in Europa supra rupes & sepimenta.

LICHEN filamentosus ramosissimus erectus, tuber-*hirtus*.
culis farinaceis sparsis. *Fl. lapp.* 989.

Usnea vulgarissima tenuior & brevior sine orbiculis.

Dill. musc. 67. t. 13. f. 12.

Muscus ramosus. *Tabern. ic.* 807.

Habitat in Europæ arboribus, sepimentis.

LICHEN filamentosus ramosissimus erectus fastigia-*vulpinus*.
tus inæquali-angulosus. *Fl. suec.* 990.

D d d d 2

Uf-

Usnea capillacea citrina, fruticuli spec. *Dill. m.*
73. t. 13. f. 16.

Habitat in Europæ tellis ligneis, maris.

articulatus. 79. Lichen filamentosus articulatus, ramulis tenuissimis punctatis. *Roy. lugdb.* 513. n. 29.

Usnea capillacea & nodosa. *Dill. musc.* 60. t. 11.

Muscus arboreus nodosus. *Baub. pin.* 361.

Musco-fungus arborum nodosus f. geniculatus. *Morif. hist.* 3. p. 634. f. 15. t. 7. f. 11.

Dendrobryon geniculatum f. nodosum. *Col. cephr.* t. 83. f. 2.

Habitat in Europæ australis sylvis.

floridus. 80. LICHEN filamentosus ramosus erectus, scutellis radiatis. *Fl. suec.* 991.

Lichen caule ramoso solido, foliis setaceis, receptaculis maximis orbiculatis peltatis ciliato-radiatis. *Hort. cl.* 477. *Roy. lugdb.* 513.

Usnea vulgarissima tenuior & brevior cum orbiculatis. *Dill. musc.* 69. t. 13. f. 13.

Muscofungus arborum capillaceus, scutellis amplius ambitum pilis radiatis. *Morif. hist.* 3. p. 635. f. 15. f. 14.

Muscus arboreus cum orbiculis. *Baub. pin.* 361.

Muscus ramosus floridus. *Tabern. ic.* 808.

Habitat in Europæ sagetis.

CHARA.

tormentosa. 1. CHARA aculeis caulinis ovatis. *Fl. suec.* 993.

Equisetum fragile majus subcinereum aquis immersum.

Morif. hist. 3. p. 621. f. 15. t. 4. f. 9.

Equisetum f. *Hippuris lacustris*, foliis mansu arendis.

Pluk. alm. 135. t. 29. f. 4.

Habitat in Europæ stagnis, mari, lacubus.

vulgaris. 2. CHARA caulibus lævibus, frondibus interne denticulatis.

Chara vulgaris. *Fl. lapp.* 469. *Fl. suec.* 992.

Chara caulibus lævibus. *Roy. lugdb.* 533.

Chara vulgaris foetida. *Vaill. act.* 1719. p. 23. t. 3. f. 1.

Equisetum foetidum sub aqua repens. *Baub. pin.* 16. p. 25. *theatr.* 251. *Baub. hist.* 3. p. 731.

Habitat in Europæ aquis pigris.

hispida. 3. CHARA aculeis caulinis capillaribus confertis. *suec.* 994. *

Chara caulibus aculeatis. Hort. cliff. 477. Roy. lugdb.
214.

Chara major, caulibus spinosis. Vaill. act. 1719. p. 18.
t. 3. f. 3.

Equisetum f. *Hippuris muscosus* sub aqua repens. *Pluk.*
alm. 135 t. 193. f. 6.

Habitat in Europæ maritimis.

CHARA caulium articulis inermibus diaphanis super-*flexilis*,
ne latioribus. *It. gotl. 215. Fl. suec. 995.*

Chara transluens minor flexilis. Raj. angl. 3. p. 133.

Habitat in Europæ maritimis.

TREMELLA. 2

TREMELLA sessilis membranacea auriformis fulva. *juniperina*,
Fl. suec. 1017.

Syffus gelatinosa fugax, junipero innascens. Fl. lapp.
531.

Habitat in Juniperetis primo vere.

TREMELLA plicata undulata. *Fl. suec. 1018. Nestor.*

Tremella terrestris sinuosa pinguis & fugax. Dill. musc.
52. t. 10. f. 14.

Syffus gelatinosa fugax terrestris. Fl. lapp. 530.

Linkia terrestris gelatinosa membranacea vulgatissima.

Mich. gen. 126. t. 67. f. 1.

Noëtic paracelsi. Act. parisi. 1708. p. 228.

Habitat in Pratis post pluvias.

TREMELLA sessilis membranacea auriformis cinea-*Auricula*,
rea. Fl. suec. 1119.

Agaricum auriculæ forma. Mich. gen. 124. t. 66. f. 1.

ungorum perniciosorum genus 1. Clas. hist. 2. p. 276.

Habitat ad Arbores putridas.

TREMELLA frondibus erectis planis: margine cri-*Lichenoides*.
spo lacinulato. *Fl. suec. 1020.*

Lichenoides pellucidum, endiviæ foliis tenuibus crispis.

Dill. musc. 143. t. 19. f. 31.

Lichen terrestris membranaceus mollior fuscus. Mich.
gen. 26. t. 38.

Musco fungus terrestris minor fuscus, foliis e latitudi-
ne crenatis musco innascens. Moris. hist. 3. p. 632. f.

15. t. 7. f. 4.

Habitat in Muscis, locis umbrosis ad montes.

- verrucosa.* 5. TREMELAA tuberculosa solida rugosa. *Fl. suec.* 102.
It. goth. 181.
 Tremella subrotunda sinuosa difformis gelatinosa. *l. w-goth.* 193.
 Tremella fluviatilis gelatinosa & uterculosa. *Dill. musc.* 54. t. 10. f. 6.
 Linkia palustris gelatinosa saxis adnascens ex obscura fulva & concava vesicam referens. *Mich. gen.* 126. t. 67. f. 2.
Habitat supra Lapides in rivulis.
- hemisphaerica.* 6. TREMELLA hemisphaerica sparsa. *It. w-goth.* 193.
Habitat in Rupibus & Confervis submarinis.
- purpurea.* 7. TREMELLA subglobosa sessilis solitaria glabra. *Fl. suec.* 1022.
 Lichenoides tuberculosum amœne purpureum. *Dill. musc.* 127. t. 18. f. 6.
Habitat in Arborum ramis moribundis & emortuis.



FUCUS.

* *Dichotomi frondescentes.*

- ferratus.* 1. FUCUS fronde plana dichotoma ferrata ad apicem tuberculata. *Hort. cliff.* 478. *Fl. suec.* 1001.
 Fucus folio dichotomo plano serrato laciniato. *Guest. stamp.* 2. p. 402. *Fl. lapp.* 462.
 Fucus f. Alga latifolia major dentata. *Morif. bist.* 3. p. 648. f. 15. t. 9. f. 1.
Habitat in Oceano.
- vesiculosus.* 2. FUCUS fronde dichotoma integra, caule medium folium transcurrente, vesiculis verrucosis terminalibus. *Fl. lapp.* 466. *Fl. suec.* 1002. *It. w-goth.* 168. *l. w-goth.* 83. *Roy. ingdb.* 514. *Guest. stamp.* 2. p. 401.
 Fucus marinus f. Quercus maritima vesiculas habens. *Banb. pin.* 363.
 Quercus marina. *Lob. ic.* 2. p. 255.
3. Fucus maritimus f. Quercus maritima foliorum eorum tumidis. *Banb. pin.* 365.
Habitat in Mari atlantico.
- ceranoides.* 3. FUCUS fronde dichotoma plana integra: apicibus fidis vesiculosis. *Fl. lapp.* 465. *Fl. suec.* 1005.
 Fucus humilis dichotomus ceranoides, latioribus foliis ut plurimum verrucosis. *Morif. bist.* 3. p. 646. f. 1. t. 8. f. 13.
Habitat in Oceano.

4. FU

4. FUCUS fronde dichotoma integra, caule fo-*spiralis*.
 lium percurrente: inferne nudo, vesiculis verruc o-
 sis terminalibus. *Fl. lapp.* 467. *Fl. succ.* 1003. *Roy.*
Ingdb. 514.
Fucus spiralis maritimus major. *Raj. angl.* 3. p. 41.
Habitat in Oceano.
5. FUCUS fronde bifida: laciniis ovato-lanceolatis in-*insens.*
 flatis apice divisilis. *Fl. lapp.* 468. *Fl. succ.* 1004.
Habitat in Oceano Atlantico.
6. FUCUS fronde membranacea lineari dichotoma di-*divaricatus.*
 varicata integra: ramificationibus vesiculosus.
Fucus bullatus frutescens, caule nudo, foliis rectis
 compressis bitidis. *Morif. hist.* 3. p. 647. f. 15. t. 8,
 f. 5.
Habitat in Anglia, Lusitania.
- * *Dichotomi caulescentes.*
7. FUCUS fronde filiformi teretiufcula dichotoma recta: *elongatus.*
 geniculis tumidiufculis.
Fucus caule lineari aphylo dichotomo. *Roy. Ingdb.* 514.
Fucus spongiosus teres ramosus viridis erectus. *Morif.*
hist. 3. p. 647. f. 15. t. 8. f. 7.
Habitat inter Angliam & Hispaniam.
8. FUCUS dichotomus linearis: axillis divaricatis. *excisus.*
Fucus pumilus dichotomus: segmentis ex una parte gib-
 bosis, ex altera excavatis. *Raj. angl.* 3. p. 43.
Fucus dichotomus membranaceus ex viridi flavescent
ceranoides, angulos rotundiusculos efformans. *Morif.*
hist. 3. p. 646. f. 15. t. 8. f. 11.
Habitat in Oceano Europeo.
9. FUCUS caule compresso dichotomo medio ramorum *nodosus.*
 in vesiculam dilatato. *Roy. Ingdb.* 514. *Fl. succ.* 1006.
Fucus caule tereti compresso dichotomo, vesicula me-
dio ramorum innata, vesiculis axillaribus laxis. *Hort.*
cliff. 479.
Fucus caule tereti compresso dichotomo, foliis oppositis
minimis, vesicula in medio singuli rami. *Fl. lapp.* 464.
Gnett. stamp. 2. p. 401.
Fucus maritimus, vesiculis majoribus singularibus per
intervalla dispositis. *Morif. hist.* 3. p. 647. f. 15. t.
 8. f. 2.
Fucus maritimus nodosus. *Dod. pempt.* 781.
Habitat in Mari Atlantico.

D d d d 4

10. FU-

- concatenatus* 10. FUCUS caule tereti ramosissimo, fructificationibus oblongis concatenatis. *Roy. lugdb.* 514.
Habitat in Oceano.

* *Ramosi foliis distinctis.*

- natus.* 11. FUCUS caule tereti ramosissimo, foliis lanceolato-serratis, fructificationibus globosis pedunculatis subaristatis *Fl. zeyl.* 369. *Roy. lugdb.* 513.
Fucus folliculaceus, serrato folio. *Bamb. pin.* 365.
Lenticula marina, serratis foliis. *Dalech. hist.* 1397.
Sargazo. *Pisf. bras.* 2. p. 266.
Habitat in Pelago libere natus, nec radicans.
Vegitabile ni fallor, inter omnia in orbe, numerosissimum.

- acinaris.* 12. FUCUS caule tereti ramoso, foliis linearibus integerrimis, fructificationibus globosis pedunculatis.
Fucus folliculaceus, linearis folio. *Bamb. pin.* 365.
Lenticula marina serapionis. *Lob. ic.* 2. p. 256.
Acinaria imperati. *Donat. marin.* 35. t. 4. f. 1.
Habitat in Italia & Oceano australiori.

- indigerus.* 13. FUCUS caule tereti corymbofo, foliis lanceolatis denticulatis alternis, fructificationibus cymosis.
Habitat ad insulam Adscensionis. Osbeck.
Caulis filiformis, fuscus, spithameus, foliis alternis, quæ deciderunt. Rami ex singulis alis simplices, caule longiores, filiformes, laxi. Folia alterna, lanceolata, densata, pellucida. Cyma pedunculata ex singulis alis superioribus, continens plurimas fructificationes subulatas, foliis breviores, fuscas, nodosas, sessiles.

* *Ramosi fronde unita.*

- turbinatus.* 14. FUCUS ramosus teres, fructificationibus turbinatis membrana cinctis. *Hort. cliff.* 478.
Fucus marinus vesiculas habens membranis extantibus alatas. *Sloan. jam.* 4. *hist.* 2. p. 58. t. 20. f. 6.
Habitat in Americæ rupibus marinis.

- glicosus.* 15. FUCUS caule tereti ramosissimo, pedunculis alternis vesiculis oblongis acuminatis. *Fl. lapp.* 463. *Fl. suec.* 1007. *Gaett stamp.* 2. p. 403.
Fucus maritimus alter, tuberculis paucissimis. *Bamb. pin.* 368.
Fucus marinus quartus. *Dod. pempt.* 480.
Habitat in Oceano.

16. FU-

16. FUCUS vesiculis oblongis pedunculatis: terminatis *barbatus*.
foliolis linearibus.
Fucus folliculaceus, foliis abrotani. *Bauh. pin.* 365?
Fucus maritimus, foliis tumidis barbatis. *Bauh. pin.* 365.
Quercus maritima barbata. *Bauh. prodr.* 154.
Habitat in Oceano.
17. FUCUS vesiculis ovatis pedunculatis, foliis multi-*senariolacens*
fidis linearibus.
Fucus folliculaceus, fœniculi folio. *Bauh. pin.* 365.
Habitat in Oceano.
18. FUCUS fronde bipinnata subfiliformi: apicibus ve-*abrotanifo-*
ficulosis dilatatis. *Læf. epist.* 1. 4. *lin.*
Habitat in mari Anglico.
- * *Fruetificationibus non vesicariis.*
19. FUCUS caule tereti ramosissimo, ramis sparsis spi- *Muscoides*.
nis mollibus alternis.
Ficus folliculaceus fuscus f. nigricans ex insula cypro.
Pet. musf. 219? *Raj. suppl.* 13.
Habitat ad Insulam Adscensionis. Osbeck.
Refert ita Muscum, ut exsiccatum nequeat a muscis
distingui. Madidus constat substantia coriacea succi,
diaphana: caulibus ramisque confertis, filiformibus,
undique adpersis rudimentis ramulorum; spinis mol-
libus, substantia & colore totius, brevissimis, inerti-
bus.
20. FUCUS cartilagineus, caule teretiufculo compres-*cartilagineus*
so: frondibus supradecompositis pinnatis: laciniis li-
nearibus coloratis. *Roy. lugdb.* 515. *Guett. stamp.* 2.
p. 404.
Fucoides rubens varie dissectum. *Raj. angl.* 8. *p.* 37.
Muscus marinus tenuissime dissectus ruber. *Bauh. pin.*
363.
Habitat in Oceano australiore.
21. FUCUS caule tereti brevissimo, folio maximo en-*saccharinu-*
siformi subsimplici. *Fl. lapp.* 460. *Fl. succ.* 1010. *Is.*
w.goth. 169.
Fucus longissimo latissimo crassoque folio. *Bauh. prodr.*
154. *Raj. hist.* 74.
Fucus alatus f. phasnagoides. *Bauh. pin.* 364.
Fucus setaceus latissimus edulis dulcis. *Sibb. scoth.* 26,
Habitat in Mari Atlantico.

- palmatus.* 22. **FUCUS** frondibus planis palmatis. *Roy. lugdb.* 515.
Fucus membranaceus ceranoides. Bauh. prodr. 155.
Fucus foliaceus humilis palmam humanam referens.
Morif. hist. 3. p. 646. f. 15. t. 8. f. 1.
Habitat in Oceano.
- rubens.* 23. **FUCUS** caule tereti ramoso, frondibus oblongis undulatis sinuatis. *Roy. lugdb.* 514.
Alga minor suave rubens varie divisa. Mart. cent. 32. t. 32.
Habitat in Oceano.
- pavonicus.* 24. **FUCUS** fronde sessili reniformi decussatim striata.
Fucus maritimus gallopavonis pennas referens. Bauh. prodr. 155. *Morif. hist.* 3. p. 645. f. 15. t. 8. f. 7.
Alga marina gallopavonis plumas referens. Bauh. pin. 364.
Fungus auricularis. Bauh. pin. 368.
Habitat in Mari Europæ australis.
- fastigiatus.* 25. **FUCUS** dichotomus ramosissimus teres uniformis fastigiatus. *Is. æl.* 120. *Fl. suec.* 1008.
Fucus caule lineari dichotomo. Roy. lugdb. 514.
Fucus palmaris tenuis in orbem expansus, in segmenta bifida breviora teretia divisus. Morif. hist. 3. p. 649. f. 15. t. 9. f. 9.
Fucus marinus polyschides. Læs. pruss. 77. t. 15.
Habitat in Oceano Balthico.
- filum.* 26. **FUCUS** filiformis simplex subfragilis opacus.
Fucus filiformis simplex. Fl. lapp. 459. *Fl. suec.* 1009. *Roy. lugdb.* 513. *Is. w-goth.* 168. *Gnest. stamp.* 2. p. 395.
Fucus f. Filum maritimum germanicum. Bocc. mus. 1. p. 271. t. 9. f. 9.
Alga nigro capillaceo folio. Bauh. pin. 364.
Filum maritimum germanicum. Bauh. prodr. 155.
Linum marinum. Bart. act. 1671. p. 118. f. 8.
Habitat in Oceano Atlantico.
- tendo.* 27. **FUCUS** filiformis simplex tenuissimus subdiaphanus.
Fucus indicus teres, setam piscatoriam referens, longissimus. Pluk. alm. 160. t. 184. f. 3.
Gramen sparteum setas equinas referens, e quibus javanenses tapetia texunt. Bauh. prodr. 11.
Habitat in China.

ULVA.

1. ULVA tubulosa simplex. *Fl. lapp.* 458. *Fl. suec.* 1013. *intestinalis.*
It. al. 33. *Hort. cliff.* 478. *Roy. lugdb.* 515. *Gron.*
virg. 204. 128.
Tremella marina tubulosa intestinorum figura. *Dill.*
musc. 47. t. 9. f. 7.
Fagus cavus. *Bauh. pin.* 364.
Habitat in Mari omni.
2. ULVA tubulosa ramosa compressa. *It. gotl.* 262. *Fl. compressa.*
suec. 1014. *
Tremella marina tenuissima & compressa. *Dill. musc.*
48. t. 9. f. 8.
Habitat in Europæ mari & testis maritimis.
3. ULVA filiformis articulata: articulis alternatim com- *confervoides*
pressis. *Hort. cliff.* 478. *
Conserua marina sinitulosa. *Dill. musc.* 34. t. 6. f. 39.
Habitat in Mari.
4. ULVA oblonga plana undulata membranacea viridis. *latissima.*
It. w:gotl. 100.
Fucus longissimo latissimo tenuique folio. *Bauh. prodr.*
154.
Habitat in Mari Europæ.
5. ULVA palmata prolifera membranacea: ramentis in- *Lactuca,*
ferne angustatis.
Ulva marina lactucæ similis. *Raj. angl.* 3. p. 62.
Tremella marina vulgaris, lactucæ similis. *Dill. musc.*
42. t. 8. f. 1.
Muscus marinus lactucæ similis. *Bauh. pin.* 364.
Habitat in Oceano.
Frondes aggregatæ, membranacæ, pallidæ, palmatæ: se-
gmentis singulis iterum in frondes palmatas enatis,
undatis, obovatis, obtusis, pellucidis.
6. ULVA fronde dilatata subfinuata, centro radicata. *umbilicatis.*
Tremella marina umbilicata. *Dill. musc.* 45. t. 8. f. 3.
Fucus umbilicus marinus dictus. *Bauh. pin.* 564.
Habitat in Oceano.
7. ULVA fronde oblonga bullata. *Linza,*
Tremella marina fasciata. *Dill. musc.* 46. t. 9. f. 6.
Fucus Linza dictus, lactucæ marinæ similis. *Bauh.*
hist. 3. p. 81.
Muscus lactucæ marinæ similis. *Bauh. pin.* 364.
Habitat in Oceano.
8. UL-

- peruviformis*. 8. *ULVA* subglobosa solitaria. *Fl. suec.* 1015. *
Fucus globosus sessilis mollis. Gleditsch. monogr.
Habitat in lacubus Sueciæ, Borussia.
- granulata*. 9. *ULVA* solitaria aggregata. *Fl. suec.* 1016. *
Tremella palustris, vesiculis sphaericis fungiformibus.
Dill. musc. 55. t. 10. f. 17.
Habitat in Europa ad ripas fluviorum.



CONFERVA.

- * *Filamentis simplicibus æqualibus geniculis destitu-*
tis.
- vicularis*. 1. *CONFERVA* filamentis simplicissimis æqualibus lon-
gissimis. *Fl. suec.* 1023.
Conferva capillaris simplicissima enodis. Roy. lugd.
511. Guett. stamp. 1. p. 45.
Conferva fluviatilis sericea vulgaris & fluitans. Dill.
musc. 12. t. 2. f. 2.
Habitat in Europæ rivulis fluvisque puerioribus.
- fontinalis*. 2. *CONFERVA* filamentis simplicissimis æqualibus di-
gito brevioribus.
Conferva minima, byssi facie. Dill. musc. 14. t. 2. f. 3.
Habitat in Europæ fontibus.
- * *Filamentis ramosis æqualibus.*
- bullosa*. 3. *CONFERVA* filamentis æqualibus ramosis, bullas
aëreas includens.
Conferva filis ramosis æqualibus radicatis. Fl. suec. 1024.
Conferva palustris bombycina. Dill. musc. 18. t. 3. f. 2.
Alge bombycina. Baub. pin. 363.
Musculus aquaticus bombycinus, tenuissimis filamentis.
Loef. pruss. 173. t. 55.
Habitat in Europæ aquis stagnantibus.
- canaliculata* 4. *CONFERVA* filamentis æqualibus basin versus ra-
mosioribus.
Conferva rivulorum capillacea, densissime congestis ra-
mulis. Dill. musc. 21. t. 4. f. 15.
Alga in tubulis aquam fontanam ducentibus. Baub. pin.
364.
Habitat in Europæ rivulis, tubulis, canalibus molen-
dinariis.
- emphibia*. 5. *CONFERVA* filamentis æqualibus ramosis, exsiccatio-
ne coeuntibus in aculeos.

con-

Conferva amphibia, fibrillosa & spongiosa. *Dill. musc.*
22. t. 4. f. 17.

Habitat in Europæ udis, fessis, umbrosis, ubi aquæ vicissitudines servant.

6. *CONFERVA* filamentis æqualibus ramosissimis e-littoralis.
longatis scabriusculis.

Conferva maritima filiformis longissima fusca, ramis alternis simplicibus brevioribus pallidis. *Fl. suec.* 1028.

It. gotb. 261. n. 3.

Conferva marina capillacea longa ramosissima. *Dill. musc.* 23. t. 4. f. 19.

Habitat in Europæ marinis rupibus.

7. *CONFERVA* filamentis ramosis mollibus digito brevioribus viridissimis. *æ-ruginosa*

Conferva marina capillacea brevis viridissima mollis.

Dill. musc. 23. t. 4. f. 20.

Habitat in Angliæ fucis.

8. *CONFERVA* filamentis æqualibus dichotomis. *dichotoma.*

Conferva dichotoma, setis porcinis similis. *Dill. musc.*

17. t. 3. f. 9.

Habitat in Angliæ fessis pratorum.

9. *CONFERVA* filamentis proliferis fastigiatis hirtis. *scoparia.*

Conferva marina pennata. *Dill. musc.* 24. f. 23.

Habitat in mari Europæo.

10. *CONFERVA* filis ramosis: filamentis alternis brevibus multipartitis digitatis. *cancellata*

Conferva marina cancellata. *Dill. musc.* 24. t. 4. f. 22.

Habitat in mari Europæo.

Filamenta lateralialia quasi intra totidem vesiculas aërem recondunt.

* *Filamentis anastomosantibus.*

11. *CONFERVA* filamentis reticulatim coadunatis. *reticulata.*

Conferva reticulata. *Dill. musc.* 20. t. 4. f. 14. *Fl. suec.*

125. *Guett. stamp.* 1. p. 45.

Conferva reticulata crispâ. *Pluk. alm.* 113. t. 24. f. 2.

Morif. bist. 3. p. 644. f. 15. t. 4. f. 4.

Muscus aquaticus bombycinus retiformis. *Læs. pruss.*

173. t. 54.

Habitat in Europæ fluviis ad eorum littora.

* *Filamentis nodosis.*

12. *CONFERVA* filis simplicissimis setiformibus rectis: *fluvialis.*
geniculis crassioribus angulatis. *Con-*

Conferva filis simplicissimis setiformibus rectis, tuberculis sparsis. *Fl. suec.* 1026.

Conferva fluviatilis lubrica setosa, equiseti facie. *Dill. musc.* 39. t. 7. f. 47.

Habitat in Europæ fluviiis confragosus.

gelatinosa. 13. *CONFERVA* filis ramosis moniliformibus, articulis globosis gelatinosis.

Conferva fontana nodosa, spermatis ranarum instar lubrica, major & fusca. *Dill. musc.* 36. t. 7. f. 42. *Guett. stamp.* 1. p. 46.

Conferva fontalis geniculata lubrica major. E. N. C. cent. 5. 6. app. 60. t. 13. f. 3.

Conferva fontana nodosa, spermatis ranarum instar lubrica, minor & viridis. *Dill. musc.* 37. t. 7. f. 43.

Corallina pingvis ramosa viridis. *Vaill. parif.* 40. t. 7. f. 6.

Habitat in Europæ fontibus præstantissimis limpidissimis.

Confer. *Conferva* stagnalis, globulis virefcentibus mucosis. *Dill. musc.* 38. t. 7. f. 44.

Conferva alpina lubrica, filamentis nodosis cæruleis. *Dill. musc.* 38. t. 7. f. 45.

Conferva fontana nodosa lubrica, filamentis tenuissimis nigris. *Dill. musc.* 39. t. 7. f. 46.

* *Filamentis geniculatis.*

capillaris. 14. *CONFERVA* filis geniculatis simplicibus: articulis alternatim compressis.

Conferva filamentis longis geniculatis simplicibus. *Dill. musc.* 25. t. 5. f. 25.

Conferva fluitans, filamentis geniculatis. *Pluk. alm.* 113. t. 84. f. 9. *Morif. hist.* 3. p. 644. f. 15. t. 4. f. 3.

Habitat in Europæ lacubus, stagnis, vadis, fossis.

corallinoides 15. *CONFERVA* filis geniculatis dichotomis.

Conferva marina gelatinosa, corallinæ instar geniculata crassior. *Dill. musc.* 33. t. 6. f. 36.

Habitat in Mari Europæo, colore albo & rubenti.

Confer. *Conferva* marina gelatinosa, corallinæ instar geniculata tenuior. *Dill. musc.* 33. t. 6. f. 37.

catenata. 26. *CONFERVA* filis geniculatis, articulis cylindricis.

Conferva ramosa, geniculis longioribus cateniformibus. *Dill. musc.* 27. t. 5. f. 27.

Ha.

Habitat in Mari Europam australem & Americam meridionalem alluente.

17. CONFERVA filamentis geniculatis, ramis fascicu- *polymorpha*.
latis.

Conferva marina articulata capillacea ramosissima viridis. *It. gotl.* 261, n. 2. *Fl. succ.* 1031.

Conferva marina geniculata nigra palmata. *Dill. musc.* 32. t. 6. f. 35.

Corallinæ affinis f. Muscus coralloides, multifido capillaceo folio, palmites pelagica nigra. *Pluk. alm.* 119. t. 47. f. 10.

Muscus tridactylites capillaceus maritimus multifidus niger, radice fibrosa. *Bocc. musc.* 1. p. 269. t. 5. f. 11.

Muscus marinus crispus pachycaulos loricatus niger. *Bocc. musc.* 315. t. 7. n. 1.

Fucus crispus loricatus nigricans, *Barr. rar.* 1323. t. 1301.

Habitat in Mari Europæo.

18. CONFERVA filamentis geniculatis flexuosis, ramis *vagabunda*.
ramulisque brevioribus.

Conferva marina trichodes, lanæ instar expansa. *Dill. musc.* 30. t. 5. f. 32.

Habitat in Mari Europæo, libera vagatur in aquarum medio.

19. CONFERVA filamentis geniculatis, ramulis brevi- *glomerata*.
oribus multifidis.

Conferva fontinalis ramosissima glomeratim congesta. *Dill. musc.* 28. t. 5. f. 34.

Conferva viridis capillacea, brevioribus setis, ramosior f. Conferva minor ramosa. *Morif. hist.* 3. p. 644. f. 15. t. 4. f. 2.

Alga fontalis trichodes. *Bauh. pin.* 364.

Habitat in Europæ fontibus, fossis, rivulis.

20. CONFERVA filamentis geniculatis ramosissimis vi- *rupestris*.
ridibus.

Conferva marina trichodes ramosior *Dill. musc.* 28. t. 5. f. 29.

Conferva marina trichodes f. Muscus marinus virens tenuifolius. *Pluk. mant.* 53. t. 182. f. 6.

Habitat in Europæ marinis rupibus copiosissima.

21. CONFERVA filamentis geniculatis ramosissimis e *aggregata*
centro confertissimis globum constituentibus.

Con-

Conserva globosa, filamentis articulatis ramosissimis e centro prodeuntibus rectis. *Fl. suec.* 1027.

Habitat in Sueciæ lacubus Dannemora, Stockholmæ. &c.

BYSSUS.

* *Filamentosa.*

- Flos aquæ.* 1. BYSSUS filamentis plumosis natantibus.
Byssus farinacea virescens aquæ insparfa. *Fl. lapp.* 532.
Fl. suec. 1128. *Guett. stamp.* 1. p. 3. *Roy. legab.* 520.
- β. Byssus membranacea aquatica. *Fl. lapp.* 529.
Byssus latissima, papyri instar supra aquam expansa. *Dill. musc.* 1.
Habitat in Mari & omni aqua, prima æstate; nocte descendit parum, die natat fere.
Constat farina cujus moleculæ, Conservæ instar, capillares, ramosæ, plumæ similes, niveæ. P. Bergius
- phosphorea.* 2. BYSSUS lanuginosa violacea lignis adnascens. *Dill. musc.* 4. t. 1. f. 6. *Fl. suec.* 1130.
Habitat in Europæ lignis putrescentibus.
- velutina.* 3. BYSSUS filamentosa viridis.
Byssus tenerissima viridis velutum referens. *Dill. musc.* 7. t. 1. f. 14.
Byssus terestris viridis herbacea & mollissima, filamentis ramosis & non ramosis. *Mich. gen.* 211. t. 89. f. 5.
Habitat in Terra.
- aurea.* 4. BYSSUS capillacea pulverulenta, fructificationibus sparsis, filamentis simplicibus ramosisque. *Gleditsch. fung.* 1.
Byssus petrææ crocea, glomerulis lanuginosis. *Dill. musc.* 8. t. 1. f. 16.
Byssus aurea derbenfis humifusa. *Pet. gaz.* 1. t. 15. f. 3.
Byssus minima saxatilis inodora, filamentis partim simplicibus partim ramosis. *Mich. gen.* 210. t. 89. f. 2.
Habitat in rupibus Arvoniz, Italiæ.
- cryptarum.* 5. BYSSUS capillacea perennis cinerea tenax rupi innata. *Fl. lapp.* 527. *Fl. suec.* 1127.
Habitat in spelunca rupestri Medelpadiæ.
- * *Pulverulenta.*
- antiquitatis.* 6. BUSSUS pulverulenta atra. *It. gosl.* 298. *Fl. suec.* 1126.
Habitat in ædium muris antiquissimis.

7. BYS-

7. BYSSUS pulverulenta cinerea rupes operiens. *saxatilis.*
Habitat in Saxo omni, diutius aeri exposito, quod perenni cinereo colore obducit, ipsa vix manifesta.
8. BYSSUS pulverulenta sanguinea faxis innascens. *Folictus.*
 Byssus perennis pulverulenta rubra lapidibus innascens.
*Fl. lapp. 528. Fl. succ. 1125. * Guss. stamp. 1. p. 3. Roy. lugdb. 520.*
 Byssus germanica minima saxatilis aurea violæ odorem spirans. *Mich. gen. 210. t. 89. f. 3.*
*Jolithus f. lapis violaceus. Swenck. foss. 382. It. w:goth. 9. 250. * 254. 263.*
Habitat in Europæ frigida sylvis opacis, supra saxa ante unum alterumve annum eversa & supinata.
9. BYSSUS pulverulenta flava lignis adnascens. *Dill. candeleris.*
musc. 3. t. 1. f. 4
 Byssus farinacea flava. *It. w:goth. 159. 52. It. al 30.*
Habitat per omnes quatuor mundi plagas in corticibus arborum, parietibus antiquis, tectis diutius venoso humido expositis.
10. BYSSUS pulverulenta viridis. *botryoides.*
 Byssus botryoides saturate virens. *Raj. angl. 3. p. 5. 6.*
Dill. musc. 3. t. 1. f. 5.
Habitat in terra diutius humida, umbrosa, ut in ollis Hortulanorum.
11. BYSSUS pulverulenta incana, farinæ instar strata. *incana.*
Dill. musc. 3. t. 1. f. 3. Fl. succ. 1131.
Habitat solo glareoso, ad latera fossularum, juxta vias publicas.
12. BYSSUS pulverulento-crustacea albissima. *lactea.*
 Byssus candidissima, calcis instar muscos vestiens. *Dill.*
musc. 2. t. 1. f. 2.
Habitat in Muscis & arborum corticibus.

SPONGIA. 7

* Tenaces.

1. SPONGIA tenax subramosa, ramis difformibus per-*officinalis.*
tusis. Roy. lugdb. 521. Mat. med. 527.
 Spongia ad usum præstantissima, foraminibus exiguis pervia. *Tournef. inst. 575.*
 Spongiæ compressæ Schiacciatæ. *Bauh. hist. 3. p. 818.*
 Spongia globosa. *Bauh. pin. 362.*
Habitat in Oceano australiore.

E c c c

2. SPON-

- oculata.* 2. SPONGIA tenax ramosissima erecta: ramis teretiusculis obrusis. *Hort. cliff.* 480. *Rev. lugdb.* 521. Spongiosus fucus. *Bauh. pin.* 369. *Habitat in Italia.*
- nodosa.* 3. SPONGIA scabra ramosa tenacissima: ramis nodosis perforatis. *Roy. lugdb.* 522. Spongia dura ramosa nigra suberis instar. *Boerb. lugdb.* 1. p. 9. *Habitat in Mari australi.*
- flabelliformis.* 4. SPONGIA flabelliformis rigidiuscula, fibris capillaribus densissime reticulatis. *Roy. lugdb.* 522. Spongia flabelliformis, caule teretiusculo, disco compresso-plano vix diviso. *Hort. cliff.* 480. Keratophyton majus nigrum, fibris tenuioribus elegantissime & densissime reticulatum. *Boerb. lugdb.* 1. p. 6. *Habitat in Indiis.*
- fistulosa.* 5. SPONGIA tubulosa simplex. *Hort. cliff.* 480. *Roy. lugdb.* 522. Spongia dura f. spuria major alba fistulosa, fibris crassioribus. *Sloan. jam.* 6. *bist.* 1. p. 62. t. 23. f. 2. *Habitat in mari ad Antillas.*
- tubulosa.* 6. SPONGIA tubulosa ramosa. *Roy. lugdb.* 522. Spongia ramosa fistulosa millepora. *Boerb. lugdb.* 1. p. 9. *Habitat in Indiis.*
- aculeata.* 7. SPONGIA cavernosa extus aculeata. *Roy. lugdb.* 522. Spongia dura f. spuria, superficie apicibus acutis exstantibus aspera, intus cavernosa. *Sloan. jam.* 7. *bist.* 1. p. 63. t. 23. f. 4. *Habitat in mari ad Americam.*
- varicata.* 8. SPONGIA ramosissima, poris cylindricis subulatis prominentibus æqualibus multitudinis hispida. *Tess. mus.* 118. t. 11. f. 1. * *Habitat in India.*
- infundibuliformis.* 9. SPONGIA turbinata cava. Spongia forma infundibuli. *Bauh. pin.* 369. Spongia elegans. *Clas. exot.* 125. *Habitat in Europa australi.*

* *Fragiles.*

10. SPONGIA repens fragilis, ramis teretibus obtusis. *lacustris.*
Hort. cliff. 480. * *Fl. succ.* 1132. *Ray. lugdb.* 522.

It. Jean. 422. *

Spongia ramis teretibus obtusis. *Fl. lapp.* 535.

Spongia ramosa fluviatilis. *Raj. hist.* 81.

Habitat in fundo Lacuum.

11. SPONGIA erecta fragilis, ramis teretiusculis. *Fl. fluviatilis.*
succ. 1133.

Spongia fluviatilis ad conservam accedens. *Rupp. jen.*
 308. t. 3. f. 1.

Spongia fluviatilis anfractuosa perfragilis ramosissima
Pluk. alm. 356. t. 112. f. 3.

Badiaga major. *Buxb. act.* 2. p. 343.

Mucos aquaticus ceratoides. *Læf. pruss.* 172. t. 52.

Habitat in Europæ fossis, paludibus.

FUNGI.

AGARICUS.

* *Stipitati, pileo orbiculato.*

1. AGARICUS stipitatus, lamellis ramosis decurrentibus. *Chentareh-
 luc.*
Fl. succ. 1049. *It. w. goth.* 215. *Dalib. paris.* 364.

Agaricus caulescens fulvus, pileo concavo: marginere-
 flexo, lamellis erectis, stipite brevissimo. *Fl. lapp.* 493.

Elvela pediculata pileata, pileo subtus reticulato. *Gyest.*
stamp. 1. p. 12.

Fungus minimus flavescens infundibuliformis. *Banh.*
pin. 373. *Vaill. paris.* t. 11. f. 9. 10.

- β. Fungus pileo per maturitatem instar agarici intybacei.
Vaill. paris. t. 11. f. 11. 12. 13.

- γ. Fungus angulosus & velut in lacinias sectus. *Vaill. pa-
 ris.* t. 11. f. 14. 15.

Habitat in Pratis.

Variat mire ex Agarico in Elvelam.

2. AGARICUS stipitatus, pileo subflavescente partito *quinquepar-
 titus.*
 lamellis albidioribus interne dentato-connexis.

Agaricus caulescens, pileo cinereo quinquepartito, la-
 mellis albis. *Fl. succ.* 1063. *

Habitat in Pratis.

3. AGARICUS stipitatus, lamellis omnibus magnitudi-*integer,*
 ne æqualibus. *Fl. succ.* 1071.

E e e e 2

Aga-

Agaricus caulescens, pileo plano-concavo purpureo, lamellis stipiteque albis. *Fl. lapp.* 486.

Agaricus major, pediculo longo modice crasso, lamellis albis creberrimis superne ad margines apparentibus. *Raj. suppl.* 18. *Buxb. cont.* 4. p. 12. 1. 19.

β. *Agaricus caulescens*, petiolo crassiusculo albo, pileo plano sanguineo, lamellis albis. *Fl. lapp.* 487.

Habitat in Sylvis.

mascorina. 4. *AGARICUS* stipitatus, lamellis dimidiatis solitariis, stipite volvato: apice dilatato, basi ovato. *Fl. succ.* 1076.

Agaricus caulescens, pileo sanguineo: verrucis lamellis-que albis, stipite albo basi globoso. *Fl. lapp.* 595.

Fungus pileo sanguineo verrucoso, lamellis albis, annulo fugaci, pediculo bulboso. *Hall. belv.* 39.

Fungorum perniciosorum genus 12. n. 4. *Clus. hist.* 2. p. 280.

Habitat in Pratis.

Cimices necat & eradicat promptissima. It. scan. 430.

dentatus. 5. *AGARICUS* stipitatus, pileo convexo, lamellis basi mucrone dentatis.

Agaricus caulescens, pileo flavo lævi margine inflexo, lamellis pallidioribus basi angulatis. *Fl. succ.* 1048.

Habitat in Sylvis.

Pileus flavescens, levis, subviscidus. Lamellæ basi a Stipite discedentes, ubi denticulo acuminato instructæ. Stipes fistulosus, angustus, cylindricus, nudus, versus pileum striatus, confertim natus.

deliciosus. 6. *AGARICUS* stipitatus, pileo testaceo succo lutescente.

Amanita lateritii coloris, croceo succo turgens. Dill. giff. 179.

Fungus perniciosus, lateritio colore varians, succum acrem & croceum fundens. *Mich. gen.* 141. *Hall. belv.* 49.

Habitat in Montibus, sterilibus, sylvestis.

lactifusus. 7. *AGARICUS* stipitatus, pileo plano carneo lactescente, lamellis rufis, stipite longo carneo. *Fl. lapp.* 484.

Fungus flavo-rufus in medio depressus, lacte non acri manans. *Hall. belv.* 50.

Habitat in Sylvis.

8. AGARICUS stipitatus, pileo planiusculo lactescente: *piperatus*,
 margine deflexo lamellis incarnato-pallidis. *Fl. suec.*
 1036.
 Fungus piperatus albus crassus, lacteo succo turgens.
Mich. gen. 141. *Hall. belv.* 34. n. 1.
 Fungus albus acris. *Bamb. pin.* 371.
Habitat in Pascuis, Sylvis.
9. AGARICUS stipitatus, pileo convexo squamato al-camp^{estris}
 bido, lamellis ruffis. *Fl. suec.* 1045.
 Agaricus caulescens, pileolo sordido lacero, petiolo al-
 bo membranaceo, lamellis rufescentibus. *Fl. lapp.* 482.
 Agaricus pediculo annulato, pileo superne albus, infer-
 ne rubens. *Guss. stamp.* 1. p. 22.
 Fungus campestris albus superne, inferne rubens. *Bamb.*
hist. 3. p. 824. *Mich. gen.* 174. *Hall. belv.* 50.
Habitat in Pratis.
10. AGARICUS stipitatus, pileo rimoso: margine vio-^{violacea}
 laceo tomentoso, stipite cærulescente lana ferruginea.
Fl. suec. 1066. *Dalib. paris.* 371.
 Fungus esculentus bulbosus dilute purpureus. *Mich. gen.*
 149. t. 49. f. 1.
 Fungus cæruleus major. *Buxb. cent.* 4. p. 7. t. 11.
Habitat ad margines Sylvarum.
11. AGARICUS stipitatus, pileo sordide flavo, lamellis ^{cinnamomeis}
 luteo-ruffis. *Fl. suec.* 1047. *
 Agaricus caulescens flavus, pileo plano. *Fl. lapp.* 594.
Habitat in Sylvis.
12. AGARICUS stipitatus, pileo purpurascente fusco viscidus.
 viscido, lamellis fusco-purpurascens.
 Agaricus caulescens, pileo convexo purpurascente, la-
 mellis ruffis, stipite brevi crassiusculo albo. *Fl. lapp.*
 485. *Fl. suec.* 1070.
Habitat in Sylvis.
13. AGARICUS stipitatus, pileo pallido: disco stella-^{equastris}
 tim luteo, lamellis sulphureis. *Fl. suec.* 1060. *
 Agaricus caulescens, pileo convexo sordido: margine
 indexo, lamellis pallidis: basi remotis, stipite cylin-
 draceo. *Fl. lapp.* 502.
Habitat in Pascuis, Sylvis.
14. AGARICUS stipitatus, pileo flavo convexo, lamel-^{gurgii}
 lis albis. *Fl. suec.* 1052.

E c c c 3

Aga-

Agaricus caulescens, pileo plano sulphureo, lamellis albis, stipite crassiusculo. *Fl. lapp.* 496.

Fungus orbicularis exalbidus pratensis. *Banb. pin.* 370. *Hall. belv.* 38.

Fungus divi georgii, coloris exalbidi cum pauca flavitie, esculenti pratensis 1. *Banb. hist.* 3. p. 824.

Fungus croceus magnus. *Sterb. fung.* 63. t. 11. f. 2. *Habitat in Sylvis.*

- mammosus.* 15. *AGARICUS* stipitatus, pileo convexo acuminato griseo, lamellis convexis griseis crenatis, stipite nudo. *Agaricus caulescens*, pileo convexo fordido: umbilico prominente, lamellis pallidis, stipite cylindraceo. *Fl. lapp.* 501.
Fungus minor, capitulo pileum nauticum referente, pediculo longiore. *Buxb. cent.* 4. p. 13. t. 21. f. 1. *Habitat in Sylvis.*

- clypatus.* 16. *AGARICUS* stipitatus, pileo hemisphærico viscido acuminato, lamellis albis, stipite longo cylindrico albo. *Fl. lapp.* 506. *Fl. succ.* 1057. *Dalib. parif.* 373. *Fungus viscidus papillaris striatus terreus*, subtus albus. *Hall. belv.* 41. *Habitat in pratis sylvaticis.*

- extinctivorus.* 17. *AGARICUS* stipitatus, pileo campaniformi albedo lacero, lamellis niveis, stipite subbulboso subulato nudo. *Agaricus caulescens albus*, pileo conico stipite cylindraceo. *Fl. lapp.* 481. *Fl. succ.* 1037. 1042. *Agaricus caulescens albus*, petiolo longissimo, pileo conico. *Fl. lapp.* 480. *Habitat in fimetis & ad pagos.*

- fimetarius.* 18. *AGARICUS* stipitatus, pileo campanulato lacero, lamellis nigris lateraliter flexuosis, stipite fistuloso. *Agaricus caulescens*, pileo conico cinereo, lamellis nigris: margine lacero, stipite longissimo albo. *Fl. lapp.* 503. *Fl. succ.* 1056. *Fungus comatus.* *Sterb. fung.* 218. t. 22. f. 7. K. L. *Fungus albus ovum* referens. *Buxb. cent.* 4. p. 16. t. 27 f. 1. *Fungorum perniciosorum genus* 14. *Spec.* 3. *Clas. hist.* 2. p. 282. *Habitat in Fimetis.*

19. A.

19. AGARICUS stipitatus, pileo campanulato striato ^{campanulatus.} pellucido, lamellis adscendentibus, stipite nudo.
 Agaricus caulescens, pileo conico cinereo lamellis albis, stipite longo. *Fl. lapp.* 507. *Fl. suec.* 1054.
 Fungus multiplex obtuse conicus, colore griseo murino. *Vaill. parif.* 71. t. 12. f. 1. 2.
Habitat in Pratis.
20. AGARICUS stipitatus, pileo lævi livido, lamellis nigricantibus separatis, stipite bulboso volvato.
Habitat in Stercoratis.
21. AGARICUS stipitatus, pileo convexo viscido pellucido lamellis luteis, stipite nudo.
 Fungus pileo croceo splendore participi *Vaill. parif.* t. 11. f. 16. 17. 18.
Habitat ad ambulacra.
22. AGARICUS stipitatus, pileo plicato membranaceo, ^{umbelliferus} lamellis basi latioribus. *Fl. suec.* 1033.
 Agaricus caulescens albus parvus, stipite longo, pileo plano pellucido: margine multifido. *Fl. lapp.* 471.
 Agaricus minimus, capitulo turbinato plano albo, lamellis margine fuscis. *It. æl.* 38.
 Fungus minimus totus albus, pileo hemisphærico utrinque striato, lamellis rarioribus. *Mich. gen.* 166. t. 80. f. 11.
Habitat inter Folia congesta, semiputrida.
23. AGARICUS stipitatus albus, pileo plicato membranaceo, stipite nigro. *Fl. suec.* 1034. *It. w. goth.* 215.
Dalib. parif. 374.
 Agaricus caulescens albus minimus, petiolo angusto, pileo plano verrucoso: margine octofido. *Fl. lapp.* 472.
 Fungus caule nigro capillari, androfaces capitulo. *Bocc. mus.* 143. t. 104.
 Fungus pileo candicante, lamellis paucis, pediculo fusco splendente. *Vaill. parif.* 69. t. 11. f. 21. 22. 23
Habitat in foliis dejectis Pini.
24. AGARICUS stipitatus, pileo luteo convexo striato, ^{clavus} lamellis stipiteque albis. *Fl. suec.* 1053.
 Fungus minimus aurantius mammillaris. *Vaill. parif.* 76. t. 11. f. 19. 20. *Hall. belv.* 38.
Habitat in Nemoribus, inter folia decidua.

* *Parasitici, acaules, dimidiati.*

- Quercinus.* 25. **AGARICUS** acaulis, lamellis labyrinthiformibus.
Fl. suec. 1082.
 Agaricus villosus, lamellis sinuosis & invicem implexis.
Buxb. cent. 5. p. 1. 4. f. 1.
 Agarico fungus lamellis crassissimis rigidis. *Hall. belv.*
 57.
Habitat in Quercubus.
- Betulinus.* 26. **AGARICUS** acaulis coriaceus villosus: margine obtuso, lamellis ramosis anastomosantibus. *Fl. suec.* 1085.*
Habitat in Betulis.
- Alnus.* 27. **AGARICUS** acaulis, lamellis bifidis pulverulentis.
Fl. suec. 1083.
 Agarico-Fungus lamellis bifidis pulverulentis. *Hall. belv.*
 58.
Habitat in Alno.
Paucas species Fungorum recenseo, ut evitem varietates, quæ plures sunt, quam vulgo creditur, & quidem sæpe aliena genera constituentes, ut in his etiamnum valde deficiat Res herbaria.

BOLETUS.* *Acaules, parasitici.*

- suberosus.* 1. **BOLETUS** acaulis pulvinatus albus lævis, poris acutis difformibus.
 Boletus acaulis coriaceo-convexus villosus albus, poris difformibus rotundis flexuosisque. *Fl. suec.* 1093.
Habitat in Betulis.
- fomentarius.* 2. **BOLETUS** acaulis pulvinatus inæqualis obtusus, poris teretibus æqualibus glaucis. *Fl. suec.* 1092.*
Habitat in Betulis.
- ignarius.* 3. **BOLETUS** acaulis pulvinatus lævis, poris tenuissimis.
Fl. suec. 1096.
 Agaricus pedis equini facie. *Tournef. inß. 56a.*
Habitat in Betulis aliisque arboribus.
- versicolor.* 4. **BOLETUS** acaulis: fasciis dicoloribus poris albis.
 Agarico-polyporus versicoloribus lineis variis, poris albis tenuissimis. *Hall. belv. 26.*
Habitat ad truncoſ Arborum antiquarum.

5. BO-

5. BOLETUS acaulis, superne lævis, salicinus. *Fl. Suec. lenc.*
lapp. 522. Fl. suec. 1095.
Habitat in Salice.

* *Stipitati.*

6. BOLETUS stipitatus perennis, pileo utrinque planius- *perennis.*
sculo. Fl. lapp. 519. Fl. suec. 1086. Dalib. paris. 383.
 Fungus lignosus fasciatus. *Vaill. paris. t. 12. f. 7.*
Habitat in sylvis super terram, subiectis candidibus ar-
borum putridis.
7. BOLETUS stipitatus, pileo pulvinato viscido, poris *viscidus.*
 teretibus convexis immersis distinctis, stipite lacero.
 Boletus caulescens, pileo luteo, poris teretibus distinctis
 lividis, stipite lacero. *Fl. suec. 1089.*
Habitat in Sylvis.
8. BOLETUS stipitatus, pileo pulvinato subviscido, po- *luteus.*
 ris rotundatis convexis flavissimis, stipite albido.
 Boletus caulescens, pileo livido subviscido, poris flavis
 convexis, stipite albido. *Fl. suec. 1088. Dalib. paris.*
381.
 Suillus annulatus terreus, inferne flavescens. *Hall.*
helv. 29.
 Fungus porosus autumnalis viscidus. *Buxb. cent. 5. p.*
7. t. 14.
Habitat in Sylvis.
9. BOLETUS stipitatus, pileo glabro pulvinato margina- *bovinus.*
 nato, poris compositis acutis: porulis angulatis bre-
 vioribus.
 Boletus caulescens, pileo testaceo, poris compositis
 angulatis griseis. *Fl. suec. 1087. Dalib. paris.*
382.
 Boletus julii mensis. *Dill. giff. 188.*
 Suillus fulvus inferne ex flavo virescens. *Hall. helv. 29.*
Habitat in Pratis.
10. BOLETUS stipitatus, pileo viscido pulvinato, po- *granulatus.*
 ris teretibus subangulatisque truncatis: angulo granu-
 lato.
Habitat in Sylvis.
 Pileus convexus, viscidus, carnosus, lividus, margine
 acuto cinctus. Pori flavi, difformes, inaequales, tere-
 tes & subangulati, truncati: margine laterali ad an-
 gulos quasi farina particula solitaria aucti. Stipes fla-
 vus, pileo brevior.

- subtomentosus*. 11. **BOLETUS stipitatus**, pileo flavo subtomentoso, poris subangulatis difformibus fulvis planis, stipite flavo. *Fl. suec.* 1091. *
Habitat in sylvis.
- subfauces*. 12. **BOLETUS stipitatus**, pileo albido, poris difformibus oblongo-flexuosis niveis. *Fl. suec.* 1090.
Habitat in Sylvis.

HYDNUM.

- imbricatum*. 1. **HYDNUM stipitatum**, pileo convexo imbricato. *Fl. lapp.* 523. *Fl. suec.* 1097. *It. W. goth.* 215. *Roy. lugdb.* 519.
Erinaceus esculentus albus crassus. Mich. gen. 132. t. 72. f. 2.
Fungus pæne candidus, præna parte erinaceus. Buxb. hist. 3. p. 818.
Habitat in Sylvis acerofis.
- repandum*. 2. **HYDNUM stipitatum**, pileo convexo lævi flexuoso. *Fl. suec.* 1098. *Dalib. parif.* 383.
Hydnum pediculatum, pileo inæqualiter laciniato scabro flavo. Guett. stamp. 1. p. 383.
Erinaceus esculentus pallide luteus. Mich. gen. 132. t. 72. f. 3.
Habitat in Sylvarum desertis.
- tomentosum* 3. **HYDNUM stipitatum**, pileo plano-infundibuliformi. *Fl. suec.* 1099.
Habitat in Sylvis acerofis.
- auriscalpium* 4. **HYDNUM stipitatum**, pileo dimidiato. *Fl. lapp.* 524. *Fl. suec.* 1100. *Roy. lugdb.* 519.
Erinaceus parvus hirsutus ex fulco fulvus, pileo semi-orbiculari, pediculo tenuiore. Mich. gen. 132. t. 72. f. 8.
Fungus erinaceus parvus in conis abietis nascens. Buxb. cens. 1. t. 57. f. 1.
Fungus erinaceus parvus, pediculo longiore auriscalpium referens buxei coloris. Buxb. bull. 129. t. 129.
Habitat in Sylvis acerofis supra terram, subjacente ramo aut strobilo.

PHALLUS.

- esculentus*. 1. **PHALLUS** pileo ovato, stipite nudo rugoso. *Fl. suec.* 1102. *Dalib. parif.* 383.

Phal-

Phallus petiolo nudo, pileo subtus undique annexo. *Hort. cliff.* 379. *Roy. lugdb.* 517.

Boletus esculentus rugosus albicans, quasi fuligine infertus. *Tournef. inst.* 561. t. 329. f. A.

Boletus esculentus rugosus 1, 2, 3. *Mich. gen.* 203. t. 85. f. 1. 2.

Fungi esculenti species tres. *Clus. hist.* 2. p. 164.

Habitat in Sylvis antiquis.

7. PHALLUS volvatus stipitatus, pileo celluloso. *impudicus.*

Phallus volvatus pileo apice clauso. *Att. Stockb.* 1732. p. 20. t. 2. f. 1. *Fl. suec.* 1101.

Phallus volva exceptus, pilei apice pervio. *Hort. cliff.* 478. *Roy. lugdb.* 517. *Guesst. stamp.* 1. p. 17.

Phallus vulgaris totus albus, volva rotunda, pileo cellutato ac summa parte umbilico pervio ornato. *Mich. gen.* 201. t. 83.

Phallus hollandicus s. batavicus. *Dalech. hist.* 1398.

Fungus foetidus penis imaginem referens. *Bauh. pin.* 374.

Habitat in Sylvis.

CLATHRUS.

* Acaules.

1. CLATHRUS acaulis subrotundus. *Hort. cliff.* 479. *cancellatus.*

Roy. lugdb. 518.

Clathrus ruber. *Mich. gen.* 214. t. 93.

Boletus cancellatus purpureus. *Tournef. inst.* 561. t. 329. f. 6.

Fungus coralloides cancellatus. *Clus. app. alt.*

Fungus rotundus cancellatus. *Bauh. pin.* 375.

Habitat in Europa australiori.

* Stipitati.

2. CLATHRUS stipitatus, capitulo oblongo volvato. *dennuatum.*

Clathrus pediculatus purpureus, capite oblongo. *Guesst.*

stamp. 1. p. 16.

Clathroides purpureum pediculo donatum. *Mich. gen.*

214. t. 94. f. 1

Habitat in Europa australiore.

3. CLATHRUS stipitatus, capitulo oblongo axi longitudinali adnato. †

Clathroidastrum. *Mich. gen.* 214. t. 94.

Habitat in Italiae lignis putridis.

FL-

ELVELA.

* *Stipitata*,*Mitra.*

1. ELVELA pileo flexo adnato lobato difformi. *Fl. succ.* 1103.
 Fungoides fungiforme crispum laciniatum & varie complicatum. *Mich. gen.* 204. t. 86. f. 7.
 Fungus autumnalis velut apex flaminis. *Menz. pug.* t. 6.
Habitat in Truncis putridis.

* *Acaulis*.*Pinet.*

2. ELVELA acaulis.
 Agaricus acaulis utrinque planiusculus. *Fl. lepp.* 517. *Fl. succ.* 1084.
Habitat in Pinu, Abiete.

PEZIZA.

* *Acaules*.*lentifera.*

1. PEZIZA campanulata lentifera.
 Peziza calyce campanulato. *Hort. cliff.* 479. *Roy. lugdb.* 518.
 Peziza caliciformis lentifera laevis. *Dill. giff.* 195. *Fl. succ.* 1109.
 Cyathoides cyathiformis cinereum & veluti sericeum. *Mich. gen.* 222. t. 102. f. 1.
 Fungoides infundibuli forma, semine foetum. *Vaill. parif.* 56. t. 11. f. 6. 7.
 Fungus campaniformis niger parvus, multa semina in se continens. *Pluk. phys.* 184. f. 9.
 Fungus pyxoides seminifer. *Laf. pruss.* 98. t. 16.
 Fungus caliciformis seminiferus. *Menz. pug.* t. 6.
 β. Peziza calice campanulato, interne striato, externe squamoso. *Guett. stamp.* 1. p. 15. *Dalib. parif.* 387.
Habitat in Agris, Lignis, Sepimentis.

punctata.

2. PEZIZA turbinata truncata, disco punctato. *Fl. succ.* 1111.
 Elvela petiolo brevissimo, pileo turbinato, disco punctato. *Hort. cliff.* 479. *Roy. lugdb.* 518.
 Fungus minimus infundibuliformis, superne nigris punctis notatus. *Raj. suppl.* 21.
 Muscus minimus lignosus, disco punctato *Bocc. musc.* 2. p. 25. t. 107.

Ha-

Habitat in Stercore equino.

Discus niveus, punctis atris eminentibus.

3. PEZIZA infundibuliformis, disco patente sinuato punctato. *cornucopioides.*

Elvela infundibuliformis: ore expanso laciniato punctato. *Guett. stamp. 1. p. 11, Dalib. parif. 384.*

Fungoides nigricans majus cornucopiæ forma. *Vaill. parif. t. 13. f. 2. 3.*

Habitat in Gallia.

4. PEZIZA cyathiformis: extus angulata venis ramosis. *Acetabulum.*

Fungoides fuscum acetabuliforme: externe ramificatum. *Vaill. parif. 57. t. 13. f. 1.*

Fungoides fuscum majus. *Buxb. act. 4. p. 283. t. 29. f. 1. 2.*

Fungoides maximum vasis fusorii forma. *Misch. gen. 206. t. 8. 6. f. 1.*

Habitat in Europa australi.

5. PEZIZA cyathiformis: margine obtuso erecto. *Fl. cyathoides.*

fucc. 1110.
Peziza lutea parva: marginibus lævibus. *Raj. angl. 3. p. 479. t. 24. f. 4.*

Habitat in Terra.

6. PEZIZA globoso-campanulata: margine crenato. *cupularis.*

Peziza calice campanulato globoso: ore crenato. *Guett. stamp. 16. Dalib. parif. 388.*

Fungoides glandis cupulam reterens, margine dentato. *Vaill. parif. 57. t. 11. f. 1. 2. 3.*

Habitat in Gallia,

7. PEZIZA plana: margine convexo piloso. *scutellata.*

Peziza lutea parva, marginibus pilosis. *Raj. angl. 3. p. 18. t. 24. f. 3.*

Elvela sessilis scutellata. *Guett. stamp. 1. p. 10.*

Fungoides, qui Fungus minimus scutellatus, coloris aurantii. *Vaill. parif. t. 13. f. 13. 14.*

Habitat in Parietibus patris.

8. PEZIZA turbinata cochleata. *cochleata.*

Elvela sessilis multis anfractibus cochleata. *Guett. stamp. 1. p. 10. Dalib. parif. 384.*

Fungoides auriculam judæ reterens, intus rufescens, extus

rus candicans & quasi farinosum. *Vaill. parif. 57. t. 11. f. 8.*

Habitat in umbrosis.

CLAVARIA.

* *Indivise.*

- pistillaris.* 1. **CLAVARIA** clavæformis simplicissima. *Fl. suec. 1104. Dalib. parif. 385.*
Clavaria clavata integerrima obtusa erecta. Hort. cliff. 479. Roy. lugdb. 517.
Clavaria alba pistilli forma. Vaill. parif. 39. t. 8. f. 5.
Clavaria major lutea. Mich. gen. 208. t. 87. f. 1.
Fungus clavatus albidus pistillaris. Bocc. mus. 1. t. 307.
Habitat in Sylvis umbrosis.

- militaris.* 2. **CLAVARIA** clavata integerrima: capite squamosa. *Guett. stamp. 1. p. 8. Dalib. parif. 385.*
Clavaria militaris crocea. Vaill. parif. 39. t. 7. f. 4.
Habitat in sylvis australibus.

- Ophioglossoides.* 3. **CLAVARIA** clavata integerrima compressa obtusa. *Guett. stamp. 1. p. 8. Dalib. parif. 386.*
Clavaria ophioglossoides nigra. Vaill. parif. 39. t. 7. f. 3.
Habitat in sylvis australibus.

* *Ramosæ*

- digitata.* 4. **CLAVARIA** ramosa lignea nigra. *Guett. stamp. 1. p. 10. Dalib. parif. 387.*
Agaricus digitatus niger. Tournes. inst. 562.
Habitat in sylvis australibus.

- Hypoxylon.* 5. **CLAVARIA** ramoso-cornuta compressa. *Fl. suec. 1105.*

Fungus ramosus niger compressus parvus: apicibus albidis Raj. angl. 3. p. 15.

- β. *Agaricus ramosus cornu rangiferi referens. Mich. Eberes. tab.*

Hypoxylon digitatum. Bruck. monogr.

Fungus cornu dorcadis facie. E. N. C. dec. 1. ann. 4. p. 195.

Habitat in Cellis, navibus, aliisque nunquam sole illustratis.

- coralloides.* 6. **CLAVARIA** ramis confertis ramosissimis inæqualibus. *Fl. suec. 1106. Dalib. parif. 386.*
Clavaria ramosissima. Roy. lugdb. 518.

Co-

Corallo-fungus flavus. Vaill. parif. 41. t. 8. f. 4.

Coralloides flava & albida. Tournesf. inst. 564. t. 332. f. B.

Barba caprina. Sterb. fung. 96. 97. t. 11. f. A. B. C. D.

Habitat in Sylvis opacis.

7. **CLAVARIA** ramis confertis ramosissimis fastigiatis *fastigiata*, obtusis luteis. *Fl. succ. 1107.*

Fungoides fungiforme luteum foetidum & minus ramosum. Raj. angl. 3. p. 479. t. 24. f. 5.

Habitat in Sylvis.

8. **CLAVARIA** ramis ramosis acutipatis inæqualibus *muscoides*, luteis. *Fl. succ. 1108.*

Fungus parvus ramosus luteus. Raj. angl. 3. p. 16. t. 24. f. 7.

Habitat inter Muscos.

LYCOPERDON.

* *Solida, subterranea absque radice.*

1. **LYCOPERDON** solidum muricatum, radice destitutum.

Lycoperdon solidum. Fl. lapp. 526. Fl. succ. 1116.

Roy. lagdb. 519. Dalib. parif. 391.

Tuber brumale, pulpa obscura odorata. Mich. gen. 221. t. 164.

Tuber. Matsh. Dissc. 544. Mich. gen. 221. t. 102.

Habitat sub Terra.

2. **LYCOPERDON** solidiusculum centro farinifero, *Cervinum* radice destitutum.

Lycoperdastrum tuberosum arrhizon fulvum, cortice duriore crasso & granulato, medulla ex albo purpurascente, semine nigro crassiore. Mich. gen. 220. t. 99. f. 4.

Tubera cervina. Bauh. pin. 376.

Tuberum genus quibusdam Cervi Boletus. Bauh. hist. 3. p. 851.

Habitat in Bohemia, Silesia. Vermelandia.

* *Puiverulenta, radicata supra terram.*

3. **LYCOPERDON** subrotundum lacerato dehiscens. *Bovista* *Fl. lapp. 525. Fl. succ. 1115. Dalib. parif. 388. Roy. lagdb. 518.*

Lycopperdon vulgare. Tournesf. inst. 563.

Fun-

Fungus maximus rotundus pulverulentus, dictus germanis Bofilt. *Baux. hist.* 3. p. 848.
Habitat in Campis sterilibus.

- aurantium.* 4. LYCOPERDON sphaeroidale basi rugosum stipitatum: laciniis obtuse emarginatis dehiscens. *Dalib. parif.* 390. n. 9.
 Lycoperdon aurantii coloris, ad basin rugosum. *Vaill. parif.* 123. t. 16. f. 9. 10.
Habitat in Gallia.
- stellatum.* 5. LYCOPERDON volva multifida patente, capitulo glabro: ore acuminato dentato. *Hort. cliff.* 479. *Fl. succ.* 1113. *Dalib. parif.* 390. *Roy. lugdb.* 519.
 Lycoperdon stellatum, calyce inverso. *Baux. cent.* 2. p. 45. t. 49. f. 3.
 Geaster major, umbilico fimbriato. *Mich. gen.* 220. t. 100. f. 1. 2. 3.
 Fungus pulverulentus, Crepitus lupidictus, coronatus & interne stellatus. *Raj. angl.* 3. p. 27. t. 1. f. 1.
 Fungus stellatus. *Bocc. mus.* 1. t. 305.
Habitat in Collibus.
- Carpobolus.* 6. LYCOPERDON volva multifida, fructu globoso ex seminibus combinatis. †
 Carpobolus. *Mich. gen.* 221. t. 101.
Habitat in Italia.
- pedunculatum.* 7. LYCOPERDON stipite longo, capitulo globoso glabro: ore cylindrico integerrimo. *Hort. cliff.* 478. *Fl. succ.* 1112. *Roy. lugdb.* 519. *Hall. helv.* 12. *Dalib. parif.* 390.
 Lycoperdon parisiense minimum pediculo donatum. *Tournef. inst.* 563. t. 331. f. E. F.
 Fungus pulverulentus minimus, pediculo longo insidens. *Raj. angl.* 3. p. 27.
Habitat in Campestribus.
- * *Parasitica, in farinam fatiscientia.*
- Epidendrum* 8. LYCOPERDON cortice farinaque purpurca. *Fl. succ.* 114.
 Lycoperdon epidendron, miniatum pulverem fundens. *Baux. hall.* 203. *Hall. helv.* 11.
 Lycoperdon sanguineum sphaericum 1. *Baux. cent.* 5. p. 15. t. 29. f. 2.
Habitat in Lignis, Parietibus antiquis.

9. LYCOPERDON aggregatum parasiticum: ore mul-epiphyllum,
tindo sacro: pulvere fulvo.

Habitat in dorso foliorum Tussilaginis.

MUCOR.

* Perennes.

- .. MUCOR perennis, stipite filiformi nigro, capitulo *sphærocephalus* globoso cinereo. *Fl. suec.* 1121. *It. scan.* 20.
Sphærocephalus niger, villo ochroleuco. *Hall. belv.* 3.
t. 1. f. 3.
Habitat in Parietibus, Lapidibus, Lignis.

- a. MUCOR perennis, stipite subulato nigro, capitulo *Lichenoides* lenticulari cinereo. *Fl. suec.* 1122. *
Coralloides fungiforme arborum nigrum vix crustosum.
Dill. musc. 78. t. 14. f. 3.
Habitat in corticibus Pini.

3. MUCOR seta nigra, villo fusco. *Embolus.*
Embolus seta nigra villo fusco. *Fl. suec.* 1136.
Embolus nigerrimus villo albo adspersus. *Hall. belv.*
8. t. 1. f. 1.
Habitat in lignis putridis.

4. MUCOR perennis pallidus, pileo fulvo. *fulvus.*
Habitat Uptaliæ D. Solander.

5. MUCOR perennis viridis, foliis furfuraceis, stipite *fil-furfuraceus* filiformi, capitulo globoso.
Habitat in Terra nuda passim in Suecia. Dan. Solander.

* Fugaces.

6. MUCOR stipitatus, capsula globosa. *Mucedo.*
Mucor filamentis granulatis fugacibus putrescentibus innatis. *Fl. lapp.* 534. *Fl. suec.* 1119. *Roy. lugdb.* 520.
Mucor vulgaris, capitulo lucido per maturitatem nigro, pediculo griseo. *Micb. gen.* 215. t. 95. f. 1.
Mucedo capitulo diaphano deinde viridi & postremo nigro. *Malp. vegetat.* t. 25. f. 108. p. 2.
Habitat in variis putridis: pane, plantis. &c.

7. MUCOR setaceus, seminibus radicalibus. *leprosus.*
Aspergillus cespitosus densissimus, initio niveus, deinde aureus, seminibus ovatis. *Micb. gen.* 213. t. 91.
f. 5.
Habitat in cavernulis autumnæ.

F f f

8. MU-

- glauens.* 8. MUCOR stipitatus, capitulo subrotundo, aggregato. Aspergillus capitatus, capitulo glauco, seminibus rotundis. *Mich. gen.* 212. *t.* 91. *f.* 1.
Habitat in Citris, Melonibus, Pomis aliisque corruptis.
- crustaceus.* 9. MUCOR stipitatus, spicis digitatis. Mucor crustaceus albus, capitulis setisque aqueis. *Fl. suec.* 1118.
Byssus filamentosa fugax alba, putrescentibus innata. *Fl. lapp.* 533. *Gnett. stamp.* 1. *p.* 3. *Roy. lugdb.* 520.
Botrytis non ramosa alba, seminibus rotundis. *Mich. gen.* 212. *t.* 91. *f.* 3.
Habitat in Cibis corruptis.
- capitosus.* 10. MUCOR stipite ramoso, spicis ternatis. Aspergillus albus tenuissimus, graminis dactyloidis facie, seminibus rotundis. *Mich. gen.* 213. *t.* 9. *f.* 3.
β. Aspergillus terrestris capitosus & ramosus albus. *Mich. gen.* 213. *t.* 93. *f.* 4.
Habitat in Putrescentibus.
- Erysiphe.* 11. MUCOR albus, capitulis fuscis sessilibus. *Fl. suec.* 1124.
Habitat in foliis Humuli, Aceris, Lamii, Galeopsidis, Lithospermi.

PALMÆ.

FLABELLIFOLIÆ.

CHAMÆROPS. 1025

1. CHAMÆROPS frondibus palmatis plicatis, stipitibus humilis.
 spinosis. *Hort. cliff.* 482. *Roy. lugdb.* 4.
 Palma minor. *Bauh. pin.* 506.
 Palma humilis f. Chamæripes. *Bauh. hist.* 1. p. 368.
 Chamæripes. *Dod. pempt.* 820.
Habitat in Europa australi, præsertim Hispania.

BORASSUS. 1026

1. BORASSUS frondibus palmatis plicatis cucullatis, flabellifer.
 stipitibus serratis.
 Borassus frondibus palmatis. *Fl. zeyl.* 395.
 Palma coccifera, folio plicatili flabelliformi; mas, fe-
 mina. *Raj. hist.* 1366.
 Ampana. *Rheed. mal.* 1. *Carimpana. Rheed. mal.* 1.
 p. 13. *t.* 10. *p.* 11. *t.* 9.
 Lontarus domestica. Rumpf.
 amb. 1. *p.* 45. *t.* 10.

Habitat in India.

CORYPHA. 1027

1. CORYPHA frondibus pinnato-palmatis plicatis filo umbraenlifo-
 interjectis. *Hort. cliff.* 482. *Fl. zeyl.* 394. *Roy. lugdb.* 4. *ra.*
 Palma montana, folio plicatili flabelliformi maximo se-
 mel tantum frugifera. *Raj. hist.* 1363.
 Codda-pana. *Rheed. mal.* 3. *p.* 1. *t.* 1. -- 12.
 Saribus. *Rumpf. amb.* 1. *p.* 42. *t.* 8.
Habitat in India.
Confer. Palma thebaica. *Pocock. orient.* 1. *p.* 281. *t.* 73. thebaica.
enj. Frondes pinnato-palmatae plicate, filo interjecto,
stipitibus ciliato-spinosis, Caudice dichotomo.

PENNATIFOLIÆ

1028

CYCAS.

- circinalis*. 1. **CYCAS** frondibus pinnatis circinnalibus: foliolis planis.
Cycas frondibus pinnatis; foliolis linearilanceolatis,
 stipitibus spinosis. *Hort. cliff.* 482. *Fl. zeyl.* 393. *Roy.*
Ingdb. 5.
Palma indica, caudice in annulos protuberante distin-
 ctio. *Raj bist.* 1360.
Arbor Zagocamboinensis. *Seb. thes.* t. p. 39, t. 25. f. 1.
Tessio. *Kämpf. jap.* 897.
Olus calappoides *Rumpf. amb.* 1. p. 86. t. 22. 23.
Todda-pana f. *Mouta-panna*. *Rheed. mal.* 3. p. 9. t.
 13. -- 21.
Habitat in India.
Foliatio circinalis more Filicum peragitur.

1028

COCOS.

- nucifera*. 1. **COCOS** frondibus pinnatis: foliolis replicatis.
Coccus frondibus pinnatis: foliolis ensiformibus mar-
 gine villosis. *Hort. cliff.* 483. *Fl. zeyl.* 391. *Roy.*
Ingdb. 4.
Palma indica coccifera angulosa. *Bauh. pin.* 502.
Palma indica nucifera. *Bauh. bist.* 1. p. 375.
Calappa. *Rumpf.* 1. p. 1. t. 1. 2.
Tenga. *Rheed. mal.* 1. p. 1. t. 1. 2. 3. 4.
Habitat in Indiæ paludosis, umbrosis.
Foliola omnia (excepto utrinque infimo) retro-plicata
sunt, contra ac in sequente.

1030

PHOENIX.

- dactylifera*. 1. **PHOENIX** frondibus pinnatis: foliolis complicatis.
Phoenix frondibus pinnatis: foliolis alternis ensiformi-
 bus basi complicatis, stipitibus compressis dorso rotun-
 datis. *Hort. cliff.* 482. *Hort. ups.* 306. *F. zeyl.* 390.
Roy. Ingdb. 5.
Palma major. *Bauh. pin.* 506.
Palma dactylifera major vulgaris. *Sloan. jam.* 174.
Palma. *Bauh. bist.* 1. p. 351. *Dod. pempt.* 819. *Raj.*
bist. 1352.
Palma hortenensis mas. | *Palma hortenensis femina.*
Kämpf. amœn. 688. t. | *Kämpf. exot.* 668. 686.
 1. 2. f. 1. 2. | t. 1. 2. f. 2. 16. 11.
Habitat in India.

ARE-

PALMÆ BIPENNATIFOLIÆ. 1189

ARECA. 1031

1. ARECA frondibus pinnatis: foliolis plicatis oppositis *Caribæo* præmorsis. *Fl. zeyl.* 392.

Areca f. Fauſel, avellana indica verſicolor. *Raj. biſt.* 1363.

Palma arecifera nucleo verſicolore moſchatæ ſimill.

Pluk. alm. 275. t. 309. f. 4.

Palma cujus fructus ſeſſilis Fauſel dicitur. *Bauh. pin.* 510.

Pinanga. *Rumpf. amb.* 1. p. 26. t. 4.

Caunga. *Rheed. mal.* 1. p. 9. t. 5. 6. 7. 8.

Habitat in India.

ELATE. 1032

1. ELATE frondibus pinnatis: foliolis oppoſitis. *Sylveſtris.*

Palma daſtylifera minor humilis ſylveſtris, fructu minore. *Herm. prodr.* 361. *Fl. zeyl.* 397.

Palma ſylveſtris malabarica, folio acuto, fructu pruniſacie. *Raj. biſt.* 1364.

Katou-indel. *Rheed. mal.* 3. p. 15. t. 22. 23. 24. 25.

Habitat in India.

BIPENNATIFOLIÆ.

CARYOTA. 1033

1. CARYOTA frondibus bipinnatis: foliolis cuneiformibus oblique præmorsis. *Fl. zeyl.* 396.

Palma indica, folio bicompoſito, fructu racemoſo. *Raj. biſt.* 1365.

Palma daſtylifera, fructu acerrimo. *Plum. ſpec.* 3.

Seguaſter major.. *Rumpf. amb.* 1. p. 64. t. 14.

Schunda panna. *Rheed. mal.* 1. p. 15. t. 11.

Habitat in India.

Palmarum omnium hiſtoria fructificationis adhuc imperfecta eſt, nec ſuffecit induſtria *Rheedi*, cui tamen debemus in his quidquid adhuc certi innotuerit. Utinam Botanici nobis traderent diſtinctas harum ſpecies ex ſtructure frondium, quam ſerius & varius apud nos floreat.

APPENDIX.

1034

ACHRAS.

Zapota.

1. ACHRAS.

Sapota fructu ovato majori. *Plum. gen.* 43.Zapotes. *Laët. amer.* 277. 332. 343.Cypote. *Cbaër. sciagr.* 24.Avellanæ indicæ 3 genus. *Baub. pin.* 418.Malus Persica maxima, foliis magnis integris longis, fructu maximo oblongo scabro, ossiculo partim rugoso partim glabro. *Sloan. jam.* 180. *hist.* 2 p. 124. t. 218.Arbor americana pomifera, frondosis ramulis, foliis amplis longioribus obtusis duris & venosis margine æquali. *Pluk. alm.* 39. t. 268. f. 2.Fructus oblongus utrinque acuminatus f. conicus lævis splendens spadiceus. *Raj. hist.* 1800.*Habitat in Jamaica.* b

1035

ADANSONIA.

digitata.

2. ADANSONIA. B. *Jussiai.*Boabab. *Alp. ægypt.* 66. t. 67.Baobab. f. Abavi. *Baub. hist.* 1. p. 110.Abavo arbor, radice tuberosa. *Baub. pin.* 434.Guanabanus scaligeri. *Baub. hist.* 1. p. 109. *Raj. angl.* 1371.*Habitat in Ægypto,* b*Arborem dixit summus Botanicus B. Jussæus a discipulo egregio, qui Africa rarissimas plantas investigavit; generis characterem non dum nobis notum ab amicissimo Auctore indefinenter expetimus.*

1036

BROSSÆA.

coccinea.

1. BROSSÆA.

Brossæa frutescens, flore coccineo, fructu nigro. *Plum. gen.* 5.*Habitat in America calidiore.*

Rami alterni. Folia alterna, ovata, serrata, petiolata.

Flores aliquot, ramos terminantes, alterni.

1037

BUCEPHALON.

racemosum.

1. BUCEPHALON.

Bucephalon fructu racemoso rubro. *Plum. gen.* 52.*Habi-*

Habitat in America.

Folia ovata, acuminata, integerrima, alterna: petiolis brevissimis. Spicæ laterales, solitariae, oppositæ.

FUCHSIA. 10 3 8

1. FUCHSIA.

triphylla.

Fuchsia triphylla, flore coccineo. Plum. gen. 14.

Habitat in America.

Caulis herbaceus, simplicissimus. Folia terna, laecolata, sessilia. Racemus terminalis, erectus.

HIPPOCRATEA. 10 3 9

1. HIPPOCRATEA. *Hort. cliff. 484.*

volubilis.

Coa scandens, fructu trigemino subrotundo. Plum. gen. 8.

β. *Coa scandens, fructu trigemino oblongo. Barr. aquin. 36.*

Habitat in America calidiori.

Caulis volubilis. Folia opposita, ovata, crenata, petiolata.

HIPPOMANE. 10 4 0

1. HIPPOMANE foliis ovatis serratis. *Hort. cliff. 484. Mancinella.*

Macanilla pyri facie. Plum. gen. 50. Catseb. car. 1.

p. 95. t. 95.

Juglandi affinis arbor julifera lactescens venenata pyrifolia Sloan. jam. 129. hist. 2 p. 3. t. 159.

Malus americana, laucerali folio, venenata. Comm. hort. 1. p. 131. t. 68.

Arbor venenata Mancinello dicta. Raj. bist. 1646.

Habitat in Caribæis sapiens inundatis.

2 HIPPOMANE foliis ovato-oblongis serratis: basi glandulosa glandulosis.

Macanilla lauri foliis oblongis. Plum. gen. 50.

Pitymalus arbor americana, mali medicæ foliis amplioribus tenuissime crenatis, succo maxime venenoso.

Pluk. alm. 369. t. 229. f. 8. Raj. suppl. 428.

Habitat in America calidiore. b.

Folia utrinque, ad latus baseos petioli, glandulam gerunt.

Amentum subulatum, longissimum, terminale.

3 HIPPOMANE foliis subovatis dentato-spinosis.

spinosi

Macanilla aquifolii foliis. Plum. gen. 50

Ilex, aquifolii folio, americana. Pluk. alm. 197. t. 196. f. 3.

F f f f 4

116-

Habitas in America calidore. ♀.

Ad classem Monœcia spectat, ordine non dum manifesto

1041

HYMENÆA.

- Courbaril.* 1. HYMENÆA. *Hort. cliff.* 484. *Hort. upf.* 305. *Mat. med.* 515.

Ceratia diphyllous antegoana, ricini majoris fructu nigro, siliqua grandi incluso. Pluk. alm. 96. t. 82. f. 2.

Arbor siliquosa ex qua Gumi Elemi elicitor. Bamb. pin. 404.

Habitas in Brasilia. ♀

1042

MATTHIOLA.

- Jobrah.* 1. MATTHIOLA.

Matthiola folio aspero subrotundo, fructu nigricante.

Plum. gen. 16.

Raregal arbor indiana. Zanow. hist. 167. t. 75?

Habitas in America.

1043

PLUKENETIA.

- volubilis.* 1. PLUKENETIA.

Plukenetia scandens, hederæ foliis serratis, fructu serrato. Plum. gen. 47.

Sajor volubilis, fructibus corniculatis. Rumpf. amb. 1. p. 194. t. 79. f. 2.

Habitas in Indiis.

Caulis volubilis. Folia cordata, serrata, acuta. Petioli laxi. Racemus compositus ex flosculis masculis numerosis, infimis semineis paucis. Ad monœciam pertinet.

1044

RANDIA.

- inermis.* 1. RANDIA inermis.

Randia. Hort. cliff. 485.

Lycium majus americanum, jasmini flore, foliis subrotundis lucidis. Pluk. alm. 234. t. 97. f. 6.

Cacao affinis frutex spinosus, lyci facie, jasmini flore albo. Sloan. jam. 135. *hist.* 2. p. 18. t. 161. f. 1.

Raj. dendr. 82.

Habitas in Jamaica. ♀

- aculeata.* 2. RANDIA aculeata.

Randia aculeata, buxi foliis majoribus. Juss. Roy. lugdb. 537.

Habitas - - - ♀

RHEE-

RHEEDIA. 1045

1. RHEEDIA.

lateriflora.

Van Rheedia folio subrotundo, fructu luteo. *Plum. gen.*
45.

Habitat in America.

Folia opposita, ovata, integerrima, petiolata. Pedunculi laterales, simplices, uniflori, numerosi.

RUMPHIA. 1046

2. RUMPHIA.

ambicincta.

Myxa pyriformis, officulo trispermo. *Raj. bist. 1556.*

Thiem-tani. *Rheed. mal. 4. p. 25. t. 11.*

Habitat in India. 3

Pertinet ad Triandriam Monogynam.

THALIA. 1047

3. THALIA. *Roy. lugdb. 11.*

geniculata.

Cortusa arundinacea, amplis cannatori foliis. *Plum. gen. 26.*

Canna indica, radice alba alexipharmaca. *Sloan. jam. 22. bist. 1. p. 253. t. 149. f. 2. Raj. suppl. 573*

Canna indica angustifolia, pediculis longis ad imum folium nodo singulari geniculato. *Pluk. alm. 79.*

Radix quædam in malacca, quæ ad versus vulnera sagittis toxico illitis facta, præsentaneum remedium.

Baub. pin. 301.

Agutigvepo-obi. *Marcgr. bras. 53. Raj. bist. 1203.*

Habitat in America.

Folia alterna, ovato-oblonga: nervis transversis, parallelis. Petioli screses, geniculo instructi; inferne membranacei vaginales. Culmus summus ramosus, exserens e singula spatula florem unum alterumve.

TREVIA. 1048

4. TREVIA.

nudiflora.

Arbor racemosa malabarica, fructu triquetra. *Comm. mal. 30.*

Causchi. *Rheed. mal. 1. p. 76. t. 42.*

Habitat in Malabariz arcuosis. 3

Ad classem Monacice pertinet.

XIMENIA. 1049

5. XIMENIA foliis solitariis.

americana.

Ximenia. *Hort. cliff. 483.*

Ximenia aculeata, flore villosa, fructu luteo. *Plam. gen. 6.*

F f f f s

Ho-

Habitat in America. b

Folia ovato-oblonga, alterna, petiolata, integerrima,
Spina ad singulam axillam foliorum. Racemus mul-
tistlorus inter spinam & ramum.

- egyptiaca.* 2. XIMENIA foliis geminis.
Lycio affinis ægyptiaca. *Banb. pin.* 479. *Raj. suppl.* 1629.
Agihalid. Alp. ægypt. 38. t. 39.
Habitat in Ægypto. b
- diandra.* 5. BOERHAVIA* glabra, floribus diandris. (*inferenda.* pag. 3. post 4.)
Habitat in India. Ott. Manchausen. L.B.
Habitus & statura omnino Boerhavia diffusa, sed Caulis & fructus glabri, pedunculi folio breviores, sæpius triflori. Flores minores, albi, pedicellati. Fructus sulcati, pentagoni, truncati, glabri.
- peltatus.* 32. CONVULVULUS foliis peltatis, pedunculis multistloris. (155. post 12.)
Convulvulus lævis indicus major. *Rumpf. amb.* 6. p. 428. t. 159.
Habitat in Amboina.
- pulemoides.* 10. VERBASCUM foliis ovatis utrinque tomentosis, inferioribus petiolatis. (178. post. 2.)
Verbascum femina, flore luteo magno. *Banb. pin.* 239.
Verbascum montanum tomentosum & incanum, folio subrotundo, caule non alato, staminibus purpureis.
Till. piss. 171.
Habitat in Europa australi. D. Monti &
Caulis erectus, spicatus, valde tomentosus. Folia ovata, minime cordata, sed basi integra, crenata, inferiora petiolata, petiolo supra plano, superiora sessilia, subamplexicaulia, non decurrentia. Racemus spiciformis ex bracteis sparsis, lanceolatis. Flores intrasingulam bracteam quatuor, quorum intermedius primum, tum infimus, demum laterales florent. Capsulæ oblongæ.
- Cervaria.* 3. SELINUM foliis radicalibus ovatis inæqualiter serratis. *Hors. cliff.* 92. *Hort. ups.* 59. (244. post 2.) †
Daucus montanus, apii folio, major. *Banb. pin.* 150.
Danci tertium genus *Fuchs. hist.* 233.
Cervaria. Riv. pent. 12.
Habitat in Helvetiæ, Sabaudia, Geneva, Alsatia, Galloprovinciæ montibus. 2

9. A-

9. **ATHAMANTA** foliis subbipinnatis: foliolis deorsum *condensa-*
imbricatis, umbella lenticulari. (244. *post* 1.)

Habitat in Sibiria. 2

Radix perennis. Caulis simplex, pedalis, levis, inferne angulatus, superne sulcatus praesertim pedunculo. Rami duo vel tres, floriferi, ex summis alis foliorum. Petioli canaliculati, basi membranacei, subtus angulati. Folia subbipinnata, foliolis alternatim pinnatifidis lobo infimo majore, unde quasi cruciata evadunt oppositiones ad basin & inferior lacinia se extendit supra folium inferius; subtus nitida. Umbella densissima. ntrinque convexa, antequam floret rufescens, Flores tamen albi, exceptis antheris fuscis. Petala inflexa. Involucrum universale saepe deficit & tantum rudimenta ostendit. Partiale longitudine florum.

31. **ALLIUM** scapo subnudo tereti, foliis linearibus, *canadense.*
capitulo bulbofero. (300. *inter* 22. - 23!)

Habitat in Canada. Kalm. 2

Radix perennis, Folia linearia, plana, levia, recta, spithamea, vix ad caulem ad basi vaginantia. Scapus nudus, teres, levis. Spatha ovata, mutica, acutiuscula. Capitulum ex bulbis; pauci flosculi pedicellati albi: Petalis ovalibus. Stamina simplicia longitudine corollae: Antheris rufescentibus.

9. **GYPHOPHILA** foliis lanceolatis laevibus, caulibus *prostrata.*
diffusis, pistillis corolla campanulata longioribus. (407.
post 2.)

Habitat - - - semina misit D. Wuchendorff.

Radix perennis protrudit Caules plurimos, undique procumbentes, diffusos, laeves, teretes, vix pedales, virides, nec ad articulos rufescentes. Folia lanceolata, glabra, vix ramosa. Panícula terminalis, brachiata, trichotoma: intermedio pedunculo simpliciore in triangulum erecto. Calyx viridis. Corolla alba: petalis obtusis, apice canaliculatis, patentibus. Stamina corolla dimidio breviora, Antheris flavis. Styli corolla paulo longiora. Habitus omnino Gypsophilæ repentis, a qua differt radice vix repente; geniculis canis viridibus nec purpurascensibus. Calycis foliolis non carinatis. Corollis minoribus albis nec subrubentibus; Antheris flavis; Pistillis petalo longioribus. Stylis apice divergentibus, Foliis minus carnosiss.

8. STEL-

arenaria.

8. STELLARIA caule erecto bitido, ramis alternis, petalis emarginatis. (423. post 7.)

Habitat in Hispania.

Radix fibrosa, annua. Caulis spithameus, erectus, teres, pilis paucis pubescens & subglutinosus. Rami alterni, longitudine fere caulis. Folia opposita, sessilia, subsus & margine pilosa, supra vero vix manifeste, ligulata apice ovato, basin versus angustiora linearia. Primus flos e dichotomia caulis, ubi caulis bifidus evadit; reliqui ex alis ramorum alterni, omnes majusculi. Pedunculus erectus, foliis longior. Calyx pentaphyllus, viscosus uti tota planta, apicibus purpurascens uti etiam folia floralia. Corolla campanulata, calyce longior, obtusissima vel vix manifeste emarginata aut retusa, alba. Stamina alba, longitudine calycis. Antheræ cerulescentes. Germen globosum. Styli 3, longitudine staminum.

hispanica.

7. COTYLEDON foliis oblongis subteretibus, floribus fasciculatis. (430. post 6.)

Cotyledon maritima, sedi folio, flore carneo, fibrosa radice. *Tournef. inst.* 90.

Cotyledon hispanica, sedi teretifolii folio, flore umbellato rubro, fibrosa radice. *Minnart. sched.* 3.

Habitat in Hispania. Læsing. ♂

Radix fibrosa. Caulis simplex digitalis, teres, erectiusculus. Folia oblonga, teretiuscula, obtusiuscula, sessilia, subpilosa, supra planiuscula, rufopunctata. Corymbus terminalis. Flores infundibuliformes. Corollæ erectæ, extus rufescentes, subpilosa: limbo plano, quinquepartito, acuto, supra incarnato striis tribus purpurascens & apice purpureo. Antheræ minime, filamenta crassiuscula, prominula. Calyx breviss. Facies *Herbæ* Sedi albi.

dubium.

9. PAPAVER capsulis oblongis glabris, caule multifloro setis adpressis foliis pinnatifidis incis. (507. post 5.)
Argemone capitulo longiore glabro. *Morif. blef.* 233. *bist.* 2. p. 279. f. 3. t. 14. f. 11.

Habitat inter Sueciæ, Angliæ segetes. ☉

Planta media inter P. argemone & Rhæas. Capsulis clavatis & pilis pedunculo adpressis cum P. Argemone convenit & a P. Rhæde differt.

squamatus.

29. CISTUS suffruticosus stipulatus, foliis obtusis squamis orbiculatis (526. post. 13.)

Hab

Habitat in Hispania. Læst. b

Suffrutex ramis erectiusculis, inferne tetragonis. Folia ovali-lanceolata, crassiuscula, petiolata, opposita aut tria utrinque. Stipulæ minutissimæ, reflexæ, marcescentes, sessiles. Pedunculi teretes. Planta tota oblecta squamis orbiculatis centro depresso.

11. RUBUS foliis ternatis subtus tomentosis, caule parvis foliis. tiolisque aculeis recurvis. (494. post. 5.)

Rubus moluccanus parvis foliis. Rumpf. amb. 5. p. 88. t. 47. f. 1.

Habitat in Ir. Osbeck. b

Caules fruticosi, teretes, incani. Folia ternata, subtus albo-tomentosa, venosa, petiolis subtus recurvato-aculeatis. Calyces tomentosi, in racemum digesti.

12. RUBUS foliis simplicibus cordatis lobatis, caule a-moluccanus. culeato decumbente. (494. post. 6.)

Rubus moluccanus latifolius. Rumpf. amb. 5. p. 88. t. 47. f. 2.

Habitat in Amboina. b

22. ANEMONE foliis reniformibus lobatis crenatis, in-malvisolia. volucro multifido. (539. post. 7.)

Anemone, cyclaminis f. malvæ folio, lutea. Baub. pin. 173.

Anemone hortensis latifolia, simplici flore. Clus. hist. 1. p. 248.

Habitat in Hispania ad vias. x

Corollæ exteriora petala subtus hirsuta, ut in Pulsatillis Tournefortii.

6. OCYIMUM caule suffruticoso, foliis ovatis, racemis gratissimum. teretibus. (597. post. 4.)

Habitat in India. b

Frutex bipedalis: Caule subserugineo, fusco. Folia opposita, ovata, acuta, profunde serrata, pallide virida, subtus scabra, serraturis 7 s. 9 utrinque ovatis. Petioli longitudine folii, supra subvillosi, plani. Racemi terminales, sessiles. Bractæ oppositæ, reflexæ, cordato-lanceolatae, nec concavæ, acuminatæ. Flosculi tres intra singulam bracteam, pedicello longitudine calycis. Calyx & Corolla Ocymi tenuiflori. Corolla minima. Odor gratiss.

29. ANTIRRHINUM foliis subulatis canaliculatis car-spartenm. notis: inferioribus ternis, caule paniculato corollisque glaberrimis. (ad pag. 612. post. 10.)

Flaki-

Habitat in Hispania. Læsing.

Radix altero anno florens, Caulis pedalis, glaberrimo, paniculato, erecto, minus stricto: ramis virgatis. Folia primordialia; antequam caulem floriferum erigit, terrena, oblonga; reliqua alterne subulata, canaliculata, glabra, carnosæ, rectæ. Flores racemosi. Calyces glabri Corollæ flavæ, palato glabro cum tinctura rubicunda in retuso gibbo palati.

molle.

30. ANTIRRHINUM corollis subcaudatis, calycibus lanceolatis pedunculatis, foliis subtomentosis ovatis (617. ante 25.)

Antirrhinum hispanicum villosum, origani folio. *Tern. inst.* 168. glutinoso. *Barr. ic.* 598. *mal.*

Habitat in Hispania.

Facies Ant. majoris, a quo differt Caulibus prostratis, Foliis caulinis oppositis. Herba tota villo vix conspicua mollissima, calcari corollæ magis protuberantæ; Pedunculis flore longioribus.

fenestralis.

31. CHEIRANTHUS foliis conferto-capitatis recurvatis undatis. (662. post. 4.)

Habitat - - - - -

Radix fibrosa. Caulis semipedalis infra flores, crassitie digiti, erectus, inferne nudus. Folia in capitulum dense congesta, lanceolata, obtusa, incana, sessilia, recurvata, undata. Racemi erecti, stricti. Flores purpurei & Siliquæ tomentosæ, curvatae ut in Ch. incano, cui maxime affinis, vix tamen varietatem dixerim.

legopodioides.

34. HEDYSARUM foliis ternatis ovatis obtusis, spicis hirsutis. (749. post 25.)

Habitat in China. Osbeck.

Folia ternata: foliolis ovatis, obtusis, nudis, minima seta terminatis. Spicæ ovata, densa, hirsutissima. Flores interstincti. Bracteis ovatis, subulato-acuminatis, longitudine florum. Pedicelli hirsuti. Calyces brevissimi, terminati setis tribus hirsutis. Corollæ parvæ. Legumina monosperma.

resedifolia.

9. SCORZONERA foliis pinnato-dentatis, caule ramossissimo subnudo divaricato (791. post 6.)

Tragopogon, resedæ minoris folio, supinum. *Barr. ic.* 800.

Habitat in Hispania. Læsing.

24. CAR-

24. **CARDUUS** acaulis, calyce glabro. *M. succ.* 656. *acaulor.*
It. gotl. 297. * *Dalib. parif.* 248. (826. *post* 23)
Carlina acaulos, minore purpureo flore. *Baub. pin.* 380
Chamæleon exiguus. *Baub. hist.* 3. p. 62.
Habitat in Europæ pascuis apricis, depressis. 2
3. **FILAGO** floribus pyramidalis pentagonis: flosculis fe- pyramidata.
mineis seriatis.
Habitat in Hispania. Læsing. ☉
Caulis pollicem unum alterumve altus, simplex, cre-
tus. Rami 4. s. 5, radicales, ascendentes, simpli-
ces. Folia lanceolata, obtusiuscula. Habitus *Gnapha-*
lis germanici. Flores caulem terminant sessiles, cincti
foliis in orbem positis, longioribus & ramis duobus
simplicibus, ut flores e dichotomia caulis quasi orian-
tur. Flores plures, pentagoni, pyramidati, tomentosi ut
tota planta, unus in medio, quem quinque cingunt, &
hos plures, compositi paleis Calyce communis quinque
farctam imbricato squamis acuminatis & inter hos flo-
sculi nudi feminei, stylo bipartito divaricato; discus
minimus, quinqueflorus, masculus absque paleis in Fi-
lagine prima calyx teres nec pentagonus est.
15. **QUERCUS** foliis lanceolato-ovatis integerrimis in- molucca,
divisis glabris. (994. *post* 1.)
Quercus molucca. Rumpf. umb. 3. p. 85. t. 56.
Habitat in Moluccis. b
14. **CROTON** foliis lanceolatis petiolatis integerrimis variegatun.
glabris. (1004. *post* 1.)
Codizum chrysolichum. Rumpf. 4. p. 65. t. 25.
C. Codizum teniosum. Rumpf. 4. p. 68. t. 26.
C. Codizum sylvestre. Rumpf. 4. p. 69. t. 27.
Habitat in India.
Opus eximium beati Rumpfi cura amplissimi D. Bur-
manni urbi botanico redditum, ad me accessit primum
absoluto a typographo opere, cujus itaque synonyma ali-
bi seorsim tradere animus est.

ADDENDA.

- P. 17. *Gratiola virginiana* (adde) *Tfiera Manga Nari*. *Rheed. mal.* 9. p. 165. t. 85.
25. *Salvia pratensis*. - *Habitat in Europæ pratis.* &
353. *Erica arborea* - - *Habitat in Europa australi.* &
494. *Rubus Chamæmorus*. *Hæc species omnino Dioica est.*
665. *Arabis canadensis*. *Habitat in America septentrionali.*
696. *Hibiscus furattensis* *Herba crinium.* *Rumpf. amb.* 4. p. 40. t. 16.
721. *Lupinus angustifolius*. *Habitat in Hispania.* ☉
781. *Medicago polymorpha*. *Habitat in Europa australi.* ☉
825. *Carduus tataricus*. - *Folia radicalia in hac specie pinnatifida sunt, non vero in præcedenti.*
479. *Pyrus communis* (rectus) *PYRUS* foliis serratis, floribus corymbosis.
- Pyrus Malus* (rectius) *PYRUS* foliis serratis, umbellis sessilibus.



INDEX GENERUM.

A.

Acalypha

Acanthus
Acer
Achillea
Achyranthes
Achras
Acnida
Aconitum
Acorus
Acrolicum
Aëlea
Adansonia
Adenanthera
Adiantum
Adonis
Adoxa
Ægilops
Æginetia
Ægopodium
Æichynomene
Æsculus
Ethusa
Agaricus
Agave
Ageratum
Agrimonia
Agrostemma
Agrostis
Aira
Ajuga
Aizoon
Alcea
Aldrovanda
Aletis
Alchemilla
Alisma
Allium
Allophylus
Aloe
Alopecurus
Alpinia
Alpine
Althæa
Alyssum
Amaranthus
Amaryllis
Ambrosia
Amethystea
Aminannia
Ammi
Amomum

1003
639
1054
896
204
1190
1037
532
324
1067
504
1190
334
1094
547
367
1050
632
265
713
344
256
1171
323
839
448
435
61
63
561
488
687
281
319
123
342
294
348
319
60
2
272
686
650
939
292
987
21
119
247
1

Amorpha
Amygdalus
Anabasis
Anacardium
Anacyclus
Anagallis
Anagyris
Anastatica
Anchuta
Andrachne
Andromeda
Andropogon
Androsace
Andryala
Anemone
Anethum
Angelica
Annona
Anchemis
Anthericum
Anthoceros
Anthospermum
Anthoxanthum
Antholyssa
Anthyllis
Antidema
Antirrhinum
Aphanes
Aphyllanthus
Apium
Apluda
Apocynum
Aquilegia
Arabis
Arachis
Aralia
Arbutus
Arctium
Arctopus
Arctotis
Areta
Arenaria
Aretia
Arethusa
Argemone
Aristida
Aristolochia
Arnica
Artemisia
Artemisia
Arum

713
472
223
383
892
148
374
641
133
1014
393
1045
141
808
538
263
250
536
893
310
1139
1158
28
37
719
1027
612
23
294
264
82
213
532
644
741
293
395
816
1058
922
1189
423
141
950
508
82
960
884
242
845
964

Arundo
Asarum
Asclepias
Ascyrum
Aspalathus
Asparagus
Asperugo
Asperula
Alphodelus
Asplenium
Aster
Astragalus
Ailanthia
Athamanta
Attaçtylis
Atragene
Atraphaxis
Atriplex
Atropa
Avena
Averrhoa
Avicennia
Axyris
Azalea

81
448
214
787
711
313
138
103
309
1078
871
755
235
244
849
542
333
1052
181
79
488
110
979
150

B.

Baccharis
Bæckea
Bailora
Bannisteria
Barleria
Barreria
Bartia
Barramia
Basella
Bauhinia
Begonia
Bellis
Bellonia
Berberis
Basleria
Beta
Betonica
Betula
Bidens
Bignonia
Biscutella
Biserrula
Bixa
Blasia
Blasia

260
358
582
427
616
274
602
389
372
1056
896
172
330
619
222
573
982
831
622
642
762
512
1138
Blasia

20

INDEX GENERUM.

<i>Aleria</i>	112	<i>Camphorolpa</i>	122	<i>Chrysobalanus</i>	511
<i>Blechnum</i>	1077	<i>Canna</i>	1	<i>Chrysocoma</i>	846
<i>Blitum</i>	4	<i>Cannabis</i>	1027	<i>Chrysogonum</i>	920
<i>Bobartia</i>	54	<i>Capparis</i>	501	<i>Chrysophyllum</i>	192
<i>Bocconia</i>	505	<i>Capraria</i>	628	<i>Chrysosplenium</i>	391
<i>Boerhavia</i>	1	<i>Capsicum</i>	188	<i>Cicer</i>	711
<i>Boerhaavia</i>	1176	<i>Cardsmine</i>	654	<i>Cichorium</i>	811
<i>Bombax</i>	511	<i>Cardiospermum</i>	366	<i>Cicuta</i>	851
<i>Bontia</i>	638	<i>Carduus</i>	820	<i>Cinchona</i>	172
<i>Borassus</i>	1187	<i>Carex</i>	971	<i>Cinna</i>	1
<i>Borbonia</i>	707	<i>Carica</i>	1036	<i>Cinrea</i>	1
<i>Borrigo</i>	137	<i>Carlina</i>	828	<i>Cissampelos</i>	1031
<i>Bosca</i>	225	<i>Carpesium</i>	557	<i>Cissus</i>	157
<i>Brahejum</i>	121	<i>Carpinus</i>	9098	<i>Cistus</i>	523
<i>Brassica</i>	666	<i>Carthamus</i>	830	<i>Cistharexylum</i>	625
<i>Breynia</i>	503	<i>Cerum</i>	263	<i>Citrus</i>	781
<i>Briza</i>	70	<i>Caryophyllus</i>	1119	<i>Clathrus</i>	1179
<i>Bromelia</i>	285	<i>Caryota</i>	376	<i>Clavaria</i>	1181
<i>Bromus</i>	76	<i>Castia</i>	268	<i>Claytonia</i>	204
<i>Brossia</i>	1190	<i>Cassine</i>	35	<i>Clematis</i>	543
<i>Browallia</i>	631	<i>Cassya</i>	812	<i>Cleome</i>	671
<i>Brunfelsia</i>	191	<i>Catananche</i>	107	<i>Clerodendrum</i>	677
<i>Brudia</i>	199	<i>Catesbea</i>	240	<i>Clethra</i>	396
<i>Bryonia</i>	1012	<i>Caulis</i>	195	<i>Cliffortia</i>	1038
<i>Bryum</i>	1115	<i>Ceanothus</i>	196	<i>Clinopodium</i>	587
<i>Bubon</i>	252	<i>Celastrus</i>	205	<i>Clitoria</i>	753
<i>Bucephalon</i>	1190	<i>Celofia</i>	621	<i>Clusia</i>	509
<i>Buchnera</i>	630	<i>Celtis</i>	1043	<i>Clypeola</i>	652
<i>Buddleja</i>	112	<i>Cenchrus</i>	1049	<i>Cneorum</i>	52
<i>Bufonia</i>	123	<i>Centaurea</i>	909	<i>Cnicus</i>	826
<i>Bulbocodium</i>	294	<i>Centunculus</i>	116	<i>Cochlearia</i>	647
<i>Buonia</i>	669	<i>Cephalanthus</i>	95	<i>Coccos</i>	1188
<i>Banum</i>	243	<i>Cerastium</i>	457	<i>Coffea</i>	171
<i>Buphralmum</i>	903	<i>Ceratocarpus</i>	969	<i>Coix</i>	971
<i>Bupleurum</i>	236	<i>Ceratonia</i>	1020	<i>Colchicum</i>	341
<i>Burmannia</i>	287	<i>Ceratophyllum</i>	998	<i>Coldenia</i>	125
<i>Bunomus</i>	372	<i>Certhera</i>	208	<i>Collinsonia</i>	78
<i>Buxus</i>	983	<i>Cercis</i>	374	<i>Columnnea</i>	682
<i>Byssie</i>	1168	<i>Cerlinthe</i>	136	<i>Colurea</i>	718
		<i>Ceropegia</i>	211	<i>Comarum</i>	508
		<i>Celstrum</i>	191	<i>Commelina</i>	40
		<i>Cachrys</i>	246	<i>Conferia</i>	1164
		<i>Cherophyllum</i>	258	<i>Conium</i>	243
		<i>Chaenactis</i>	1187	<i>Connarus</i>	675
		<i>Chamaecypariss</i>	1156	<i>Conocarpus</i>	176
		<i>Chelidonium</i>	661	<i>Convallaria</i>	314
		<i>Chelone</i>	505	<i>Convolvulus</i>	153
		<i>Chenopodium</i>	611	<i>Conyza</i>	861
		<i>Cherleria</i>	218	<i>Corchorus</i>	529
		<i>Chionanthus</i>	425	<i>Cordia</i>	190
		<i>Chironia</i>	8	<i>Coreopsis</i>	905
		<i>Chondrilla</i>	189	<i>Coriandrum</i>	256
		<i>Chrysanthemum</i>	706	<i>Coriaria</i>	1027
			887		

C.

Cacalis
Cacta
Catalpa
Celastrus
Calceolus
Calla
Callicarpa
Callitriche
Calophyllum
Caltha
Camellia
Cameraria
Campanula

314
466
380
921
325
968
111
969
513
558
698
210
263

Cassia
Cassine
Cassya
Catananche
Catesbea
Caulis
Ceanothus
Celastrus
Celofia
Celtis
Cenchrus
Centaurea
Centunculus
Cephalanthus
Cerastium
Ceratocarpus
Ceratonia
Ceratophyllum
Certhera
Cercis
Cerlinthe
Ceropegia
Celstrum
Cachrys
Cherophyllum
Chaenactis
Chamaecypariss
Chelidonium
Chelone
Chenopodium
Cherleria
Chionanthus
Chironia
Chondrilla
Chrysanthemum

511
846
920
192
391
711
811
851
172
1
1
1031
157
523
625
781
1179
1181
204
543
671
677
396
1038
587
753
509
1042
652
52
826
647
1188
171
971
341
125
78
682
718
508
40
1164
243
675
176
314
153
861
529
190
905
256
1027

Coria

INDEX GENERUM.

[illegible]

INDEX GENERUM.

Alberia	218	Camphorolipa	122	Chrysobalanus	512
Blechnum	1077	Canna	1	Chrysocoma	840
Blirum	4	Cannabis	1027	Chrysogonum	920
Bobartia	54	Capparis	503	Chrysophyllum	192
Bocconia	504	Capraria	628	Chrysopsplenium	396
Boerhavia	1	Caplicum	188	Cicer	718
Boletus	1178	Cardsmine	654	Cichorium	813
Bombax	511	Cardioidpermum	366	Cicuta	256
Bontia	638	Carduus	820	Cinchona	172
Borassus	1187	Carex	973	Cinna	5
Borbonia	707	Carica	1036	Circua	9
Borrage	337	Carina	828	Cissampelos	1031
Bosca	225	Carlinia	857	Cissus	157
Brachyum	121	Carpeium	909	Cistus	523
Brafica	666	Carpinus	830	Citharexylum	625
Breynia	503	Carthamus	263	Citrus	722
Briza	70	Casum	515	Clathrus	1179
Bromelia	285	Caryophyllus	1119	Clavaria	1122
Bromus	76	Caryota	376	Claytonia	204
Brossca	1190	Callia	268	Clematis	543
Browallia	631	Calline	35	Cleome	678
Brunsfelfia	191	Callitha	812	Clerodendrum	637
Brudia	199	Carananche	109	Clethra	396
Bryonia	1012	Catesbea	240	Cliffortia	1023
Bryum	1115	Caulalis	195	Clinopodium	527
Budon	252	Ceanothus	196	Clitoria	753
Bucephalon	1190	Celastrus	205	Clusia	509
Buchnera	650	Celofia	621	Clusia	1042
Buddleia	112	Celsia	1043	Clypeola	652
Bufonia	122	Celtis	1049	Cneorum	52
Bulbocodium	294	Cenchrus	909	Cnicus	826
Buonia	669	Centaurea	116	Cochlearia	647
Bunium	243	Centunculus	95	Coccolaria	2188
Buphralmum	903	Cephalanthus	457	Coffea	178
Bupleurum	236	Cerastium	969	Coix	978
Burmiania	237	Ceratocarpus	1020	Colchicum	341
Butomus	372	Ceratonion	998	Coldenia	125
Buxus	983	Ceratophyllum	208	Collinsonia	28
Bysshe	2148	Cercis	374	Columnnea	682
		Cerinth	136	Colurea	723
		Ceropegia	211	Comarum	508
		Cestrum	193	Commelina	40
		Cachrys	246	Conferua	1164
		Cherophyllum	218	Conium	245
		Chaetrops	1187	Connarus	671
		Chara	1116	Conocarpus	176
		Cheiranthus	661	Convallaria	314
		Chelidonium	505	Convolvulus	153
		Chelone	611	Conyza	861
		Chenopodium	218	Corchorus	529
		Cherleria	425	Cordia	190
		Chionanthus	8	Coreopsis	907
		Chironia	189	Coriandrum	256
		Chondrilla	706	Coriaria	1027
		Chrysanthemum	882		

Coris

INDEX GENERUM.

Coris	177	Datura	1037
Corypha	1187	Daucus	179
Corispermum	4	Delphinium	242
Cornucopia	54	Dalibarda	491
Cornus	117	Dentaria	530
Cornutia	628	Dianthera	653
Coronilla	742	Dianthus	27
Corrigiola	271	Diapensia	409
Cortula	144	Dictamnus	141
Corvulus	998	Digitalis	383
Corymbium	928	Dillenia	611
Costus	2	Diodia	535
Cotula	891	Dioscorea	104
Cotyledon	429	Diosma	1032
Cracca	752	Ditrea	198
Crambe	673	Diospyros	358
Craniolaria	618	Diptacus	1057
Crassula	282	Dodaria	97
Crassus	475	Dodecatheon	633
Crateva	444	Dolichos	144
Crepis	805	Doronicum	725
Crescentia	626	Dorstenia	885
Cressa	223	Draba	181
Crinum	291	Dracocephalum	642
Crithmum	246	Dracontium	594
Crocus	36	Drosera	967
Crotalaria	714	Dryas	281
Croton	1024	Drypis	501
Crucianella	108	Duranta	413
Cucubalus	414		637
Cucumis	1011		
Cucurbita	1010		
Cuminum	254		
Cupania	200		
Cupressus	1001		
Curcuma	2		
Cuscuta	124		
Cycas	1188		
Cyclamen	145		
Cymbaria	618		
Cynanchum	212		
Cynara	827		
Cynofurus	72		
Cynoglossum	134		
Cynometra	382		
Cynomorium	970		
Cyperus	74		
Cypripedium	911		
Cytisus	739		

E.

Ebenus	254
Echinophora	200
Echinops	1001
Echium	2
Elaeagnus	124
Elatine	1188
Elate	145
Eleocarpus	618
Elephantopus	212
Elveia	827
Elymus	72
Empetrum	134
Ephedra	382
Epidendrum	970
Epigea	74
Epilobium	911
Epimedium	739
Equisetum	
Eranthemum	
Erica	
Erigeron	
Eriurus	
Eriocaulon	

Eriocaulon	87
Eriophorum	52
Ervum	738
Eryngium	332
Erysimum	660
Erythrina	706
Erythronium	305
Eugenia	470
Evonymus	197
Eupatorium	836
Euphorbia	450
Euphrasia	604
Exacum	212

F.

Fagonia	386
Fagus	997
Ferula	246
Festuca	13
Fevilla	1013
Ficus	1059
Filago	927
Flagellaria	313
Fontinalis	1107
Fraxia	494
Frankenia	331
Fraxinus	1057
Fritillaria	303
Fuchsia	1191
Fucus	1158
Fumaria	699

G.

Galanthus	288
Galax	200
Galega	714
Galenia	319
Galeopsis	579
Galium	105
Garcinia	443
Garidella	435
Gaura	347
Gaultheria	395
Genista	709
Gentiana	227
Geranium	675
Gerardia	610
Gesneria	612
Geryllis	442
Geum	500
Gladiolus	36
Glaux	207
Glechoma	578
Gleditsia	1056

cli.

INDEX GENERUM.

Clinus 462
 Globularia 95
 Gloriosa 305
 Glycine 753
 Glycirrhize 741
 Gmelina 626
 Gnaphalium 850
 Gnidia 358
 Gomphrena 224
 Gostypium 793
 Gratiola 17
 Grevia 965
 Grillea 348
 Gronovia 202
 Gvajacum 381
 Guettarda 991
 Guilandina 381
 Gundelia 814
 Gypsophila 406

H.

Heimanthus 325
 Hemathoxylum 324
 Hælleria 625
 Hamamelis 124
 Hebenstretia 629
 Hedera 202
 Hedyotis 101
 Hedyfarum 745
 Helenium 886
 Helianthus 904
 Heliocarpus 448
 Helicteres 963
 Heliotropium 130
 Helonias 348
 Helleborus 557
 Hemerocallis 324
 Hemionitis 1077
 Heracleum 249
 Hermannia 673
 Hetnandia 981
 Herniaria 218
 Hesperis 663
 Heuchera 226
 Hibiscus 693
 Hieracium 792
 Hippocratea 1191
 Hippocrepis 744
 Hippomane 1191
 Hippophaë 3023
 Hippuris 4
 Mirrella 32
 Holcus 1017
 Holosteum 78

Hordeum 84
 Horminum 596
 Hottonia 585
 Hugonia 675
 Humulus 1028
 Hura 1008
 Hyacinthus 316
 Hydnum 1178
 Hydrangea 397
 Hydrocharis 1036
 Hydrocotyle 284
 Hydrophyllum 146
 Hymenæa 1192
 Myofcyamus 179
 Hyoseris 808
 Hypocoum 124
 Hypericum 783
 Hypnum 1122
 Hypocheris 810
 Hylopus 569

I.

Jambolifera 349
 Jastione 928
 Jasminum 7
 Jatropha 1006
 Iberis 648
 Ilex 125
 Iliecebrum 806
 Impatiens 937
 Imperatoria 259
 Indigofera 751
 Inula 881
 Ipomœa 159
 Iris 38
 Ixiatis 670
 Ilichæmum 1049
 Inardia 120
 Isoetes 1100
 Isopyrum 357
 Itea 199
 Iva 983
 Juglans 997
 Juncus 325
 Jungermannia 1131
 Juniperus 1039
 Justicia 388
 Julticia 15
 Ixia 36
 Ixora 110
 Kalmia 391

K.

Kampharis
 Kiggelaria
 Knauria
 Knoxia
 L.
 Lachnea
 Lactuca
 Lagœcia
 Legurus
 Lamium
 Lantana
 Lapsana
 Lalerpitium
 Lathræa
 Lathyrus
 Lavandula
 Lavatera
 Laurus
 Laufonia
 Lechea
 Ledum
 Lemna
 Leptnice
 Leonodon
 Leonurus
 Lepidium
 Leucadendron
 Leucojum
 Lichen
 Ligusticum
 Ligustrum
 Lilium
 Limodorum
 Limofella
 Linnaea
 Linum
 Lippia
 Liquidambar
 Liriodendrum
 Lithospermum
 Lobelia
 Lœdingia
 Lœelia
 Lolium
 Lonchitis
 Lonicea
 Loranthus
 Lotus
 Ludwigia
 Lunaria
 Lupinus
 Lychnis
 Lycium

INDEX GENERUM.

ycoperdon
ycopodium
ycopis
ycopus
ymachia
ythrum

M.

agnolia
alope
alpighia
alva
ammea
andragora
angifera
aranta
arcgravia
archanria
arrubium
arsilea
artynia
aricaria
atchiola
edeola
edicago
clampodium
clampyrum
clanthium
clastoma
clia
clianthus
cliffa
clitris
lochis
lochia
lochia
luncylon
lispermum
lntha
lnzelia
lnyanthes
lrcurialis
lrmhryanthemum
lpilus
lua
lchelia
lucos
leropus
lium
leria
lofa
lulus
lusops
luarta
labilis

1183 Mitchella
1100 Mitella
138 Mnium
21 Muehringia
146 Mollugo
440 Moluccella
Momordica
Monarda
Monotropa
535 Montia
692 Morina
425 Morinda
687 Morisona
512 Morus
281 Mucor
200 Muntingia
2 Musa
503 Mussenda
1137 Myagrum
582 Myosotis
1099 Myosurus
618 Myrica
890 Myriophyllum
1192 Myrsine
Myrtus

N.

339 Najas
389 Nana
384 Napea
639 Narcissus
592 Nardus
597 Nepenthes
674 Nepetha
35 Nerium
349 Neurada
310 Nicotiana
576 Nigella
516 Nyctanthes
145 Nymphaea
1035 Nyssa
Obolaria
Ochna
Ocyrum
Oenanthe
Oenothera
Olax
634 Oldenlandia
349 Olea
89 Onoclea
177 Ononis

O.

111 Onopordon
406 Ophioxylon
1109 Ophioglossum
359 Ophiorrhiza
89 Ophrys
587 Orchis
1009 Origanum
22 Ornithogalum
387 Ornithopus
87 Orobanche
28 Orobus
176 Orontium
503 Ortega
986 Orvala
1185 Oryza
509 Osbeckia
1043 Osmunda
277 Osteospermum
640 Ostrya
121 Othonna
284 Oviada
1024 Oxalis

P.

Paeonia
Panax
Pancratium
Panicum
Papaver
Parietaria
Paris
Parkinsonia
Parnassia
Parthenium
209 Passerina
441 Passiflora
180 Paltinaca
534 Paragonula
6 Pavetta
510 Paullinia
1038 Pedicularis
Peganum
Penaea
Pentapetes
Penthorum
518 Peplis
597 Periploca
254 Petiveria
346 Petrea
32 Peucedanum
119 Peziza
8 Phaca
1062 Phalaris
716 Phallus

827
1043
1062
150
945
939
588
306
743
632
728
324
560
578
333
345
1063
923
1022
924
637
433

Phar -

INDEX GENERUM.

Pharnaceum	272	Primula	143	Ruppiea	12
Phatium	1106	Prinos	330	Rufus	104
Phaseolus	723	Proserpinaca	88	Ruta	38
Phelandrium	255	Protea	94		
Philadelphus	470	Prunella	600		
Phillyrea	7	Prunus	472		
Phileum	59	Psidium	470		
Phionis	584	Pteralea	762	Saccharum	5
Phlox	151	Prelea	118	Sagina	18
Phoenix	1188	Pteris	1073	Sagittaria	99
Phryma	601	Pulmonaria	135	Salicornia	101
Phylla	195	Punica	472	Saxif	21
Phyllanthus	981	Pyrula	596	Salfola	12
Phyllis	232	Pyrus	479	Salvadora	2
Phyllis	182			Salvia	26
Phytocoma	170			Sambucus	17
Phytolacca	441			Samolus	44
Picris	792	Quercus	994	Samyda	50
Pilularia	1100	Queria	90	Sanguinaria	10
Pimpinella	163			Sanguisorba	21
Pinguicula	17			Sanicula	34
Pinus	1000			Sanctum	84
Piper	28	Rajania	994	Santolina	31
Pisonia	1020	Randia	1192	Sapindus	4
Pistacia	1025	Ranunculus	534	Saponaria	51
Pistia	963	Raphanus	659	Sarracena	2
Pisum	727	Rauwolfia	208	Sarothra	51
Plantago	112	Renalmia	286	Satureja	94
Platanus	958	Releda	448	Strygium	30
Plinia	916	Rhamnus	193	Saururus	21
Plukenetia	1192	Rheedia	1193	Sauvagesia	31
Plumbago	217	Rheum	371	Saxifraga	31
Plumeria	209	Rhexia	346	Scabiola	1
Poa	67	Rhinanthus	603	Scandix	31
Podophyllum	105	Rhizophora	443	Scheuchzeria	31
Poinciana	380	Rhodiola	1035	Schinus	31
Polemonium	160	Rhododendron	392	Schonus	6
Polyanthes	316	Rhus	265	Schwalbea	31
Polycnemum	35	Ribes	200	Scilla	31
Polygala	701	Riccia	1138	Scirpus	4
Polygonum	350	Richardia	330	Scleranthus	4
Polymnia	926	Ricinus	1007	Scolymus	1
Polypodium	1082	Rivinia	121	Scoparia	1
Polypremum	111	Robinia	723	Scorpiurus	2
Polytrichum	1109	Rocella	170	Scorzonera	2
Pomederia	288	Rondeletia	172	Scrophularia	1
Populus	1034	Rosa	491	Scutellaria	1
Porella	1106	Rosmarinus	23	Secale	1
Portulaca	445	Royena	397	Securidaca	1
Potamogeton	126	Rubia	109	Sedum	1
Potentilla	495	Rubus	492	Selago	1
Poterium	994	Rudbeckia	905	Selinum	1
Pothos	968	Ruellia	634	Sempervivum	1
Potium	601	Rumex	333	Senecio	1
Prenanthes	797	Rumphia	1193		

INDEX GENERUM

Serapias	949	Syrax	444	Tribulus	386
Seriphium	928	Subularia	442	Trichomanes	1097
Serrania	816	Suriana	284	Trichosanthes	1008
Sesamum	634	Svertia	226	Trichostema	598
Seseli	219	Symphytum	136	Tridax	900
Sherardia	102	Syringa	9	Trientalis	344
Sibbaldia	284			Trifolium	764
Sibthorpia	631			Triglochin	338
Sicyos	1013			Trigonella	776
Sida	683	Tabernaemontana	210	Trillium	319
Sideritis	574	Tagetes	887	Triopteris	418
Sideroxylum	192	Tamarindus	32	Triosteum	176
Sigesbeckia	900	Tamarix	270	Triticum	85
Silene	416	Tamus	1028	Triumfetta	444
Silphium	919	Tanacetum	843	Trollius	556
Sinapis	668	Tarhonanthus	842	Tropaeolum	345
Siphonanthus	109	Targionia	1136	Tulipa	305
Sison	252	Taxus	1040	Turnera	271
Silybrium	657	Telephium	271	Turritis	666
Silyrinchium	954	Tetracera	533	Tussilago	865
Sium	251	Terragonia	480	Typha	972
Sluanea	512	Terragonotheca	903		
Smilax	1028	Teucrium	562		
Smymnium	462	Thalia	1193		
Solanum	184	Thalictrum	545	Ulex	741
Soldanella	145	Thapsia	261	Ulmus	225
Solidago	878	Thea	515	Uva	1163
Sonchus	793	Theligonum	993	Uniola	71
Sophora	373	Theobroma	782	Urena	692
Sorbus	477	Theophrasta	149	Urtica	923
Sorganium	971	Thesium	207	Utricularia	18
Spartium	708	Thlaspi	645	Uvaria	536
Spergula	440	Thya	1002	Uvularia	304
Spermacoce	102	Thymbra	569		
Spiranthes	927	Thymus	590		
Phagnum	1106	Tiarella	405		
Pigelia	149	Tilia	514		
Pinacia	1027	Tillia	128	Vaccinium	349
Piraea	439	Tillandria	286	Valantia	1053
Plachnum	1108	Toluidifera	384	Valeriana	51
Pondias	371	Tomex	118	Vallisneria	1015
Pongia	1169	Tordylium	239	Vateria	515
Pachys	580	Torenia	619	Velexia	332
Phelina	840	Tormentilla	500	Velva	641
Phelia	217	Tournefortia	140	Veratrum	1044
Phyllaea	270	Tozzia	607	Verbascum	177
Picea	274	Trachelium	171	Verbena	18
Pellera	421	Tradescantia	288	Verbesina	901
Pellera	519	Tragacantha		Veronica	9
Perculia	1007	Tragia	980	Viburnum	267
Pervaria	698	Tragopogon	789	Vicia	734
Pea	78	Trapa	120	Vinca	209
Pebe	831	Tremella	1157	Viola	933
Parnotes	535	Trevis	1193	Viscum	1023
Pachno.	189	Trianthema	223	Vitex	638
				Viris	

INDEX GENERUM.

Viris	203	Xeranthemum	857	Zanonia	1028
Volkameria	637	Ximenia	1193	Zanthoxylum	270
		Xyris	42	Zea	971
				Zizania	958
Waltheria	673	Yucca	319	Ziziphora	21
				Zosteria	694
Xanthium	987	Zanichellia	969	Zygophyllum	331



INDEX SYNONYMORUM.

A	A.					
Abavo	1190	Acinos	21. 592	Agnanthus	64	
Abies	1001. 1002	Aconitifolia	505	Agrimonia	389. 44	
Abrotanoides	93. 845. 928	Aconitum	367. 504. 531.	Agrimonia similis	44	
Abrotaunum	842. 844. 845.		550. 557. 900.	Agrimonioides	44	
	846. 849. 927	Acorus	39	Agul	741	
Abrus	754	Acriviole	345	Agutizvepo-obi	1191	
Ablinthium	844. 845. 846.	Acrosticum	1074	Argyrocome	851	
	847. 848. 849. 898.	Actea	548	Ahouaj	208. 209	
	900. 988.	Aculeosa	170	Aira	1044	
Abulito affinis	963	Adaca-Manjen	927	Aizoon	431	
Abutilon	675. 685. 686.	Adamboe	160	Alaternoides	195. 196	
	688.	Adhatoda	15. 639		1025. 1042	
Acacia	380. 382. 517. 519.	Adiantum	1065. 1067.	Alaternus	19	
	523. 709. 722. 1057		1068. 1072. 1075.	Alcea	529. 530. 675. 684	
Acacia accedens	517		1078. 1080. 1081.		685. 686. 687. 688	
	allinis		1092. 1098. 1099		690. 691. 693. 694	
	similis		1112. 1122		695. 696. 697. 698	
Acacia	371	Admirabilis	177		783.	
Acalypha	980	Aegilops	77. 1045	Alcea affinis	68	
Acanthium	827	Aegoceras	675	Alchemilla	123. 207. 39	
Acanthoides	828	Aegopodium	256. 263		406	
Acaricoba	834	Adhatoda	635	Alectorolophus	602. 60	
Acarna	820. 823. 828. 829.	Aelchynomene	518		610	
	830	Aethiopsis	27	Alga	968. 1167. 116	
Acathia-valli	35	Afra	382		1164. 1167	
Acer	427. 428	Agarico fungus	1176	Algoides	94	
Aceri affinis	428		polyporus	1176	Alhenna	24
	cognata	Agaricum	1157	Alifanus	34	
Acetosa	336. 337. 338. 372	Agaricus	1176. 1180. 1182	Alifina	34	
Achyrantha	205	Agary	728	Alkekengi	181. 18	
Achyrophorus	810	Ageratum	630. 843. 853.		183.	
Acinaria	1160		897. 900	Alliaria	64	
Acini species	591	Agerato affinis	989	Allium	24	
Acinodendrum	390	Agihalid	1194	Alni effigie	476	

INDEX SYNONYMORUM.

Alnifolia	396	Anacardium	110	Apiofenodon	444
Alni fructu	176	Anagallidistrum	116	Apium 244. 246. 250. 253.	
Alnus	176. 193. 983	Anagallis 12. 116. 147.		260. 264 265.	
Aloe 282. 319. 323. 325.		148 157. 171		Apocynum 209. 210. 211.	
635		Anagyris	733	217	
Alopecuros	573	Ananas	285	Aquifolium	125
Alopecurus	167	Anandria	865	Aquilegia	557
Alpina	2	Anantali-maravara	952	Arabis	660
Alfina	993	Ananthocyclus 891. 892		Araca i puda	282
Alfinastrum	368	Anapodophyllum	505	Arachnoides	741
Alfina 13. 14. 88. 89.	116.	Anascovadi	814	Arachidua	730. 741
119. 120 123. 128.		Anblatum	606	Aracus	730
142. 169. 172. 207.		Anchusa 132. 133. 137		Araliastrum	1059
226. 272. 332. 344.		Anchusa degeneris facie		Arapabaca	149. 150
359 407. 414. 421.		137		Arbor egocerathos	675
422. 423. 424. 425.		Andropogon	78	æthiopica	121
438. 439. 440 441.		Andromeda	291	africana	512
463. 464. 631. 632		Androsamum 783. 784.		alni fructu	782
Alfina affinis	141	786. 787.		americana 208. 265.	
Alfines facie	993	Androface	141	626. 284. 370.	
Alfinoides	82. 123	Anemone	543	390. 533. 1190	
Althæa 673. 674. 675. 684.		Anemonospermus 922. 923		baccifera 371. 426	
685. 686. 687. 688.		Anerifolius	94	brasiliانا	471
690. 691. 693. 694.		Angelica 248. 256. 259.		cacavifera	782
695. 696		261. 263. 265. 273		camphorifera	369
Althæa similis	673	Angeli-maravara	953	crepitans	1008
Aly-alu	2060	Angiopteris	1062	cucurbitifera	626
Alypum	96	Anguina	1008	exotica	381
Alyfoides	651	Angurek Warua	9 2	ferens folia argen-	
Alyilon. 582. 642. 651.		Anguria	101.	rea	94
652		Anil	751	fraxini folio	384
Alyffum 574. 642. 656		Anisum	253. 264	Gvajaci	625
Anialago	29	Anora	192. 512. 537	Indica 385. 512.	
Amanja	1172	Anonis 716. 717. 718. 754.		523. 1017	
Amaracus	590	767. 775.		Judæ	374
Amarantho affinis	224	Anonyma 391. 395. 607		ligni sancti	382
Amaranthoides 205. 224.		Anonymos 102. 111. 120.		maderaspatana	361
225		200 207. 704		malabarina	118
Amaranthus 122. 204. 205		Anthanisophyllum	3	malabarensium	
206. 225		AntEuphonium	834	110	
Ambapaya	1035	Anthemis	891	mexicana	512
Amberbot	910	Anthora	538	negundo	938
Ambrosia	648. 846	Anthyllis 89. 122. 223.		platani folio	1036
Ambuyambo	960	331. 332. 563. 764		Poyfon- vood-Tree	
Amelanchier	8	Antirrhinum	603	266	
Amellus	864	Aparine 18. 105. 106. 108.		racemosa 390. 1193	
Ameri	751	1052		Salvifolia	1044
Amerhyffina	21	Aparines folio	120	sebifera	1004
Ammi 252. 253. 255		Apate	796	S. Thomæ	375
Ammoides	260	Apaiiba	512	Siliquosa 516. 519.	
Ampans	1187	Aphaca	739	625. 1192	
Ampeloprasium	297	Aphyllanthes 96. 929		stercoraria	31
Amygdalus	121	Aphyllon	633	trifolia	266
Amygdalis similis	782	Apialtrum	593	tritiss	6
Anacampferos 430. 431.		Apios	733. 713	venenata	1201

INDEX SYNONYMORUM.

Arbor virginiana 370. 371	Astragaloides 754. 759	Bellon-parfia 1108
vitz 1002	Astragalus 719. 728. 751.	Bem-curini 15
zagoe 1188	753. 754. 762.	Bennet 938
Zeylanica 8	Arthanasia 845	Bemfcherri 110
Arbuscula afra 1038	Attractylis 829. 830	Bem-Tamara 511
afriana 338	Atraphaxis 359	Bermudiana 36. 957
baccifera 117	Atroplex 4. 5. 219. 220. 353.	Beta 337
jamaicensis 390	359. 979.	Betonica 15. 413. 570. 578.
mariana 393	Avellana 999. 1005	620. 673
nerii foliis 860	Avellane indica 3 gener	Betonie folio 573
viscosa 118	1190.	Beella 29
Arbutus 395. 397	Avena 94	Bidens 814. 900. 901. 908.
Arcahu 1059	Aurantia 782	919
Archangelica 251	Aureliana 1058	Bifolium 246
Arcrotheca 923	Auricula 143. 144. 145.	Bihei 1043
Arenaria 422. 425. 440	179. 421.	Bilimbi 428
Aretia 141. 143	Azadirachte 385	Bistorja 360
Argemone 506. 508.	Azalea 392	Biri-maram-maravara 953
1196	Azedarach 184. 385	Blairia 19
Arguzia 141		Blattaria 178. 179. 446.
Argyracome 861	B.	621. 698
Argyrodendros 24	Baccharis 864	Blennochas 120
Aira 475	Baccifera 111. 182. 1037	Blinbingum 428
Aria Bepou 385	Badiaga 1171	Blitum 204. 230. 790
Aria-vela 672	Baducca 504	Boabab 1190
Arifarum 966. 967	Balam-pulli 54	Bobarrta 907
Armenicata 473. 474	Balaustia 472	Bolerus 1179
Armeria 410	Baleincanda-schularmandi 36	Bon 173
Armerius 409. 412. 421	Ballel 158	Bema 737
Aroeira 389	Ballote 582. 584.	Bonarota 11
Artemisia 344. 345. 928	Bandura 955	Bonduc 381
Arum 324. 967. 968	Balfamina 937. 938. 959.	Bontia 953
Aruncus 490	1009	Bonus Henricus 218
Arundo L. 54. 55. 325. 391.	Balsamica 897	Borsgo 133. 115
1050	Balsamum alpinum 393	Borrys 219. 220. 561
Afarina 612	Tolutanum 384	Borrtis 1186
Aclepias 218	Bandura 955	Brassica 159. 665. 666. 671.
Alcyon 785	Baobab 1190	965
Alcyrum 784. 786. 788	Barba Capre 490	Breytia 928
Aspalathus 708. 709. 722.	Caprina 1183	Briza 86
727.	hirci 789	Brunella 570. 600. 601
Asparagus 339	Bardana 816	Bryonia 140. 1012. 1028.
Aspergillus 1185. 1186	Baratas 154	1032
Asperula 103	Batti-Scherigeram 985	Bryonie similis 1033
Aphodelus 310. 311	Beetle-codi 29	Bryonioides 1013
Aplenium 1070	Behen album 414. 914.	Bryum 1108. 1109. 1110.
Asta freida 248	Beid-el-ollat 214	1111. 1112. 1113. 1114.
Aster 841. 859. 861. 864.	Belilla 177	Bubonium 883
865. 869. 881. 882. 883.	Belladonna 182	Bucanephyllum 510
884. 885. 886. 888. 900.	Bellidialtrum 886	Buccaferrea 128
902. 904. 926.	Bellis 96. 97. 658. 843. 876.	Buchnera 640
Asteri similis 864	886. 888. 889. 890. 892.	Buglossum 133. 134. 137.
Asterias 227	982	138. 139. 792.
Asteriscus 896. 903. 920		Bugula 561. 601
Alterocephalus 100		Bulbine 311
		Bul-

INDEX SYNONYMORUM

Bulbocestanum	343	Canna Indica	1193	Cataria	570. 573
Bulbocodium	36	Cannoides	1	Catiline-vegou	361
Bulbus	309	Canschi	1193	Cattu-gasturi	696
Bunium	253	Canabrica	158	Cattu-irpali	29
Bupariti	694	Capnos	701	Carulli-pola	190
Buphthalmum	548. 895.	Capo-molago	189	Caucalis 259. 240. 242. 243	
896. 902.		Capparis	385	Cavalam	1008
Buphthalmum simile	893	Capraria	90. 116	Cauda muris	284
Bupleuroides	252	Caprificus	1059	vulpis	61
Bupleurifolia	928	Caprifolium	173	Caulis-Rapum	667
Bupleuri similis	928	Caput Gallinaceum	751	Caapeba	1031. 1032
Burau	1008	Cara	1033	Caunga	1139
Bursa Pastoris	642. 643.	Caragana	722	Ceanorhus	820
647		Caraginata	286	Cedronella	594. 596
Bursa Pastoris similis	665	Carambu	119. 388	Cedrus	1001. 1039. 1040.
Buxus	196	Caranofi	922	Celastrus	195. 196. 197
Byssus	1141. 1157. 1186	Caraschulli	636	Cenchrameda	509. 782
		Caraveza	671	Centaurea	819
		Caraxeron	224	Centaurium	112. 190. 204.
		Cardamine	644	208. 230. 232. 272. 818.	
		Cardaminidum	345	826. 839. 910. 915. 918.	
		Cardiaca	524. 586. 587	Centaurii species	912
		Cardioperium	921	Cepa	293. 300. 301.
		Carduo-cirsium	817	Cepaea	431
		Carduus	285. 413. 639.	Cerafa	474. 475
		814. 815. 816. 817. 820.		Cerastis similis	473
		826. 827. 828. 829. 830.		Cerastis affinis	474
		831. 917. 919.		Cerastus	269. 425. 426. 473.
		Carduus-acanthus	939	474.	
		Carduncellus	831	Ceraria	374. 653. 706. 1192
		Carretti	381	Ceratocephalus	833. 908.
		Carim-curini	15	Ceratoideis	979
		Carimgola	288	Ceratonie affinis	503
		Carlina	826. 829. 1199	Cerefolium	257. 258. 259
		Carou-melli	193	Coreus	466. 467. 469
		Carpinus	118	Cerinth	136
		Carpobolus	1184	Cerri glans	996
		Carthamoides	831	Cervaria	171. 1194
		Cartnamus	829	Cerviana	272
		Carua	369	Cervicaria	167
		Carum	253	Cervispina	193
		Carvi	253. 261.	Ceterach	1080
		Carvifolia	261	Chaa	515
		Caryophyllata	500. 501	Chareciolum	257
		Caryophyllus	88. 191. 274.	Cherophyllo-similis	123
		294. 406. --- 413. 415.		Cherophyllum	257
		421. 422. 436. 438. 439.		Chalceos	815
		472. 827.		Chamaebuxus	704
		Caschon	383	cerastus	173. 174
		Castia	1022	cistus	152. 396. 302.
		Castia	369. 371	525. 527. 528.	
		Castida	598. 599. 600	678.	
		Castine	125	crista	379. 380
		Castanea	344. 994	daphne	112. 391.
		Castorea	637	394	
				Chame	

INDEX SYNONYMORUM.

<i>Chemadrifolia</i>	441	869. 886. 892. 895.	<i>Codda-pans</i>	1187	
drya 11. 13. 502.		896. 901. 902. 903.	<i>Codium</i>	1199	
564. 566. 603.		904. 905. 906. 907.	<i>Codi-avanacu</i>	981	
<i>filix</i>	1081	908. 909. 920. 923.	<i>Coilantha</i>	227	
<i>genista</i>	710	924. 927	<i>Coilophyllum</i>	510	
<i>jafme</i>	559	<i>Chrysis</i>	905	<i>Colchicum</i>	292. 294
<i>iris</i>	38	<i>Chrylocome</i> 841. 853. 858.	<i>Colletta-veetla</i>	636	
<i>larix</i>	711	859	<i>Colocasia</i>	965	
<i>lea</i> 35. 195. 981.		<i>Chrysolplenium</i>	631	<i>Colocynthis</i>	1008. 1011
1042		<i>Cicer</i> 719. 738. 752. 756.	<i>Colubrin liagnigenus</i>	189	
<i>leagnus</i>	1024	757	<i>Coluppa</i>	225	
<i>leon</i> 815. 820. 828.		<i>Ciceri affinis</i>	759	<i>Colutea</i> 373. 381. 742. 743.	
915. 1199		<i>Cichonium</i> 792. 805. 806.	751		
<i>linum</i>	281	807. 811. 900	<i>Coma</i> 840. 841. 843. 854		
<i>melum</i> 888. 889.		<i>Cicuta</i>	243. 256.	<i>Condonium</i>	513
891. 892. 891.		259	<i>Conna</i>	378	
894. 895. 896.		<i>Cicutaria</i> 243. 255. 256.	<i>Conocarpodendron</i>	93. 94.	
<i>melo affine</i>	893	258. 259	95		
<i>melpilus</i>	579	<i>Cimicifuga</i>	504	<i>Conserva</i>	1163
<i>moly</i>	302	<i>Cinara</i>	827. 828	<i>Confolida media</i>	561
<i>nerion</i>	348	<i>Cinaroides</i>	25	regalis	531
<i>orchis</i>	948	<i>Cineraria</i>	992	<i>Convolvulus</i>	160. 161. 162.
<i>peuce</i>	819	<i>Cinnamomum</i>	369. 370.	164. 1031	
<i>pitys</i> 356. 562. 595.		371	<i>Convolvulo-Tithymalus</i>		
<i>rhododendron</i>	150	<i>Cirium</i> 817. 818. 819.	1054		
<i>rhododendros</i>	150.	820. 821. 822. 823.	<i>Conyza</i> 836. 840. 841. 842.		
151		824. 825. 826	851. 860. 861. 863.		
<i>riphes</i>	1187	<i>Cistoides.</i>	864. 865. 870. 881.		
<i>rubus</i>	494	<i>Cistus</i> 150. 151. 198. 271.	882. 925		
<i>svce</i>	455	391. 392. 394	<i>Coyza affinis</i>	882. 883	
<i>Chamejasine</i>	105	<i>Citrullus</i>	1011	<i>Coral</i>	706. 707
<i>Chamelza</i>	357	<i>Clandellina</i>	606	<i>Corallina</i> 1154. 1166. 1167	
<i>Chamelco</i>	829	<i>Clatroides</i>	1179	<i>Corallodendron</i>	706
<i>Champsam</i>	536	<i>Clatroidaltrum</i>	1179	<i>Corallofungus</i>	1183
<i>Chanfchena-pou</i>	375	<i>Clematis</i> 189. 209. 210.	<i>Coralloides</i>	1141. 1145.	
<i>Charantia</i>	1009	366. 374. 427. 543.	1151. 1154. 1183. 1185.		
<i>Cheiranthus</i>	664. 665.	543. 623. 624. 625.	<i>Corchoro affinis</i>	529	
913		955. 956. 957. 958.	<i>Corchorus</i>	803	
<i>Chelidonia</i>	550	959. 960. 1031. 1043.	<i>Cordis indi folio</i>	366	
<i>Chelidonium</i>	505. 550	<i>Clematitris</i>	544. 836	<i>Corduba</i>	314
<i>Chenopodium</i>	35. 204	<i>Clinopodium</i> 21. 21. 23.	<i>Cordylone</i>	319	
<i>China</i>	1029. 1031	567. 568. 571. 591.	<i>Corindum</i>	367	
<i>Chlora</i>	232	601. 602.	<i>Coriotragematodendros</i>		
<i>Chondrilla</i>	790. 792. 793.	<i>Clitoria</i>	753	1025	
794. 796. 797. 798.		<i>Clitorius</i>	753	<i>Coris</i>	787
799. 806. 811. 812.		<i>Clymenum</i>	732	<i>Corispermum</i>	969
813. 835. 864. 865.		<i>Cneorum</i>	357	<i>Corni species</i>	1042
909		<i>Cnicus</i>	829. 830	<i>Cornus</i>	349. 371
<i>Chovanna-mandaru</i>	375	<i>Coa</i>	1191	<i>Corona</i>	303. 904. 905. 906
<i>Chritophoriana</i>	273. 274.	<i>Cocculus</i>	241	<i>Coronaria</i>	436
504. 548		<i>Cocconilea</i>	267	<i>Coronilla</i>	752
<i>Chritophoriana facie</i>	504	<i>Coccus</i>	1188	<i>Coronopus</i>	114. 115.
<i>Chrysantheinoides</i>	923	<i>Codagan</i>	234	<i>Corrua</i>	179. 226. 406. 1193
<i>Chrysanthemum</i>	95. 832.	<i>Codaga-pala</i>	209	<i>Cotulie species</i>	630
833. 835. 842. 860.		<i>Coda-pileva</i>	176	<i>Corymbifera</i>	898

Coryza

INDEX SYNONYMORUM.

Coryza	831	Cynosurus	71	Draconi affinis	319
Costus	845	Cynoxylon	1058	Dracocephalo affinis	594
Corinus	267	Cyperella	43	Dracontium	764
Coronea	480	Cyperoides	972. 974. 975.	Dracunculus	849. 898.
Cortum	597	976. 977. 978. 979.		907. 964	965. 968
Corula	892	Cyperus	42. 44. 49. 50. 51.	Drakena	121
Coryledon	234. 282. 283.	52. 121.		Dryopteris	1081. 1089
399. 431		Cyporel	1190	Drupis	414
Covalum	444	Cypripedium	948. 951	Dugglassia	67
Craeca	735. 738	Cyrti apnos	701	E.	
Crambe	640. 670	Cyrti facie	641		
Craffula	129	Cyrtus	373. 709. 710. 713.		
Cratægus	479	720. 764. 778			
Crepis	792. 798			Ebenus	193
Crista Galli	603			Ebulus	369
Pavonis	380			Ecbulium	15
				Echino-Melocactus	466
				Echinophora	241. 242.
Crithmum	239. 883. 884.	Dactylis	55. 57.		
Crocodylodes	829. 830	Dalea	764	Echinophora affinis	816
Crocodylium	815. 914.	Damafonium	343	Echinus	276
Croton	980. 1043	Dantia	120	Echmi facie	133
Cruciata	149. 1051. 1052	Daphnoides	356	Echioides	132. 138
Cucubalus	418. 419. 437	Darea	1097. 1098. 1099	Echium	131
Cucullaria	699	Dalystephana	228	Edera	202
Cucumis	35. 1009. 1010.	Dauci tertium genus	1194	Ela Calli	452
1013.		Daucus	243. 244. 245. 252.	Elajocarpus	575
Cucurbita	1008	253. 260. 264. 1194.		Elatine	613
Cucurbitifera	444	Dens Canis	305	Eleschnus	121
Cudupariti	693	Leonis	798. 799. 808.	Elengi	349
Culiramar	993	809. 885.		Elephas	603
Cuminum	124. 103	Dentaria	606. 607	Electari	1
Cunila	574. 575	Deutarie-facie	146	Elleborus	558
Cunonia	37	affinis	633	Elleborus	558. 1044
Cupamen	1003	Dentellaria	104. 151	Elichryso-affinis	205. 842.
Cupi	177	Defcurea	659	861. 989	
Cupressopinus	199	Diapensia	235	Elichrysum	840. 842.
Curuni	366	Diconangia	179	843. 844. 850. 851.	
Cururu-ape	366	Dictamnium	589	852. 853. 854. 855.	
Curutu-pala	211	Dictamnus	584. 589	857. 858. 859.	
Cuscuta	35. 272	Dierville	175	Ellicrylon	844
Cyanus	93. 909. 910. 911.	Digitalis	610. 611. 612.	Elicarmos	307
912. 913. 916. 919		634.		Eluteria	1042
Cyanoides	918	Dimorphotheca	922	Elvela	1171. 1180. 1181
Cythoides	1180	Diuretica	884	Flymus	84. 752
Cyclaminus	145	Dodonæa	118. 125	Embolus	1185
Cydonia	144	Doria	871	Endivia	96. 813
Cymbalaria	612	Doronici species	885	Entada	518
Cynanchum	211. 212	Doronicum	880. 881. 884.	Ephemerum	9. 41. 42.
Cynocrambe	993. 1035	906		282. 342	
Cynoglossa	134	Dortmanna	989	Epidendron	953
Cynoglossoides	137. 138	Dorycnium	158. 712. 720.	Epipachis	945. 949
Cynoglossum	131. 136.	742. 762. 764. 776.		Epipogium	945
118.		Draba	645. 649. 660. 664.	Epithyrium	124
Cynomorium	382. 383	931.		Equiletrum	4. 368. 992.
Cynoforchis	940. 941	Draco	849	1156. 1157	

Еран-

INDEX SYNONYMORUM.

Erangelica	288	Falsel - Taril	452	Frustrax cinereus	978
Erefia	149	Ferrum equinum	744	coronarius	470
Erica 112. 198. 199. 393.		Ferula 246. 253. 254		foliis oblongis	394
560. 651. 1022.		Fetulego	247	ferratis	394
Ericaformis	199	Festuca 76. 77. 78. 79. 82.		Indiæ orientalis	834
Ericoila	229	1046. 1051.		indicus	639. 963.
Ericu	214. 274	Ficoidea	488	pavoninus	380
Erigeron	808. 883	Ficoides 466. 467. 468.		peregrinus	225
Erigerum	867	481 - 488		peruvianus	29
Erinacea	720	Ficus 468. 469. 481. 483.		scandens	203
Erinaceus	1178	488. 1036. 1043		spicatus	489
Erini genus	169	Filago	856	spinofus	110
Erioccephalus	354	Filicifolia	981	viminibus lentis	196
Eriophorum	808	Filicula 1068. 1071. 1081.		virginianus	118
Eriothrea	230	1089. 1091. 1096. 1098		vilcosus	536
Eruca 657. 658. 659. 660.		Filipendula 490. 608. 610		Fruciculus capsularis	981.
665. 667. 668. 670.		Filix 1062. 1063. 1066. --		982.	
Eruago	450. 670	1078. 1082. 1083.		convolvulaceus	
Erva	224	1085. 187. -- 1094.		1029	
Ervilia	727	1096. 1098. 1099		foliis rufci	339
Ervum	373	Filum	1162	sinensis	373
Eryngium 252. 814. 830.		Fistularia	607	Fruclus	1190
831.		Flammula 543. 544. 548		Fucoides	1161
Erysimum 364. 657. 658.		Fles adonis	547	Fucus 128. 1147. 1154.	
659. 661		africanus	887	1163. 1164. 1167.	
Erythimo similis	666	clitoridis	753	Fumaria	362
Esula 213. 456. 461. 462.		cuculi	656	Fungi esculenti species	
Eupatoria 818. 831. 837.		constantinopolitanus		1179	
839. 862.		436		Fungilli	1147
Eupatorio affinis	819	moscatus	696	Fungoides 1180. 1181. 1183	
Eupatorioides 831. 852.		pavonis	380	Fungus 970. 1141. 1151.	
853.		pallionis 956. 957.		1162. 1171. 1175. 1177.	
Eupatorio-phalacron 902.		958. 959. 960.		180. 1182. 1184.	
920.		folis	905. 906	Fungorum perniciosum	
Eupatorium 448. 832. 839.		Fluvialis	1015	genus 1172. 1174.	
860. 862. 909. 921.		Foeniculum 246. 260. 263.		Fu-Ran	954
Euphorbium 450. 451. 452.		Foenum Burgundicum 779			
Euphrasia 15. 16. 602. 603.		Græcum 718. 766.			
606. 607. 630.		777. 778.			
Euphrasia affinis	16	Fontinalis	128	Gaidarothymo	582
Evonymoides	196	Fragaria	490. 500	Gale	1024. 1025
Evonymo affinis 197. 430.		Fragaria affinis	284	Galega	714
515. 1037.		Franca	331. 332	Galega affinis	722
Evonymo similis	172	Frangula	193. 269	Galeopsis 102. 579. 580.	
Evonymus 195. 196. 392.		Fraxinella	333	586	
490.		Fructus	968	Galium	1052
Eyrythalia	231	canne	325	Gallitrichum	25. 596
		orbicularis	189	Gallium 104. 105. 106.	
		Frumentum ladicum 171		107	
Faba	578. 737	972. 1047.		Gealter	1184
Fabago	385. 386	Frutex æthiopicus 94. 196.		Gelfeminum	623. 625
Fagopyrum 363. 364. 365		269. 707. 708. 853.		Genista 375. 707. 712. 715.	
Fagotritica similis	364	1038. 1042.		741. 745. 746.	
Fagotriticum	364. 365	africanus 354. 1058		Genista spartium 708. 712.	
Fagus	533. 1163	Recoitar 111. 365.		720. 745. 763.	

G.

Genista

INDEX SYNONYMORUM.

<i>Geniste affinis</i> 762	<i>Gramen cryophyllum</i> 975	<i>Gramen panicula pendula</i> aurea 72
<i>Gentifella</i> 710. 711.	<i>creticum</i> 84. 1051	<i>paniculatum</i> 55. 61.
<i>Gentiana</i> 190. 226. 273.	<i>cyperoides</i> 43. 44.	62. 64. 66. 68. 69.
409. 633.	45. 46. 47. 50.	74. 75. 1048.
<i>Gentianella</i> 229. 231. 635.	51. 52. 58. 972.	<i>paniculis atrifimis</i>
<i>Gentianoides</i> 231	979	68. 71
<i>Gerbera</i> 888	<i>Dactyloides</i> 73	<i>parnassi</i> 273. 316
<i>Gesnera</i> 618	<i>Dactylon</i> 57. 58. 73.	<i>parvum</i> 66. 71
<i>Gesneria</i> 622	73. 972. 1045.	<i>phalaroides</i> 60. 66.
<i>Geum</i> 401. 402	1246. 1947.	68. 76
<i>Ghandiroba</i> 1014	<i>echinatum</i> 1050	<i>piperium</i> 1100
<i>Gingidium</i> 239. 242	<i>Eranthemum</i> 71	<i>pratense</i> 28. 52. 63.
<i>Gladiolo æthiopico simili</i>	<i>eriphorum</i> 76	64. 67. 68. 72. 74.
<i>lis</i> 37	<i>festuaceum</i> 74. 75	75. 1048.
<i>lacustri accedens</i> 43	<i>floridum</i> 423	<i>pusillum</i> 50
<i>Gladiolus</i> 37. 39. 288. 372.	<i>fluviatile</i> 44	<i>sciurum</i> 73
929	<i>foliis junceis</i> 85. 74.	<i>secalinum</i> 83. 84.
<i>Glacium</i> 5106	<i>geniculatum</i> 55	1055
<i>Glaucoides</i> 332	<i>glumis variis</i> 72	<i>segetum</i> 61. 65
<i>Glaux</i> 120. 332. 759. 760	<i>hirtum</i> 329. 330	<i>serotinum</i> 62
1014	<i>holosteum</i> 328	<i>sparreum</i> 28. 53. 78.
<i>Globularia</i> 94	<i>hordeaceum</i> 84. 85.	82. 1162
<i>Globularia affinis</i> 87	<i>Indicum</i> 54	<i>spica Brizæ</i> 78
<i>Glycyrrhiza</i> 373. 758	<i>ischæmum</i> 72	<i>echinata</i> 1049
<i>Graphalium</i> 840. 927	<i>junceum</i> 43. 44. 48.	<i>triticea</i> 87
<i>Gnaphalodes</i> 852	50. 87. 115. 327	<i>spicatum</i> 42. 71. 78
<i>Gossypium</i> 511. 512	328. 338.	81. 82. 84. 85.
<i>Gramen</i> 63. 422	<i>juncoides</i> 328	339. 1051
<i>Gramen aculeatum</i> 42	<i>lanatum</i> 1048	<i>supinum</i> 62
<i>alopecuroides</i> 56. 71	<i>legitimum</i> 58	<i>sylvaticum</i> 61. 69.
81	<i>locustis echinatis</i>	77.
<i>alopecuro simile</i> 60	1050	<i>tenuæ</i> 74
<i>alopecurum</i> 61. 72	<i>loliaceum</i> 53. 69. 80.	<i>tomentosum</i> 52. 53
<i>alpinum</i> 65. 67. 74	83. 86.	<i>tremulum</i> 70
975	<i>Maris Borussorum</i>	<i>tribuloides</i> 1050
<i>americanum</i> 1050	1048	<i>triticeum</i> 78
<i>aquaticum</i> 55. 60.	<i>maritimum</i> 60. 71.	<i>typhinum</i> 60
64. 67. 75	1050	<i>typhoides</i> 55. 59. 60
<i>aromaticum</i> 1046	<i>niliaceum</i> 59	<i>vernum</i> 70
<i>arundinaceum</i> 52.	<i>minum</i> 63. 66. 68.	<i>virginianum</i> 59
55. 54. 82. 1048	1050	<i>xerampelinum</i> 70
<i>avense</i> 70. 72	<i>montanum</i> 62. 66.	<i>Graminitolia</i> 642. 1100
<i>avenaceum</i> 61. 62.	74. 75. 1050	<i>Grana paradisi</i> 2
64. 65. 66. 67.	<i>neurorum</i> 69. 74	<i>Gramadilla</i> 955. 956. 957.
75. 76. 77. 79.	<i>myloicophorum</i> 71	958. 959.
80. 82. 560. 1045	<i>myuros</i> 54	<i>Gratiola</i> 634
<i>Barcinonense</i> 73	<i>neurorolium</i> 65. 328.	<i>Gratiolæ affinis</i> 628
<i>bermudense</i> 57	329. 974.	<i>Grossularia</i> 200. 201. 390
<i>bromoides</i> 76. 78	<i>nodosum</i> 79	<i>Guajacana</i> 1058
<i>bulonium</i> 328	<i>orientale</i> 54. 64. 84.	<i>Guajava</i> 470
<i>bulbosum</i> 993	1050	<i>Guanabanus</i> 537. 1190
<i>Caninum</i> 62. 64. 83.	<i>palustre</i> 55. 67	<i>Guazuma</i> 530. 782
86. 1049.	<i>panicum</i> 56. 57. 73	<i>Guidonia</i> 443
<i>capillatum</i> 61		<i>Gislandinoides</i> 382

H. Ha-

INDEX SYNONYMORUM.

H.		809. 810. 811. 812.	Icaeo	513
Hacub	814	Hieracloe	1048	Ilex
Halicacabum	183. 367	Hinapariti	694	994- 995- 1191
Halimus	1053	Hippolapathum	366	Illecebra
Hamdamanias	514	Hippophæstum	917	Illecebrum
Heiera 266. 340. 578. 612.		Hippofelinum	262	Ily
1041		Hirculus	403	Imperatoria
Hedera species	619	Holochænus	49	Indigophora
Hederula	970	Holoiteo-affinis	284	Indiamie
Hediunda	191	Holoiteum	88. 114. 115.	1034
Hedypnois	798. 809	Hordeum	84. 86	Inrybum
Hedyisrum	713. 743	Horminum	23 - 27. 571.	Intybus
Heliteria	704	573. 603		Jolithus
Helenium	831. 905	Hummatu	179	Jon - Draba
Helianthemoides	271	Hyacintho affinis	292. 325	chlespi
Helianthemum	585-529	Hyacinthus	292. 308. 309.	652. 653
Helianthus	904	316. 319.		654
Helichrysoides	831	Hydroceratophyllum	992	Ina C. Balari
Heliotropii flore	141	Hydropiper	361. 367	Irio
Heliotropium	140. 578.	Hyoferis	811	658. 659
1004		Hypericonides	787	Irion
Helix	202	Hypericum	198	Lripa
Helleboraster	558	Hypocittis	442	Ischemum
Helleborine	949. 950. 951	Hypoglossum	1041	Isophyllum
Helleborus	235. 548. 556.	Hypophyllocarpodendron	93.	Isora Murri
557. 558. 1004.		93.		963
Helminthotheca	793	Hypoxis	306	Irticanni
Helxine	363. 364. 1052	Hypoxyton	1182	Juglandi affinis
Helichrysom	842. 851. 853.	Hyssopifolia	447	Jujuba
857		Hyssopus	569. 595. 603	194. 195
Hemeroallis	291. 303.	Hyssopum	569	Jujube
Hemionitis	1075. 1078.			195
1088. 1094				388. 389
Hepatica	273. 538. 1132.			Juncaria
1136-1139				Juncellus
Hepatorium	832. 909			48. 49
Herba britannica	336			Junco affinis
crinium	1200			Juncoides
gerardi	265			326. 327. 329.
indica	937			330
maxima	905			Juncoidi adfinis
mimosa	434			Juncus
non nominata	162			43. 44. 47. -- 50. 52
S. Jacobi	870			53. 54. 328. 372. 1046.
stella	115			1068
viva	434			Jungermannia
Hermidium	947			1114
Hermadactylus	40			1007
Herniaria	35. 123. 206. 207			
Hesperis	660. 618. 662			
Hieracioides	806. 807			
Hieracium	790. 792. 793.			
792. 806. 807. 808.				

Kazon

INDEX SYNONYMORUM.- 2

Katou indel	1189	Laurus 267. 268. 304. 536.	Limnædorum	944	
Katou - Kadali	391	1025. 1037. 1041	Limon	782	
Katou Thobam	991	Lavandula	19	274. -- 277.	
Katfil - kelengu	1033	Lavendula	572	510	
Katjiletti - pullu	42	Ledon 523. 524. 525. 529	Linagrostis	53	
Katu - kara - valli	1027	Ledum 392. 523. 524	Linaria 18. 207. 221. 278.		
Katu - katfil	1033	Lemna	1099	559. 613. -- 617. 841	
Katu - kurka	573	Lens 738. 970 1099. 1100	Lingua Cervina 1067. 1069.		
Katu - pithegamulla	6	Lentibularia 18. 970	1070. 1073. 1074.		
Katu - tagera	751	Lenticula 281. 963. 969.	1079. 1080. 1082. --		
Katu - thica	95	970. 971. 1099. 1100	1084		
Katu - kelengu	155	Lentiscus 388. 389. 1026	Linkia	1157. 1158	
Kauki	349	Leonrodoides	808	Linocarpum	281
Kavara	622	Leonorodon	806	Linofyris	841
Kempheria	19	Leontopetaloides	313	Linum 111. 148. 1163	
Kerstophyron	1170	Leontopetal affinis	312	Lithonchlaspi	646
Ketmia 673. 674. 695. 696.	697	Leontopetalon	312	Lithospermum 131. 134.	
		Leontopodium 114. 855	279 559. 972		
Kirganelli	982	Leonurus 580. 586. 587	Lobus	952	
Kleinia	834. 835	Lepidium 151. 648. 649	Locusta	34	
Knavei	406	Lepicarpodendron 91. 92	Lolium	435	
Kobbe	267	Leptostachya	601	Lonchitis 1063. 1064. 1067.	
Kodda - pail	963	Leucadendros 93. 94	1069 1071. 1074. 1076.		
Kolongging	952	Leucanthemum 888. 889.	1077. 1080. -- 1082.		
Kondampallu	937	896	1087		
Konni	753	Leucas	571	Lonicera 111. 176. 331.	
		Leucojum 288. 642. 643.	625 626	Lori similis	1058
		648. 651. 652. 661.		Lori arboris folio	509
		662. 663. 664. 665		Lorus 511. 712. 719. 740.	
		Leucoxylos	624	764. 765. 771. 776.	
		Levisanus	199	778. 1043. 1044. 1057	
		Leviticum	250	Loro affinis 719. 754. 778.	
		Libanotis 244. 246. 247.	376		
		248. 260		Ludvigia	120. 388
		Lichen 1135. -- 1140. 1157		Lunaria 336. 651. 652. 778.	
		Lichenastrum 1114. 1131.		779. 1064. 1065. 1137	
		-- 1136. 1139		Lupinaster	766
		Lichenoides 1140. 1142. -		Lupulinom	772
		- 1151. 1157. 1158		Lupulo simile	985
		Ligno brasiliiano simile 382		Lupulus 437. 1028. 1054	
		Lignum campechianum		Luteola 448. 449. 1037	
		384		Lutum	448
		moluccence 1004		Lychnidea 151. 153. 630	
		peregrinum 381		Lychnis 88. 151. 331. 332.	
		Ligustroides	637	408. 409. 413. -- 421.	
		Ligustrum 8. 9. 349. 637.		425. 436. 438. 439. 853	
		1043		Lychni - (cabiofa	101
		Lilio - alphodelus 292. 324		Lychnoides	152. 424
		Narcissus 291. -- 294		Lycio affinis	1194
		Lilium 292. 294. 303. 305.		Lycium 197. 331. 626. 636.	
		324. 339		1192	
		Convallium 315.		Lycopersicon	186
		316		Lycopodioides 1100. 1102.	
		Limeum	550	-- 1106	
		Limnia	204		

L.

Laburnum	739	Labiata	739	Labiata	739
Lactryma Jobii	792	Labiata	739	Labiata	739
Lactuca	795. 797	Labiata	739	Labiata	739
Ladanum	579	Labiata	739	Labiata	739
Lagopodium	719	Labiata	739	Labiata	739
Lagopogon	769	Labiata	739	Labiata	739
Lagopogus 767. 768. 769		Labiata	739	Labiata	739
Lagurus 1045. 1046		Labiata	739	Labiata	739
Lamium 578. 580. 597. 598		Labiata	739	Labiata	739
601. 620		Labiata	739	Labiata	739
Lampfana	811	Labiata	739	Labiata	739
Lantana	268	Labiata	739	Labiata	739
Lapathum 218. 333. -- 337.		Labiata	739	Labiata	739
359. 372. 1027		Labiata	739	Labiata	739
Lappa	816. 987.	Labiata	739	Labiata	739
Lappula	240. 389. 444	Labiata	739	Labiata	739
Laplana	803. 810	Labiata	739	Labiata	739
Larix	1000. 1001	Labiata	739	Labiata	739
Laferpitium	247	Labiata	739	Labiata	739
Larhyroides	728	Labiata	739	Labiata	739
Larhyrus	457. 728	Labiata	739	Labiata	739
Laurentia	931	Labiata	739	Labiata	739
Laureola	191. 356. 357	Labiata	739	Labiata	739
Laurifolia	444	Labiata	739	Labiata	739
Lauro affinis	389	Labiata	739	Labiata	739
Lauro - cerasus	473. 474	Labiata	739	Labiata	739

10()0(

Ly-

INDEX SYNONYMORUM.

Lycopsis	139	Melanium	935	Morus Gallinæ	579
Lyfimachia 119. 151. 189.		Melampyro cognata	636	Ranae	1036
223. 346. 347. 348.		Melampyrum	556. 629	Morungu	381
388. 446. 447. 594.		Melandrium	437	Morus norvegica	494
598. 624		Melanomphale	307	Moscatella	367
Lyfimachia affinis 119.		Melanthium	584	Moul-elavou	511
628		Meleagris	304	Mouricou	706
		Melilobus	1056	Mucedo	185
		Melilorus 765. 766. 776.		Mu-ke-tengu	1033
		777.		Muiva	390
		Melissa 568 587. 594-597.		Munda-valli	161
		601.		Mundubiguacu	1006
		Melo	1011. 1012	Muriguti	103
		cactus	466. 468	Muscari	317
		Melochia	673. 675	Musco-fungus 1143. 1144	
		Melopepo	1010	1146 - 1149. 1152 -	
		Memecylum	395	1154 1156. 1157.	
		Mendon	305	Muscoides 1110. 1134	
		Mentha 570. 571. 578. 588.		Muscus adiant. aurei facie	
		596. 845.		1112.	
		Menthallrum	576	alpinus 415. 1119.	
		Mercurialis	989 1003	1120.	
		Mespilus 268. 426. 476.		amarus	1146
		177. 509.		apocarpus	1115
		Messerichmidia	141	aquaticus 1107 1164	
		Methonica	305	165. 1171	
		Meum 245. 255. 259.		arboreus 1129 1130.	
		Michelia	626	1144. 1154. 1156.	
		Milium 59. 1017		aureus 1100. 1108	
		Millefolio adfinis	629	bisfolius	1131
		Millefolium 18. 144. 146.		capillaceus 1108 -	
		556. 854. 897. 899.		1110. 1112. 1115.	
		900. 992. 998 1070		1119. 1121. 1155.	
		Mimola	713	capillaris 1116. 1117.	
		Mirella	401. 406	1121. 1122	
		Mitra	150	ceranoides	1154
		Mitreola	150	clavatus 1103. 1104	
		Mnium	1138	coralloides	1153
		Modjra-canni	675	corallina +	1153
		Modira-camirani	189	corniculatus	1153
		Moringia	422	coronatus 1108. 1113	
		Moldavica	595. 596	1116. 1119.	
		Molle	389	cristam castrisem	
		Mollugo	90 107	representans 1126	
		Molucca	187	cupressiformis	1154
		Moly 295. 296. 298. 300.		dendroides	1128
		301		denticulatus 1104 -	
		Mombin	371	1106 1114. 1124	
		Monilifera	924	denticulato similis	
		Monorchis	947. 948	1128	
		Monospermalthæa	673	erectus 1102. 1103.	
		Monrana	392	1109. 1117 1118.	
		Morinda	76	silicinus 1103. 1123.	
		Moringha	381	1125.	
		Moro-ficus	1059	fistulosus	1132
				Muscus	

M.

Macanilla	1191
Mahaleb	474
Mail-anfschi	349
Majorana	588. 590
Mela 186. 428. 474. 479	
Malacodendron	698
Malacca-schambu	470
Malago-codi	88
Malus 389. 469. 472. 474.	
479. 480. 782. 783	
1190. 1191	
Malva 684. -- 687 690-692	
Malvaviscus	695
Malvinda	684. 692
Mallam roddali	1044
Mammay	512
Mammei	513
Mancanilla	1191
Mandragoras	181
Manfiadi	384
Manelli	712
Manga	200
Manghas	208
Mangles	443
Mangostana	444
Mangostans	444
Manja-kua	2
Manja-pumeram	6
Manihot	1007
Mao	200
Mareta-mala-maravara	
1067	
Marquias	956
Marrubialtrum	584
Marrubium 21. 580. 582.	
584. 586	
Marfilea 1100. 1135. 1136	
Martynia	618
Marum 27. 564. 567. 590.	
Matricaria 889. 893. 894.	
897. 899.	
Maurocchia	269
Meadia	144
Medica 778. 779. 780. 781	
Medicago 766. 777. 778.	

INDEX SYNONYMORUM.

Muscus foliis scurellatis nit	Mutallina	255	Nigellastrum	425
folio lato 1113	Myagrum	661	Nirvala	444
frutescens 1100	Myofotus	437-439	Niota-niodem valli	211
helveticus 1119	Myofurus	556	Nir-pulli	42751
humilis 1115	Myrrhis 245-251. 256. 257.		Niruri	982
Islandicus 1145	259		Noelitali	1027
juniperifolius 1109	Myrica	1038	Noli me tangere	938
lactuca similis 1163	Myrobalanus	513-982	Nostoc paracelsi	1157
Lichendides 1132	Myrobatindum	626. 627	Nuciperfica	479
lycopodioides 1106	Myrtifolia	196	Nuciprunifera	367
marinus 1161. 1163	Myrtillus	350	Nucula	243
1167	Myrtoides	472. 1024	Nummularia	148. 631
maximus 1100. 1103	Myrto-Cistus	783	Nuxeni-kelangu	1032
montanus 1098	Myrtus	471	Nux avellana	1006
1127. 1128	Myxa	1193	americana	167
minimus 1116 1121	Myxstax	675	juglans	997
1180			malabarica	1007
muralis 1111 1134			vesicaria	981
myofuroides 1124			vomica	189
norvegicus 1108			zeylanica	1008
palustis 1106 1110			Nyctage	177
1113. 1120 1134			Nymphæa	145 1036
parvus 1110			Nymphæa affinis	511
pendulus 1121				
pennatus 1122 1126				
polytrichoides 1112				
1113 1118 1122				
pulmonarius 1145				
pyxidatus 1151				
pyxoides 1151				
zsmofus 1109. 1124				
1155. 1156				
repens 1105. 1127.				
1130				
roseus 1113				
rupestris 1101				
saxatilis 1118 1129				
squamosus 1105.				
1107. 1115 1121.				
1127-1130.				
stellaris 1110. 1123				
1119. 1122.				
saxiformis 1123				
terrestris 1101.				
1102-1104 1106.				
1108. 1120 1123-				
1130. 1153.				
trichoides 1107.				
1110. 1111. 1116.				
1118. 1121. 1130.				
trichomanoides				
1123				
tridactylites 1167				
vulgaris 1117 1118				
1131				

N.

Naga-mu-valli	374
Naghas	515
Nalagu	441
Napellus	532
Napo brassica	667
Napus	666
Narcissus	290-294
Nardo celtica adfinis	33
Nardo celtica similis	32
Nardus	32 33. 65. 835
Narinam-poulli	696
Narthesium	311
Narum-panel	536
Narukila	288
Nasturtium	353. 641. 643.
	644 645 654. 655. 656.
	657. 659 1059
Nari-schambu	470
Nattisiam	341
Natrix	718
Nedel-ambel	145
Nelam mari	747
Nelam-parenda	937
Nelensichena	966
Neli-pouli	428
Nelipu	18
Neli-tali	714
Nella-randale-Corti	716
Nelliha	982
Neotia	945. 948
Nepera	169
Nerio affinis	109. 624
Nerium	210. 211
Nhandiroba	1013
Nidus avis	945
Nigella	425

O.

Obeliscotheca	907
Ochrus	727
Ocinum	597
Ocymastrum	580
Ocymoides	417. 419
Ocymophyllum	120
Ocymum	364
Odollam	208
Odontires	436
Oel-ambel	535
Oenarthe	152. 253. 260.
	545
Oepata	111
Olea	121. 185. 638.
Oliganthomum	289
Olus calappoides	1188
Omphalodes	135
Onagra	246. 247 386 516
Onobrychis	168. 207. 746.
	747-748. 749. 750. 751.
	760
Ononis	754
Ophioxylon	112
Ophris	946. 947
Opiantha	230
Opulus	268
Opunria	468. 469
Orchiastrum	946

Orchis

)o()o(2

INDEX SYNONYMORUM.

Orchis 325. 944. 945. 947. 948. 949. 950. 951. 953	Panover-Tsierana 121	Perficaria 360. 383
Orchidi affinis 910	Papaver 414. 506. 508. 509	Perfici officulo fructus 470
Orega 589	Papaya 1036	Personata 816
Oreoselinum 244	Papaya-maram 1036	Pes aserinus 219
Origanum 588	Papia 578	Petalites 866
Origano cognata 22. 588	Papyrus 47	Petilium 303
Orleans 512	Parietaria 121. 605. 985	Petræa 533
Ornithogalo affinis 204	Parietariae affinis 993	Petroselinum 252. 253
Ornithogalum 339	Paris 340	Phaca 761
Ornithopodium 743. 746	Pariti 694	Phagus 997
Ornithopodii affinis 744. 759.	Paronychia 205. 206. 207. 404. 1081	Phælypæa 606
Orobanche 387. 388. 606. 607. 945.	Paronychiae facie 105	Phalangium 310. 312. 316 324.
Orobis 738. 752	Parqui 191	Phaeoloides 753
Oryza 55	Parthenium 899. 989	Phaeolo affinis 741
Osinurda 1063	Partheniastrum 988	Phaeolus 516. 675. 725 - 727. 739. 753. 754.
Ostrya 998	Palserina 279. 559	Philyrea 296
Osyris 221. 616. 841	Pastinaca 239. 242. 243.	Phlomis 173
Oxalidi affinis 433	Paragonica 149	Phylla 194
Oxalis 337. 338	Paullinia 427	Phyllanthos 469
Oxycantha 477. 478	Pavate 110	Phyllis 1068
Oxycedrus 1038	Pavel 1009	Phyllitis 1067. 1069. 1070. 1073. 1077. --- 1080. 1082. -- 1084. 1097
Oxyoides 434	Pavia 544	Phyllochnois 561
Oxys 434. 435	Pecten 257	Phyllon 1035. 1036
	Pedicularis 602. 611. 629	Phyllum 1035
	Pee-inota-modien 184	Phyteuoides 116
	Peekandel 443	Picea 1002
	Pee-mottenga 52	Picnoman 820
	Peetandali-cotti 715	Picris 790. 799
	Peetjanga-pulpani 635	Pilosella 574. 642. 665. 800 807. 850
	Peetumba 16	Pimpinella 116. 117. 252. 261. 994
	Penza 703	Pinanga 1189
	Penar-valli 1028	Pinafter 1000
	Pentaphylloides 284. 495 - 497. 499.	Pinguicula 17
	Pentraphyllum 497. 498. 500. 503	Pingvin 285
	Pentstemon 612	Pinus 1004
	Peplis 455	Piper 189
	Peplon 455	Piperi similis 638
	Peplus 456	Pistachia 270
	Pepo 1009. 1010	Pistacia 124. 270. 397
	Peragu 637	Pistacio-vitex 638
	Peralu 1060	Pistulochia 699. 961. 962
	Pereskia 469	Pisum 366. 737. 754
	Perfoliata 236. 237. 238	Pitronia 140. 141
	Perfoliatum 238	Piryhusa 457. 458
	Periclymenum 118. 173 - 176. 627. 628.	Planra 631
	Periclymeri similis 637	africana 385
	Perinkara 515	annua 112
	Periploca 212	lactaria 452
	Perlea 370	leguminosa 708. 753
	Perfica 472	
	Perfica similis 200	

P.

Pachyphylla 180
Paco-teroca 2
Pada-valam 1008
Padus 193. 473. 474
Paderota 11
Penoc 515
Pajaneli 623
Painafchylli 639
Paiparoca 964
Pala 210
Palega 625
Paliuro affinis 637. 1026
Paliurus 194
Palma 944. 1043. 1187 - 1189
Palmata 944
Palmodecca 156
Panaces 249. 274
Panambu-valli 333
Panax 247. 249. 262.
Pandipavel 1009
Panicastrella 1050
Panicum 75
Panna-kelengo-marava 1087

Planta

INDEX SYNONYMORUM.

)()(3

INDEX SYNONYMORUM.

Ricinoides 196 1004-1006	Scabiosa 87. 95-97. 101.	Selinum 244
1042	102. 815. 902. 913.	Semen sanctum 245
Ricinus 195. 1004-1007	927	Sempervivum 129 401. 412
1042	Scabiosa affinis 159. 588	Sendra-claudi 157
Ricophora 1032	Scammonia 212	Senecio 808. 814. 835. 860.
Rirro 815	Scammonia 193	861. 862. 880. 902.
Romphal 965	Scammonie affinis 212	carduus 909
Ros folis 281. 282	Sceptrum Carolinum 608	Senecioides 862
Rosa hierocuntea 282	Schadidacali 460	Seena 176. 381. 741
sinensis 694	Schageri-Cottam 514	Senna affinis 375
Rosmarinus 391. 563	Scherubala 804	Serapis 950. 952
Royena 628	Scherunam-Cottam 1042	Seriana 365. 366
Royce 176	Scherupauri 694	Seriphium 659
Rubeola 102. 104. 107. 1051	Scherti 110	Serpentaria 341. 567.
Rubia 102-105. 107-109.	Schinoides 389	Serpentina 115.
560.	Schit-edu 634	Serpyllum 591.
Rudbeckia 176. 177. 908	Schrenanbus 82	Serpyllifolia 631
Ruellia 17	Schono-prafum 301	Sesamoides 207. 415.
Ruppia 968	Schorigeram 980	449. 459
Rus 675	Schovanna-adambee 159	Sesamum 972
Ruscum 1041	Schunambu-valli 117	Sesban 722
Ruta 444. 546. 621. 1087	Schunda-panna 1189	Seseli 238. 239. 240. 244.
Ruychiana 595	Schunga-Cuspi 753	246. 248. 250. 259.
	Scirpo-Cyperus 50. 51	260. 264
	Scirpoides 49	Sherardia 19. 359
	Scirpus 43. 54	Sicyoides 1013
	Sciarca 25. 26	Sideritis 574. 578. 579. 581.
	Scolopendria 1079	584. 588. 596
	Scolopendriae facie 885	Silaum 246
	Scolymoccephalus 93	Silene 392. 414. 415. 416.
	Scolymus 827. 828	437
	Scoparia 90	Siler 246. 249
	Scordio affinis 603	Siliqua 34. 374. 1036
	Scordium 565. 567. 587	Silybum 814
	Scorodonia 19. 564	Simaruba 1026
	Scorodoprasum 898	Sinapi 659. 662. 669
	Scorodo-Thlaspi 646	Sinapis 640
	Scorpii spina 232	Sinapistrum 672
	Scorpioides 744-746	Sivanna-pola-rali 391
	Scorpius 711	Sisfarum 251
	Scorzonera 813	Sisymbria 657
	Scotellaria 588	Sisymbrium 576. 655. 660.
	Sebestena 190	667
	Secale 83	Sisyrinchium 36. 40. 306
	Securidaca 743. 752. 758.	Sitz f Sits-dlin 266
	762	Sio affinis 655
	Sedo-affinis 372	Sium 252. 253. 255. 262
	Sedum 129. 142. 144.	Smilax 155. 304. 723. 994.
	221. 236. 398-400	1032
	402-405. 425. 429.	Soda 243
	464. 465. 642.	Solanifolia 9
	Seguaster 1189	Solano-eogener 340
	Seleginoides 1102	Solanoides 122
	Selago 422. 1100. 1101.	Solanum 3. 122. 177. 179.
	1102,	

S.

Stammona 344
Sabina 1039
Sagina 422
Sagina affinis 128
Sagitta 993
Sagitta similis 288
Sagittaria 343
Sajor 1192
Salicaria 447
Saliflore 281. 282
Salfolia 35
Salvia 585. 627
Salvinia 1099
Sambucus 268
Samoloides 116
Sarapfacus 567
Samitravadi 471
Sanemunda 956. 560
Sanicula 17. 142-144. 179.
226. 399. 400. 401. 403
Santolina 840. 843. 927
Sapote 1190
Saponaria 406-408. 417
Sargazo 1160
Saribus 1187.
Sassafras 371
Saryrium 942. 946. 950
Saururus 29. 30
Saxifraga 128. 260. 398. 421

181.

INDEX SYNONYMORUM.

181. 182. - 184. 340.		T.			
367. 441. 635					
Soldanella 159	Tabaxir	81	Tilia	225	
Solidago 870. 871. 924. --	Tagadi	1049	Tilia affinis	426	
936	Tagera	376	Tinda-pana	986	
Soncho - edfinia 811	Tagetes	835	Tinus 267	269	
Sonchus 790. 792. 796.	Tagolina	835	Tiri panua	1067	
797. 798. 808	Tala-neli	156	Tirucalli	452	
Sorbus 476. 478	Taludama	3	Tithymali myrsinias 111.		
Sparreum 707	Tamara	428. 511	112 1025		
Spartium 720. 762	Tamariscus 1999. 271. 928		Tithymalus 213. 450- 463.		
Spathula 99	Tanacerum 887. 890. 899		1197		
Speculum veneris 168	Tana- pouel- paatliamara-		Todda-pana f. monta-pan-		
169. 635	vara	1101	na	1188	
Spergula 88. 272. 281. 408.	Tandali-Corri	715	Todda-vaddi	434	
424. 632	Tanghekolli	792	Tormentilla	123	
Spherocephalus 1185	Taraxaconoides	299	Tota-pivi	1008	
Sphagnum 1106. -- 1108.	Taraxacomastrum 808. 809		Toxicodendron 265. 266		
1115	Taraxacum	798	Trachelium 164. 166. 167.		
Sphondylium 249. 250	Tarchonanthos	989	168. 171		
Sphondylococcus 111	Tarton-Raire	356	Tragacantha	761	
Spongiosus fucus 1170	Tatula	179	Tragacantha affinis	762	
Spica celien	Telephiastrum	446	Tragium	784	
trifolia	Telephoides	1014	Tragopogon 791. 1198		
Spina	Telephium 423. 430. 1035		Tragorchis	740	
alba 815. 827	Tenga	1188	Tragoriganum	591	
tomentosa	Terabintus 509. 1025. 1026		Tragoselinum	264	
Spirma 198	Testiculi species	939	Tragum	1040	
Strachyoides	Testiculus 941. 946		Tremella 1163. 1164.		
Strachys 573. - 575. 587	Tetragonocarpus	480	Tretorrhiza	231	
Strachas 572. 573	Teucri facie	125	Tribulus	120	
Strachadi affinis	Teucrium 561. 598. 600.		Trichomanes 1074. 1080-		
Strachelina	602		1082. 1088. 1089. 1094.		
Staphisagria	602		1095		
Staphyloendron 270. 397	Thalictrides	504	Trifoliastrum	766	
Stellaria	Thalictrum	542. 557	Trifolii spica	225	
Storbe 813. 818. 819. 912.	Thapfia	242	Trifolio affinis	692	
914-916. 919. 928. 994	Thee	515	Trifolium 145. 386. 433.		
Storhas 853. 854	Thelypteris	1072	435. 538. 672. 719. 740.		
Storhadi citrinæ affinis	Thevetia	208	749. 763. 764. 776-784		
855	Thlaspi 642. 643. 545. 647.		Trilopus	124	
Steramonioides	653		Trinitatis herba	936	
Stratiotes 897. 963	Thlaspidium	649. 652	Trionum	697	
Seyrax	Thora	350	Triopteris	428	
Suber	Thrafi	45	Triorchis 940. 946. 348		
Sabularia	Thuris folium	510	Triosteospermum	176	
Succisa	Thylacitis	228	Tripolium 151. 873		
Suffrutex ericetiformis	Thymbra	567. 704	Trineum 83. 601		
704	Thymelæa 96 207. 356.		Triumfetta	389	
Suillus 1177	358 539. 560. 629		Trixago	601	
Sumach	Thymelæa facie	284	Trixis	18	
Syalina	affinis	1038	Tierou-katou - nagegam		
Sycomorus	Thymo simile	591	389		
Symphoricarpos	Thymum	567. 568	Tiageri-nuren	1022	
Symphytum 135. 177	Thymus	567- 569. 591	Tiama-pullu	58. 69	
Syringa 112. 191. 470.	Tiera-kuren-pullu.	54	Tiam-pangam	381	
623			Tiela	1060	
			Tieli		

INDEX SYNONYMORUM.

Theli	27	Urtica	579. 580. 979. 980.	Virga sanguinea	118
Thien-cumulu	632	1003		Viscago	416. 417. 479. 480
Thiercammulla	6	Urticu	512	Viscum	110. 286. 288. 953
Thiem-Tani	1193	Urtica	1154-1156	Vistna-clandi	157
Thiera Maga Nari	1200	Utricsaria	955	Vitalba	541
Thiera-famstravadi	471	Uva urli	395. 478	Vitaliana	144
Therou-mau-maravara	912	Uvifera	365	Vitex	255. 267
Therou-ponna	514			Viricelle	200
Theru-kendel	443			Visis	202. 1012. 1013
Theru-kirganelli	982			idza	175. 196. 349 -
Theru-korhijelleri-pullu	87	Vaccaria	409	352. 395. 397. 478	
Theru-valli-panna	1063	Valdia	637	Volubilis	162. 212. 1033
Theru veela	672	Valeriana	162. 840	Volli-panna	1013
Theru-uren	675	Valerianella	3. 34. 101.	Vulnaria	719
Thetti-mandam	380	102. 232. 234. 629.		Vulvaria	220
Tsubakki	698	Valerianoides	629. 1058		
Tsutufu	150	Vallia-capo-molago	189		
Tuber	1183	Valli-onapu	937		
Tubera	1183	Vallisperioides	1015		
Tuberaria	526	Vanillas	952		
Tuberum	1183	Vanille	952		
Tulipifera	535	Van Rheedia	1193		
Tuna	468. 469	Velutia-mandaru	375		
Tunica	408. 410-412	Veratrum	235		
Turpethum	155	Verbasci tolio	112		
Turrita	666	Verbasculum	143		
Turritis	661. 664. 665	Verbasceum	585. 621		
Tusei	303	Verbena	601. 660		
Tussilago	336. 881	Verbenæ affinis	342		
Tuzpatia	121	Verbesina	901		
Typha	324	Vermicularis	432		
Tydelinum	244	Vermiculata	406		
		Veronica	116. 613		
		Veronice similis	19		
		Vesicaria	651		
		Viburnum	268. 627		
		Vicia	729. 734. 738. 759		
		Vicia similis	730		
		Vidi-maram	190		
		Vincetoxicum	216		
		Viola	176. 167. 345. 653.		
		662. 663.			
		Virga aurea	140. 863. 871.		
		378. - 881			



NO.

NOMINA TRIVIALIA.

NOMINA TRIVIALIA.

A.

Abelmoschus	696
Abies	1002
Abrotanum	845
Abrus	753
Abrothium	848
Abus	376
Acarua	820
Acanthium	327
Acetabulum	1181
Acerosa	317
Acetosella	338. 433
Acinodendrum	389
Acinos	591
Acnella	901
Adhatoda	15
Adiant. nigrum	1081
Egilops	996
Ethiopsis	25
Ageratum	897
Agnus Castus	638
Ahouai	208
Aizoon	430
Aramasca	292
Alaternus	193
Alcea	689
Alhagi	745
Alkekengi	183
Alopecuros	573
Alfinsastrum	368
Alypum	95
Alysson	582
Amalago	29
Amarella	230
Amelachier	478
Amellus	873
Ammi	252
Ammoides	260
Annomum	252
Ampeloprasum	294
Anacampseros	430. 445
Anagallis	12
Ananas	285
Anandria	865
Anblatum	606
Androsæmum	784
Anguria	1011
Angvina	1008
Anisum	264
Anserina	495
Antheuphorbium	335

Anthora	532
Antrifcus	240. 257
Aperine	108. 1052
Aphaca	729
Apios	457. 753
Aquifolium	125
Arbor triflis	6
Archangelica	250
Arctostaphylos	351
Arcturus	178
Arcira	389
Argemone	506
Aria	475
Arisarum	966
Armeria	274. 410. 420
Armoracia	648
Arunce	490
Aclepium	261
Alfa fœtida	248
Afcyron	789
Alterifcus	920
Auricula	143. 800. 1157
Auricularia	101
Auriscalpium	1178
Auranthium	782
Avellana	998
Azrolus	477
Azedarac	384
Azediracta	335

B.

Baducca	504
Belanghas	1007
Balsamina	983. 1009
Balsamita	846
Balsamum	384
Barnbos	81
Basilicum	597
Barba Jovis	720
Barbarea	659
Barometz	1092
Batatas	154
Beccabunga	12
Behen	418. 914. 414
Bella donna	181. 293
Bellidjastrum	886
Bénzoin	370
Bermudiana	954
Beule	28

Betonica	15
Bidens	908
Bihai	1042
Bilimbi	428
Bistorta	360
Blattaria	178
Blitum	990
Bonarota	11
Bonduc	381
Bonus henricus	218
Bortys	219. 562
Bovista	1183
Brignola	475
Britannica	334
Bryopteris	1103
Bulbocastanum	243
Bulbocodium	36. 294
Burfia pastoris	647

C.

Caceo	788
Cajan	739
Calnito	192
Cakile	670
Calaba	514
Calebura	509
Calamintha	595
Calamus	324
Calainagrostis	82
Calceolus	951
Calendula	922
Calitrapa	917
Calomelanos	1072
Camara	627
Campechianum	384
Camphora	369
Candel	443
Candelabrum	211
Caprifolium	173
Capillus Veneris	1096
Caput Galli	751
Medusa	84. 452
Caracalla	725
Caragana	722
Carambola	428
Cardamominum	1
Cardiaca	484
Car-	

NOMINA TRIVIALIA.

Cardinalis	930	Citrullus	1010	Diclamnus	189
Cardunculus.	827. 831	Clava Herculis	170	Dierville	175
Carica	1059	Clymenum	730	Dodonæa	125
Cariophyllus	410	Cneorum	157. 357	Donax	81
Carota	242	Cobbe	267	Doronicum	880
Carpobolus	1184	Cocculus	340	Dortmanna	929
Carvi	263	Cœli rosa	436	Dorychnium	776
Cascarilla	1042	Colocasia	965	Draba	646
Cassia	369	Colocynthis	1011	Dracontium	964
Cassina	125	Coma aurea	840	Dracunculus	850. 964
Castanea	997	Conocarpodendrum	93	Dryopteris	1093
Catalpa	622	Consolida	530	Eudaïm	1011
Catechu	1185	Contrajerva	121	Dulcamara	185
Cedrus	1001	Copallium	266		
Ceiba	511	Convulvulus	364		
Cemhra	1000	Coralloriza	945		
Centaurium	229. 910	Coris	787		
Cepa	301	Coronaria	436		
Cepza	431	Coronopus	115. 648		
Cephalotes	582	Cota	894		
Cerasus	474	Cotinus	267		
Ceratonia	523	Coroneaster	479		
Ceretolium	257	Cotula	894		
Cerris	997	Courbaril	1192		
Cervaria	167. 1194	Coryledon	398		
Cerviana	272	Cracca	735		
Ceterach	4080	Crista	380		
Chama-Buxus	704	Galli	603		
Cifus	392	Crocodyllum	919		
Crista	379	Cruciata	1052		
Cyparissias	842	Crupina	909		
drys	12. 565	Crux Galli	56. 476		
Jaline	559	Crux andrææ	787		
Iea	981	Cucullaria	699		
Mespilus	981	Cujete	626		
Moly	301	Cunonia	37		
Morus	494	Curcas	1006		
pitys	562	Cururu	365		
peuce	819	Cyanus	911		
Syce	455	Cydonia	481		
Cherophyllos	555	Cymbalariz	405. 613		
Chamomilla	891	Cyminum	854		
Chempaca	536	Cynapium	256		
Characias	463	Cynops	116		
Charantia	1009	Cynocrambe	993		
Cheiranthus	933	Cynosbatus	202		
Chairi	661				
Chelidonia	655				
China	894				
Chironium	1029				
Chryfogonum	249				
Cicer	312				
Cicera	757				
Cinamomum	730				
	369				

NOMINA TRIVIALIA.

opyrum
 fara
 ulago
 aria
 idea
 s indica
 pendula
 ten
 mula
 s aeris
 aque
 cuculi
 Jovis
 uni graecum
 ngula
 naria

36
 865
 247
 530
 225
 468
 490
 1162
 544. 548
 952
 1168
 436
 436
 777
 193
 525

Holoschænus
 Horminum
 361. 367
 809
 442
 1041
 93
 1041
 387
 344
 447. 647
 988
 560
 1182
 Laburnum
 Lacryma Jobi
 Lactuca
 Ladanum
 Lagopus
 Lantana
 Lappa
 Lappula
 Lathanthus
 Lamx
 Lathyrus
 Laurentia
 Laureola
 Laurocerasus
 Lavenia
 Lebeck
 Lentago
 Lentiscus
 Leontopodium
 Leonthopetalon
 Leonurus
 Lepidocarpodendron
 Leptostachya
 Leucanthemum
 Leucoxydon
 Levisanus
 Levillicum
 Libanotis
 Liljago
 Liljaltrum
 Liljo Asphodelus
 Hyacinthus
 Lima
 Limon
 Limonia
 Limonium
 Linaria
 Lingua
 Linophyllon
 Linofyris
 Linum stellatum
 Linza
 Locusta
 Lonchitis
 Lophanthus
 Lotus
 Luffa
 Lunaria
 Lureola
 Lycopersicum
 Lupinaster
 Lupulus

739
 962
 1163
 579
 114
 268
 816
 444. 137
 783
 1001
 457
 231
 857
 474
 902
 516
 268
 1026
 855
 312
 587
 94
 601
 388
 624
 199
 250
 244. 246
 310
 324
 324
 308
 72
 782
 389
 272
 616
 545 950
 207
 841
 148
 1163
 33
 1088
 509
 157. 194. 311
 1009
 336. 656. 1064
 448
 185
 766
 1028

G.

anga
 banum
 e
 eobdolon
 bera
 m
 gidium
 nago
 ucium
 ux
 idium
 na parad.
 narum
 stularia
 ajava
 izuma
 idonia

2
 253
 1024
 586
 885
 401
 242
 435
 508
 759
 357
 2
 472
 201
 470
 781
 443
 Jabotapita
 Jacca
 Jacobæa
 Jalappa
 Jambos
 Jberis
 Icaco
 Ilex
 Illecebrum
 Inga
 Infisia
 Ineybus
 Jolirhus
 Jonquilla
 Jonehlaapi
 Ipecacuanha
 Iria
 Irio
 Isora
 Jujuba
 Jungermannia

I.

H.

imus
 icacabum
 mala
 Heria
 enium
 leborine
 lanchemum
 k
 atica
 pa venti
 pionitis
 ulus

1052
 366
 444
 704
 881
 949
 528
 102. 1017
 538
 586
 1078
 402
 Kali
 Karatas
 Kauki
 Kleinia
 Labrusca

K.

L.

222
 285
 349
 834
 783
 203

Ma

NOMINA TRIVIALIA.

[illegible]

NOMINA TRIVIALIA

[illegible]

NOMINA TRIVIALIA.

Virgaurea	880		X.		Zapota	11
Vitisga	848				Zacintha	8
Vitalba	544	Xiphium		40	Zeocriton	
Vitaliana	143	Xylotium		174	Zerumber	
Viticella	548		T.		Zingiber	
Vitis idæa	351	Yervamora		225	Ziziphus	11
Volema	479		Z.		Zygis	55
Vulneraria	719					
Vulvaria	220	Zenonia		41		



ADDENDA.

- P. 17. Gratiola virginiana (adde) Tſieria Manga Nari. *Rheed mal. g. p. 165. t. 85.*
25. Salvia pratensis. - *Habitat in Europæ pratis. 2*
353. Erica arborea. - *Habitat in Europa australi. 1*
494. Rubus Chamæmorus - *Hæc species omnino Dioica est*
665. Arabis canadensis - *Habitat in America septentrionali.*
696. Hibiscus surattensis - *Herba crinium. Rumpf. amb g. p. 40. t. 16.*
721. Lupinus angustifolius - *Habitat in Hispania ☉*
781. Medicago polymorpha - *Habitat in Europa australi. ☉*
825. Carduus tataricus - *Folia radicalia in hac specie pinnatifida sunt, non ut in precedenti.*
479. Pyrus communis (rectius) PYRUS foliis serratis, floribus corymbosis.
- Pyrus malus - - PYRUS foliis serratis, umbellulis sessilibus.
858. Xeranthem: speciosissimum. *Habitat in Æthiopia. 3*
348. § 4. EPILOBIUM: major. *Bauh. pin. 303.*
927. MICROPUS (*supinus*) procumbens, foliis geminis.
- 2 MICROPUS erectus, foliis solitariis. *erectus.*
- Habitat in Hispania. ☉*
1199. 3. Filago (*pyramidata*) conjugatur Gnaphalio germanico p. 857.
4. FILAGO floribus conicis terminalibus axillari-busque, caule subdichotomo erecto. *est Gnaphalium gallicum p. 857. n. 33.*
5. FILAGO floribus conicis lateralibus, caule paniculato. *est Gnaphalium arvense p. 856. n. 32.*
6. FILAGO floribus subulatis axillaribus, caule dichotomo erecto. *est Gnaphalium gallicum p. 857. n. 34.*



ERRATA.

6. n. 3. <i>Virginia</i> lege <i>virginica</i> .	P. 169. n. 34 adde <i>betophylla</i> ,
30. n. 14. <i>petatum</i> - <i>petatum</i> .	296. n. 6. - <i>magicum</i> .
59. n. 9. <i>filiforme</i> - <i>filiforme</i> .	538. n. 1. - <i>Heratice</i> .
108. n. 19. <i>apavina</i> - <i>Apavine</i> .	652. n. 3. - <i>campestris</i> .
246. n. 2. <i>vaicum</i> - <i>pyrenaicum</i> .	869. n. 15. - <i>crutifolia</i> .
312. n. 2. <i>Lam:ropetatum</i> <i>Leontopetatum</i> ,	99. <i>Scabiosa</i> 9. <i>corollis trifidis error</i>
345. n. 1. <i>majus</i> - <i>minus</i> ,	<i>typographi: secundum</i>
2. <i>minus</i> - <i>majus</i> .	<i>Cl. Austorem.</i>
373. n. 1. <i>alo pedes</i> - <i>alopecuroides</i> ,	300. <i>Allium</i> 23. <i>Synonymon ex Hort</i>
408. n. 6. <i>hybrina</i> - <i>hybrida</i> .	<i>cliff excludatur.</i>
447. n. 6. <i>inylimifolia</i> , <i>thymifolia</i> .	420. <i>Silene</i> 23. <i>LYCHNIS d:icatur.</i>
514. n. 1. <i>aniculata</i> , <i>paniculata</i> .	570. <i>Nepeta</i> 5. <i>linea tertia deleatur.</i>
2. <i>ateriflora</i> - <i>lateriflora</i> .	844. <i>Tanacetum</i> 2 - 4. * <i>fruticosa de-</i>
540. n. * <i>Anemoneide</i> - <i>Anemonoides</i> .	<i>leatur.</i>
557. n. 1. <i>umariondes</i> - <i>Fumarioides</i> .	<i>Tanacetum</i> 3 - 4. * <i>Herbacea ad-</i>
596. n. 11. <i>triflora</i> - <i>triflorum</i> .	<i>datur.</i>
608. n. 7. <i>recucisa</i> - <i>recutita</i> .	
708. n. 2. <i>radiafum</i> - <i>radiatum</i> .	
798. n. 7. <i>altissima</i> - <i>repens</i> .	
896. n. 16. <i>triumfeti</i> - <i>triumfeti</i> .	
1006. n. 3. <i>Gurcas</i> - <i>Curcas</i> .	
P. 61. <i>MILIUM</i> 1. <i>foliis dispermis</i> - lege <i>floribus dispermis</i> .	
83. <i>LOLIUM</i> 1. <i>⊙</i> - <i>2. ♀</i> - <i>⊙</i> .	
93. <i>LEUCADENDROS</i> 12. 13. - <i>LEUCADENDRON</i> .	
193. <i>RHAMNUS</i> 1. <i>spicis</i> - <i>spinis</i> .	
221. <i>CHENOPODIUM</i> 18. <i>⊙</i> . <i>Chenopodium</i> , - <i>Salsola</i> .	
268. <i>VIBURNUM</i> 4. <i>ovatis</i> - <i>plicaris</i> .	
436. <i>AGROSTEMMA</i> 3. <i>laciniatis</i> - <i>lanceolatis</i> .	
<i>LYCHNIS</i> 4. <i>bifidis corymbosis</i> - <i>bifidis, floribus corymbosis</i> .	
479. <i>PYRUS</i> 1. <i>foliis basi</i> - <i>pomis basi</i> .	
524. <i>CISTUS</i> 5. <i>trinervius</i> - <i>trinerviis</i> .	
527. <i>CISTUS</i> 19. <i>foliariis subfessilibus bractea tri-</i>	
<i>plici</i> - <i>erectis, pedicellis horizonta-</i>	
<i>libus</i> .	
598. <i>SCUTELLARIA</i> 3. <i>floribus lævibus</i> - <i>foliis lævibus</i> .	
777. <i>TRIGONELLA</i> 3. <i>peduncularis</i> - <i>fessilibus</i> .	
824. <i>CARDUUS</i> 18. <i>fessilibus</i> - <i>petiolatis</i> .	
843. <i>TANACETUM</i> 1. * <i>Herbacea</i> - <i>* Fruticosa</i> .	
916. <i>CFNTAUREA</i> 37. <i>sinuatis</i> - <i>lyratis</i> .	
1077. <i>BLECHNUM</i> 2. <i>alteanis</i> - <i>alternis</i> .	
859. §. 7. <i>HERANTHEMUM</i> - <i>XERANTHEMUM</i>	
828. §. 3. <i>CARLINA vulgaris</i> - <i>corymbosa</i>	
§. 4. <i>corymbosa</i> - <i>vulgaris</i>	
1199. §. 3. <i>absque paleis in</i> - <i>absque paleis. In</i>	
1135. §. 19. <i>gulacea</i> - <i>julacea</i>	
1011. §. 5. <i>foliorum angulis</i> - <i>foliis dentatis integria</i>	
63. §. 1. <i>spicatum</i> - <i>indicum</i>	
598. §. 1. <i>America</i> - <i>Armeria</i>	
192. <i>Plukenetia fr. serrato</i> - <i>fructu tetragono</i>	
420. §. 21. <i>SILENE portensis excludatur, cultura enim constitit eandem esse cum</i>	
<i>SILENE</i> 19 <i>in aperta. que in hortis corollam non protrudit.</i>	
418. §. 10. <i>SILENE gigantea</i> - <i>CUCUBALUS giganteus ad genus itaque an</i>	
<i>recedens amandetur.</i>	

Paginas & Paragraphos allegatas cum libris conferre otia non admiseri, quas L. B. emendes, usque dum altera prouideat editio.



